

POSTER PRESENTATIONS

Open Access

Design of a Phase IIb, randomized, controlled, double-blind, multi-centre study to evaluate the efficacy, safety, and immunogenicity of the GMZ2 candidate malaria vaccine in Ugandan, Ghanaian, Burkinabe and Gabonese children aged 12-60 months

Ramadhani A Noor^{1*}, Michael Thiesen², Benjamin Mordmuller³, Paul Milligan⁴, Frank Atuguba⁵, Sodiomon Sirima⁶, Saadou Issifou⁷, Kalifa Bojang⁸, Fred Kironde⁹, Alfred Tiono⁶, Brenda Okech¹, Roma Chilengi¹⁰, Ateba Ngoa⁷, Kaddu Mukasa⁹, Soren Jepsen²

From Parasite to Prevention: Advances in the understanding of malaria
Edinburgh, UK. 20-22 October 2010

The malaria vaccine candidate GMZ2 is a recombinant fusion protein of Plasmodium falciparum Glutamate Rich Protein and Merozoite Surface Protein 3, adjuvanted with aluminium hydroxide. Trials in malaria naive adults in Germany and in Gabon in adults exposed to malaria and subsequently in children 1-5yrs showed that three doses of 100ug GMZ2 are well tolerated and immunogenic. The aim of this phase 2b trial is to determine whether the GZ2 vaccine can protect against clinical attacks of malaria in children aged 1-5yrs, to obtain more extensive data on safety, tolerability and immunogenicity, and to determine if the vaccine can prevent anaemia and reduce the parasite density and gametocyte carriage.

The primary endpoint is the incidence of clinical malaria defined as fever or history of fever in the previous 24 hours with parasite density of 5000 asexual parasites per ul or more, detected by passive surveillance over a 6 month period from the third vaccination. All malaria episodes will be included in the calculation of vaccine efficacy. This endpoint is chosen in order to have the optimum power to detect a protective effect; efficacy will also be calculated using a range of different

parasite density cutoff values. Children will be followed for a total of 22 months from the third vaccination. Immune responses to the vaccine antigens GMZ2, GLURP and MSP3 will be assessed by measuring antigen specific IgG by ELISA, and antigen specific memory B-cell responses by ELISPOT. Functionality of the immune response will be assessed by growth inhibition of P. falciparum in the presence or absence of monocytes. Cell mediated immunogenicity will be assessed by cytokine profiling and intracellular cell staining following stimulation with the vaccine antigen, and the quality of the antibody response by type and subclass-specific ELISA.

The planning of the trial was complicated by the need to take account of the changing epidemiology of malaria. Baseline studies were conducted in five potential trial sites to determine the current incidence of malaria; in each site two cohorts of children were followed to estimate the incidence of malaria by active and passive surveillance and by passive surveillance alone. Four sites with higher malaria incidence were chosen for recruitment. A total sample size of 1840 will be enrolled in each of the four sites in 2010/11.

¹African Malaria Network Trust (AMANET), Dar Es Salaam, Tanzania, United Republic of

Full list of author information is available at the end of the article

Author details

¹African Malaria Network Trust (AMANET), Dar Es Salaam, Tanzania, United Republic of. ²Statens Serum Institut (SSI) Denmark, Copenhagen, Denmark. ³Universitätsklinikum Tübingen (UKT), Tübingen, Germany. ⁴London School of Hygiene and Tropical Medicine (LSHTM), London, UK. ⁵Navrongo Health Research Centre (NHRC), Navrongo, Ghana. ⁶CNRFP, Burkina Faso, Ouagadougou, Burkina Faso. ⁷Medical Research Unit-Albert Schweitzer Hospital, Lambarene, Gabon. ⁸Medical Research Council (UK)The Gambia, Fajara/Banjul, Gambia. ⁹Makerere University, Kampala, Uganda. ¹⁰KEMRI-Wellcome Trust Research Programme, Mombasa, Kenya.

Published: 20 October 2010

doi:10.1186/1475-2875-9-S2-P26

Cite this article as: Noor *et al.*: Design of a Phase IIb, randomized, controlled, double-blind, multi-centre study to evaluate the efficacy, safety, and immunogenicity of the GMZ2 candidate malaria vaccine in Ugandan, Ghanaian, Burkinabe and Gabonese children aged 12-60 months. *Malaria Journal* 2010 **9**(Suppl 2):P26.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

