



Secular changes in bone mineral density of adult Japanese women from 1995 to 2013

Hiroaki Watanabe¹⁾²⁾, Yasuko Minagawa³⁾, Ichiro Suzuki³⁾, Kaori Kitamura¹⁾, Yumi Watanabe¹⁾, Keiko Kabasawa⁴⁾, Kseniia Platonova¹⁾, Aya Hinata¹⁾ and Kazutoshi Nakamura¹⁾

¹⁾Department of Preventive Medicine, Niigata University Graduate School of Medical and Dental Sciences, Niigata, Japan, ²⁾Department of Clinical Engineering and Medical Technology, Niigata University of Health and Welfare, Niigata, Japan, ³⁾Shibata Comprehensive Health Care Service Center, Shibata, Niigata, Japan, ⁴⁾Department of Health Promotion Medicine, Niigata University Graduate School of Medical and Dental Sciences, Niigata, Japan

(Received June 16, 2021, accepted October 14, 2021)

Abstract

Introduction : Secular changes in hip fracture incidence have been reported in the last few decades in Japan, but whether long-term bone mineral density (BMD) is also changing is unclear. This study aimed to determine whether BMD of Japanese women has changed over time.

Methods : Subjects were 10,649 adult women who underwent BMD measurement in a health check-up population in Niigata, Japan, between 1995 and 2013. BMD of the distal, non-dominant forearm was measured by dual-energy X-ray absorptiometry. Demographic information and BMI were also obtained. Secular trends were determined by linear regression analysis.

Results : BMD of subjects in their 40's decreased significantly in the age-adjusted model (P for trend=0.0162), but not in the age- and BMI-adjusted model (P for trend=0.2171). BMD of subjects in their 50's decreased marginally in the age-adjusted model (P for trend=0.0535), but not in the age- and BMI-adjusted model (P for trend=0.6601). BMDs of subjects in their 30's and 60's did not significantly change, while BMIs of subjects in their 40's-60's decreased significantly.

Conclusions : A secular decrease in BMD, partly attributed to decreases in BMI, was observed in middle-aged Japanese women from 1995 to 2013. Measures to help maintain suitable BMI will be necessary to prevent a decrease in BMD among women.

Key words : BMI, bone density, osteoporosis, secular change, women

Introduction

Osteoporosis is a skeletal disorder involving compromised bone strength which predisposes individuals to an increased risk of fracture¹⁾. The burden of osteoporosis and osteoporotic fracture is enormous. The global number of hip fractures in 2010 was estimated to be 2.7 million²⁾. In Japan, there were an estimated 175,700 hip fractures in 2012³⁾, representing an increase over the last few decades^{3,4)}. This increase may partly be explained by aging of the population. While hip fracture inci-

dence rates in European and North American countries generally stabilized with age-adjusted decreases in the last few decades^{5,6)}, one review⁵⁾ reported that the incidence rate rose in Asia, and Japan was not an exception. Orimo *et al.*³⁾ reported that hip fracture incidence rates in Japan for people in their 80's and 90's increased from 1992 to 2012, and Miyasaka *et al.*⁴⁾ similarly reported that age-adjusted incidence rates increased from 1985 to 2010 in Niigata, Japan.

The reasons underlying the long-term increase in hip fracture incidence in Japan are unclear, but

Corresponding author : Kazutoshi Nakamura, MD E-mail : kazun@med.nigata-u.ac.jp

©2021 The Fukushima Society of Medical Science. This article is licensed under a Creative Commons [Attribution-NonCommercial-ShareAlike 4.0 International] license.

<https://creativecommons.org/licenses/by-nc-sa/4.0/>

may be related to an increased prevalence of osteoporosis, generally diagnosed based on bone mineral density (BMD)¹¹. One question that arises is whether BMD of community-dwelling people changes over time in the long term. While some have reported a secular change in BMD, others have reported conflicting results. For instance, Xu and Wu reported a decreasing trend of BMD in a US population of the National Health and Nutrition Examination Survey (NHANES) 2005–2014⁷, whereas Cheung *et al.*⁸ reported an increasing trend of BMD in Hong Kong women. Thus, further studies on secular changes in BMD in a general population are needed.

Female sex is a risk factor for low BMD due not only to biological factors but also to behavioral ones. For instance, dieting to lose weight is a typical, female-related risk factor. According to previous studies, nearly half of adult females had a history of dieting in Japan⁹ and worldwide¹⁰. The National Health and Nutrition Survey in Japan reported that total energy intake (men and women combined) showed a decreasing trend from 2,088 kcal/day in 1985 to 1,897 kcal/day in 2017, and that the proportion of underweight (body mass index [BMI] <18.5 kg/m²) women showed an increasing trend from 8.1% in 1985 to 10.3% in 2017¹¹. These data suggest that BMD of Japanese women is likely decreasing.

This study aimed to determine whether BMD of Japanese women decreased in the long term, while also considering BMI changes. To this end, we analyzed a large dataset of BMD measurements from a health check-up program for community-dwelling people between 1995 and 2013.

Subjects and methods

Subjects

We targeted 12,799 women, aged between 20 and 79 years, who underwent BMD measurement as part of a health check-up examination at Shibata Comprehensive Health Care Service Center (Niigata, Japan) between 1995 and 2013, when BMD measurements using the DTX-200 osteometer were terminated. All women whose BMD was measured during this period were eligible for inclusion in this study. However, women who did not have concomitant measurements of BMI (an important determinant of BMD) were excluded. For statistical analysis, we used only data from the first BMD measurement for those who underwent two or more BMD measure-

ments during the study period. Of the 12,799 women, 10,649 were analyzed after excluding 2,150 who had missing BMI data. Men were not included in the analysis given the limited number of men with BMD measurements ($n=2,250$) relative to women. This study used data obtained from an anonymized database of Shibata Comprehensive Health Care Service Center. The protocol of this study was approved by the Ethics Committee of Niigata University (No. 2017-0396).

Measurements

BMD of the distal, non-dominant forearm was measured with the dual-energy X-ray absorptiometry (DXA) method using the DTX-200 osteometer (Osteometer MediTech A/S, Rødovre, Denmark). The DTX-200 system automatically scans 24 mm of the radius and the ulna, proximal from the point with an 8 mm radius-ulna gap. Details regarding BMD measurements have been described previously¹². The Shibata Comprehensive Health Care Service Center had checked the intra-coefficient of variation (CV) of DTX-200 measurements using a standard phantom every morning during health check-up examinations throughout the study period and confirmed that values were within $\pm 2.0\%$. The long-term CV of the Shibata Comprehensive Health Care Service Center is reported to be 0.8%¹³. Age, sex, weight, and height were obtained from health check-up records. BMI was calculated as weight (kg) divided by the square of height (m²).

Statistical analysis

Mean and standard deviation (SD) were calculated to characterize BMD and BMI by year. Mean BMDs by age group and year were then subjected to a trend test using simple linear regression analysis. The statistical significance of secular changes in BMD from 1995 to 2013 was tested with simple and multiple linear regression analyses using the following models: 1) unadjusted, 2) age-adjusted, and 3) age- and BMI-adjusted models. In addition, we conducted the same analysis for secular changes from 1995 to 2008 as a sensitivity analysis. From 2009, the local government canceled subsidies for BMD measurements, and thus the number of subjects with BMD measurements from 2009 onward steeply decreased. Under these circumstances, subjects might have had more interest in their bone health after 2009 compared to before 2009 (i.e., potential self-selection bias). Statistical analyses

were conducted with the SAS statistical package (release 9.4, SAS Institute Inc., Cary, NC, USA). $P < 0.05$ was considered statistically significant.

Results

Table 1 shows age, BMI, and BMD by year. The number of subjects was the largest in the first year of the study period (1995), and decreased gradually thereafter. The number of subjects in 2009

Table 1. Subject age, body mass index (BMI), and forearm bone mineral density (BMD) by year

	N	Mean (SD)		
		Age	BMI (kg/m ²)	BMD (g/cm ²)
1995	1,134	49.1 (11.2)	22.3 (3.0)	0.441 (0.072)
1996	1,026	50.1 (11.9)	22.6 (3.0)	0.437 (0.075)
1997	1,053	46.5 (13.3)	22.2 (2.9)	0.439 (0.074)
1998	977	48.6 (13.1)	22.3 (3.1)	0.433 (0.074)
1999	682	53.5 (10.7)	22.4 (3.0)	0.421 (0.075)
2000	624	51.9 (10.0)	22.4 (2.9)	0.437 (0.076)
2001	681	49.7 (10.3)	22.4 (3.2)	0.442 (0.075)
2002	602	51.5 (9.5)	22.2 (3.1)	0.434 (0.072)
2003	588	52.1 (10.4)	22.5 (2.9)	0.430 (0.073)
2004	488	52.4 (10.1)	22.4 (3.1)	0.429 (0.078)
2005	440	53.9 (10.2)	22.2 (3.2)	0.421 (0.081)
2006	521	56.8 (10.0)	22.4 (3.2)	0.409 (0.081)
2007	585	56.0 (10.1)	22.4 (3.0)	0.407 (0.077)
2008	518	56.5 (9.8)	22.5 (3.3)	0.405 (0.079)
2009	204	59.3 (10.1)	22.2 (3.3)	0.384 (0.085)
2010	153	58.1 (9.5)	22.3 (3.0)	0.400 (0.093)
2011	146	58.5 (10.5)	22.0 (2.8)	0.399 (0.082)
2012	133	56.9 (9.2)	21.9 (3.2)	0.417 (0.077)
2013	94	58.1 (9.3)	21.9 (3.1)	0.411 (0.082)

sharply decreased from the number in 2008 (a decrease of 60.6%). Mean BMIs were 20.4 (SD, 2.6, $N=430$), 21.1 (SD, 2.9, $N=1,197$), 21.9 (SD, 3.0, $N=2,257$), 22.7 (SD, 3.1, $N=3,342$), 22.9 (SD, 3.0, $N=3,048$), and 23.1 kg/m² (SD, 3.0, $N=375$) for women in their 20's, 30's, 40's, 50's, 60's, and 70's, respectively.

Table 2 shows secular changes in mean BMD by age group, with a graphical representation presented in Figure 1. Data for subjects in their 20's and 70's are not included given the small sample size. BMDs of subjects in their 40's (P for trend=0.0054) and 70's (P for trend=0.0225) decreased significantly from 1995 to 2013.

Secular changes in BMD from 1995 to 2013 by age group were examined by regression coefficients (β) in linear regression analyses (Table 3). The BMD of subjects in their 40's decreased significantly in unadjusted (P for trend=0.0054) and age-adjusted (P for trend=0.0162) models, but not in the age- and BMI-adjusted model (P for trend=0.2171). The BMD of subjects in their 50's decreased marginally in the age-adjusted model (P for trend=0.0535), but not in the age- and BMI-adjusted model (P for trend=0.6601).

Secular changes in BMD from 1995 to 2008 by age group are also shown in Table 3. The BMD of subjects in their 40's decreased significantly in unadjusted (P for trend=0.0161) and age-adjusted models (P for trend=0.0445), but not in the age- and BMI-adjusted model (P for trend=0.4186). The BMD of subjects in their 50's decreased significantly in the age-adjusted model (P for trend=0.0090), but not in the age- and BMI-adjusted model (P for trend=0.4648). Significant changes in BMD were

Table 2. Forearm bone mineral density of women by age group and year

Year	20's			30's			40's			50's			60's			70's		
	n	Mean	SD	n	Mean	SD	n	Mean	SD	n	Mean	SD	n	Mean	SD	n	Mean	SD
1995	59	0.447	0.049	156	0.464	0.045	365	0.481	0.051	318	0.440	0.068	221	0.370	0.065	15	0.315	0.032
1996	60	0.474	0.051	136	0.467	0.048	250	0.483	0.052	280	0.440	0.068	289	0.377	0.069	11	0.334	0.101
1997	124	0.466	0.048	244	0.469	0.047	210	0.480	0.055	204	0.447	0.065	259	0.365	0.065	12	0.278	0.060
1998	82	0.475	0.045	207	0.467	0.051	157	0.480	0.051	244	0.439	0.065	278	0.368	0.065	9	0.316	0.038
1999	22	0.475	0.062	46	0.460	0.049	124	0.475	0.045	237	0.439	0.064	239	0.370	0.063	14	0.300	0.062
2000	16	0.469	0.058	51	0.473	0.043	138	0.489	0.048	259	0.441	0.069	150	0.375	0.069	10	0.308	0.057
2001	27	0.456	0.040	87	0.465	0.047	174	0.486	0.058	255	0.446	0.067	136	0.361	0.066	2	0.376	0.062
2002	14	0.446	0.035	48	0.447	0.050	134	0.475	0.050	264	0.443	0.070	137	0.376	0.066	5	0.325	0.059
2003	14	0.463	0.054	48	0.461	0.048	136	0.471	0.049	226	0.440	0.065	143	0.376	0.069	21	0.320	0.061
2004	2	0.494	0.011	55	0.458	0.051	104	0.478	0.052	193	0.444	0.064	119	0.362	0.076	15	0.325	0.050
2005	2	0.386	0.026	37	0.467	0.055	91	0.474	0.054	174	0.438	0.066	124	0.356	0.073	12	0.305	0.060
2006	2	0.450	0.059	24	0.472	0.054	78	0.471	0.061	159	0.439	0.065	206	0.374	0.070	52	0.331	0.071
2007	2	0.521	0.019	19	0.458	0.053	112	0.474	0.056	163	0.430	0.057	232	0.371	0.065	57	0.340	0.074
2008	2	0.510	0.096	22	0.467	0.047	78	0.474	0.049	150	0.435	0.066	223	0.366	0.068	43	0.334	0.052
2009	0	-	-	5	0.447	0.064	29	0.466	0.052	55	0.414	0.072	83	0.363	0.068	32	0.301	0.076
2010	0	-	-	5	0.540	0.068	18	0.472	0.052	50	0.452	0.071	62	0.356	0.072	18	0.301	0.055
2011	1	0.485	-	4	0.484	0.061	24	0.472	0.052	31	0.445	0.064	65	0.362	0.071	21	0.341	0.062
2012	1	0.365	-	2	0.515	0.036	18	0.475	0.059	51	0.441	0.070	51	0.373	0.067	10	0.395	0.074
2013	0	-	-	1	0.478	-	17	0.474	0.068	29	0.455	0.063	31	0.377	0.060	16	0.329	0.063




Fig. 1. Secular trends in mean forearm bone mineral density (BMD) by age group. Results of statistical tests are presented in Table 3. The number of subjects with BMD measurements from 2009 onward steeply decreased.

Table 3. Regression coefficients (β) of year for forearm bone mineral density as an outcome variable calculated by simple and multiple linear regression analyses between 1995 and 2013

Age group	Predictor variable, manner of adjustment	1995-2013		1995-2008	
		Regression Coefficient	P for trend	Regression coefficient	P for trend
30's	Year, unadjusted	-0.000018	0.9642	-0.000435	0.2990
	Year, age-adjusted	-0.000084	0.8309	-0.000492	0.2412
	Year, age- & BMI-adjusted	-0.000232	0.5173	-0.000627	0.1030
40's	Year, unadjusted	-0.000672	0.0054	-0.000673	0.0161
	Year, age-adjusted	-0.000587	0.0162	-0.000569	0.0445
	Year, age- & BMI-adjusted	-0.000274	0.2171	-0.000208	0.4186
50's	Year, unadjusted	-0.000279	0.2718	-0.000399	0.1868
	Year, age-adjusted	-0.000453	0.0535	-0.000729	0.0090
	Year, age- & BMI-adjusted	0.000097	0.6601	-0.000191	0.4648
60's	Year, unadjusted	-0.000279	0.2541	-0.000200	0.5016
	Year, age-adjusted	-0.000144	0.5490	-0.000120	0.6804
	Year, age- & BMI-adjusted	0.000234	0.2890	0.000033	0.9017

not observed for subjects in their 30's or 60's.

Secular changes in mean BMI from 1995 to 2013 by age group are shown in Figure 2. A simple linear regression analysis showed that BMI decreased significantly for subjects in their 40's ($\beta = -0.050$, P for trend = 0.0002), 50's ($\beta = -0.076$, P for trend < 0.0001), and 60's ($\beta = -0.042$, P for trend < 0.0001). Significant changes in BMI were not observed for subjects in their 30's ($\beta = 0.027$, $P = 0.2399$).

Discussion

The present study investigated secular changes in forearm BMD and BMI of adult Japanese women from 1995 to 2013, with the following findings: 1) age-adjusted BMD of subjects in their 40's and 50's decreased, 2) their age- and BMI-adjusted BMD did not decrease, and 3) BMI of subjects in their 40's, 50's, and 60's decreased.

Only a few studies have reported on secular changes in BMD. Xu and Wu⁷ studied hip BMD in




Fig. 2. Secular trends in mean body mass index (BMI) by age group.

BMI decreased significantly in subjects in their 40's ($\beta = -0.050$, $P = 0.0002$), 50's ($\beta = -0.076$, P for trend < 0.0001), and 60's ($\beta = -0.042$, P for trend < 0.0001), but not in subjects in their 30's ($\beta = 0.027$, P for trend $= 0.2399$). The number of subjects with BMD measurements from 2009 onward steeply decreased.

US subjects (age ≥ 30 years [mean age, 53 years]) who participated in the NHANES from 2005 to 2014, and found that their BMD decreased during that period, particularly between 2009 and 2014. The authors suggested that this decrease in BMD corresponded to an increased trend of hip fracture in women aged ≥ 65 years between 2012 and 2015⁶, and that BMD is a strong predictor of hip fracture⁷. While our findings are consistent with this previous report⁷, our study has the advantage of having conducted age-stratified analyses of BMD changes. Contrary to the results of Xu and Wu's study⁷ and ours, Cheung *et al.*⁸ reported an increase in hip BMD in Hong Kong women aged ≥ 20 years (mean age, 47 years) across the entire adult age range from 1995–2000 to 2005–2010. Their finding was partly consistent with the long-term reduction in hip fracture incidence in Hong Kong¹⁴. In sum, these previous studies showed that long-term changes in adult BMD are associated with changes in hip fracture occurrence. In our study, women in their 40's and 50's showed a decreasing BMD trend between 1995 and 2008. It will be informative to observe the changing trend of osteoporotic fracture, including hip fracture, in the next few decades (from 2021) and determine whether the decreasing BMD trend reflects the changes in fracture trend.

Although age-adjusted BMDs of subjects in their 40's and 50's decreased during our study period, they did not significantly decrease when further

adjusted for BMI. This suggests that the decrease in BMD was due in part to the decrease in BMI. Although the distal forearm, which we used to measure BMD, is not a weight-bearing site, forearm BMD has been reported to correlate well with BMI and body weight^{15,16}.

The secular decrease in BMI is not limited to the present population, but rather is a phenomenon observed throughout Japan. According to the National Health and Nutrition Survey in Japan, the BMI of middle-aged women apparently decreased during our study period¹⁷. The long-term decrease in BMI was associated with a combination of decreased energy intake and worsened exercise habits, as well as breakfast skipping¹⁷. Important factors associated with low BMI in women include a desire for thinness and dieting behavior¹⁸, which are prevalent not only among young women but also among middle-aged and older women¹⁹. Moreover, dieting behavior itself has been suggested to have an unfavorable effect on bone strength, including BMD¹⁹. The National Health and Nutrition Survey also reported a long-term decrease in calcium intake in middle-aged women, which may be involved in the decrease in BMD we observed¹⁷. Thus, controlling dietary factors is a good strategy for bone health maintenance.

We did not observe a significant long-term decrease in BMD of subjects in their 60's, although their BMIs decreased significantly. While the rea-

son for this inconsistency is unclear, we speculate that menopausal status and healthy lifestyles may have played a role. Women in their 60's are mostly post-menopausal and experience estrogen insufficiency. However, the extent of decrease in estrogen in post-menopausal women varies, which may have impacted their long-term BMD changes.

Notably, there were some differences between the present female study population and the general Japanese female population with respect to BMI. Mean BMIs reported by the National Health and Nutrition Survey, Japan, 2004 (corresponding to the midpoint of the present study period) were 21.0 (SD 3.0, $N=121$), 22.6 (SD 3.6, $N=127$), 23.0 (SD 3.2, $N=167$), and 23.4 kg/m² (SD 3.5, $N=170$) for women in their 30's, 40's, 50's, and 60's, respectively²⁰. In comparison, BMIs in the present study for women in their 40's (22.7 kg/m²) and 60's (22.9 kg/m²) were slightly, but significantly, lower than the national average. These differences may have resulted from the present study population being a health-check population (i.e., those with potentially healthier lifestyles²¹). Thus, caution should be exercised when interpreting the results of the present study.

This study has some limitations. First, the decreasing trend in BMD may have been due to selection bias because the number of participants decreased over time during the study period. It is also possible that more women who were worried about their low BMD may have participated in the health check-up examinations later in the study period. Nonetheless, even if there was selection bias, it might not be associated with decreased BMD because the decline in number of subjects after 2008 did not bring about a corresponding decrease in BMD, but rather an increase in BMD. Second, we did not assess secular changes in some age groups given the limited sample size. In particular, changes in BMD of subjects in their 30's tended to decrease, but not significantly. The 20's and 30's are important ages in terms of bone acquisition and maintenance, and thus should be investigated further. Fourth, we could not assess factors other than BMI, such as lifestyles and anti-osteoporotic medications. Nevertheless, their confounding effects may be limited, as evidenced by Japanese community studies^{22,23} reporting that lifestyle factors are much less influential on adult BMD than BMI (or body weight). With regard to anti-osteoporotic medications that affect BMD, such as bisphosphonates, we previously reported in a community study of BMD²² that 11 of 461 (2.4%) women aged between 60 and 75 years used such medications,

whereas none of the women in their 50's (213 women) used them. Therefore, our results may be biased in terms of the use of anti-osteoporotic medications in women aged over 60 years. Fifth, we conducted several trend tests, which may have resulted in Type I error (i.e., false positives). Finally, we measured cortical BMD, but not spongy BMD. Therefore, while our results apply to osteoporosis of long bones, they may not apply to vertebral bones.

Our study has several implications. First, BMD is an important predictor of osteoporotic fracture, and a good indicator that reflects lifestyle factors. However, long-term trends of BMD have rarely been reported worldwide, and no reports on this topic from Japan have been published. Second, we observed a long-term decrease in BMD in middle-aged women during the period spanning 1995–2013, which could serve as an alert for a potential increase in osteoporosis risk several decades later because hip fracture typically occurs in older adults aged ≥ 80 years. Finally, although we observed a long-term decrease in BMD being mediated by BMI, the present study design did not allow us to determine a causal association. Nevertheless, such an association has been suggested by previous Japanese cohort studies^{16,24}, and the results of the present study underscore the importance of BMI as a bone mass indicator.

Conclusions

The secular decrease in BMD in middle-aged Japanese women during the period spanning 1995–2013 was partly attributed to decreases in BMI. Measures to help maintain suitable BMI will be necessary to prevent a decrease in BMD among women.

Acknowledgements

The authors thank the staff of Shibata Comprehensive Health Care Service Center for their assistance with data collection. None of the authors have conflicts of interest to report.

Disclosure

The authors declare no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

References

1. NIH Consensus Development Panel on Osteoporosis

- sis Prevention, Diagnosis, and Therapy. Osteoporosis prevention, diagnosis, and therapy. *JAMA*, **28** : 785-795, 2001.
2. Odén A, McCloskey EV, Johansson H, Kanis JA. Assessing the impact of osteoporosis on the burden of hip fractures. *Calcif Tissue Int*, **92** : 42-49, 2013.
 3. Orimo H, Yaegashi Y, Hosoi T, *et al.* Hip fracture incidence in Japan : Estimates of new patients in 2012 and 25-year trends. *Osteoporos Int*, **27** : 1777-1784, 2016.
 4. Miyasaka D, Endo N, Endo E, *et al.* Incidence of hip fracture in Niigata, Japan in 2004 and 2010 and the long-term trends from 1985 to 2010. *J Bone Miner Metab*, **34** : 92-98, 2016.
 5. Cooper C, Cole ZA, Holroyd CR, *et al.* Secular trends in the incidence of hip and other osteoporotic fractures. *Osteoporos Int*, **22** : 1277-1288, 2011.
 6. Lewiecki EM, Wright NC, Curtis JR, *et al.* Hip fracture trends in the United States, 2002 to 2015. *Osteoporos Int*, **29** : 717-722, 2018.
 7. Xu Y, Wu Q. Decreasing trend of bone mineral density in US multiethnic population : analysis of continuous NHANES 2005-2014. *Osteoporos Int*, **29** : 2437-2446, 2018.
 8. Cheung E, Bow C, Loong C, *et al.* A secular increase in BMD in Chinese women. *J Bone Miner Metab*, **32** : 48-55, 2014.
 9. Nakamura K, Hoshino Y, Watanabe A, *et al.* Eating problems and related weight control behaviour in adult Japanese women. *Psychother Psychosom*, **68** : 51-55, 1999.
 10. Santos I, Sniehotta FF, Marques MM, Carraça EV, Teixeira PJ. Prevalence of personal weight control attempts in adults : a systematic review and meta-analysis. *Obes Rev*, **18** : 32-50, 2017.
 11. Ministry of Health, Labour, and Welfare, Japan : The National Health and Nutrition Survey in Japan, 2017. Ministry of Health, Labour, and Welfare, Tokyo, 2018. (in Japanese)
 12. Nakamura K, Tanaka Y, Saitou K, Nashimoto M, Yamamoto M. Age and sex differences in the bone mineral density of the distal forearm based on health check-up data of 6343 Japanese. *Osteoporos Int*, **11** : 772-777, 2000.
 13. Nakamura K, Kazama JJ, Tanaka Y, *et al.* Microscopic hematuria is associated with low bone mineral density in aged women and men. *J Bone Miner Metab*, **27** : 251-254, 2009.
 14. Kung AWC, Yates S, Wong V. Changing epidemiology in hip fracture rate in Hong Kong. *Arch Osteoporos*, **2** : 43-47, 2007.
 15. Nakamura K, Saito T, Nishiwaki T, *et al.* Correlations between bone mineral density and demographic, lifestyle, and biochemical variables in community-dwelling Japanese women 69 years of age and over. *Osteoporos Int*, **17** : 1202-1207, 2006.
 16. Nakamura K, Oyama M, Saito T, *et al.* Nutritional and biochemical parameters associated with 6-year change in bone mineral density in community-dwelling Japanese women aged 69 years and older : The Muramatsu Study. *Nutrition*, **28** : 357-361, 2012.
 17. Ministry of Health, Labour and Welfare, Japan : The National Health and Nutrition Survey in Japan, 2013. Ministry of Health, Labour, and Welfare, Tokyo, 2015. (in Japanese)
 18. Hayashi F, Takimoto H, Yoshita K, Yoshiike N. Perceived body size and desire for thinness of young Japanese women : a population-based survey. *Br J Nutr*, **96** : 1154-1162, 2006.
 19. Tatsuno I, Terano T, Nakamura M, *et al.* Lifestyle and osteoporosis in middle-aged and elderly women : Chiba bone survey. *Endocr J*, **60** : 643-650, 2013.
 20. Ministry of Health, Labour and Welfare, Japan : The National Health and Nutrition Survey in Japan, 2004. Ministry of Health, Labour, and Welfare, Tokyo, 2006. (in Japanese)
 21. Iwasaki M, Yamamoto S, Otani T, *et al.* Generalizability of relative risk estimates from a well-defined population to a general population. *Eur J Epidemiol*, **21** : 253-262, 2006.
 22. Nakamura K, Tsugawa N, Saito T, *et al.* Vitamin D status, bone mass, and bone metabolism in home-dwelling postmenopausal Japanese women : Yokogoshi Study. *Bone*, **42** : 271-277, 2008.
 23. Nakamura K, Ueno K, Nishiwaki T, *et al.* Nutrition, mild hyperparathyroidism, and bone mineral density in young Japanese women. *Am J Clin Nutr*, **82** : 1127-1133, 2005.
 24. Kitamura K, Nakamura K, Saito T, *et al.* High serum 25-hydroxyvitamin D levels do not retard postmenopausal bone loss in Japanese women : the Yokogoshi Study. *Arch Osteoporos*, **8** : 153, 2013.