

Research Article

Genetic Variability of *Aspergillus flavus* Isolates from a Mississippi Corn Field

Cesar D. Solorzano,¹ Hamed K. Abbas,¹ Robert M. Zablotowicz,²
Perng-Kuang Chang,³ and Walker A. Jones¹

¹ Biological Control of Pests Research Unit, United States Department of Agriculture, Agricultural Research Service, Stoneville, MS 38776, USA

² Crop Production Systems Research Unit, United States Department of Agriculture, Agricultural Research Service, Stoneville, MS 38776, USA

³ Food and Feed Safety Research Unit, United States Department of Agriculture, Agricultural Research Service, New Orleans, LA 70124, USA

Correspondence should be addressed to Hamed K. Abbas; hamed.abbas@ars.usda.gov

Received 24 July 2014; Accepted 5 October 2014; Published 12 November 2014

Academic Editor: Dongsheng Zhou

Copyright © 2014 Cesar D. Solorzano et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

A nontoxigenic *Aspergillus flavus* strain, K49, is currently being tested as a biological control agent in corn fields in the Mississippi Delta. However, little is known about the overall genetic diversity of *A. flavus* from year to year in corn fields and specifically in Mississippi. Our objective was to assess the genetic variability of *A. flavus* isolates from different seasons, inoculum sources, and years, from a no-till corn field. Of the 175 *A. flavus* isolates examined, 74 and 97 had the typical *norB-cypA* type I (1.5 kb) and type II (1.0 kb) deletion patterns, respectively. Variability in the sequence of the *omtA* gene of the majority of the field isolates ($n = 118$) was compared to strain K49. High levels of haplotypic diversity (24 *omtA* haplotypes; $Hd = 0.61 \pm 0.04$) were found. Among the 24 haplotypes, two were predominant, H1 ($n = 71$), which consists of mostly toxigenic isolates, and H49 ($n = 18$), which consists of mostly atoxigenic isolates including K49. Toxigenic isolates were prevalent (60%) in this natural population. Nonetheless, about 15% of the population likely shared the same ancestral origin with K49. This study provides valuable information on the diversity of *A. flavus*. This knowledge can be further used to develop additional biological control strains.

1. Introduction

Corn, *Zea mays* L., is a major agricultural crop in the state of Mississippi and in the U.S., with an estimated acreage of 770,000 and 91,897,000 acres planted in 2011, respectively [1]. One major food safety concern for corn, peanuts, rice, and edible tree nuts is the contamination with toxins produced by *Aspergillus flavus* Link [2]. *Aspergillus flavus* is a soil-borne haploid fungus, known to asexually produce conidia and sclerotia and recently reported to have a sexual stage [3, 4]. The fungus and other *Aspergillus* spp. in section *Flavi* can synthesize aflatoxins and cyclopiazonic acid (CPA) [5, 6]. Cyclopiazonic acid is able to inhibit mammalian calcium-ATPase and interfere with Ca^{2+} level in animal systems [7].

Aflatoxins B₁ (AFB₁) and B₂ (AFB₂) are produced by *A. flavus*, and aflatoxins G₁ (AFG₁) and G₂ (AFG₂) in addition to the B aflatoxins are produced by *A. parasiticus* [8–10]. *A. flavus* isolates with large deletions of the aflatoxin gene cluster and portions of the subtelomeric region generally result in no aflatoxin and/or CPA production [11].

Aflatoxins can be more problematic for corn grain and residues in the southern U.S. than other regions due to subtropical climate and summer precipitation deficit of the former region [12, 13]. In addition, 70% of corn production uses some form of reduced tillage, therefore increasing the corn residues contaminated with *A. flavus* left in the field for the next growing season. The U.S. Food and Drug Administration (FDA) regulatory limit for aflatoxin levels in

corn feed used in finishing beef cattle is 300 ng/g and for corn used for human consumption is 20 ng/g [14, 15].

A biological control approach to reduce the impact that toxigenic *A. flavus* strains have on corn production is to use nontoxigenic *A. flavus* strains to displace the toxigenic strains in the field [16]. These nontoxigenic strains include K49 (NRRL 30797) and the commercially available AF36 (NRRL 18543) and afla-guard (NRRL21882) which are approved by U.S. Environmental Protection Agency (EPA) for use in a variety of crops in the U.S. These strains competitively exclude aflatoxin producers by applying them to soil and plant parts during the vegetative growth stages [17]. International efforts in Argentina [18], Nigeria [19], and China [20] are currently underway to identify prospective biocontrol strains.

Aflatoxins in *A. flavus* strains are synthesized by a group of approximately 25 genes that form the aflatoxin gene cluster, a region spanning 65–70 kb [10, 25]. Genetic variability occurs in the aflatoxin gene cluster of *A. flavus* isolates. Partial gene deletions in the *norB-cypA* gene region result in known deletion patterns I and II [26]. The biocontrol strain AF36 contains all genes in the aflatoxin biosynthesis gene cluster, with a 90% homology to that of aflatoxin producing *A. flavus* [27]. The basis for its nontoxigenicity is a single nucleotide mutation that results in a stop codon near the beginning of the coding sequence of *pksA* which encodes a polyketide synthase [28]. Previous studies have confirmed that K49 also contains the whole aflatoxin gene cluster [29]. A recent study reports that the same *pksA* missense mutation occurs in K49 [30].

A variety of techniques have been used to examine population diversity of toxigenic and nontoxigenic strains present in various crops and geographical locations. They include amplified fragment length polymorphism (AFLP) [31], random amplified fragment polymorphism (RAPD) [32, 33], direct sequencing, intersimple sequence repeats (ISSR) [34], and microsatellite markers [20, 35, 36]. Eight deletion patterns in the aflatoxin gene cluster have been reported for a group of 38 nonaflatoxigenic *A. flavus* isolates [37]. In a group of soil field isolates from Mississippi corn fields, 60% of *A. flavus* isolated had the potential to produce aflatoxins and all isolates expressed *aflD*, *aflG*, *aflP*, *aflR*, and *aflS* aflatoxin genes [29]. Recently, the presence of mating type genes has been determined (*MAT1-1* or *MAT1-2*) in *A. flavus* [38]. Moreover, another important target of research is the quantification of *A. flavus* by real-time PCR that targets the *nor-1* gene from the aflatoxin gene cluster [39].

The objective of this research was to assess the temporal genetic variability of *A. flavus* isolates from a no-tilled corn field in Mississippi using two markers, *norB-cypA* and *omtA* genes from the aflatoxin gene cluster. Single nucleotide polymorphism (SNP) in the *omtA* gene has been shown to have a sufficient discriminatory power in typing strains of *A. flavus* and closely related *A. oryzae* [37]. Knowledge of this genetic information will help to better understand the community of isolates in a given corn field with no previous exposure to nontoxigenic strains and to determine if additional local isolates could be developed for biological control programs.

2. Materials and Methods

2.1. Collection of *Aspergillus flavus* Isolates and Confirmation of Aflatoxin Production. One hundred-seventy five *A. flavus* isolates were obtained from soil, plant parts (e.g. silk, leaves, cobs, seeds, and tassels), and adult beetles collected randomly from a no-till Bt hybrid corn field in Elizabeth, MS, during the growing (May–October) and overwintering (December–April) seasons of 2006–2009. Soil isolates were obtained as described elsewhere by diluting 10 g of soil into 150 mL of a solution of 2 g of agar/L of water [40]. 100 μ L of each sample was transferred to modified dichloronitroaniline rose bengal agar (MDRB) with 3% NaCl and the inoculum was distributed uniformly with a sterile spreader and incubated at 37°C for 5 days [41, 42]. Putative *A. flavus* isolates were transferred into potato dextrose agar (PDA) plates amended with 0.3% β -cyclodextrin to enhance fluorescence and incubated at 28°C for 5 days in the dark. *Aspergillus flavus* identification was based on morphological characteristics [43]. Isolates were examined for fluorescence under UV light (365 nm) and subjected to biochemical aflatoxin confirmation with the ELISA Veratox test kit [12, 44].

2.2. DNA Extraction, PCR Amplification, and Sequencing. A single agar plug containing spores was transferred into a sterile 2 mL centrifuge tube containing 1 mL of 0.05% Triton X-100 solution. The tube was vortexed vigorously and 100 μ L of the spore suspension was inoculated into 25 mL of potato dextrose broth (PDB) in a 50 mL plastic tube. Tubes were incubated horizontally for 4-5 days at 30°C at 150 rpm.

After visible fungal growth, mycelia was vacuum-dried. Approximately 100–150 mg of wet mycelia was used for DNA extraction using the ZR Fungal/Bacterial DNA kit (Zymo Research, Orange County, CA). DNA extraction followed the manufacturer's instructions in which cells were lysed by bead beating for 5 min with a Disruptor Genie (Zymo Research, Orange County, CA). Eluted DNA was the template for PCR amplification of partial fragments of *norB-cypA* (300/800 bp) and *omtA* (594 bp) genes of the aflatoxin gene cluster and *ITS* (600 bp) gene regions. Platinum Taq DNA Polymerase High Fidelity (Life Technologies, 5791 Van Allen Way, Carlsbad, CA, 92008) was used for the PCR reaction.

Primer pairs were *norB-cypA*-F 5'-GTGCCAGCA-TCTTGGTCCA-3', *norB-cypA*-R 5'-AGGACTTGATGAT-TCCTCGTC-3' [27], *omtA*-F 5'-CAGGATATCATTTGTG-GACGG-3', *omtA*-R 5'-CTCCTCTACCAGTGGCTTCG-3' [36], *ITS1* 5'-TCCGTAGGTGAACCTGCGG-3', and *ITS4* 5'-TCCTCCGCTTATTGATATGC-3' [20]. ITS sequences of selected samples were used for species confirmation. PCR was conducted following the profile of 2 min at 95°C, followed by 30 cycles of 94°C for 60 s, 45°C for 60 s, and 72°C for 90 s, with a final extension of 72°C for 5 min. Amplicons were separated in 1.5% agarose gels in TBE and photographed using a BioDoc-It Imaging System (UVP Inc., Upland, CA). PCR products for sequencing were gel extracted using Quantum Prep Freeze 'N Squeeze DNA Gel extraction spin columns (Biorad Laboratories, Hercules, CA) and sequenced at the

TABLE 1: *Aspergillus flavus* isolates collected in Elizabeth, MS, from 2006 through 2009, UV fluorescence and ELISA aflatoxin detection (+/-), *norB-cypA* amplicon fragment size, and *omtA* haplotype frequency.

Sources	Time period	AF (+)	AF (-)	<i>norB-cypA</i> patterns	<i>omtA</i> haplotypes (frequency)
Soil	Jan.-Apr. 2006	140, 161, 163-172	159, 160, 162, 173	4 ^b 12 ^c	H1 (10) ^d , H6 (1), H14 (1), H17 (1), H49 (2)
	May-Oct. 2006	93-99	90-92	4 ^b 6 ^c	H1 (5), H7 (1), H11 (1), H49 (1)
	Jan.-Apr. 2007	56-58, 60, 76-80, 141, 143-146, 148, 149, 153-155	59, 81, 82, 142, 147, 150-152, 156	7 ^b 21 ^c	H1 (15), H16 (1), H19 (1), H49 (6)
	May-Oct. 2007	101, 105, 107, 110, 115, 123, 138, 139	100, 108	2 ^b 8 ^c	H1 (7), H8 (1), H18 (1), H49 (1)
	Jan.-Apr. 2008	61, 62, 64, 83-88, 157, 158, 175-177, 180	63, 65-67, 174, 178, 179	6 ^b 14 ^c	H1 (15), H49 (2)
	May-Oct. 2008	111, 113, 116, 117, 120, 121, 127, 136	132	3 ^b 6 ^c	H1 (4), H5 (1), H15 (1), H21 (1), H49 (1)
	Jan.-Apr. 2009	55, 73	50-54, 68-72, 74, 75, 89	8 ^b 7 ^c	H1 (2), H2 (1), H49 (2)
Above ground ^a	May-Oct. 2006	124, 128, 134, 137	102, 106, 114, 122, 131, 135	6 ^b 4 ^c	H1 (3), H3 (1), H4 (1), H7 (2), H49 (2)
	May-Oct. 2008	104, 109, 112, 118, 125, 126, 129, 133	103, 119	2 ^b 8 ^c	H1 (6), H20 (1), H22 (1), H23 (1), H49 (1)
Plant debris	Jan.-Apr. 2006	1-8	9-14	7 ^b 7 ^c	H1 (4), H12 (1)
	Jan.-Apr. 2007	19, 22, 24-26, 28, 29	16-18, 20, 21, 23, 27, 30	15 ^b	H8 (2), H9 (1), H10 (1)
	Jan.-Apr. 2008	34, 36-38, 40-45	31-33, 35, 39	11 ^b 4 ^c	H8 (2), H13 (2)
Combined		105	66	74 ^b , 97 ^c	118 ^e

^a Above ground corresponding to isolates from silk, tassel, ear, and insect.

^b 300 bp fragment, deletion pattern I.

^c 800 bp fragment, deletion pattern II.

^d Number of isolates belonging to the indicated haplotype.

^e Number of isolates examined for *omtA* amplicon.

Genomics and Bioinformatics Research Unit of the USDA-ARS in Stoneville, Mississippi.

2.3. Alignment, Population Genetic Analysis, and Population Structure. DNA sequences were aligned with SeqMan Pro in DNASTAR (Madison, WI). *omtA* haplotypes were determined manually. The program DNAsp version 5.10.01 [45] was used to estimate population genetic parameters: haplotype diversity (Hd), nucleotide diversity (π), mean number of pair wise nucleotide differences (K), theta per site (θ_s), and theta per gene (θ_g). Levels of gene flow were determined through the effective number of migrants (N_m) according to the formula of Nei [24] across years and inoculum source. Parameters that test for population expansion or selection: D^+ and F^+ statistics [21], Fu's F_s statistic, Strobeck S statistic [22], and Tajima's D [23] statistic, were also calculated with DNAsp.

The program Arlequin version 3.5.1.2 [46] calculated nucleotide differences between years and inoculum sources by estimating F_{ST} values and determined genetic structure with an analysis of molecular variance (AMOVA) with the

methodology from Excoffier et al. [47]. Phylogenetic relationships among isolates were established using the Maximum Likelihood (ML) analysis method in MEGA version 5 [48].

3. Results

3.1. Confirmation of Isolates and Deletion Patterns of the *norB-cypA* Gene Region. All 175 isolates were confirmed with morphological characteristics and also by *ITS* sequence data (>99.5% homology) to be *A. flavus*. Conidia of isolates appeared with a characteristic yellowish green to darker green. A total of 105 isolates that were positive for UV fluorescence were also positive for ELISA aflatoxin detection, and 66 were not positive for UV fluorescence and ELISA (Table 1). Amplification of the partial *norB-cypA* gene region also confirmed that isolates were *A. flavus*. From a total of 175 isolates, 171 were amplified for the *norB-cypA* gene region. The *norB-cypA* F/R primer pair generated a 300 bp and an 800 bp fragment for 74 and 97 *A. flavus* isolates that correspond to the 1.5 kb (type I) and a 1.0 kb (type II) deletion patterns, respectively (Table 1).

TABLE 2: *omtA* gene polymorphism statistics for *Aspergillus flavus* from a single Mississippi field.

Season/Source/Year	Sequence and haplotype				Nucleotide diversity				Population expansion				
	<i>N</i>	<i>S</i>	<i>H</i>	Hd ± sd	π	<i>K</i>	θ_s	θ_g	<i>D</i> ⁺	<i>F</i> ⁺	<i>F</i> _s	<i>S</i>	<i>D</i>
May–Oct./soil/2006	8	28	4	0.64 ± 0.18	0.020	12.17	0.018	10.79	0.96	0.99	4.86	0.05	0.67
May–Oct./soil/2007	10	26	4	0.53 ± 0.18	0.014	8.77	0.015	9.19	1.11	0.88	5.02	0.03	–0.21
May–Oct./soil/2008	8	39	5	0.78 ± 0.15	0.023	14.25	0.025	15.42	0.15	0.02	3.32	0.16	–0.40
May–Oct./above ground/2006	9	28	5	0.86 ± 0.08	0.022	13.27	0.017	10.30	1.32	1.51	3.92	0.09	1.44
May–Oct./above ground/2008	10	32	5	0.66 ± 0.16	0.011	6.88	0.020	12.01	–2.36*	–2.58*	2.45	0.23	–2.05*
Jan.–April/soil/2006	15	28	5	0.59 ± 0.14	0.015	9.27	0.014	8.80	0.92	0.84	5.34	0.02	0.22
Jan.–April/soil/2007	23	26	4	0.52 ± 0.09	0.016	9.84	0.011	7.04	1.20	1.51	11.4	0.00	1.49
Jan.–April/soil/2008	17	24	2	0.22 ± 0.12	0.008	5.29	0.011	7.09	1.59*	0.98	10.3	0.00	–1.02
Jan.–April/soil/2009	5	25	3	0.80 ± 0.16	0.024	14.80	0.020	12.00	1.73*	1.86*	4.74	0.09	1.73
Jan.–April/plant debris/2006	5	23	2	0.40 ± 0.23	0.015	9.20	0.018	11.04	–1.23*	–1.32	6.64	0.02	–1.23
Jan.–April/plant debris/2007	4	23	3	0.83 ± 0.22	0.001	1.00	0.001	1.09	–0.70	–0.60	–0.88	0.95	–0.70
Jan.–April/plant debris/2008	4	23	2	0.66 ± 0.20	0.025	15.33	0.021	12.54	2.28*	2.38*	6.76	0.03	2.28*
Combined	118	68	24	0.61 ± 0.04	0.018	11.08	0.023	13.94	–4.07*	–3.13*	1.83	0.20	–0.65

Note: *N* = number of sequences; *S* = number of variable sites; *H* = number of haplotypes; Hd = haplotype diversity ± standard deviation; π = nucleotide diversity; *K* = mean number of pairwise nucleotide differences; θ_s = theta per site; θ_g = theta per gene; *D*⁺ and *F*⁺ statistics (Fu and Li 1993) [21]; *F*_s Fu's *F*_s statistic; Strobeck *S* statistic (Strobeck 1987) [22]; *D* = Tajima's (1989) [23] statistic; * *P* < 0.05.

3.2. Genetic Analysis of *omtA* Gene Region. A total of 118 *A. flavus* isolates (67% of *n* = 175 isolates) were selected for *omtA* sequence analysis (Table 1). These isolates were selected because their *omtA* gene fragments were clearly amplified as revealed by gel electrophoresis. The subsequent SNP analysis identified 24 *omtA* haplotypes. Seventy-one of the 118 isolates belonged to a single haplotype designated as haplotype “1” (H1). This haplotype was observed in 10 of the 12 season/source/year locations. Likewise, haplotype H49, designated “49” because its sequence is exactly the same as K49, was found in 18 isolates in 9 of the 12 season/source/year locations but was not found in plant debris 2006–2008.

Haplotype diversity was high (Hd = 0.61) across sites and ranged from 0.22 to 0.86 (Table 2). The variable number of nucleotide sites (*S*) totaled 68 from the 597 nucleotide sites analyzed. Of those, singleton variable sites were 31 and parsimony informative sites were 37. Nucleotide diversity (π) was low, ranging between 0.001 and 0.025 with an average of 0.018. Other nucleotide diversity parameters were *K* = 11.08, θ_s = 0.023, and θ_g = 13.94 for the total population. The hypothesis of neutral mutation was tested with statistical parameters *D*⁺, *F*⁺, *F*_s, Strobeck's *S*, and Tajima's *D*. Except for a few parameters, most were not significant at *P* < 0.05, and hence no assumptions on population growth or selection can be made. Plant debris isolates from the overwintering season of 2007 appear to be most genetically distinct when compared to all other populations. *N*_M in these comparisons was low (≤ 1.29), which demonstrates low levels of gene flow, and 8 of the 11 *F*_{ST} comparisons of this group of isolates were significantly different from all other populations tested (Table 3).

3.3. Population Structure. AMOVA partitioned the variation among years, among populations within years and within populations. Variation was mostly explained by within populations (84.92%) (df = 106, *F*_{ST} = 0.15, *P* < 0.0009). Variation among populations within years accounted for 20.63% of the

variation (df = 8, *F*_{SC} = 0.19, *P* < 0.0009) and there was no variation among years (0%) (df = 3, *F*_{CT} = –0.05, *P* < 0.56).

3.4. Phylogenetic Analysis. Our ML phylogenetic analysis showed that isolates were not clustered based on the season, inoculum source, or year (Figure 1). For example, isolates 159 to 172 were collected from soil between January and April in 2006 (Table 1), but isolates of 161 and 163–172 are in a clade different from that of isolates 159, 160, and 162. Nonetheless, the 118 isolates formed two major clades based on the aflatoxin producing ability. H1 (isolates 1 to 5) is toxigenic and found to be most distantly related to H49 (H49 to 72). The branch that contains H1, at the upper part of the genealogy tree, includes the group of isolates that were confirmed to be toxigenic. Only 6 isolates within this group were nontoxigenic (isolates 66, 67, 70, 89, 90, 132, and 178) and however closely related to toxigenic isolates by analysis of the *omtA* gene region. Likewise, there were 7 toxigenic isolates (42, 73, 94, 97, 113, 137, and 148) among the 34 isolates that grouped within the nontoxigenic branch where H49 was positioned. These isolates are closely related to H49 but result in production of aflatoxin.

4. Discussion

In this study we assessed the population genetic diversity of *A. flavus* across years from a single field in Mississippi and determined how this population of isolates is related to the native Mississippi nontoxigenic isolate K49. H49 was present in 10 of the 12 locations, and therefore these isolates could present a shared origin. The other six isolates that were closely related to H49 were nontoxigenic and they may have mutations or deletions in other genes with the aflatoxin gene cluster that result in nontoxigenicity. We found that a high proportion (60%) of the populations are toxigenic isolates of

TABLE 3: *omtA* observed F_{ST} values (below) and gene flow (N_M) calculated with Nei (1982) [24] for *Aspergillus flavus* isolates collected from twelve spatiotemporal locations from a single corn field in Mississippi.

$F_{ST} N_{M \rightarrow}$	1	2	3	4	5	6	7	8	9	10	11	12
1	—	1.72	2.62	11.73	10.95	29.37	0.34	10.88	3.26	4.10	1.68	2.42
2	0.12	—	0.62	0.98	0.93	1.58	1.29	1.35	0.57	0.57	3.00	22.19
3	0.00	0.37*	—	4.51	3.78	4.77	0.11	3.11	3.86	4.74	0.58	0.65
4	-0.05	0.29*	-0.00	—	87.94	42.29	0.32	15.52	9.36	18.02	1.25	1.57
5	-0.07	0.27*	-0.02	-0.08	—	39.32	0.23	12.87	9.04	18.43	1.05	1.30
6	-0.06	0.22*	0.04	-0.04	-0.06	—	0.60	22.67	6.48	7.77	2.22	2.99
7	0.54*	0.17	0.77*	0.65*	0.67*	0.57*	—	0.34	0.15	0.18	0.54	0.70
8	-0.09	0.17	-0.01	-0.06	-0.08	-0.06	0.54*	—	5.18	6.05	1.42	1.94
9	0.02	0.40*	-0.02	-0.03	-0.05	0.01	0.77*	-0.02	—	17.39	0.63	0.74
10	0.03	0.44*	0.00	-0.03	-0.05	0.01	0.79*	-0.00	-0.05	—	0.73	0.83
11	0.08	-0.01	0.31	0.27*	0.24*	0.21	0.30	0.12	0.39*	0.45*	—	2.65
12	0.02	-0.15	0.29	0.19	0.18	0.12	0.25	0.06	0.33*	0.39*	-0.08	—

1 = growing season soil 2006; 2 = growing season above ground 2006; 3 = overwintering plant debris 2006; 4 = overwintering soil 2006; 5 = growing season soil 2007; 6 = overwintering soil 2007; 7 = overwintering plant debris 2007; 8 = growing season soil 2008; 9 = growing season above ground 2008; 10 = overwintering soil 2008; 11 = overwintering plant debris 2008; 12 = overwintering soil 2009; *significant at $P < 0.05$.

haplotype H1. These H1 toxigenic isolates are present in all years and inoculum sources. In contrast, Abbas et al. [40] observed a higher percent of toxigenic *A. flavus* isolates in soil than stover and cobs with grain; cobs had the lowest percent of toxigenic *A. flavus*. We speculate that this no-till field with continuous corn production has supported a buildup of a higher number of H1 toxigenic isolates over time probably in part due to their better competitiveness against *A. flavus* isolates of other haplotypes.

Aspergillus flavus populations from a corn field community can be diverse due to large amounts of propagule materials [49]. Conidia can be carried through air movement, and sclerotia can remain in the soil for the next growing season. However, we found that, in addition to H1, haplotype H49 is prevalent (15%) in soil and above ground sources but not in plant debris. The biocontrol strain K49 also belongs to the H49 haplotype. Hence, we can assume that H49 isolates from this Mississippi field share an ancestral origin with K49, which was isolated from corn in Mississippi [41]. H49 isolates are present in 10 of the 12 locations examined, which suggests their adaptability and survivability in the field. Other biocontrol candidate strains may be further selected from this particular haplotype. Application of a mixture of highly competitive biocontrol *A. flavus* strains has become a new paradigm for mitigating preharvest aflatoxin problems in some parts of world, especially in Nigeria [Joseph Atehnkeng personal communication, also see <http://www.aflasafe.com/> for details].

Our somewhat limited sample size might explain why a large number of haplotypes were found only once as evidence by the high levels of haplotype diversity. The importance and relative frequency of these single-isolate haplotypes cannot be ascertained unless a larger sample size is taken into consideration. Isolates, detected only once, might have an origin beyond this field and could have been carried in by wind or insects, thereby increasing the genetic diversity of this field. Besides H1 and H49, H7 and H8 are the only other

haplotypes found in more than one source. H7 was found in the growing season of 2006 in both soil and above ground inoculum, and H8 was found during the overwintering seasons of 2007 and 2008. Given that there was no structure found across years, variation in the populations cannot be explained as a year effect, but within a year, populations are variable according to their source.

Bayman and Cotty [50] found that 66/105 of *A. flavus* soil isolates comprised 13 vegetative compatibility groups in a single Arizona cotton field over a 3-year period and one VCG includes 20% of all isolates from the field. Unlike Gashgari et al. [51], who used RAPD analysis, we are able to differentiate toxigenic from nontoxigenic isolates using both analytical methods of UV detection and ELISA and, to a lesser extent, with the partial *omtA* gene sequence. Others like Midorikawa et al. [33] have been able to characterize *A. flavus* populations based on host plant. In the context of *A. flavus* diversity from a single corn field, the *omtA* marker provided a satisfactory level of resolution. Geiser et al. [52, 53] and Chang et al. [37] included this genetic marker as part of their studies when comparing *A. flavus* and *A. oryzae* isolates. Yin et al. [20] also included this genetic marker in the study that examined nontoxigenic peanut *A. flavus* isolates from a field in China. They found five deletion patterns in 56 toxigenic and nontoxigenic isolates from peanut fields and included 11 genes from the aflatoxin gene cluster in their analysis. The *omtA* and *norB-cypA* gene regions can be more commonly used by researchers, as a way to genetically characterize *A. flavus* isolates. The majority of genes in the aflatoxin gene cluster are highly conserved, as demonstrated by Ehrlich et al. [54] for *A. flavus* and the related aflatoxin producers of *A. nomius* and *A. parasiticus*. The atoxigenic isolates that are clustered with other toxigenic isolates (Figure 1) may have mutations or deletions in genes in the aflatoxin gene cluster, which results in nontoxigenicity [26].

This study provides useful information to the scientific community that studies *A. flavus* populations. The knowledge




FIGURE 1: Maximum likelihood analysis of *omtA* gene sequences for 118 *A. flavus* collected from a single field in Mississippi 2006–2009. (Scale is 0.005.)

of *A. flavus* genetic diversity can be used to develop additional biocontrol agents native to the local agricultural areas. Further work will determine if these atoxigenic HI isolates are genetically similar to other genes within the aflatoxin gene cluster, specially the *pkSA* gene that contains the missense

mutation that results in nontoxigenicity in the biocontrol strains K49 and AF36.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Acknowledgments

The authors thank Ms. Bobbie Johnson, Ms. Carol Benson, and Mr. Leon Hicks for technical scientific assistance. This research is a contribution to USDA National Program 303.

References

- [1] USDA-NASS, "USDA National Agricultural Statistics Service," Grain crops national and state data, <http://www.nass.usda.gov/>.
- [2] H. K. Abbas, *Aflatoxin and Food Safety*, CRC Press, Boca Raton, Fla, USA, 2005.
- [3] S. Amaike and N. P. Keller, "*Aspergillus flavus*," *Annual Review of Phytopathology*, vol. 49, pp. 10.1–10.27, 2011.
- [4] B. W. Horn, G. G. Moore, and I. Carbone, "Sexual reproduction in *Aspergillus flavus*," *Mycologia*, vol. 101, no. 3, pp. 423–429, 2009.
- [5] H. K. Abbas, R. M. Zablutowicz, B. W. Horn et al., "Comparison of major biocontrol strains of non-aflatoxigenic *Aspergillus flavus* for the reduction of aflatoxins and cyclopiazonic acid in maize," *Food Additives and Contaminants*, vol. 28, no. 2, pp. 198–208, 2011.
- [6] H. K. Abbas, M. A. Weaver, B. W. Horn, I. Carbone, J. T. Monacell, and W. T. Shier, "Selection of *Aspergillus flavus* isolates for biological control of aflatoxins in corn," *Toxin Reviews*, vol. 30, no. 2-3, pp. 59–70, 2011.
- [7] E. D. King, A. B. Bassi Jr., D. C. Ross, and B. Druebbisch, "An industry perspective on the use of "atoxicogenic" strains of *Aspergillus flavus* as biological control agents and the significance of cyclopiazonic acid," *Toxin Reviews*, vol. 30, no. 2-3, pp. 33–41, 2011.
- [8] H. K. Abbas, J. R. Wilkinson, R. M. Zablutowicz et al., "Ecology of *Aspergillus flavus*, regulation of aflatoxin production, and management strategies to reduce aflatoxin contamination of corn," *Toxin Reviews*, vol. 28, no. 2-3, pp. 142–153, 2009.
- [9] U. L. Diener, R. J. Cole, T. H. Sanders, G. A. Payne, L. S. Lee, and M. A. Klich, "Epidemiology of aflatoxin formation by *Aspergillus flavus*," *Annual Review of Phytopathology*, vol. 25, pp. 249–270, 1987.
- [10] J. Yu, P.-K. Chang, K. C. Ehrlich et al., "Clustered pathway genes in aflatoxin biosynthesis," *Applied and Environmental Microbiology*, vol. 70, no. 3, pp. 1253–1262, 2004.
- [11] P.-K. Chang, B. W. Horn, and J. W. Dorner, "Clustered genes involved in cyclopiazonic acid production are next to the aflatoxin biosynthesis gene cluster in *Aspergillus flavus*," *Fungal Genetics and Biology*, vol. 46, no. 2, pp. 176–182, 2009.
- [12] H. K. Abbas, W. T. Shier, and R. D. Cartwright, "Effect of temperature, rainfall and planting date on aflatoxin and fumonisin contamination in commercial Bt and non-Bt corn hybrids in Arkansas," *Phytoprotection*, vol. 88, no. 2, pp. 41–50, 2007.
- [13] H. A. Bruns, "Controlling aflatoxin and fumonisin in maize by crop management," *Journal of Toxicology—Toxin Reviews*, vol. 22, no. 2-3, pp. 153–173, 2003.

- [14] US-FDA (Food and Drug Administration), "Action levels for poisonous or deleterious substances in human food and animal feed," <http://www.fda.gov/food/guidanceregulation/guidancedocumentsregulatoryinformation/ucm077969.htm>.
- [15] H. P. van Egmond, R. C. Schothorst, and M. A. Jonker, "Regulations relating to mycotoxins in food. Perspectives in a global and European context," *Analytical and Bioanalytical Chemistry*, vol. 389, no. 1, pp. 147–157, 2007.
- [16] D. Bhatnagar, J. W. Cary, K. Ehrlich, J. Yu, and T. E. Cleveland, "Understanding the genetics of regulation of aflatoxin production and *Aspergillus flavus* development," *Mycopathologia*, vol. 162, no. 3, pp. 155–166, 2006.
- [17] H. K. Abbas, R. M. Zablutowicz, and H. A. Bruns, "Modeling the colonization of maize by toxigenic and non-toxigenic *Aspergillus flavus* strains: implications for biological control," *World Mycotoxin Journal*, vol. 1, no. 3, pp. 333–340, 2008.
- [18] G. G. Barros, A. M. Torres, M. I. Rodriguez, and S. N. Chulze, "Genetic diversity within *Aspergillus flavus* strains isolated from peanut-cropped soils in Argentina," *Soil Biology and Biochemistry*, vol. 38, no. 1, pp. 145–152, 2006.
- [19] M. Donner, J. Atehnkeng, R. A. Sikora, R. Bandyopadhyay, and P. J. Cotty, "Molecular characterization of atoxigenic strains for biological control of aflatoxins in Nigeria," *Food Additives and Contaminants A: Chemistry, Analysis, Control, Exposure and Risk Assessment*, vol. 27, no. 5, pp. 576–590, 2010.
- [20] Y. Yin, T. Lou, L. Yan, T. J. Michailides, and Z. Ma, "Molecular characterization of toxigenic and atoxigenic *Aspergillus flavus* isolates, collected from peanut fields in China," *Journal of Applied Microbiology*, vol. 107, no. 6, pp. 1857–1865, 2009.
- [21] Y.-X. Fu and W.-H. Li, "Statistical tests of neutrality of mutations," *Genetics*, vol. 133, no. 3, pp. 693–709, 1993.
- [22] C. Strobeck, "Average number of nucleotide differences in a sample from a single subpopulation: a test for population subdivision," *Genetics*, vol. 117, pp. 149–153, 1987.
- [23] F. Tajima, "Statistical method for testing the neutral mutation hypothesis by DNA polymorphism," *Genetics*, vol. 123, no. 3, pp. 585–595, 1989.
- [24] M. Nei, "Evolution of human races at the gene level," in *Human Genetics Part A: The Unfolding Genome*, B. Bonne-Tamir, T. Cohen, and R. M. Goodman, Eds., pp. 167–181, Alan R. Liss, New York, NY, USA, 1982.
- [25] G. A. Payne, J. Yu, W. C. Nierman et al., "A first glance into the genome sequence of *Aspergillus flavus*," in *The Aspergilli: Genomics, Medical Aspects, Biotechnology, and Research Methods*, S. A. Osmani and G. H. Goldman, Eds., pp. 15–23, CRC Press, Boca Raton, Fla, USA, 2008.
- [26] P.-K. Chang, K. C. Ehrlich, and S.-S. T. Hua, "Cladal relatedness among *Aspergillus oryzae* isolates and *Aspergillus flavus* S and L morphotype isolates," *International Journal of Food Microbiology*, vol. 108, no. 2, pp. 172–177, 2006.
- [27] M. K. Das, K. C. Ehrlich, and P. J. Cotty, "Use of pyrosequencing to quantify incidence of a specific *Aspergillus flavus* strain within complex fungal communities associated with commercial cotton crops," *Phytopathology*, vol. 98, no. 3, pp. 282–288, 2008.
- [28] K. C. Ehrlich and P. J. Cotty, "An isolate of *Aspergillus flavus* used to reduce aflatoxin contamination in cottonseed has a defective polyketide synthase gene," *Applied Microbiology and Biotechnology*, vol. 65, no. 4, pp. 473–478, 2004.
- [29] C. Accinelli, H. K. Abbas, R. M. Zablutowicz, and J. R. Wilkinson, "*Aspergillus flavus* aflatoxin occurrence and expression of aflatoxin biosynthesis genes in soil," *Canadian Journal of Microbiology*, vol. 54, no. 5, pp. 371–379, 2008.
- [30] P.-K. Chang, H. K. Abbas, M. A. Weaver, K. C. Ehrlich, L. L. Scharfenstein, and P. J. Cotty, "Identification of genetic defects in the atoxigenic biocontrol strain *Aspergillus flavus* K49 reveals the presence of a competitive recombinant group in field populations," *International Journal of Food Microbiology*, vol. 154, no. 3, pp. 192–196, 2012.
- [31] D. T. Wicklow, C. E. McAlpin, and C. E. Platis, "Characterization of the *Aspergillus flavus* population within an Illinois maize field," *Mycological Research*, vol. 102, no. 3, pp. 263–268, 1998.
- [32] A. Lourenço, E. L. Durigon, P. Zanotto, J. E. G. C. Madeira, A. P. De Almeida, and B. Correa, "Genetic diversity of environmental *Aspergillus flavus* strains in the state of São Paulo, Brazil by random amplified polymorphic DNA," *Memorias do Instituto Oswaldo Cruz*, vol. 102, no. 6, pp. 687–692, 2007.
- [33] G. E. O. Midorikawa, M. R. R. Pinheiro, B. S. Vidigal et al., "Characterization of *Aspergillus flavus* strains from Brazilian Brazil nuts and cashew by RAPD and ribosomal DNA analysis," *Letters in Applied Microbiology*, vol. 47, no. 1, pp. 12–18, 2008.
- [34] P. P. Batista, J. F. Santos, N. T. Oliveira, A. P. D. Pires, C. M. S. Motta, and E. A. Luna-Alves Lima, "Genetic characterization of Brazilian strains of *Aspergillus flavus* using DNA markers," *Genetics and Molecular Research*, vol. 7, no. 3, pp. 706–717, 2008.
- [35] L. C. Grubisha and P. J. Cotty, "Twenty-four microsatellite markers for the aflatoxin-producing fungus *Aspergillus flavus*," *Molecular Ecology Resources*, vol. 9, no. 1, pp. 264–267, 2009.
- [36] A. D. Hatti, S. D. Taware, A. S. Taware, P. P. Pangrikar, A. M. Chavan, and D. S. Mukadam, "Genetic diversity of toxigenic and non-toxigenic *Aspergillus flavus* strains using ISSR markers," *International Journal of Current Research*, vol. 5, pp. 61–66, 2010.
- [37] P.-K. Chang, B. W. Horn, and J. W. Dorner, "Sequence breakpoints in the aflatoxin biosynthesis gene cluster and flanking regions in nonaflatoxigenic *Aspergillus flavus* isolates," *Fungal Genetics and Biology*, vol. 42, no. 11, pp. 914–923, 2005.
- [38] J. H. Ramirez-Prado, G. G. Moore, B. W. Horn, and I. Carbone, "Characterization and population analysis of the mating-type genes in *Aspergillus flavus* and *Aspergillus parasiticus*," *Fungal Genetics and Biology*, vol. 45, no. 9, pp. 1292–1299, 2008.
- [39] M. A. Passone, L. C. Rosso, A. Ciancio, and M. Etcheverry, "Detection and quantification of *Aspergillus* section Flavi spp. in stored peanuts by real-time PCR of *nor-1* gene, and effects of storage conditions on aflatoxin production," *International Journal of Food Microbiology*, vol. 138, no. 3, pp. 276–281, 2010.
- [40] H. K. Abbas, C. Accinelli, R. M. Zablutowicz et al., "Dynamics of mycotoxin and *Aspergillus flavus* levels in aging Bt and non-Bt corn residues under Mississippi no-till conditions," *Journal of Agricultural and Food Chemistry*, vol. 56, no. 16, pp. 7578–7585, 2008.
- [41] H. K. Abbas, R. M. Zablutowicz, M. A. Weaver, B. W. Horn, W. Xie, and W. T. Shier, "Comparison of cultural and analytical methods for determination of aflatoxin production by Mississippi Delta *Aspergillus* isolates," *Canadian Journal of Microbiology*, vol. 50, no. 3, pp. 193–199, 2004.
- [42] B. W. Horn and J. W. Dorner, "Soil populations of *Aspergillus* species from section Flavi along a transect through peanut-growing regions of the United States," *Mycologia*, vol. 90, no. 5, pp. 767–776, 1998.
- [43] M. A. Klich, *Identification of Common Aspergillus Species*, Centraalbureau voor Schimmelcultures, Utrecht, The Netherlands, 2002.

- [44] H. K. Abbas, W. P. Williams, G. L. Windham, H. C. Pringle III, W. Xie, and W. T. Shier, "Aflatoxin and fumonisin contamination of commercial corn (*Zea mays*) hybrids in Mississippi," *Journal of Agricultural and Food Chemistry*, vol. 50, no. 18, pp. 5246–5254, 2002.
- [45] P. Librado and J. Rozas, "DnaSP v5: a software for comprehensive analysis of DNA polymorphism data," *Bioinformatics*, vol. 25, no. 11, pp. 1451–1452, 2009.
- [46] L. Excoffier and H. E. L. Lischer, "Arlequin suite ver 3.5: a new series of programs to perform population genetics analyses under Linux and Windows," *Molecular Ecology Resources*, vol. 10, no. 3, pp. 564–567, 2010.
- [47] L. Excoffier, P. E. Smouse, and J. M. Quattro, "Analysis of molecular variance inferred from metric distances among DNA haplotypes: application to human mitochondrial DNA restriction data," *Genetics*, vol. 131, no. 2, pp. 479–491, 1992.
- [48] K. Tamura, D. Peterson, N. Peterson, G. Stecher, M. Nei, and S. Kumar, "MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods," *Molecular Biology and Evolution*, vol. 28, no. 10, pp. 2731–2739, 2011.
- [49] R. M. Zablotowicz, H. K. Abbas, and M. A. Locke, "Population ecology of *Aspergillus flavus* associated with Mississippi Delta soils," *Food Additives and Contaminants*, vol. 24, no. 10, pp. 1102–1108, 2007.
- [50] P. Bayman and P. J. Cotty, "Vegetative compatibility and genetic diversity in the *Aspergillus flavus* population of a single field," *Canadian Journal of Botany*, vol. 69, no. 8, pp. 1707–1711, 1991.
- [51] R. M. Gashgari, Y. M. Shebany, and Y. A. Gherbawy, "Molecular characterization of mycobiota and aflatoxin contamination of retail wheat flours from jeddah markets," *Foodborne Pathogens and Disease*, vol. 7, no. 9, pp. 1047–1054, 2010.
- [52] D. M. Geiser, J. I. Pitt, and J. W. Taylor, "Cryptic speciation and recombination in the aflatoxin-producing fungus *Aspergillus flavus*," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 95, no. 1, pp. 388–393, 1998.
- [53] D. M. Geiser, J. W. Dorner, B. W. Horn, and J. W. Taylor, "The phylogenetics of mycotoxin and sclerotium production in *Aspergillus flavus* and *Aspergillus oryzae*," *Fungal Genetics and Biology*, vol. 31, no. 3, pp. 169–179, 2000.
- [54] K. C. Ehrlich, J. Yu, and P. J. Cotty, "Aflatoxin biosynthesis gene clusters and flanking regions," *Journal of Applied Microbiology*, vol. 99, no. 3, pp. 518–527, 2005.