



Enhanced Production of Docosahexaenoic Acid in Mammalian Cells

Guiming Zhu*, Xudong Jiang, Qin Ou, Tao Zhang, Mingfu Wang, Guozhi Sun, Zhao Wang, Jie Sun, Tangdong Ge

Laboratory of Biochemistry and Molecular Biology, College of Basic Medicine, Jiamusi University, Jiamusi, Heilongjiang, China

Abstract

Docosahexaenoic acid (DHA), one of the important polyunsaturated fatty acids (PUFA) with pharmaceutical and nutraceutical effects, may be obtained through diet or synthesized in vivo from dietary α -linolenic acid (ALA). However, the accumulation of DHA in human body or other mammals relies on the intake of high dose of DHA for a certain period of time, and the bioconversion of dietary ALA to DHA is very limited. Therefore the mammalian cells are not rich in DHA. Here, we report a new technology for increased production of DHA in mammalian cells. By using transient transfection method, *Siganus canaliculatus* $\Delta 4$ desaturase was heterologously expressed in chinese hamster ovary (CHO) cells, and simultaneously, mouse $\Delta 6$ -desaturase and $\Delta 5$ -desaturase were overexpressed. The results demonstrated that the overexpression of $\Delta 6/\Delta 5$ -desaturases significantly enhanced the ability of transfected cells to convert the added ALA to docosapentaenoic acid (DPA) which in turn get converted into DHA directly and efficiently by the heterologously expressed $\Delta 4$ desaturase. This technology provides the basis for potential utility of these gene constructs in the creation of transgenic livestock for increased production of DHA/related products to meet the growing demand of this important PUFA.

Citation: Zhu G, Jiang X, Ou Q, Zhang T, Wang M, et al. (2014) Enhanced Production of Docosahexaenoic Acid in Mammalian Cells. PLoS ONE 9(5): e96503. doi:10.1371/journal.pone.0096503

Editor: Ayyalasomayajula Vajreswari, National Institute of Nutrition, India

Received: November 29, 2013; **Accepted:** April 8, 2014; **Published:** May 2, 2014

Copyright: © 2014 Zhu et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by grants from the National Natural Science Foundation of China (No. 31000984) and Science and Technology Innovation Team Research Project of Jiamusi University (No. 04099904). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: xi78926@sina.com

Introduction

Docosahexaenoic acid (DHA; 22:6n-3) is an n-3 polyunsaturated fatty acid (PUFA) that is particularly enriched in mammalian tissues such as brain, testis, and retina. Clinical studies have demonstrated the crucial role of DHA in the development and functions of these tissues [1,2]. DHA enhances membrane elasticity and molecular motion and thus promotes signal transduction via enhanced protein/receptor interactions [3–6]. DHA is also the activating ligand for multiple transcriptional factors that control the expression of enzymes involved in fatty acid synthesis and β -oxidation [7]. Depletion of brain DHA in some mammals leads to distinct impairments in brain and neural function [8–13]. In adult humans, low levels of DHA in blood have been correlated with psychological disturbances such as alcoholism, major depression (nonpsychotic), postpartum depression, and senile dementia [19–21]. The essentiality of DHA for human infant nutrition in support of neuronal function has been shown by DHA supplementation, enhancing visual acuity and cognition-related test scores in human infants [14–18].

It is well established that DHA can be biosynthesized from α -linolenic acid (ALA, 18:3n-3), a shorter chain n-3 fatty acid precursor, through chain elongation and desaturation processes. α -linolenic acid is desaturated to 18:4n-3 by $\Delta 6$ -desaturase, chain-elongated to 20:4n-3, and subsequently converted to eicosapentaenoic acid (EPA, 20:5n-3), also a biologically active n-3 fatty acids, by $\Delta 5$ -desaturase in the endoplasmic reticulum. Mammalian

$\Delta 5$ - and $\Delta 6$ -desaturases have been identified and cloned [22]. However, $\Delta 4$ -desaturase, responsible for making 22:6n-3 directly from docosapentaenoic acid (DPA, 22:5n-3), an elongation product of 20:5n-3, has been identified only in a few of organisms such as microalgae [23]. In mammals, 22:5n-3 is further elongated to 24:5n-3 followed by desaturation by $\Delta 6$ -desaturase to 24:6n-3. Subsequently, 24:6n-3 is transferred to peroxisomes and converted to 22:6n-3 by removing two carbon chains by β -oxidation. Long-chain n-6 fatty acids, such as arachidonic acid (ARA, 20:4n-6), the commonly observed n-6 fatty acids in most tissues, are biosynthesized from linoleic acid (LA, 18:2n-6) using the analogous pathway and the same enzyme system.

Although dietary ALA can be converted to DHA in vivo, the efficiency of conversion is limited [24]. It was shown that the mature rat brain has little capacity to synthesize its own DHA, even when stimulated to do so, under dietary conditions of feeding ALA alone or during n-3 PUFA dietary deprivation [25,26]. In contrast, the adult rat liver does convert ALA to DHA, but at a rate sufficient to convert <2% of the total ALA that enters the liver per unit time, with the majority (>70%) of the entering ALA being lost to β -oxidation. Therefore, mammals are unable to produce high levels of DHA in their bodies. Algae are the primary producers of DHA and EPA in the ecosystem, and several refined algal oils are rich sources of DHA. Fish consume micro algae (phyto plankton) and therefore are rich in DHA and EPA, thus fish become the main dietary source of these PUFA. Nevertheless, marine products such as encapsulated fish oil are not suitable for

daily use. Furthermore, the decline in marine fish stocks and the potential contamination of marine products with mercury and other chemicals call for land-based dietary sources of DHA. One strategy is to create genetically modified land animals by making use of the desaturases and elongases involved in DHA biosynthesis. In this paper, we report the successful production of DHA in chinese hamster ovary (CHO) cells by heterologous expression of *Siganus canaliculatus* $\Delta 4$ desaturases after overexpression of mice $\Delta 6/\Delta 5$ -desaturase. The results provide a basis for potential applications of this technology in creating of transgenic livestock for enhanced production of DHA in the related products.

Materials and Methods

Construction of Recombinant Plasmids

The $\Delta 4$ desaturase gene from *Siganus canaliculatus*, *sScD4*, was artificially synthesized (with genetic codons optimized) from its original sequence (GenBank accession number: GU594278). Then, the synthetic gene (1352 bp) was inserted into the pcDNA3.1 expression vector with EcoRI/HindIII double digestion. The expression vectors for *fads2* and *fads1* genes (cDNAs), coding for mouse $\Delta 6$ -desaturase and $\Delta 5$ -desaturase respectively, were constructed previously in our laboratory, with the cDNAs cloned into pcDNA3.1. The double-gene expression vector pcDNA3.1-F2F1 was constructed as follows: based on the enzyme sites BglIII/MluI carried on pcDNA3.1-fad1, PCR primers were designed to amplify the pCMV-fad2-polyA DNA fragment on pcDNA3.1-fad2, and then the amplicon was inserted into pcDNA3.1-fad1 by BglIII/MluI double digestion and ligation.

Cell Cultures and transfection

Transient transfections in chinese hamster ovary (CHO) cells were carried out using LipofectamineTM 2000 (Invitrogen Inc., USA) as described in the manufacturer's manual. One day prior to transfection, cells were plated onto 60-mm culture dishes in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% fetal bovine serum (FBS) and 10 μ M PUFA substrates (DPA for *sScD4* transfections, LA and ALA for F2F1 transfections). Cells were transfected with 8 μ g of the plasmids. To monitor transfection efficiencies, cells were transfected with an expression vector containing the enhanced green fluorescent protein (EGFP) cDNA, pcDNA3.1-EGFP. The day following transfection, media were replaced with fresh media containing the same PUFA substrates as above. Two days after transfection, cells were harvested and used for analysis of gene expression or fatty acid composition.

RT-PCR

RT-PCR was performed to analyze transcripts in transgenic cells. Total RNA was extracted from the harvested cells, using TRIzol Reagent (Invitrogen) followed by DNase treatment to remove genomic DNA contamination. cDNA was prepared using the TaKaRa RNA PCR Kit 3.0 (TaKaRa). The following primer pairs were used to detect the target genes—*sScD4*: 5'-TCGGGCATCTGAGCGTGTTTC-3' and 5'-ATACTCCACGGCTGGGTCTCC-3'; F2: 5'-TAGTCTAGAGCATGGGGAAGGGAGGTAAC-3' and 5'-AGCAAGCTTCATTTATGAGGT AAGCATCCAG-3'; F1: 5'-TACTCTAGAGCTTGCTATGGCTCCCGACC-3', 5'-ATGAAGCTTGCTATTGTGAAGGTAAGCGTCC-3'. The PCR conditions were as follows: 94°C for 2 min for desaturation, 35 cycles of 94°C for 30 s for desaturation, 60°C for 30 s for annealing, and 72°C for 30 s for extension, followed by a final extension period of 72°C for 10 min. The amplified products were subjected to electrophoresis




Figure 1. *sScD4*, *fads2* and *fads1* transcripts in transfected cells were analyzed by RT-PCR. Lane 1 were cells transfected with pcDNA3.1-EGFP; Lane 2 were cells transfected with pcDNA3.1-F2F1; Lane 3 were cells co-transfected with pcDNA3.1-F2F1 and pcDNA3.1-*sScD4*.

doi:10.1371/journal.pone.0096503.g001

on a 2% agarose gel. The amplification of beta-actin gene was carried out as control.

Lipid Analysis

The fatty acid composition of total cellular lipids was analyzed as following: an aliquot of cell pellet homogenate in a micro reaction vessel (Supelco) was mixed with 1 ml of 14% BF₃/MeOH reagent containing 0.005% butylated hydroxytoluene (as antioxidant). After blanketed with nitrogen, the mixture was heated at 95–100°C for 1 h, then cooled to room temperature and methyl esters extracted in the hexane phase following addition of 1 ml H₂O and 1 ml of hexane. The samples were centrifuged for 5 min at 3000 rpm, then the upper hexane layer was removed and concentrated under nitrogen. Fatty acid methyl esters were quantified with GC-MS by using an HP-5890 Series II gas chromatograph equipped with an Omegawax 250 capillary column (Supelco, Bellefonte, PA) attached to an HP-5971 mass spectrometer. The oven program was maintained initially at 150°C for 2 min, then ramped to 200°C at 10°C/min and held for 4 min, ramped again at 5°C/min to 240°C, held for 3 min, and finally, ramped to 270°C at 10°C/min and maintained for 5 min. Peaks were identified by comparison with fatty acid standards (Sigma), then area and percentage for each resolved peak were analyzed.

Statistical Analysis

Statistical analysis was performed with SPSS version 18.0 for Windows, using one-way ANOVA followed by LSD comparison test for posthoc analyses. Results were expressed as the mean \pm SD. A p-value of <0.05 was considered to be statistically significant.

Results

To test whether pcDNA3.1-*sScD4* could be expressed in mammalian cells, the vector was introduced into CHO cells, by using liposome transfection method. DPA was added to the cell culture medium for 3 days. The RT-PCR results showed mRNA of this synthetic gene could be greatly transcribed, while the cells transfected with the control plasmid pcDNA3.1-EGFP, which carries the enhanced green fluorescent protein as reporter gene, detected no *sScD4* mRNA (Fig.1). The transient transfections of pcDNA3.1-*sScD4* in CHO cells confirmed *sScD4* can directly and efficiently convert DPA to DHA, as shown in Fig 2. Table 1 shows fatty acids composition of total cellular lipids from the CHO cells transfected with *EGFP* or *sScD4*. None of the percentage distribution of a fatty acid varied between the CHO cells transfected with *EGFP* and *sScD4* except DPA and DHA. The




Figure 2. Partial gas chromatograph traces showing fatty acid profiles of total cellular lipids extracted from the control cells transfected with pcDNA3.1-EGFP (A), and the cells infected with pcDNA3.1-sScD4 (B).
doi:10.1371/journal.pone.0096503.g002

percentage distribution of DPA decreased from 20.2 ± 1.62 in CHO cells transfected with *EGFP* to 14.4 ± 1.46 in the cells transfected with *sScD4*, while the percentage distribution of DHA increased from 4.0 ± 0.95 in cells transfected with *EGFP* to 9.2 ± 0.94 in the cells transfected with *sScD4*.

CHO cells transfected with pcDNA3.1-F2F1 have the ability to overexpress of *fads2* and *fads1* genes as shown in Fig.1. The fatty acid profiles were remarkably different between the control cells transfected with pcDNA3.1-EGFP and the cells transfected with pcDNA3.1-F2F1 (Fig.3). In the cells overexpressing the $\Delta 6/\Delta 5$ desaturase, the percentage distribution of 18:2n-6 and 18:3n-3 were significantly decreased, while 20:3n-6, 20:4n-6, 20:5n-3, 22:5n-3 were significantly increased, as compared to the control cells. This means the substrates (18:2n-6 and 18:3n-3) added to the transfected cells were efficiently converted into longer PUFA. However, the DHA level increased slightly but was not statistically significant, indicating that the overexpression of the $\Delta 6/\Delta 5$ desaturases did not contribute to the elevation of DHA level. An

interesting phenomenon is that the percentage composition of 20:3n-3 and 20:4n-3 were also significantly decreased in the cells overexpressing the $\Delta 6/\Delta 5$ desaturase when compared to the control cells. Obviously, the cells tend to convert them to longer chain PUFA, such as 20:5n-3, 22:5n-3.

The co-transfection of pcDNA3.1-F2F1 and pcDNA3.1-sScD4 into CHO cells also showed similar overexpression of *fads2* and *fads1* genes as cells transfected with pcDNA3.1-F2F1, but only the former detected the expression of *sScD4* gene (Fig.1). The $\Delta 4$ desaturase (*sScD4* gene product) confer the cells co-transfected with pcDNA3.1-F2F1 and pcDNA3.1-sScD4 the ability of efficiently converting the accumulated DPA to DHA (Fig.3). The level of DHA in the cells co-transfected with pcDNA3.1-F2F1 and pcDNA3.1-sScD4 was more than twofold higher than in the cells transfected with pcDNA3.1-F2F1, and threefold higher than in the cells transfected with pcDNA3.1-EGFP (Table 2). In addition, compared to the cells transfected with pcDNA3.1-F2F1, the percentage distribution of 20:5n-3, 22:5n-3 in CHO cells co-

Table 1. PUFA composition of total cellular lipids from the CHO cells transfected with EGFP or sScD4 genes.

Mol % of fatty acids	EGFP	sScD4
14:0	4.6±0.42	4.5±0.31
16:0	2.1±0.34	1.6±0.22
18:0	19.2±2.13	20.3±2.51
18:1n-9	24.4±3.35	23.6±2.87
18:1n-7	9.1±1.86	10.9±1.94
18:2n-6	4.2±0.93	4.5±1.17
20:3n-6	1.4±0.38	1.6±0.35
20:4n-6	5.8±1.69	6.3±2.12
20:5n-3	3.3±0.89	2.3±0.81
22:5n-3	20.2±1.62 ^a	14.4±1.46 ^b
22:6n-3	4.0±0.95 ^b	9.2±0.94 ^a

Values are means of three measurements. Values for each fatty acid with different superscripts differ significantly ($P < 0.05$) between control (EGFP), sEgD4 and sScD4.
doi:10.1371/journal.pone.0096503.t001




Figure 3. Partial gas chromatograph traces showing fatty acid profiles of total cellular lipids extracted from the control cells transfected with pcDNA3.1-EGFP (A), the cells transfected with pcDNA3.1-F2F1 (B) and the cells co-transfected with pcDNA3.1-F2F1 and pcDNA3.1-sScD4 (C).

doi:10.1371/journal.pone.0096503.g003

transfection of pcDNA3.1-F2F1 and pcDNA3.1-sScD4 decreased significantly in that they were converted to DHA.

Discussion

Here, we reported a new technology for enhanced production of DHA in mammalian cells. Mammals lack the $\Delta 12$ and $\Delta 15$ fatty acid desaturases responsible for converting oleic acid (18:1n-9) into linoleic acid (18:2n-6) and α -linolenic acid (18:3n-3) and thus, are unable to biosynthesize polyunsaturated fatty acids (PUFA) *de novo* [27,28]. Therefore, the ingested plant-derived 18:2n-6 and 18:3n-3 are essential dietary nutrients for mammals [27,28]. Nevertheless, most of the ingested plant-derived 18:2n-6 and 18:3n-3 can not be converted to ARA, EPA or DHA but are lost to β -oxidation. Our results demonstrated that, by overexpression of $\Delta 6$ and $\Delta 5$ fatty

acid desaturases, the ability of converting 18:2n-6 to 20:4n-6, 18:3n-3 to EPA and DPA in mammalian cells was significantly elevated. This may be attributed to the fact that the overexpression of these desaturases disequibrated the regulatory system of PUFA biosynthesis. We had studied that the overexpression of $\Delta 6$ and $\Delta 5$ fatty acid elongases (the most two important elongases involved in PUFA biosynthesis) in CHO cells, and found that they contributed little to elevating the level of any longer PUFA biosynthesized from 18:2n-6 or 18:3n-3 (unpublished data). We conclude that, among the fatty acid desaturases and elongases involved in PUFA biosynthesis in mammals, $\Delta 6$ and $\Delta 5$ desaturases are the key ones that contribute to production of longer PUFA.

The conversion of DPA to DHA is the rate-limiting step in the pathway of biosynthesis of DHA from ALA [29]. As shown in our results, even the overexpression of $\Delta 6$ and $\Delta 5$ fatty acid

Table 2. PUFA composition of total cellular lipids from the CHO cells transfected with EGFP, F2F1 or F2F1+D4.

Mol % of fatty acids	EGFP	F2F1	F2F1+D4
14:0	4.2±0.83	3.8±0.81	3.2±0.78
16:0	0.7±0.03	0.8±0.04	2.1±0.11
18:0	18.8±2.12	19.6±2.33	18.2±1.98
18:1n-9	20.3±1.18	20.3±1.12	20.7±1.27
18:1n-7	10.9±0.91	10.3±0.95	8.2±0.78
n-6 PUFA			
18:2n-6	16.1±1.23 ^a	12.6±1.07 ^b	12.3±0.92 ^b
20:3n-6	0.9±0.07 ^b	1.7±0.21 ^a	1.9±0.20 ^a
20:4n-6	5.6±0.88 ^b	7.1±0.69 ^a	6.8±0.70 ^a
22:4n-6	1.5±0.09	1.4±0.14	1.5±0.21
Total	24.0±1.91	22.7±1.76	22.5±1.69
n-3 PUFA			
18:3n-3	6.9±0.84 ^a	4.0±0.37 ^b	3.4±0.29 ^b
20:3n-3	3.0±0.23 ^a	2.1±0.31 ^b	1.8±0.13 ^b
20:4n-3	1.3±0.12 ^a	0.9±0.13 ^b	0.9±0.09 ^b
20:5n-3	0.6±0.08 ^c	2.6±0.29 ^a	1.8±0.15 ^b
22:5n-3	3.9±0.71 ^b	7.2±0.99 ^a	4.2±0.73 ^b
22:6n-3	4.7±0.48 ^b	5.3±0.66 ^b	12.5±0.85 ^a
Total	20.4±1.30 ^b	22.0±1.07 ^b	24.6±1.51 ^a

Values are means of three measurements. Values for each fatty acid with different superscripts differ significantly ($P < 0.05$) between control (EGFP), sN3 and N3D4. doi:10.1371/journal.pone.0096503.t002

desaturases did not promote the conversion of DPA to DHA. They only led to an accumulation of high level of DPA in CHO cells. The results also confirmed that the Sprecher pathway in mammals had very low efficiency to convert DPA to DHA. A few of lower organisms can directly convert DPA to DHA by utilizing their $\Delta 4$ desaturases activities. Some researchers believe that the rate of DHA synthesis could be faster in the more direct $\Delta 4$ pathway and that only requires endoplasmic reticulum than in the Sprecher pathway that involves peroxisomes, translocation of PUFA intermediates and limited fatty acid oxidation (a catabolic step). $\Delta 4$ desaturases have been demonstrated in some organisms including protozoan trypanosomes [30], the photosynthetic freshwater protist *Euglena gracilis* [31], marine microalgae *Pavlova lutheria* and *Thraustochytrids* [32,33], and herbivorous, marine teleost fish, *Siganus canaliculatus* [34]. By heterologous expression in yeast, these $\Delta 4$ desaturases conferred on the yeast, the ability to convert DPA (22:5n-3) to DHA. This paper has clearly demonstrated that, when heterologously expressed in CHO cells, the *S. canaliculatus* $\Delta 4$ -desaturase can efficiently convert supplied DPA into DHA. More important, the proportions of DPA converted to DHA were about 36.3% in CHO cells expressing *S. canaliculatus* $\Delta 4$ -desaturase, which is even much higher than that in transgenic yeast expressing this desaturase. This indicates the mammalian cells or even mammals are likely to accept these $\Delta 4$ -desaturase activities from lower organisms. Significantly *S. canaliculatus* $\Delta 4$ -desaturase activity could cooperate with $\Delta 6$ and $\Delta 5$ -desaturase activities in co-transfected CHO cells with their gene constructs, directly and efficiently converting the accumulated DPA to DHA. Therefore, our strategy, by heterologous expression of *Siganus canaliculatus* $\Delta 4$ desaturase with overexpression of mice $\Delta 6/\Delta 5$ -desaturase in mammalian cells, is very effective in high-level production of DHA from dietary ALA.

References

- Muskiet FA, Fokkema MR, Schaafsma A, Boersma ER, Crawford MA (2004) Is docosahexaenoic acid (DHA) essential? Lessons from DHA status regulation, our ancient diet, epidemiology and randomized controlled trials. *J. Nutr.* 134: 183–186.
- Giusto NM, Pasquare SJ, Salvador GA, Castagnet PI, Roque ME, et al. (2000) Lipid metabolism in vertebrate retinal rod outer segments. *Prog Lipid Res*;39:315–91
- Gawrisch K, Eldho NV, Holte LL (2003) The structure of DHA in phospholipid membranes. *Lipids.* 38: 445–452.
- Mitchell DC, Niu SL, Litman BJ (2003) DHA-rich phospholipids optimize G-protein-coupled signaling. *J. Pediatr.* 143 (Suppl.): 80–86.
- Mitchell DC, Niu SL, Litman BJ (2003) Enhancement of G protein-coupled signaling by DHA phospholipids. *Lipids.* 38:437–443.
- Niu SL, Mitchell DC, Lim SY, Wen ZM, Kim HY, et al. (2004) Reduced G protein-coupled signaling efficiency in retinal rod outer segments in response to n-3 fatty acid deficiency. *J. Biol. Chem.* 279: 31098–31104.
- Clarke SD (2001) Polyunsaturated fatty acid regulation of gene transcription: a molecular mechanism to improve the metabolic syndrome. *J. Nutr.* 131: 1129–1132.
- Moriguchi T, Greiner RS, Salem NJr (2000) Behavioral deficits associated with dietary induction of decreased brain docosahexaenoic acid concentration. *J. Neurochem.* 75: 2563–2573.
- Carrie I, Smirnova M, Clement M, De Javel D, Frances H et al. (2002) Docosahexaenoic acid-rich phospholipid supplementation: effect on behavior, learning ability, and retinal function in control and n-3 polyunsaturated fatty acid deficient old mice. *Nutr. Neurosci.* 5: 43–52.
- Catalan J, Moriguchi T, Slotnick B, Murthy M, Greiner RS, et al. (2002) Cognitive deficits in docosahexaenoic acid deficient rats. *Behav. Neurosci.* 116: 1022–1031.
- Fedorova I, Salem NJr (2006) Omega-3 fatty acids and rodent behavior. *Prostaglandins Leukot. Essent. Fatty Acids.* 75: 271–289.
- Greiner RS, Moriguchi T, Hutton A, Slotnick BM, Salem NJr (1999) Rats with low levels of brain docosahexaenoic acid show impaired performance in olfactory-based and spatial learning tasks. *Lipids.* 34 (Suppl): 239–243.
- Champoux M, Hibbeln JR, Shannon C, Majchrzak S, Suomi SJ, et al. (2002) Fatty acid formula supplementation and neuromotor development in rhesus monkey neonates. *Pediatr. Res.* 51: 273–281.
- Innis SM (2000) The role of dietary n-6 and n-3 fatty acids in the developing brain. *Dev. Neurosci.* 22: 474–480.
- Neuringer M (2000) Infant vision and retinal function in studies of dietary long-chain polyunsaturated fatty acids: methods, results, and implications. *Am. J. Clin. Nutr.* 71 (Suppl): 256–267.
- Birch EE, Castaneda YS, Wheaton DH, Birch DG, Uauy RD, et al. (2005) Visual maturation of term infants fed long-chain polyunsaturated fatty acid-supplemented or control formula for 12 mo. *Am. J. Clin. Nutr.* 81: 871–879.
- Birch EE, Garfield S, Castaneda Y, Hughbanks-Wheaton D, Uauy R, et al. (2007) Visual acuity and cognitive outcomes at 4 years of age in a double-blind, randomized trial of long-chain polyunsaturated fatty acid-supplemented infant formula. *Early Hum. Dev.* 83: 279–284.
- Gibson RA, Chen W, Makrides M (2001) Randomized trials with polyunsaturated fatty acid interventions in preterm and term infants: functional and clinical outcomes. *Lipids.* 36: 873–883.
- Marangell LB, Martinez JM, Zboyan HA, Kertz B, Kim HF, et al. (2003) A cerebrole-blind, placebo-controlled study of the omega-3 fatty acid docosahexaenoic acid in the treatment of major depression. *Am. J. Psychiatry.* 160: 996–998.
- Otto SJ, Groot RH, Hornstra G (2003) Increased risk of postpartum depressive symptoms is associated with slower normalization after pregnancy of the functional docosahexaenoic acid status. *Prostaglandins Leukot. Essent. Fatty Acids.* 69: 237–243.
- Hibbeln JR, Linnoila M, Umhau JC, Rawlings R, George DT, et al. (1998) Essential fatty acids predict metabolites of serotonin and dopamine in cerebrospinal fluid among healthy control subjects, and early- and late-onset alcoholics. *Biol. Psychiatry.* 44:235–242.
- Nakamura MT, Nara TY (2004) Structure, function, and dietary regulation of delta6, delta5, and delta9 desaturases. *Annu. Rev. Nutr.* 24, 345–376
- Pereira SL, Leonard AE, Huang YS, Chuang LT, Mukerji P (2004) *Biochem. J.* 384, 357–366
- Salem NJr, Wegher JB, Mena P, Uauy R (1996) Arachidonic and docosahexaenoic acids are biosynthesized from their 18-carbon precursors in human infants. *Proc. Natl. Acad. Sci. USA.* 93: 49–54.
- DeMar JC, Ma JK, Chang L, Bell JM, Rapoport SI (2005) Alpha-linolenic acid does not contribute appreciably to docosahexaenoic acid within brain phospholipids of adult rats fed a diet enriched in docosahexaenoic acid. *J. Neurochem.* 94: 1063–1076.
- Igarashi M, DeMar JC, Ma JK, Chang L, Bell JM, Rapoport SI (2007) Docosahexaenoic acid synthesis from alpha-linolenic acid by rat brain is unaffected by dietary n-3 PUFA deprivation. *J. Lipid Res.* 48: 1150–1158.

Production of DHA in land animals may be a good choice for mankind to meet the growing demand for this important LC-PUFA, and our results provide potential solution on this regard. Like humans, most land animals can accumulate DHA through their diet, but usually there is no adequate DHA in animal feeds. Adding DHA to animal feeds for the accumulation of it in animal bodies is expensive and unadvisable. Mammalian cells in culture, to a large extent, can simulate the process of PUFA biosynthesis in living animals. So our technology, by heterologous expression of *Siganus canaliculatus* $\Delta 4$ desaturases and overexpression of mice $\Delta 6/\Delta 5$ -desaturases simultaneously, will probably produce high level of DHA in transgenic livestock carrying these gene constructs as in transfected CHO cells reported here. If such genetically modified land animals are successfully created and widely accepted by the public, DHA can be provided abundantly by these animals, which will probably change our views on the ordinary 'unhealthy' animal fat.

In conclusion, this work has demonstrated that mammalian cells can be genetically modified by transfection of an delta-4 desaturase that is heterologously expressed with concomitant overexpression of delta-5 and delta-6 desaturases to convert ALA to DHA efficiently, and provide a potential for creation of new land animal breeds who can produce abundant DHA in their related products.

Author Contributions

Conceived and designed the experiments: GZ TZ. Performed the experiments: GZ XJ JS TG MW ZW. Analyzed the data: GZ XJ QO GS. Contributed reagents/materials/analysis tools: QO TZ GS. Wrote the paper: GZ.

27. Tinoco J (1982) Dietary requirements and functions of alpha-linolenic acid in animals. *Prog. Lipid Res.* 21, 1–45.
28. Holman RT (1986) Control of polyunsaturated acids in tissue lipids. *J. Am. Coll. Nutr.* 5, 183–211.
29. Pawlosky RJ, Hibblen JR, Lin Y, Goodson S, Riggs P, et al. Effects of beef- and fish-based diets on the kinetics of n₃ fatty acid metabolism in human subjects. *Am J Clin Nutr* 2003;77:565–72.
30. Tripodi KEJ, Buttiglierio LV, Altabe SG, Uttaro AD (2006) Functional characterization of front-end desaturases from trypanosomatids depicts the first polyunsaturated fatty acid biosynthetic pathway from a parasitic protozoan. *FEBS J* 273:271–280.
31. Meyer A, Cirpus P, Ott C, Schlecker R, Zähringer U, et al. (2003) Biosynthesis of docosahexaenoic acid in *Euglena gracilis*: Biochemical and molecular evidence for the involvement of a Δ^4 -fatty acyl group desaturase. *Biochemistry* 42:9779–9788.
32. Tonon T, Harvey D, Larson TR, Graham IA (2003) Identification of a very long chain polyunsaturated fatty acid Δ^4 -desaturase from the microalga *Pavlova lutheri*. *FEBS Lett* 553:440–444.
33. Qiu X, Hong H, MacKenzie SL (2001) Identification of a Δ^4 fatty acid desaturase from *Thraustochytrium* sp involved in the biosynthesis of docosahexaenoic acid by heterologous expression in *Saccharomyces cerevisiae* and *Brassica juncea*. *J Biol Chem* 276:31561–31566.
34. Li Y, Monroig O, Zhang L, Wang S, Zheng X, et al (2010) Vertebrate fatty acyl desaturase with Δ^4 activity. *Proc. Natl. Acad. Sci. USA.* 107(39):16840–16845.