













Supp. Fig. 1. Staged differential expression analyses identify E14.5-E15.5 transition as greatest overall shift in gene expression. Up-set plot of all staged comparisons conducted for differential expression analysis (n=3 biological replicates and n=3 technical replicates for each developmental stage analyzed). Closer investigation of the stage of palatal fusion (E14.5-E15.5) demonstrated the greatest shift in overall gene expression compared to all other staged comparisons.


*Intersection size indicates number of differentially expressed genes in each staged comparison.



Supp. Fig. 2. Functional Enrichment Analysis from E14.5 vs E15.5 highlights bone morphogenesis and cilia development as key processes at this timepoint. Functional enrichment analyses from GO Biological Processes database corroborated gene patterning associations based on stage-specific signatures identified from single genes.



Supp. Fig. 4. Sub-clustering of mesenchymal cells from single-nucleus RNA+ATAC-seq experiments revealed novel *Pax9*⁺ cell type. (A-B). Initial UMAP projection of re-clustering based on separate analysis of isolated mesenchymal cells. Overlaid with cluster annotation (A) and contributing sample (B). (C-D) UMAP projection as in (A) overlaid with deviation scores of enrichment of *Pax9* motif within marker peaks for each cell type. (E-F) Ridgeplot of chondrogenic, mesenchyme, osteogenic, and *Pax9*⁺ mesenchyme cell types. G) UMAP projection as in (A) overlaid with cluster annotations. H) Pseudo-replicate cell type proportion plot of ATAC output from all samples analyzed.



Tissue Expression Database: **Osteoblast [BTO:0001593]**
(Col1a1, Sparc, Runx2)

Node Color



colored nodes:
query proteins and first shell of interactors



white nodes:
second shell of interactors

Node Content



empty nodes:
proteins of unknown 3D structure



filled nodes:
some 3D structure is known or predicted

Known Interactions



from curated databases



experimentally determined

Predicted Interactions



gene neighborhood



gene fusions



gene co-occurrence

Others



textmining




co-expression





protein homology

Supp. Fig. 5. From String Consortium Database, K-means clustering highlights osteoblast markers with common interactions with high strength of association (2.41) and low false discovery rate (1.20e-06).



Supp. Fig. 6. *In vivo* spRNAseq confirms temporal increase in osteogenic markers and provides spatial quantification of lineage-specific differentiation of condensed mesenchyme progenitors of palatal bone. Markers of osteogenic lineage induction included the earliest **a** osteogenic progenitors *Six2*, *Erg*, and *Bmp7*, then **b** pre-osteoblast markers *Runx2*, *Sp7*, and *Mmp13*, followed by **c** more mature, functional osteoblast markers *Ifitm5*, *Ibsp*, and *Spp1*; finally, **d** markers of osteocyte activity *Sost*, *Dmp1*, and *Phex*, increase at this stage, yet to a lesser extent than earlier markers. Notably, markers from all early osteogenic cell types (osteoprogenitors, pre-osteoblasts, and osteoblasts) demonstrated some expression at E14.5; however, all osteocyte conserved markers only appeared in substantial expression by E15.5.



Supp. Fig. 7. Spatial RNA-seq-derived novel marker genes differentially expressed at the stage of palatal fusion. a Spatial *in vivo* expression, UMAPs, and violin plots from the top 5 marker genes from unbiased differential analysis of selective palate barcoding sequencing capture areas from E14.5 – E15.5 mid-palate coronal sections. **b** Gene plots from previously curated bulk RNA-seq analyses corroborate above spRNA-seq data, further evidence of the potential important functional roles for *Cdc20b*, *Deup1*, *Dynlrb2*, *Lrrc23*, and *Tnn* (aka *Tn-W*) in secondary palate development.