



Research Article

Diol-ginsenosides from Korean Red Ginseng delay the development of type 1 diabetes in diabetes-prone biobreeding rats



Chung Ju ^{1, ☆, ☆☆}, Sang-Min Jeon ^{2, ☆}, Hee-Sook Jun ^{3, 4}, Chang-Kiu Moon ^{1, 2, *}

¹ College of Pharmacy, Seoul National University, Seoul, Republic of Korea

² College of Pharmacy and Research Institute of Pharmaceutical Science and Technology (RIPST), Ajou University, Suwon, Gyeonggi-do Republic of Korea

³ College of Pharmacy, Gachon University, Incheon, Republic of Korea

⁴ Lee Gil Ya Cancer and Diabetes Institute, Gachon University, Incheon, Republic of Korea

ARTICLE INFO

Article history:

Received 27 September 2018

Received in Revised form

16 April 2019

Accepted 3 June 2019

Available online 12 June 2019

Keywords:

β-cell cytotoxicity

Cytokines

Diol-ginsenosides

Type 1 diabetes

ABSTRACT

Background: The effects of diol-ginsenoside fraction (Diol-GF) and triol-ginsenoside fraction (Triol-GF) from Korean Red Ginseng on the development of type 1 diabetes (T1D) were examined in diabetes-prone biobreeding (DP-BB) rats that spontaneously develop T1D through an autoimmune process.

Methods: DP-BB female rats were treated with Diol-GF or Triol-GF daily from the age of 3–4 weeks up to 11–12 weeks (1 mg/g body weight).

Results: Diol-GF delayed the onset, and reduced the incidence, of T1D. Islets of Diol-GF-treated DP-BB rats showed significantly lower insulinitis and preserved higher plasma and pancreatic insulin levels. Diol-GF failed to change the proportion of lymphocyte subsets such as T cells, natural killer cells, and macrophages in the spleen and blood. Diol-GF had no effect on the ability of DP-BB rat splenocytes to induce diabetes in recipients. Diol-GF and diol-ginsenoside Rb1 significantly decreased tumor necrosis factor α production, whereas diol-ginsenosides Rb1 and Rd decreased interleukin 1β production in RAW264.7 cells. Furthermore, mixed cytokine- and chemical-induced β-cell cytotoxicity was greatly inhibited by Diol-GF and diol-ginsenosides Rc and Rd in RIN5mF cells. However, nitric oxide production in RAW264.7 cells was unaffected by diol-ginsenosides.

Conclusion: Diol-GF, but not Triol-GF, significantly delayed the development of insulinitis and T1D in DP-BB rats. The antidiabetogenic action of Diol-GF may result from the decrease in cytokine production and increase in β-cell resistance to cytokine/free radical-induced cytotoxicity.

© 2019 The Korean Society of Ginseng. Publishing services by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Autoimmune or type 1 diabetes (T1D) is thought to result from cell-mediated immune destruction of the pancreatic β cells of the islets of Langerhans [1]. In both human and spontaneously diabetic animal models, islets are infiltrated by inflammatory immune cells that cause the destruction of β cells (insulinitis) either through major histocompatibility–restricted autoimmune events via autoreactive CD8⁺ lymphocytes or nonspecific inflammatory reactions [1–4]. Cytokines produced by autoreactive effector cells, especially by macrophages or T cells, also destroy β cells [5]. Cytokines are capable of recruiting and activating autoreactive effector cells and

injuring β cells directly or indirectly through the production of reactive oxygen intermediates and nitrogen intermediates [6–8]. Among animal models of T1D, diabetes-prone biobreeding (DP-BB) rats exhibit several clinical and histopathological characteristics of human T1D, including the spontaneous onset of diabetes and insulinitis at the age of 8–16 weeks, selective β-cell destruction, requirement of exogenous insulin to sustain life, and severe ketosis [9].

Although the prevalence of T1D (5~10%) is lower than that of type 2 diabetes in diabetic patients, the complications of T1D can be more severe and detrimental to the patients because T1D commonly occurs in children and young adults [10]. Thus, it is

* Corresponding author. College of Pharmacy, Seoul National University, 1 Gwanak-ro, Gwanak-gu, Seoul 08826, Korea, and College of Pharmacy, Ajou University, 206 Worldcup-ro, Yeongtong-gu, Suwon, Gyeonggi-do 16499, Republic of Korea.

E-mail addresses: moonck@snu.ac.kr, moonck21@ajou.ac.kr (C.-K. Moon).

☆ These authors contributed equally to this study.

☆☆ Present address: Research Headquarters, Shin Poong Pharm. Co., Ltd., Seoul, 06242, Korea.

clinically important to identify biologically effective substances that help prevent or delay the onset of T1D for improving quality of life. One such candidate would be *Panax ginseng* C.A. Meyer (Araliaceae), which is one of the most popular natural tonics used to manage diabetes mellitus in southeastern Asia for hundreds of years [11]. Ginsenosides (ginseng saponins) are known as the major components of ginseng [12,13] and may be classified into diol- and triol-saponins based on the structures of aglycones (protopanaxadiol and protopanaxatriol) [12,14] (Fig. 1). Diol-ginsenoside fraction (Diol-GF) and triol-ginsenoside fraction (Triol-GF), which are prepared from the radix of Korean Red Ginseng (KRG) as per a well-defined procedure, contain several ginsenosides with different sugar moieties in the molecules [15,16].

Ginseng extracts/fractions or individual ginsenosides have been reported to exhibit various physiological actions. Ginseng is effective in the treatment of type 2 diabetes in human and animal models [12,17–19]. It was also reported that islet pretreatment with red ginseng attenuated cytokine-induced islet damage [20]. However, the active components and their hypoglycemic mechanisms are questionable. Furthermore, the effects of ginsenosides on the development of T1D and their mechanisms are not yet clear [21,22].

In this study, we examined the effects of ginsenoside fractions (Diol-GF and Triol-GF) on the development of T1D in DP-BB rats. The mechanisms underlying the antidiabetogenic effect of Diol-GF were investigated with Diol-GF and purified diol-ginsenosides (Rb1, Rb2, Rc, and Rd) using macrophages (RAW264.7) and β cells (RIN5mF).

2. Materials and methods

2.1. Animals and cell lines

Three- or four-week-old female DP-BB rats were purchased from the University of Massachusetts Medical School (Worcester, MA, USA) and maintained under the standard nonspecific pathogen-free conditions with regular rat chow and water ad libitum. All the experiments were performed under the Principles of Laboratory Animal Care (NIH publication no. 85-23, revised 1985).

The rat insulinoma β -cell line, RINm5F, was kindly provided by Dr K.-S. Suh, Kyung Hee University, Seoul, Korea. RINm5F cells were maintained in RPMI-1640 medium supplemented with 10% heat-inactivated fetal bovine serum (FBS) and L-glutamine. Mouse

Table 1

The major constituents of Diol-GF and Triol-GF used in this study [16].

Fractions	Saponins							Other constituents
	Rg1	Re	Rf	Rb1	Rb2	Rc	Rd	
Triol-GF	50.22	35.28	4.5	-	-	-	-	10
Diol-GF	-	-	10.7	33.7	21.7	18.9	13.7	1.3

Diol-GF, diol-ginsenoside fraction; Triol-GF, triol-ginsenoside fraction.

Unit: relative content (%) of individual saponins in each saponin fraction.

macrophage cell line, RAW264.7, was obtained from the Korean Cell Line Bank (Seoul, Korea) and cultured in Dulbecco's modified Eagle's medium supplemented with 10% heat-inactivated FBS.

2.2. Reagents and chemicals

Diol-GF, Triol-GF, and purified diol-ginsenosides Rb1, Rb2, Rc, and Rd (Fig. 1) were provided by the Korea Ginseng and Tobacco Research Institute (Daejeon, Korea), where Diol-GF and Triol-GF were fractionated from total crude KRG saponin extracts as described previously [15,16]. The major constituents of each fraction were reported previously [16] and are summarized in Table 1. Voucher specimens have been deposited in the laboratory of the KT&G Central Research Institute in Korea as described before [23]. Mouse fluorescein isothiocyanate (FITC)-conjugated monoclonal antibodies OX19, W3/25, and OX8 were purchased from Sera Lab (Crawley Down, UK). Mouse FITC-conjugated monoclonal antibody (mAb) OX41 was supplied by Serotec (Oxford, UK). Human interleukin 1 β (hIL-1 β), recombinant mouse interferon γ (mIFN- γ), and recombinant mouse tumor necrosis factor α (mTNF- α) were procured from Sigma (St. Louis, MO, USA). Ficoll-Hypaque, RPMI-1640 medium, DMEM, FBS, and L-glutamine were purchased from Gibco (Paisley, UK). Mouse TNF- α and IL-1 β enzyme-linked immunosorbent assay kits were obtained from R&D System (Minneapolis, MN, USA). Other reagents were purchased from Sigma.

2.3. Treatment of animals with ginsenosides

Female DP-BB rats were orally treated with 1 mg ginsenoside/g body weight in 10% of ethanol/phosphate-buffered saline (PBS) once every day from the age of 3–4 weeks up to the age of 11–12




Fig. 1. Chemical structures of ginsenosides. (A) 20(S)-protopanaxadiol. (B) 20(S)-protopanaxatriol. Ara(fur), α -l-arabinofuranosyl; Ara(pyr), α -l-arabinopyranosyl; Glc, β -d-glucopyranosyl.




Fig. 2. Effects of ginsenosides on the development of T1D in DP-BB rats. (A) DP-BB rats were treated with 1 mg/g body weight of Diol-GF or Triol-GF every day for 8 weeks from the age of 3–4 weeks. The onset of diabetes was monitored by measuring urine and blood glucose levels after every 2–3 days from the age of 7 weeks. The number of animals in each group was 12. $p < 0.05$, PBS versus Diol-GF by the log-rank test. (B) At the age of 20 weeks, DP-BB rats were sacrificed, and the plasma and pancreatic immunoreactive insulin (IRI) levels were measured. Bar represents the mean ($n = 12$). Wist, PBS, Diol-GF, and Triol-GF denote Wistar Furth rat, PBS-treated DP-BB rat, Diol-GF-treated DP-BB rat, Triol-GF-treated DP-BB rat, respectively. Diol-GF, diol-ginsenoside fraction; DP-BB, diabetes-prone biobreeding; PBS, phosphate-buffered saline; T1D, type 1 diabetes; Triol-GF, triol-ginsenoside fraction.

weeks. Control rats received 10% ethanol in PBS. From 7 weeks of age, urine glucose and ketone levels were determined every 2–3 days with Diastix and Ketostix reagent strips (Ames, Miles, Ontario, Canada). If the reading of the urine glucose level was 500 mg/dL or higher, a second measurement was performed on the following day. Animals with values higher than 500 mg/dL on consecutive days were considered as diabetic. Development of diabetes was also confirmed by measuring blood glucose levels. Animals with non-fasting blood glucose levels greater than 250 mg/dL were considered as diabetic [24].

2.4. Examination of insulinitis and insulin contents

DP-BB rats were sacrificed at the age of 20 weeks, and each pancreas was excised and immediately frozen at -70°C until use. Sections ($4.5\ \mu\text{m}$) of the frozen pancreas were air-dried for 30 min, fixed with acetone for 10 min, air-dried again for 15 min, and stored at -20°C until use. For the examination of the degree of insulinitis, the sections of each pancreas were stained with hematoxylin and eosin and examined under a light microscope. Insulinitis lesions were arbitrarily classified as none (score 0), early (score 1), intermediate (score 2), late (score 3), and end-stage (score 4) insulinitis, according to the morphological criteria previously described [25].

Immunoreactive insulin contents in plasma and the pancreas were measured using a radioimmunoassay with a rat insulin antibody, as previously described [26].

2.5. Fluorescence-activated cell sorting analysis

The splenocytes and peripheral blood mononuclear cells isolated from DP-BB rats (at 20 weeks) using Ficoll-Hypaque were incubated with an appropriate amount of FITC-conjugated mouse monoclonal antibody OX19, W3/25, OX8, or OX41. The cells were fixed in 1% paraformaldehyde and analyzed using fluorescence-activated cell sorting (Beckton Dickinson, Sunnyvale, CA, USA) [27].

2.6. Adoptive transfer of diabetes

Splenocytes were isolated from nondiabetic PBS-treated or ginsenoside-treated rats at the age of 8–10 weeks and suspended

at a density of 2×10^6 cells/mL in a medium comprising 80 parts of RPMI-1640 with 10% of FBS, 20 parts of concanavalin A (ConA)-conditioned medium, and $5\ \mu\text{g}/\text{mL}$ ConA. The ConA-conditioned medium was obtained from the splenocytes of Sprague-Dawley rats incubated for 48 h in the presence of $2.5\ \mu\text{g}/\text{mL}$ of ConA. DP-BB neonates (within 24 h after birth) were injected with splenocytes (3×10^7 cells/rat), prepared as described previously, into the superficial orbital vein [28]. Development of diabetes was monitored as described previously.

2.7. Measurement of cytokines

RAW264.7 cells (1×10^6 cells/mL) were seeded in a 96-well plate and treated with diol-ginsenosides ($500\ \mu\text{g}/\text{mL}$ of Diol-GF and $100\ \mu\text{M}$ of Rb1, Rb2, Rc, and Rd) in complete Dulbecco's modified Eagle's medium for 24 h. The cells were stimulated with lipopolysaccharide (LPS) ($10\ \mu\text{g}/\text{mL}$) for another 24 h. The supernatants were collected, and cytokine productions (TNF- α and IL-1 β) were measured by enzyme-linked immunosorbent assays according to the manufacturer's description.

2.8. Measurement of nitric oxide production

The levels of nitrite, the stable end product of nitric oxide (NO) in aqueous solution, were determined as described by Green et al [29] with some modifications. RAW264.7 cells (1×10^6 cells/mL) were seeded in a 96-well plate and treated with diol-ginsenosides in complete Dulbecco's modified Eagle's medium for 24 h. The cells were washed and further treated with IFN- γ ($100\ \text{U}/\text{mL}$) in modified Eagle's medium for 24 h. After incubation, $50\ \mu\text{L}$ of the supernatant was removed and incubated with an equal volume of the Griess reagent at room temperature for 10 min. The absorbance of the solution was measured at 550-nm wavelength using a Titertek microplate reader (Flow Laboratories, Herts, U.K.).

2.9. Measurement of cell viability

The cell viability and mitochondrial activity of RINm5F cells were determined by 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium assay. RINm5F cells (1×10^6 cells/mL)




Fig. 3. Effects of ginsenosides on insulinitis development in DP-BB rats. The pancreas was removed from each rat (20-week-old) after treatment, fixed in formalin, and stained with hematoxylin and eosin. (A) Collective insulinitis score: 0, none; 1, early stage; 2, intermediate stage; 3, late stage; and 4, end-stage ($n = 5$). (B) Representative images of pancreatic islets from (a) normal Wistar Furth rat, (b) PBS-treated DP-BB rat, (c) Diol-GF-treated DP-BB rat, and (d) Triol-GF-treated DP-BB rat. Original magnification is $\times 100$. Diol-GF, diol-ginsenoside fraction; DP-BB, diabetes-prone biobreeding; PBS, phosphate-buffered saline; Triol-GF, triol-ginsenoside fraction.

seeded in 96-well plates were preincubated for 24 h at 37°C in complete RPMI medium with or without diol-ginsenosides. After washing, the cells were incubated in phenol red-free Modified Eagle's medium with or without 500 U/mL of mIFN- γ , 1,000 U/mL of mTNF- α , and 10 U/mL of hIL-1 β for another 24 h. In some experiments, sodium nitroprusside was added instead of cytokines. At the end of the incubation, 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium solution (1 mg/mL) was added, and the cells were incubated for another 3 h at 37°C. The formazan crystals were dissolved in a solution containing isopropanol and 1 N hydrochloric acid (HCl) (1,000:3, v/v) by vigorous shaking of the plates for 15 min. The optical density was determined at 540-nm wavelength, with 630 nm as a reference wavelength.

2.10. Statistical analysis

Data are expressed as means \pm standard error of mean and analyzed for statistical significance using the Student *t* test. Statistical analysis of incidence of diabetes was performed using the log-rank test. A value of $p < 0.05$ was regarded as significantly different.

3. Results

3.1. Effects of ginsenosides on the development of T1D in DP-BB rats

Treatments with Diol-GF or Triol-GF were well tolerated by DP-BB rats, as evident from their behaviors and general appearances. We observed no histopathological changes in tissues of major organs, including the spleen, lung, liver, kidney, heart, brain, and stomach after ginsenoside treatment (data not shown). As shown in Fig. 2A, PBS-treated control DP-BB rats developed an acute form of diabetes with glycosuria, hyperglycemia, and marked (>10%) weight loss by the age of 8 weeks. In contrast, Diol-GF treatment significantly delayed the onset of T1D (mean age of diabetes onset: 11.2 weeks in the PBS-treated group versus 17.7 weeks in the Diol-GF-treated group, $p < 0.05$). Moreover, the disease incidence was significantly reduced among rats treated with Diol-GF at the end of the experiment (83.3% in the PBS-treated group versus 50% in the Diol-GF-treated group). At 12 weeks, blood glucose levels of DP-BB rats treated with Diol-GF (137.67 ± 12.47 mg/dL) were significantly lower than those of control rats (311.33 ± 6.96 mg/dL). Triol-GF slightly delayed the onset of diabetes (mean age of diabetes onset: 11.2 weeks versus 14.4 weeks).

3.2. Effects of ginsenosides on insulinitis and the insulin level in DP-BB rats

To investigate if ginsenoside treatment prevents the destruction of pancreatic β cells, the DP-BB rats were sacrificed at 20 weeks, and insulin levels in plasma and the pancreas and the development of insulinitis were examined in treated animals. Immunoreactive insulin contents in the plasma and pancreas were significantly low in PBS-treated DP-BB rats than in control nondiabetic Wistar rats (Fig. 2B; 6.5 versus 23.1 μ mol/mL in plasma and 0.64 versus 2.50 U/g in the pancreas). In comparison with PBS treatment, Diol-GF treatment increased the insulin levels in plasma (12.65 μ mol/mL) and the pancreas (1.53 U/g) of the DP-BB rats. Triol-GF, however, had no effect on insulin levels in both plasma and the pancreas.

Treatment with Diol-GF significantly reduced insulinitis in the DP-BB rats (Fig. 3A). The islets from the PBS-treated DP-BB rats showed massive infiltration of lymphocytes and apparent β -cell destruction (Fig. 3B, Panel b) as compared with the islets from normal Wistar rats (Panel a). On the other hand, the islets from the Diol-GF-treated rats showed reduced insulinitis (Panel c). Although the islets from the Triol-GF-treated DP-BB rats showed less β -cell destruction, the level of insulinitis observed in the islets from the Triol-GF-treated DP-BB rats was similar to that observed in the PBS-treated rats (Fig. 3A and B, Panel d).

3.3. Effect of ginsenosides on lymphocyte subsets and adoptive transfer of diabetes

To study the effect of ginsenosides on changes in immune cell populations, we examined the proportions of T cells, natural killer cells, and macrophages in the spleen or peripheral blood after Diol-GF treatment. The number and the proportion of CD4⁺ cells (W3/25; CD4⁺ T cells and macrophages), CD8⁺ cells (OX8; CD8⁺ T cells and natural killer cells), and total T cells (OX19) were unchanged after Diol-GF treatment. No change in the proportion of CD172a⁺ cells (OX41; macrophages and monocytes) was observed (Fig. 4A and B).

To examine the generation of immunoregulatory/suppressor cells that mediate the prevention of T1D in the DP-BB rats, we adoptively transferred the ConA-activated splenocytes from nondiabetic PBS-, Diol-GF-, or Triol-GF-treated rats to DP-BB neonate recipients. The ConA-activated spleen cells prepared from the PBS-treated DP-BB rats induced insulinitis and diabetes in the DP-BB neonates (Table 2). Diol-GF or Triol-GF failed to alter the




Fig. 4. Lymphocyte subsets in ginsenoside-treated DP-BB rats. (A) At the age of 20 weeks, DP-BB rats were sacrificed, and lymphocyte subsets were examined by fluorescence-activated cell sorting analysis. Values are mean \pm SEM and represent the absolute number of the lymphocyte subpopulation per spleen and should be multiplied by 10^6 to obtain the actual number. The total number of lymphocytes per spleen after Ficoll-Hypaque isolation was $4.3 \pm 1.1 \times 10^7$ cells. (B) Values are mean \pm SEM and represent the absolute number of lymphocyte subpopulation per microliter of peripheral blood. The total number of lymphocytes was 3848 ± 730 cells/ μ L of peripheral blood. Diol-GF and Triol-GF represent Diol-GF-treated DP-BB rat and Triol-GF-treated DP-BB rat, respectively. Diol-GF, diol-ginsenoside fraction; DP-BB, diabetes-prone biobreeding; SEM, standard error of mean; Triol-GF, triol-ginsenoside fraction.

ability of splenocytes to transfer insulinitis and diabetes in the recipients.

3.4. Effects of diol-ginsenosides on cytokine and NO productions in RAW264.7 cells

To evaluate the effect of diol-ginsenosides on macrophage functions, cytokine and NO productions were measured in the mouse macrophage cell line, RAW264.7, after LPS or IFN- γ stimulation for 24 h. TNF- α and IL-1 β productions were significantly increased after stimulation of RAW264.7 cells with LPS (Fig. 5A and B). Diol-GF treatment decreased the production of TNF- α , while Rb1 greatly decreased TNF- α levels. Although Diol-GF had no effect on IL-1 β production, Rb1 and Rd decreased IL-1 β production in RAW264.7 cells (Fig. 5B). NO production increased in cells treated with IFN- γ , and diol-ginsenosides failed to reduce the NO level increased by IFN- γ treatment (Fig. 5C).

3.5. Effects of diol-ginsenosides on cytokine- or sodium nitroprusside-induced cytotoxicity in RIN5mF cells

To investigate whether diol-ginsenosides induce β -cell resistance to cytotoxicity, cell viability was measured in the rat

insulinoma β -cell line, RIN5mF, after treatment with cytokines or sodium nitroprusside for 24 h. The combined treatment of hIL-1 β (10 U/mL), mTNF- α (1,000 U/mL), and mIFN- γ (500 U/mL) resulted in a significant decrease in cell viability and mitochondrial function of RIN5mF cells (Fig. 6A). Cytokine-induced cell death was confirmed by the examination of cellular morphology under a light microscope (data not shown). Preincubation of RIN5mF cells with Diol-GF, Rc, and Rd for 24 h significantly decreased the cytokine-induced cell death (Fig. 6A). The effect of diol-ginsenosides on exogenous NO-induced cytotoxicity in RIN5mF cells was also studied in the presence of sodium nitroprusside. The decomposition of sodium nitroprusside results in the generation of NO in the medium. Sodium nitroprusside treatment caused 95% cytotoxicity in RIN5mF cells (Fig. 6B). However, treatment of RIN5mF cells with Diol-GF, Rc, and Rd for 24 h significantly decreased the NO-induced toxicity (Fig. 6B).

4. Discussion

This is the first study to demonstrate that Diol-GF obtained from the KRG may significantly delay the onset of T1D, reduce the disease incidence, and decrease insulinitis in DP-BB rats. Although obtained from the same plant, Triol-GF failed to effectively delay the onset of T1D or reduce the incidence of the disease. Ginsenosides are classified into Diol-GF and Triol-GF according to the structure of aglycones (Fig. 1). Although there are minor differences in the number of hydroxyl groups and sugar moieties between Diol-GF and Triol-GF, the two compounds showed different, and sometimes opposite, physiological functions [30,31]. Different effects of Diol-GF and Triol-GF were also observed in the experiment on antidiabetogenesis. Although Diol-GF exhibited remarkable antidiabetogenic effects, Triol-GF showed none.

To elucidate the inhibitory mechanism of Diol-GF on T1D, we first examined the changes in immune cell balance in DP-BB rats. Macrophages, CD4⁺ T cells, and CD8⁺ T cells are involved in the destruction of β cells in nonobese diabetic (NOD) mice and BB rats [2]. The onset of T1D in the DP-BB rats and NOD mice was prevented by the depletion of macrophages after treatment with silica [32] or the monoclonal antibody (OX19), directed against antigens

Table 2

Effects of ginsenosides on adoptive transfer of insulinitis and diabetes to neonate DP-BB rats with ConA-activated splenocytes.

Donor of ConA-activated splenocytes	Incidence	
	Diabetes	Insulinitis
PBS-treated DP-BB rats	8 of 10 (80%)	5 of 5 (100%)
Diol-GF-treated DP-BB rats	7 of 10 (70%)	5 of 5 (100%)
Triol-GF-treated DP-BB rats	8 of 10 (80%)	5 of 5 (100%)

ConA, concanavalin A; Diol-GF, diol-ginsenoside fraction; DP-BB, diabetes-prone biobreeding; PBS, phosphate-buffered saline; Triol-GF, triol-ginsenoside fraction. Splenocytes were isolated from nondiabetic PBS-treated, Diol-GF-treated, or Triol-GF-treated rats at the age of 8–10 weeks and activated with ConA (5 μ g/mL). DP-BB rat neonates (within 24 h after birth) were injected with ConA-activated splenocytes (3×10^7 cells/rat) into the superficial orbital vein. Animals with immunocytes in pancreatic islets (from early to late insulinitis) were considered as insulinitis positive.




Fig. 5. Effects of diol-ginsenosides on cytokine and NO productions in RAW264.7 cells. RAW264.7 cells (1×10^6 cells/mL) were preincubated with or without diol-ginsenosides (Diol-GF: 500 μ g/mL; Rb1, Rb2, Rc, and Rd: 100 μ M) for 24 h. (A–B) Cells were further incubated with lipopolysaccharide (LPS) (10 μ g/mL) for 24 h. TNF- α (A) and IL-1 β (B) in the supernatants were measured by ELISA. (C) Cells were further incubated with IFN- γ (100 U/mL) for another 24 h. At the end of the incubation, NO release was measured by the Griess reagent. The results are expressed as mean \pm SEM of five independent experiments. * $p < 0.05$ as compared with PBS-LPS or PBS-IFN- γ group. Diol-GF, diol-ginsenoside fraction; ELISA, enzyme-linked immunosorbent assay; IFN- γ , interferon γ ; IL-1 β , interleukin 1 β ; PBS, phosphate-buffered saline; SEM, standard error of mean; TNF- α , tumor necrosis factor α .

expressed on the surface of all T cells [33]. Furthermore, the generation of immunoregulatory/suppressor cells may mediate the prevention of T1D in DP-BB rats [28]. Thus, the changes in the proportion of effector immune cells/regulatory cells may affect the development of insulinitis and T1D in DP-BB rats. However, we failed to observe any change in the proportion of immune cells in the spleen and peripheral blood, and adoptive transfer experiments of ConA-activated splenocytes from Diol-GF-treated DP-BB rats to neonate recipients showed no protective effect on the development of diabetes. These results indicate that the preventive effect of Diol-GF on T1D may not result from the alteration in the immune cell populations or generation of regulatory cell populations.

As Diol-GF treatment failed to change the immune cell populations, we investigated the effect of Diol-GF and individual diol-ginsenosides on cytokine production and β -cell viability. Cytokines produced by Type 1 T helper cells and macrophages exert cytotoxicity on β cells. IL-1 β , TNF- α , and IFN- γ expressed in the insulinitic lesions of NOD mice and DP-BB rats inhibited insulin

synthesis and secretion and exerted cytotoxicity to the rodent pancreatic islet β cells [34]. Moreover, these cytokines are usually more potent when present in combination to destroy human islet β cells [7,35,36]. In addition, these cytokines increase the production of inducible nitric oxide synthase (iNOS), which generates NO from L-arginine. NO exerts toxic effects on β cells by inhibiting iron-dependent enzymes, thereby impairing the cellular mitochondrial function and DNA synthesis. NO also generates cytotoxic hydroxyl radicals. Macrophages and islets of animals with advanced infiltration show high expression of iNOS, suggesting that the expression of iNOS may be associated with the development of diabetes in BB rats and NOD mice [34,36–39].

Diol-GF and Rb1 significantly decreased LPS-induced TNF- α production, whereas Rb1 and Rd decreased IL-1 β production induced by LPS in RAW264.7 cells. The evaluation of cytotoxicity induced by the combination of cytokines in β cells revealed that diol-ginsenosides (Diol-GF, Rc, and Rd) significantly reduced the cytotoxic effects of these cytokines. It is unlikely that Diol-GF




Fig. 6. Effects of ginsenosides on cytokine- or sodium nitroprusside-induced cytotoxicity in RINm5F cells. (A) RINm5F cells (1×10^6 cells/mL) were preincubated with or without diol-ginsenosides (Diol-GF: 500 μ g/mL; Rb1, Rb2, Rc, and Rd: 100 μ M) for 24 h. After washing, the cells were incubated in phenol red-free Modified Eagle's medium with or without 500 U/mL of mIFN- γ , 1,000 U/mL of mTNF- α , and 10 U/mL of hIL-1 β for another 24 h. At the end of incubation, cell viability was measured by MTT assay. The results are expressed as mean \pm SEM of five independent experiments. (B) RINm5F cells (1×10^6 cells/mL) were preincubated with or without diol-ginsenosides (Diol-GF: 500 μ g/mL; Rb1, Rb2, Rc, and Rd: 100 μ M) for 24 h. After washing, the cells were treated with 2 mM sodium nitroprusside for 18 h. Cell viability was measured by MTT assay. The results are expressed as mean \pm SEM of five independent experiments. * $p < 0.05$ as compared with the PBS-cytokine mixture or PBS-sodium nitroprusside (SNP) group. Diol-GF, diol-ginsenoside fraction; hIL-1 β , human interleukin 1 β ; mIFN- γ , mouse interferon γ ; mTNF- α , mouse tumor necrosis factor α ; MTT, PBS, phosphate-buffered saline; SEM, standard error of mean.

interacts directly with cytokines and interferes with their binding to receptors as cells were pretreated with diol-ginsenosides, and diol-ginsenosides were removed before the addition of cytokines. Diol-GF, Rc, and Rd also reduced exogenous NO-induced cell death in RINm5F cells. Diol-ginsenosides, however, had no effect on the cytokine-induced NO production in RAW264.7 cells. Preliminary experiments in a cell-free system showed that diol-ginsenosides had no direct effect on the reduction of NO/nitrite production induced by sodium nitroprusside (data not shown). This result suggests that the inhibitory effects of diol-ginsenosides on RINm5F cytotoxicity are not associated with the direct scavenging of NO from the medium [40]. These results strongly suggest that the antidiabetogenic effect of Diol-GF is caused by the decrease in cytokine productions by ginsenoside Rb1 and Rd and the increase in β -cell resistance to cellular stress in response to the ginsenoside Rc and Rd treatment.

It should be noted that those diol ginsenosides have the same chemical structure except that a single sugar residue is attached to the β -OH at the C20 position (Fig. 1). Currently, it remains unknown how the single sugar residue such as glucose, arabinopyranose, and arabinofuranose can differentially regulate cytokine production and β -cell viability. Previous studies have shown that individual components of ginsenosides can exert specific biological functions through binding to different nuclear receptors or regulating different intracellular signaling pathways although their structure–function relationships are largely unknown [41–43]. Thus, it would be plausible that each sugar residue attached to the β -OH at the C20 position may differentially affect the interaction between the ginsenosides and nuclear receptors or signaling proteins involved in cytokine production and cell viability. Further studies are needed to elucidate the underlying mechanisms behind selective effectiveness of individual ginsenosides.

It is reported that the increased mitochondrial energy supply [44] or elimination of the leukocyte 12-lipoxygenase gene [45] increases the resistance of β cells to streptozotocin or cytokines. The inhibition of protein kinase C δ was shown to protect rat INS-1 cells from IL-1 β - and streptozotocin-induced apoptosis [46]. It is unclear whether diol-ginsenosides exert effects on the factors previously reported. However, Diol-GF was shown to increase Cu/Zn superoxide dismutase, an oxygen free radical-scavenging enzyme, in rat hepatoma cell lines. The transgenic expression of Cu/Zn superoxide dismutase partially protected mice from alloxan- and streptozotocin-induced diabetes [30,31,44]. It is also reported that cytokine-induced aldehyde production, such as malondialdehyde, is associated with islet β -cell destruction and that oral administration of ginsenosides decreases malondialdehyde levels [47,48]. However, the mechanism underlying diol-ginsenoside-mediated increase in the β -cell resistance is elusive.

In conclusion, we demonstrated that Diol-GF delayed the development of T1D in the DP-BB rats without any apparent toxicity. The protective effect of Diol-GF may be mediated through the decreased production of cytokines in macrophages and increased resistance of β cells to cytokine/free radical-induced cytotoxicity.

Conflicts of interest

The authors declare no conflicts of interest.

Acknowledgments

This work was supported by a grant from the Korea Ginseng & Tobacco Research Institute.

References

- [1] Clark M, Kroger CJ, Tisch RM. Type 1 diabetes: a chronic anti-self-inflammatory response. *Front Immunol* 2017;8:1898.
- [2] Yoon JW, Jun HS. Cellular and molecular pathogenic mechanisms of insulin-dependent diabetes mellitus. *Ann N Y Acad Sci* 2001;928:200–11.
- [3] Wallberg M, Cooke A. Immune mechanisms in type 1 diabetes. *Trends Immunol* 2013;34(12):583–91.
- [4] Li M, Song LJ, Qin XY. Advances in the cellular immunological pathogenesis of type 1 diabetes. *J Cell Mol Med* 2014;18(5):749–58.
- [5] Thomas HE, Graham KL, Chee J, Thomas R, Kay TW, Krishnamurthy B. Proinflammatory cytokines contribute to development and function of regulatory T cells in type 1 diabetes. *Ann N Y Acad Sci* 2013;1283(1):81–6.
- [6] Suarez-Pinzon WL, Mabley JG, Strynadka K, Power RF, Szabo C, Rabinovitch A. An inhibitor of inducible nitric oxide synthase and scavenger of peroxynitrite prevents diabetes development in NOD mice. *J Autoimmun* 2001;16(4):449–55.
- [7] Rabinovitch A. Immunoregulation by cytokines in autoimmune diabetes. *Adv Exp Med Biol* 2003;520:159–93.
- [8] Padgett LE, Broniowska KA, Hansen PA, Corbett JA, Tse HM. The role of reactive oxygen species and proinflammatory cytokines in type 1 diabetes pathogenesis. *Ann N Y Acad Sci* 2013;1281(1):16–35.
- [9] Mordes JP, Bortell R, Blankenhorn EP, Rossini AA, Greiner DL. Rat models of type 1 diabetes: genetics, environment, and autoimmunity. *ILAR J* 2004;45(3):278–91.
- [10] Maahs DM, West NA, Lawrence JM, Mayer-Davis EJ. Epidemiology of type 1 diabetes. *Endocrinol Metab Clin North Am* 2010;39(3):481–97.
- [11] Lee K-H, Morris-Natschke S, Qian K, Dong Y, Yang X, Zhou T, Belding E, Wu S-F, Wada K, Akiyama T. Recent progress of research on herbal products used in traditional chinese medicine: the herbs belonging to the divine husbandman's herbal foundation canon (Shén Nóng Bèn Cǎo Jīng). *J Trad Complement Med* 2012;2(1):6–26.
- [12] Attele AS, Wu JA, Yuan CS. Ginseng pharmacology: multiple constituents and multiple actions. *Biochem Pharmacol* 1999;58(11):1685–93.
- [13] Sotaniemi EA, Haapakoski E, Rautio A. Ginseng therapy in non-insulin-dependent diabetic patients. *Diabetes Care* 1995;18(10):1373–5.
- [14] Shin BK, Kwon SW, Park JH. Chemical diversity of ginseng saponins from *Panax ginseng*. *J Ginseng Res* 2015;39(4):287–98.
- [15] Kim SK, Kwak YS, Kim SW, Hwang SY, Ko YS, Yoo CM. Improved method for the preparation of crude ginseng saponin. *J Ginseng Res* 1998;22(3):155–60.
- [16] Kim CS, Jeong SL, Lee YG. A rapid separation of an edible panaxadiol and panaxatriol in ginseng saponins by benzene ethylene resin adsorption. *J Ginseng Res* 1998;22(3):211–5.
- [17] Kimura I, Nakashima N, Sugihara Y, Fu-jun C, Kimura M. The anti-hyperglycaemic blend effect of traditional chinese medicine byakko-ka-ninjin-to on alloxan and diabetic KK-CaY mice. *Phytother Res* 1999;13(6):484–8.
- [18] Kim JH, Yi YS, Kim MY, Cho JY. Role of ginsenosides, the main active components of *Panax ginseng*, in inflammatory responses and diseases. *J Ginseng Res* 2017;41(4):435–43.
- [19] Gui QF, Xu ZR, Xu KY, Yang YM. The efficacy of ginseng-related therapies in type 2 diabetes mellitus: an updated systematic review and meta-analysis. *Medicine (Baltimore)* 2016;95(6):e2584.
- [20] Kim JS, Jang HJ, Kim SS, Oh MY, Kim HJ, Lee SY, Eom DW, Ham JY, Han DJ. Red ginseng administration before islet isolation attenuates apoptosis and improves islet function and transplant outcome in a syngeneic mouse marginal islet mass model. *Transplant Proc* 2016;48(4):1258–65.
- [21] Hong YJ, Kim N, Lee K, Hee Sonn C, Eun Lee J, Tae Kim S, Ho Baeg I, Lee KM. Korean red ginseng (*Panax ginseng*) ameliorates type 1 diabetes and restores immune cell compartments. *J Ethnopharmacol* 2012;144(2):225–33.
- [22] Hong BN, Ji MG, Kang TH. The efficacy of red ginseng in type 1 and type 2 diabetes in animals. *Evid Based Complement Alternat Med* 2013;2013:593181.
- [23] Kim JH, Kang SA, Han SM, Shim I. Comparison of the antiobesity effects of the protopanaxadiol- and protopanaxatriol-type saponins of red ginseng. *Phytother Res* 2009;23(1):78–85.
- [24] Nicoletti F, Di Marco R, Morrone S, Zaccone P, Lembo D, Grasso S, Santoni A, Meroni PL, Bendtzen K. Reduction of spontaneous autoimmune diabetes in diabetes-prone BB rats with the novel immunosuppressant fusidic acid. Effect on T-cell proliferation and production of interferon-gamma. *Immunology* 1994;81(2):317–21.
- [25] Chung YH, Jun HS, Kang Y, Hirasawa K, Lee BR, Van Rooijen N, Yoon JW. Role of macrophages and macrophage-derived cytokines in the pathogenesis of Kilham rat virus-induced autoimmune diabetes in diabetes-resistant Bio-Breeding rats. *J Immunol* 1997;159(1):466–71.
- [26] Rabinovitch A, Suarez-Pinzon W, El-Sheikh A, Sorensen O, Power RF. Cytokine gene expression in pancreatic islet-infiltrating leukocytes of BB rats: expression of Th1 cytokines correlates with beta-cell destructive insulinitis and IDDM. *Diabetes* 1996;45(6):749–54.
- [27] Nicoletti F, Zaccone P, Di Marco R, Lunetta M, Magro G, Grasso S, Meroni P, Garotta G. Prevention of spontaneous autoimmune diabetes in diabetes-prone BB rats by prophylactic treatment with antirat interferon-gamma antibody. *Endocrinology* 1997;138(1):281–8.
- [28] Sobel DO, Goyal D, Ahvazi B, Yoon JW, Chung YH, Bagg A, Harlan DM. Low dose poly I:C prevents diabetes in the diabetes prone BB rat. *J Autoimmun* 1998;11(4):343–52.

- [29] Green LC, Wagner DA, Glogowski J, Skipper PL, Wishnok JS, Tannenbaum SR. Analysis of nitrate, nitrite, and [15N]nitrate in biological fluids. *Anal Biochem* 1982;126(1):131–8.
- [30] Kim YH, Park KH, Rho HM. Transcriptional activation of the Cu,Zn-superoxide dismutase gene through the AP2 site by ginsenoside Rb2 extracted from a medicinal plant, *Panax ginseng*. *J Biol Chem* 1996;271(40):24539–43.
- [31] Chang MS, Lee SG, Rho HM. Transcriptional activation of Cu/Zn superoxide dismutase and catalase genes by panaxadiol ginsenosides extracted from *Panax ginseng*. *Phytother Res* 1999;13(8):641–4.
- [32] Lee KU, Amano K, Yoon JW. Evidence for initial involvement of macrophage in development of insulinitis in NOD mice. *Diabetes* 1988;37(7):989–91.
- [33] Like AA, Biron CA, Weringer EJ, Byman K, Sroczynski E, Guberski DL. Prevention of diabetes in BioBreeding/Worcester rats with monoclonal antibodies that recognize T lymphocytes or natural killer cells. *J Exp Med* 1986;164(4):1145–59.
- [34] Eizirik DL, Sandler S, Welsh N, Cetkovic-Cvrlje M, Nieman A, Geller DA, Pipeleers DG, Bendtzen K, Hellerström C. Cytokines suppress human islet function irrespective of their effects on nitric oxide generation. *J Clin Invest* 1994;93(5):1968–74.
- [35] Rabinovitch A. Immunoregulatory and cytokine imbalances in the pathogenesis of IDDM. Therapeutic intervention by immunostimulation? *Diabetes* 1994;43(5):613–21.
- [36] Nicoletti F, Meroni PL, Landolfo S, Gariglio M, Guzzardi S, Barcellini W, Lunetta M, Mughini L, Zanussi C. Prevention of diabetes in BB/Wor rats treated with monoclonal antibodies to interferon-gamma. *Lancet* 1990;336(8710):319.
- [37] Andersen NA, Larsen CM, Mandrup-Poulsen T. TNFalpha and IFNgamma potentiate IL-1beta induced mitogen activated protein kinase activity in rat pancreatic islets of Langerhans. *Diabetologia* 2000;43(11):1389–96.
- [38] Eizirik DL, Flodstrom M, Karlsen AE, Welsh N. The harmony of the spheres: inducible nitric oxide synthase and related genes in pancreatic beta cells. *Diabetologia* 1996;39(8):875–90.
- [39] Rabinovitch A, Baquerizo H, Pukel C, Sumoski W. Effects of cytokines on rat pancreatic islet cell monolayer cultures: distinction between functional and cytotoxic effects on islet beta-cells. *Reg Immunol* 1989;2(2):77–82.
- [40] Kim YK, Guo Q, Packer L. Free radical scavenging activity of red ginseng aqueous extracts. *Toxicology* 2002;172(2):149–56.
- [41] Ahuja A, Kim JH, Kim JH, Yi YS, Cho JY. Functional role of ginseng-derived compounds in cancer. *J Ginseng Res* 2018;42(3):248–54.
- [42] Mohanan P, Subramaniam S, Mathiyalagan R, Yang DC. Molecular signaling of ginsenosides Rb1, Rg1, and Rg3 and their mode of actions. *J Ginseng Res* 2018;42(2):123–32.
- [43] Leung KW, Wong AS. Pharmacology of ginsenosides: a literature review. *Chin Med* 2010;5:20.
- [44] Burkart V, Brenner HH, Hartmann B, Kolb H. Metabolic activation of islet cells improves resistance against oxygen radicals or streptozocin, but not nitric oxide. *J Clin Endocrinol Metab* 1996;81(11):3966–71.
- [45] Bleich D, Chen S, Zipser B, Sun D, Funk CD, Nadler JL. Resistance to type 1 diabetes induction in 12-lipoxygenase knockout mice. *J Clin Invest* 1999;103(10):1431–6.
- [46] Carpenter L, Cordery D, Biden TJ. Inhibition of protein kinase C delta protects rat INS-1 cells against interleukin-1beta and streptozotocin-induced apoptosis. *Diabetes* 2002;51(2):317–24.
- [47] Ismail MF, Gad MZ, Hamdy MA. Study of the hypolipidemic properties of pectin, garlic and ginseng in hypercholesterolemic rabbits. *Pharmacol Res* 1999;39(2):157–66.
- [48] Deng HL, Zhang JT. Anti-lipid peroxidative effect of ginsenoside Rb1 and Rg1. *Chin Med J (Engl)* 1991;104(5):395–8.