



# Two new species of *Limbodessus* diving beetles from New Guinea - short verbal descriptions flanked by online content (digital photography, $\mu$ CT scans, drawings and DNA sequence data)

Michael Balke<sup>‡</sup>, Bernhard Ruthensteiner<sup>‡</sup>, Evie Lilly Warikar<sup>§</sup>, Katja Neven<sup>‡</sup>, Lars Hendrich<sup>‡</sup>

<sup>‡</sup> Zoologische Staatssammlung, Munich, Germany

<sup>§</sup> Department of Biology, Universitas Cendrawasih, Jayapura, Indonesia

Corresponding author: Michael Balke ([coleoptera-zsm@zsm.mwn.de](mailto:coleoptera-zsm@zsm.mwn.de))

Academic editor: Miguel Alonso-Zarazaga

Received: 05 Nov 2015 | Accepted: 18 Dec 2015 | Published: 22 Dec 2015

Citation: Balke M, Ruthensteiner B, Warikar E, Neven K, Hendrich L (2015) Two new species of *Limbodessus* diving beetles from New Guinea - short verbal descriptions flanked by online content (digital photography,  $\mu$ CT scans, drawings and DNA sequence data). Biodiversity Data Journal 3: e7096. doi: [10.3897/BDJ.3.e7096](https://doi.org/10.3897/BDJ.3.e7096)

ZooBank: [urn:lsid:zoobank.org/pub:B84A30C1-77BE-4C49-9B60-7C0AE708D21E](https://urn:lsid:zoobank.org/pub:B84A30C1-77BE-4C49-9B60-7C0AE708D21E)

## Abstract

### Background

To date only one species of *Limbodessus* diving beetles has been reported from the Island of New Guinea, *L. compactus* (Clark, 1862), which is widespread in the Australian region.

### New information

We describe two new species of microendemic New Guinea *Limbodessus* and use a compact descriptive format flanked by enriched online content in wiki powered species pages. *Limbodessus baliem* **sp.n.** is described from ca. 1,600 m altitude in the Baliem Valley of Papua and *Limbodessus alexanderi* **sp.n.** from >3,000 m altitude north of Sugapa, Papua.

Based on our analysis, we also transfer three species from other genera to *Limbodessus* Guignot, 1939, with the following changes: *Limbodessus deflectus* (Ordish, 1966), **new combination**; *Limbodessus leverii* (J. Balfour-Browne, 1944), **new combination**; and *Limbodessus plicatus* (Sharp, 1882), **new combination**.

## Keywords

Dytiscidae, *Limbodessus*, new species, high resolution photography,  $\mu$ CT scans, DNA sequence data, minimalistic descriptions, New Guinea

## Introduction

*Limbodessus* Guignot (1939) contains 71 described species mostly distributed in the Australasian region, few species are Oriental or reach the Palearctic in Japan (Nilsson (2001), Nilsson (2015)). The majority of species was described from underground waters in Australia, the otherwise epigeal species prefer stagnant water habitats or densely vegetated backflows or marginal areas of slowly flowing streams. Underground and interstitial species show adaptations to their environment, such as reduction or loss of the eyes as well as reduced wings (Watts and Humphreys (1999), Larson (1994)). Based on their morphology being different from the epigeal species (Watts and Leijds (2005)), new genera such as *Boongurrus* Larson and *Tjirtudessus* Watts & Humphreys were suggested, but later synonymised with *Limbodessus* by Balke and Ribera (2004) based on molecular phylogenetic data and a morphological apomorphy (male paramere with hook or bent finger like apical part).

Here we describe two new microendemic species from New Guinea. One occurs in an intramontane depression at 1,600–1,700 m and the other one in the tropical-montane to alpine habitat above 3,000 m. We utilize a compact, integrative descriptive format following Riedel et al. (2013a), Riedel et al. (2013b) combining morphological and molecular evidence. We provide high resolution online digital imaging resources, as well as  $\mu$ CT data illustrating characters not explicitly described here but anticipating potential interest in certain structures in the future (e.g. sculpture of metacoxa, width and structure of lateral wings of metaventrite).

## Materials and methods

Specimens are in the following collections:

**CLH** Collection of Lars Hendrich, Munich, Germany (property of NHMW)

[MZB Museum Zoologicum Bogoriense, now LIPI RC Biology, Division of Zoology, Cibinong, West Java, Indonesia](#)

[NHMW Naturhistorisches Museum Wien, Vienna, Austria](#)

[NMPC National Museum, Prague, Czech Republic](#)

[ZSM Zoologische Staatssammlung München, Munich, Germany](#)

Measurements were taken with a Leica M205 C stereomicroscope. The following abbreviations were used: TL (total body length), TL-H (total body length without head), and MW (maximum body width). UNCEN legit indicates specimens collected during a field course with the Cendrawasih University, Jayapura, Papua (UNCEN).

DNA sequence data were generated using standard methods described in detail in our laboratory wiki: [http://zsm-entomology.de/wiki/The Beetle D N A Lab](http://zsm-entomology.de/wiki/The_Beetle_D_N_A_Lab)

Digital images were taken with a Nikon D3X equipped with a bellows or expansion rings and lenses: Mitutoyo 10x ELWD Planapo or Leitz Photar 25/2.8. Illumination came from three compact Nikon flashes, and the instrument was moved on an Isel linear drive or Cognisys Stackshot (for very small steps 1–7 µm). Image stacks were combined using the method A in Helicon focus software.

For micro-CT scanning, the specimens were attached with soft dental wax to tips of glass Pasteur pipettes. Scanning was performed with a Phoenix Nanotom M (GE Measurement & Control, Wunstorf, Germany) cone beam CT scanner at a voltage of 60 kV and a current of 240mA (*Limbodessus baliem*, *Kaef2-gr*) or a voltage of 50 kV and a current of 275 mA (*Kaef2-kl*) respectively, using a molybdenum target. Each scan took 144 minutes. 1.440 projections were prepared per scan. The 3D datasets (prepared with the `datos|x` reconstruction software, GE Measurement & Control) were examined by volume rendering with Drishti 2.3.2 Limaye (2012) and VGStudioMax 2.2 (Visual Graphics GmbH, Heidelberg, Germany) software.

In addition, surface meshes were generated using the threshold tool in the segmentation editor of the software Amira 5.4.5 (FEI VisualizationSciences Group, Burlington MA, USA). For varying intensities in the volume data, the threshold was slightly locally adjusted. The attachment wax, which has X-ray absorption nearly as intense as that of the specimens' skeletons, was removed during segmentation by a combination of threshold and manual (lasso and brush tools) segmentation. To constrain total mesh complexity (eventually to ca. one million faces) and reduce file size to a tolerable level, all internal structures were removed.

The PDF 3D models were prepared largely following the procedures outlined by Ruthensteiner and Heß (2008).

## Data resources

DNA sequences were uploaded to EMBL and are available under accession numbers [LN884305–LN884314](#).

The *cox1* sequences are available in alignment format on Dryad: doi:[10.5061/dryad.q1b24](#). Newly generated data are appended here in fasta format Suppl. material 1.

Animated videos of the  $\mu$ CT scans are available at YouTube tagged as "*Limbodessus*". The  $\mu$ CT data were deposited in the [Morphosource](#) database in their own [project](#).

## Taxon treatments

### *Limbodessus alexanderi* Balke & Hendrich, sp. n.

- ZooBank [urn:lsid:zoobank.org:act:9C940C82-646F-4075-86B9-3E88D397ABEA](https://zoobank.org/act:9C940C82-646F-4075-86B9-3E88D397ABEA)

#### Materials

##### *Holotype:*

- a. scientificName: *Limbodessus alexanderi*; taxonID: [urn:lsid:zoobank.org:act:9C940C82-646F-4075-86B9-3E88D397ABEA](https://zoobank.org/act:9C940C82-646F-4075-86B9-3E88D397ABEA); order: Coleoptera; family: Dytiscidae; island: New Guinea; country: Indonesia; stateProvince: Papua; locality: N Sugapa; verbatimElevation: 3000 m; verbatimCoordinates: 3°40'11.90"S, 137°4'51.31"E; decimalLatitude: -3.669972; decimalLongitude: 137.080919; samplingProtocol: collected with strainer; eventDate: 27.xii.1995; sex: male; lifeStage: adult; recordedBy: Riedel; institutionCode: NHMW; collectionCode: Insects

##### *Paratype:*

- a. scientificName: *Limbodessus alexanderi*; taxonID: [urn:lsid:zoobank.org:act:9C940C82-646F-4075-86B9-3E88D397ABEA](https://zoobank.org/act:9C940C82-646F-4075-86B9-3E88D397ABEA); order: Coleoptera; family: Dytiscidae; island: New Guinea; country: Indonesia; stateProvince: Papua; locality: N Sugapa; verbatimElevation: 3000 m; verbatimCoordinates: 3°40'11.90"S, 137°4'51.31"E; decimalLatitude: -3.669972; decimalLongitude: 137.080919; samplingProtocol: collected with strainer; eventDate: 27.xii.1995; individualID: four of them with green extraction voucher labels: M. Balke 4424, 4425, 6423, 6424; individualCount: 130; lifeStage: adult; recordedBy: Riedel; institutionCode: MZB, NHMW, ZSM; collectionCode: Insects

#### Description

A large, dark brown to black *Limbodessus*: length of body 3.0–3.5 mm (N=20); with pronounced habitus disruption between pronotum and elytron Figs 1, 2, 3, 4, 5, 6, 37, 39; dorsoventrally rather thick Figs 7, 8; cervical line present but faint in male or partly dissolved into punctures in females, distinct pronotal plica present, elytral plica absent; flight wings vestigial; metacoxa and metaventrite with few punctures only Fig. 22; ventral side with notable modifications: elytral epipleuron apically modified: slightly concave, with inner margin dilated, its form reminding of a spear tip or shallow spoon Figs 7, 16, 17, 37, 39.



Figure 1.

*Limbodessus alexanderi*, male, dorsal habitus. Length of beetle: 3.1 mm.



Figure 2.

*Limbodessus alexanderi*, female, dorsal habitus. Length of beetle: 3.1 mm.




Figure 3.

*Limbodessus alexanderi*, male, dorsal habitus.  $\mu$ CT scan. Length of beetle: 3.1 mm.



Figure 4.

*Limbodessus alexanderi*, female, dorsal habitus.  $\mu$ CT scan. Length of beetle: 3.1 mm.




Figure 5.

*Limbodessus alexanderi*, male, frontal habitus.  $\mu$ CT scan. Length of beetle: 3.1 mm.



Figure 6.

*Limbodessus alexanderi*, female, frontal habitus.  $\mu$ CT scan. Length of beetle: 3.1 mm.



Figure 7.

*Limbodessus alexanderi*, female, lateral habitus. Length of beetle: 3.1 mm.




Figure 8.

*Limbodessus alexanderi*, male, lateral habitus.  $\mu$ CT scan. Length of beetle: 3.1 mm.



Figure 9.

*Limbodessus baliem*, female, lateral habitus. Length of beetle: 2.8 mm.



Figure 10.

*Limbodessus alexanderi*, female, detailed elytral and pronotal sculpture. Lens: Mitutoyo ELWD 10x.





Figure 11.

*Limbodessus baliem*, male, lateral habitus.  $\mu$ CT scan. Length of beetle: 2.8 mm.



Figure 12.

*Limbodessus baliem*, female, detailed elytral and pronotal sculpture. Lens: Mitutoyo ELWD 10x.



Figure 13.

*Limbodessus baliem*, male, habitus dorsally. Length of beetle: 2.8 mm.



Figure 14.

*Limbodessus baliem*, female, habitus dorsally. Length of beetle: 2.8 mm.




Figure 15.

*Limbodessus baliem*, male, habitus dorsally.  $\mu$ CT scan. Length of beetle: 2.8 mm.



Figure 16.

*Limbodessus alexanderi*, male, habitus ventrally.  $\mu$ CT scan. Length of beetle: 3.1 mm.



Figure 17.

*Limbodessus alexanderi*, female, habitus ventrally.  $\mu$ CT scan. Length of beetle: 3.1 mm.



Figure 18.

*Limbodessus baliem*, male, habitus frontally.  $\mu$ CT scan. Length of beetle: 2.8 mm.



Figure 19.

*Limbodessus baliem*, male, detailed elytral and pronotal sculpture. Lens: Mitutoyo ELWD 10x.



Figure 20.

*Limbodessus baliem*, female, habitus ventrally. Length of beetle: 2.8 mm.



Figure 21.

*Limbodessus baliem*, male, habitus ventrally. Length of beetle: 2.8 mm.




Figure 22.

*Limbodessus alexanderi*, ventral aspect, metacoxa and metaventricle.



Figure 23.

Distribution of *Limbodessus baliem* and *L. alexanderi* in New Guinea. *Limbodessus compactus* is widespread especially in the lowlands, it is absent from the Baliem Valley and altitudes above 2000 m.



Figure 24.

*Limbodessus baliem*, male, habitus ventrally.  $\mu$ CT scan. Length of beetle: 2.8 mm.




Figure 25.

*Limbodessus alexanderi*, male, median lobe in ventral view.




Figure 26.

*Limbodessus alexanderi*, male, median lobe in lateral view.






Figure 27.

*Limbodessus baliem*, ventral aspect, metacoxa and metaventrite.




Figure 28.

*Limbodessus compactus*, ventral aspect, metacoxa and metaventrite.




Figure 29.

*Limbodessus baliem*, male, median lobe in ventral view.




Figure 30.

*Limbodessus baliem*, male, median lobe in lateral view.




Figure 31.

*Limbodessus alexanderi*, male, paramere. Total length 0.46 mm.




Figure 32.

*Limbodessus compactus*, male, median lobe in ventral view (after Biström 1988).




Figure 33.  
*Limbodessus compactus*, male, median lobe in lateral view (after Biström, 1988).




Figure 34.  
*Limbodessus baliem*, male, paramere. Total length 0.33 mm.




Figure 35.

*Limbodessus compactus*, male, paramere. Total length 0.3 mm




Figure 36.

Elytral epipleuron of *Limbodessus compactus*, arrows point to base where there is a carina.




Figure 37.

*Limbodessus alexanderi*, male, habitus. Animated  $\mu$ CT scan. Length of beetle: 3.1 mm.




Figure 38.

Elytral epipleuron of *Allodessus megacephalus* (Gschwendtner 1931), arrow points to base where there is no carina.



Figure 39.

*Limbodessus alexanderi*, female, habitus. Animated  $\mu$ CT scan. Length of beetle: 3.1 mm.




Figure 40.

Neighbour joining tree using 3' cox1 sequence data for New Guinea *Limbodessus* with other epigeal species from neighbouring areas. *Limbodessus capeensis* Watts and Leijs (2005) was not available.

Sexes dimorphic, see below.

**Male:** Antenna filiform Figs 1, 3, 5. Dorsal surface with coarse punctures but otherwise with shiny surface. Median lobe of aedeagus as in Figs 25, 26 with tiny setae on tip, paramere as in Fig. 31.

**Female:** Antenna with strongly enlarged antennomeres forming a conspicuous club Figs 2, 4, 6, 17. Dorsal surface with very fine microreticulation between surface punctures and surfaces therefore appearing matt Figs 2, 10.

### Etymology

Named after Alexander Riedel who discovered this species. The species name is a noun in the genitive case.

### Distribution

Indonesian New Guinea, known only from the type locality which is the mountain range north of Sugapa, Papua (Fig. 23).

### Ecology

Puddles in high altitude grassland in the tropical montane / subalpine habitat. Here, a second, much smaller and black *Limbodessus* was also collected which is the female of an undescribed species.

### Conservation

The species is most likely not threatened due to its occurrence on remote high altitude plateau.

### Online resources

Higher resolution digital images,  $\mu$ CT data as well as sequence data have been deposited in public databases. The Species-ID species page is a versioned wiki site and can be enhanced through community contributions Hendrich and Balke (2011).

**Species-ID species page:** [http://species-id.net/wiki/Limbodessus\\_alexanderi](http://species-id.net/wiki/Limbodessus_alexanderi)

**DNA sequences:** Partial 3' *cox1* sequence deposited at: LN884309–LN884312.

**$\mu$ CT data:** In two Morphosource projects [here](#) and [here](#).



***Limbodessus baliem* Balke & Hendrich, sp. n.**

- ZooBank [urn:lsid:zoobank.org:act:3EC40F14-C7D9-42C8-84EB-91ADF93C9F7D](https://doi.org/10.3896/abx-3EC40F14-C7D9-42C8-84EB-91ADF93C9F7D)

**Materials***Holotype:*

- a. scientificName: *Limbodessus baliem*; taxonID: [urn:lsid:zoobank.org:act:3EC40F14-C7D9-42C8-84EB-91ADF93C9F7D](https://doi.org/10.3896/abx-3EC40F14-C7D9-42C8-84EB-91ADF93C9F7D); order: Coleoptera; family: Dytiscidae; island: New Guinea; country: Indonesia; stateProvince: Papua; locality: Wamena; verbatimElevation: 1600 m; locationRemarks: water holes near runway; verbatimCoordinates: 4° 6'16.76"S, 138° 57'37.96"E; samplingProtocol: collected with strainer; eventDate: 31.viii. & 6.ix.1990; sex: male; lifeStage: adult; recordNumber: IR1; recordedBy: Balke & Hendrich; institutionCode: NHMW; collectionCode: Insects

*Paratype:*

- a. scientificName: *Limbodessus baliem*; taxonID: [urn:lsid:zoobank.org:act:3EC40F14-C7D9-42C8-84EB-91ADF93C9F7D](https://doi.org/10.3896/abx-3EC40F14-C7D9-42C8-84EB-91ADF93C9F7D); order: Coleoptera; family: Dytiscidae; island: New Guinea; country: Indonesia; stateProvince: Papua; locality: Wamena; verbatimElevation: 1600 m; locationRemarks: water holes near runway; verbatimCoordinates: 4° 6'16.76"S, 138° 57'37.96"E; samplingProtocol: collected with strainer; eventDate: 31.viii. & 6.ix.1990; individualCount: 15; lifeStage: adult; recordNumber: IR1; recordedBy: Balke & Hendrich; institutionCode: NHMW; collectionCode: Insects
- b. scientificName: *Limbodessus baliem*; taxonID: [urn:lsid:zoobank.org:act:3EC40F14-C7D9-42C8-84EB-91ADF93C9F7D](https://doi.org/10.3896/abx-3EC40F14-C7D9-42C8-84EB-91ADF93C9F7D); order: Coleoptera; family: Dytiscidae; island: New Guinea; country: Indonesia; stateProvince: Papua; locality: Wamena; verbatimElevation: 1700 m; locationRemarks: water holes near runway; verbatimCoordinates: 4° 6'16.76"S, 138° 57'37.96"E; samplingProtocol: collected with strainer; eventDate: 20.-27.ix.1992; individualCount: 3; lifeStage: adult; recordNumber: IR54A = 57; recordedBy: Balke; institutionCode: NHMW; collectionCode: Insects
- c. scientificName: *Limbodessus baliem*; taxonID: [urn:lsid:zoobank.org:act:3EC40F14-C7D9-42C8-84EB-91ADF93C9F7D](https://doi.org/10.3896/abx-3EC40F14-C7D9-42C8-84EB-91ADF93C9F7D); order: Coleoptera; family: Dytiscidae; island: New Guinea; country: Indonesia; stateProvince: Papua; locality: Wamena; verbatimElevation: 1700 m; locationRemarks: water holes near runway; verbatimCoordinates: 4° 6'16.76"S, 138° 57'37.96"E; samplingProtocol: collected with strainer; eventDate: 15.x.1993; individualCount: 1; lifeStage: adult; recordedBy: Riedel; institutionCode: ZSM; collectionCode: Insects
- d. scientificName: *Limbodessus baliem*; taxonID: [urn:lsid:zoobank.org:act:3EC40F14-C7D9-42C8-84EB-91ADF93C9F7D](https://doi.org/10.3896/abx-3EC40F14-C7D9-42C8-84EB-91ADF93C9F7D); order: Coleoptera; family: Dytiscidae; island: New Guinea; country: Indonesia; stateProvince: Papua; locality: Wamena; verbatimElevation: 1600 m; locationRemarks: water holes near runway; verbatimCoordinates: 4° 6'16.76"S, 138° 57'37.96"E; samplingProtocol: collected with strainer; eventDate: 19.ix.1992; individualCount: 39; lifeStage: adult; recordNumber: IR52; recordedBy: Balke; institutionCode: NHMW; collectionCode: Insects
- e. scientificName: *Limbodessus baliem*; taxonID: [urn:lsid:zoobank.org:act:3EC40F14-C7D9-42C8-84EB-91ADF93C9F7D](https://doi.org/10.3896/abx-3EC40F14-C7D9-42C8-84EB-91ADF93C9F7D); order: Coleoptera; family: Dytiscidae; island: New Guinea; country: Indonesia; stateProvince: Papua; locality: Wamena; verbatimElevation: 1600 m; locationRemarks: water holes near runway; verbatimCoordinates: 4° 6'16.76"S, 138° 57'37.96"E; samplingProtocol: collected with strainer; eventDate: 19.ix.1992; individualCount: 3; lifeStage: adult; recordNumber: IR53; recordedBy: Balke; institutionCode: NHMW; collectionCode: Insects

- f. scientificName: *Limbodessus baliem*; taxonID: [urn:lsid:zoobank.org:act:3EC40F14-C7D9-42C8-84EB-91ADF93C9F7D](https://doi.org/10.3896/abx3-3EC40F14-C7D9-42C8-84EB-91ADF93C9F7D); order: Coleoptera; family: Dytiscidae; island: New Guinea; country: Indonesia; stateProvince: Papua; locality: Wamena; verbatimElevation: 1600 m; locationRemarks: water holes near runway; verbatimCoordinates: 04 06.323S, 138 57.693E; samplingProtocol: collected with strainer; eventDate: 18.x.2011; individualID: one of them with green extraction voucher label M. Balke 5082 and one with 5083; individualCount: 37; lifeStage: adult; recordNumber: PAP04; recordedBy: UNCEN; institutionCode: MZB, ZSM; collectionCode: Insects
- g. scientificName: *Limbodessus baliem*; taxonID: [urn:lsid:zoobank.org:act:3EC40F14-C7D9-42C8-84EB-91ADF93C9F7D](https://doi.org/10.3896/abx3-3EC40F14-C7D9-42C8-84EB-91ADF93C9F7D); order: Coleoptera; family: Dytiscidae; island: New Guinea; country: Indonesia; stateProvince: Papua; locality: Wamena, 20 mins towards Jiwika; verbatimElevation: 1620 m; locationRemarks: limestone creek; verbatimCoordinates: 03 56.953S, 138 54.375E; samplingProtocol: collected with strainer; eventDate: 18.x.2011; individualCount: 10; lifeStage: adult; recordNumber: PAP05; recordedBy: UNCEN; institutionCode: MZB, ZSM; collectionCode: Insects
- h. scientificName: *Limbodessus baliem*; taxonID: [urn:lsid:zoobank.org:act:3EC40F14-C7D9-42C8-84EB-91ADF93C9F7D](https://doi.org/10.3896/abx3-3EC40F14-C7D9-42C8-84EB-91ADF93C9F7D); order: Coleoptera; family: Dytiscidae; island: New Guinea; country: Indonesia; stateProvince: Papua; locality: Jiwika; verbatimLocality: INDONESIA, Papua: Jayawijaya Distr., Baliem valley, 14 km NNE of Wamena, wetland & gardens nr Jiwika [Kuru], 03°58.0-5'S, 138°55.8 56.3'E; 1660 m, J. Hájek & J. Šumpich leg., 6.ii.2015; verbatimElevation: 1660 m; locationRemarks: wetland; verbatimCoordinates: 03°58.0'S, 138°55.8'E; samplingProtocol: collected with strainer; eventDate: 6.ii.2015; individualCount: 8; lifeStage: adult; recordedBy: J. Hájek & J. Šumpich; institutionCode: NMPC; collectionCode: Insects
- i. scientificName: *Limbodessus baliem*; taxonID: [urn:lsid:zoobank.org:act:3EC40F14-C7D9-42C8-84EB-91ADF93C9F7D](https://doi.org/10.3896/abx3-3EC40F14-C7D9-42C8-84EB-91ADF93C9F7D); order: Coleoptera; family: Dytiscidae; island: New Guinea; country: Indonesia; stateProvince: Papua; locality: Wamena; verbatimLocality: INDONESIA, Papua: Jayawijaya Distr., Baliem valley, Wamena env., ruderal & open marsh nr Baliem river, 04°04.9'S, 138°57.3'E; 1650 m, J. Hájek & J. Šumpich leg., 31.i.+8.ii.2015; verbatimElevation: 1650 m; locationRemarks: wetland; verbatimCoordinates: 04°04.9'S, 138°57.3'E; samplingProtocol: collected with strainer; eventDate: 31.i.+8.ii.2015; individualCount: 31; lifeStage: adult; recordedBy: J. Hájek & J. Šumpich; institutionCode: NMPC; collectionCode: Insects

## Description

A large, mainly yellow to orange *Limbodessus*: length of body 2.6–3.0 mm (N=20); with slight habitus disruption between pronotum and elytron Figs 13, 14, 15, 18; dorso ventrally rather flattened Figs 9, 11; cervical line present (sometimes faint or partly dissolved into punctures in females), distinct pronotal and elytral plicae present Figs 13, 14, 15, 18, 41; fully developed flight wings present. Metacoxa and metaventrite with few punctures only Fig. 27. Ventral side without modified elytral epipleuron.

Sexes dimorphic, see below.

**Male:** Antenna filiform Figs 13, 15, 18, 21, 24. Dorsal surface with coarse punctures but otherwise with shiny surface Figs 13, 19. Ventrally shiny, dark orange to dark brown Fig. 21. Median lobe of aedeagus as in Figs 29, 30 with tiny setae on tip, paramere as in Fig. 34.

**Female:** Antenna moniliform Fig. 14. Dorsal surface with very fine microreticulation between surface punctures and surfaces therefore appearing matt Figs 12, 14. Ventrally matt, darker, metacoxa and metaventricle blackish Fig. 20.



Figure 41.

*Limbodessus baliem*, female, habitus. Animated  $\mu$ CT scan. Length of beetle: 2.8 mm.

### Etymology

The species is named after the type locality, the Baliem River Valley. The name is a noun in the nominative singular standing in apposition.

### Distribution

Indonesian New Guinea, known only from the Baliem Valley which also contains the type locality Wamena (Fig. 23). The name is a noun in the nominative singular standing in apposition.

### Ecology

The species was collected from small bodies of stagnant water around Wamena in the vast, flat valley floor of the Baliem Valley Figs 42, 43. This species seems to prefer sunny or partly shaded habitat with sand or clay on the bottom and not too strongly vegetated. Other species it was associated with were: *Exocelina baliem* Shaverdo, Hendrich & Balke, 2013, *Hydrovatus enigmaticus* Biström, 1997, *Hyphydrus dani* Biström, Balke & Hendrich, 1993, *Hydaticus okalehubyi* Balke & Hendrich, 1992, *H. rivanolis* Wewalka, 1979, and *Rhantus dani* Balke, 2001 (Shaverdo et al. (2013), Biström et al. (1993), Balke and Hendrich (1992), Balke (2001).



Figure 42.  
Habitat of *Limnodessus baliem* near Wamena.



Figure 43.  
Habitat of *Limnodessus baliem* near Jiwika.

**Conservation**

Aquatic habitats in the Baliem Valley in general are threatened by eutrophication from domestic animals, fish farming to some degree and increasing intensity of gardening throughout the valley.

## Online resources

Higher resolution digital images,  $\mu$ CT data as well as sequence data have been deposited in public databases. The Species-ID species page is a versioned wiki site and can be enhanced through community contributions Hendrich and Balke (2011).

**Species-ID species page:** [http://species-id.net/wiki/Limbodessus\\_baliem](http://species-id.net/wiki/Limbodessus_baliem)

**DNA sequences:** Partial 3' *cox1* sequence deposited at: LN884313–LN884314.

**$\mu$ CT data:** Deposited in [Morphosource](#).

## Identification keys

<b>Key to New Guinea <i>Limbodessus</i></b>		
Based on Watts and Leijds (2005).		
1	Base of elytral epipleuron with raised transverse carina delineating a basal pit, see Fig. 36 (widespread in the Australian region; ventral aspect as in Fig. 28, with metaventrite and metacoxa rather smooth; male genital as in Figs 32, 33, 35).	<i>Limbodessus compactus</i> (Clark, 1862)
–	Base of elytral epipleuron without raised transverse carina delineating a basal pit as in Fig. 38	2
2	Elytral plica present; beetle mainly yellowish; microendemic of the Baliem Valley of Papua	<i>Limbodessus baliem</i> <b>sp.n.</b>
–	Elytral plica absent; beetle dark brown to blackish; only known from high altitude grassland of the range north of Sugapa, Papua	<i>Limbodessus alexanderi</i> <b>sp.n.</b>

## Analysis

We obtained sequence data for the 3' end of the mitochondrial cytochrome oxidase I gene and analyzed these in the context of our database of Australasian diving beetles Hendrich et al. (2010) by constructing neighbour joining trees (HKY model as implemented in Geneious 8.1.6 software) as well as by clustering the data in SpeciesIdentifier module of Taxon DNA software (v1.6.; Meier et al. (2006). Both species described here differ from known extant taxa (Fig. 40). We queried the two new species against the other *Limbodessus* in our dataset and found uncorrected *p*-distances for *L. baliem* sp.n. of 6.68–12.23% (*L. gemellus* (Clark 1862)/ *L. praelargus* (Lea 1899)) and for *L. alexanderi* sp.n. of 8.83–13.96% (*L. inornatus* (Sharp 1882)/ *L. cheesmanae* (Balfour-Browne 1939)).

As another result of our analysis, we propose new combinations for the following three species:

*Limbodessus deflectus* (Ordish 1966), **new combination**

- *Liodessus deflectus* Ordish 1966: 229; Nilsson 2015: 115.

*Limbodessus leverii* (Balfour-Browne 1944), **new combination**

- *Bidessus leverii* Balfour-Browne 1944: 98; Nilsson 2015: 126.

*Limbodessus plicatus* (Sharp 1882), **new combination**

- *Bidessus plicatus* Sharp 1882: 360

- *Liodessus plicatus* (Sharp 1882): Nilsson 2015: 116.

## Discussion

The merits and possible applications of (X-ray) Micro-CT for arthropod systematics have recently been discussed by Friedrich et al. (2013) and Simonsen and Kitching (2014). The strength of this approach lies in the non-destructive examination. Data acquisition is relatively straight forward and data cover a wide range of information including that of internal structures.

In our examples the CT examination provided comprehensive information on overall morphology; general proportions as well as external surface details could be assessed. Regarding surface details, Micro-CT complements conventional ways of examinations, such as light microscopic photography and SEM. Both methods are superior concerning structural surface details. In addition, light microscopy provides true colours. In the specimens examined in the present study, the X-ray absorption was relatively poor, this might be the reason for some deficiencies in structural resolution. In places, internal structures cannot be discerned. For example, the elytra could not be separated from the underlying material.

The external surface renderings appear to be a very useful approach for distributing the CT data. By interactive manipulation, these polygonal mesh surfaces enable intuitive understanding of overall proportions of the specimens. They also contain an enormous amount of geometric information that could easily be extracted and used for e.g., morphometric studies. The 3D PDF models make the entire external morphological information available to “readers”, who might use this information in further comparative studies.

In order to also provide information on specimen coloration and surface microsculpture, we provided digital photographs of different aspects of the beetles using a high throughput, automated imaging approach that could easily be implemented by a technician. The goal

was to optimize image quality and time spent on each image so that taxonomically relevant or potentially interesting details can be recognized easily. Digital imaging does, in our opinion, still outperform CT examination for routine work and in terms of documentation of taxonomically relevant structures in the beetles studied here.

## Acknowledgements

We thank Alexander Riedel for sharing specimens. Fieldwork of the Cendrawasih University UNCEN was organized by Dr S. Surbakti which is greatly appreciated here. Dr Matthew van Dam kindly helped with the linguistic check of the paper which is highly appreciated here.

## Author contributions

MB and LH designed the study and conducted the taxonomic work and fieldwork. MB took high resolution digital images and SEMs and conducted the molecular work. BR conducted the uCT work and wrote the corresponding part of the paper. ELW organized and conducted fieldwork. KN performed illustration work.

## References

- Balfour-Browne J (1939) On the aquatic Coleoptera of the New Hebrides and Banks Islands. Dytiscidae, Gyrinidae, and Palpicornia. The Annals and Magazine of Natural History 3 (11): 459-479. DOI: [10.1080/03745481.1939.9723627](https://doi.org/10.1080/03745481.1939.9723627)
- Balfour-Browne J (1944) New and interesting Dytiscidae (Coleoptera) from Fiji. Proceedings of the Royal Entomological Society of London (B) 13: 97-100. DOI: [10.1111/1/j.1365-3113.1944.tb00796.x](https://doi.org/10.1111/1/j.1365-3113.1944.tb00796.x)
- Balke M (2001) Biogeography and classification of New Guinean Colymbetini (Coleoptera: Dytiscidae: Colymbetinae). Invertebrate Systematics 15 (2): 259-275. DOI: [10.1071/IT98008](https://doi.org/10.1071/IT98008)
- Balke M, Hendrich L (1992) Ein neuer Schwimmkafer der Gattung *Hydaticus* Leach (Coleoptera: Dytiscidae) aus dem Hochland von West Papua. Mitteilungen der Schweizerischen Entomologischen Gesellschaft 65 (3): 297-302.
- Balke M, Ribera I (2004) Jumping across Wallace's line: *Allodessus* Guignot and *Limbodessus* Guignot revisited (Coleoptera: Dytiscidae, Bidessini) based on molecular-phylogenetic and morphological data. Australian Journal of Entomology 43 (2): 114-128. DOI: [10.1111/j.1440-6055.2004.00415.x](https://doi.org/10.1111/j.1440-6055.2004.00415.x)
- Biström O (1988) Generic review of the Bidessini (Coleoptera, Dytiscidae). Acta Zoologica Fennica 184: 1-41.
- Biström O, Balke M, Hendrich L (1993) A new species of *Hyphydrus* Illiger 1802 (Coleoptera Dytiscidae) from West New Guinea, and notes on other species of the genus. Tropical Zoology 6 (2): 287-298. DOI: [10.1080/03946975.1993.10539229](https://doi.org/10.1080/03946975.1993.10539229)

- Clark H (1862) Catalogue of the Dytiscidae and Gyrinidae of Australasia, with descriptions of new species. The Journal of Entomology. Descriptive and Geographical 1: 399-421.
- Friedrich F, Matsumura Y, Pohl H, Bai M, Hörnschemeyer T, Beutel R (2013) Insect morphology in the age of phylogenomics: innovative techniques and its future role in systematics. Entomological Science 17 (1): 1-24. DOI: [10.1111/ens.12053](https://doi.org/10.1111/ens.12053)
- Gschwendtner L (1931) Neue Dytiscidae aus Formosa. Entomologischer Anzeiger 11: 21.
- Guignot F (1939) Contribution à l'étude des Bidessus. Bulletin de la Société d'Étude des Sciences Naturelles de Vaucluse Bulletin de la Société d'Étude des Sciences Naturelles de Vaucluse 10 (4): 51-61.
- Hendrich L, Balke M (2011) A simultaneous journal / wiki publication and dissemination of a new species description: *Neobidessodes darwiniensis* sp. n. from northern Australia (Coleoptera, Dytiscidae, Bidessini). ZooKeys 79: 11-20. DOI: [10.3897/zookeys.79.803](https://doi.org/10.3897/zookeys.79.803)
- Hendrich L, Pons J, Ribera I, Balke M (2010) Mitochondrial Cox1 Sequence Data Reliably Uncover Patterns of Insect Diversity But Suffer from High Lineage-Idiosyncratic Error Rates. PLoS ONE 5 (12): e14448. DOI: [10.1371/journal.pone.0014448](https://doi.org/10.1371/journal.pone.0014448)
- Larson DJ (1994) Boongurrus rivulus, a New Genus and Species of Water Beetle (Coleoptera: Dytiscidae: Bidessini) from Northern Queensland, Australia. Australian Journal of Entomology 33 (3): 217-221. DOI: [10.1111/j.1440-6055.1994.tb01222.x](https://doi.org/10.1111/j.1440-6055.1994.tb01222.x)
- Lea AM (1899) Descriptions of new species of Australian Coleoptera. Part V. Proceedings of the Linnean Society of New South Wales 23: 521-645.
- Limaye A (2012) Drishti: a volume exploration and presentation tool. Proc. SPIE 8506, Developments in X-Ray Tomography VIII, 85060X October 17, 2012: 1. DOI: [10.1117/1.2.935640](https://doi.org/10.1117/1.2.935640)
- Meier R, Shiyang K, Vaidya G, Ng P (2006) DNA Barcoding and Taxonomy in Diptera: A Tale of High Intraspecific Variability and Low Identification Success. Systematic Biology 55 (5): 715-728. DOI: [10.1080/10635150600969864](https://doi.org/10.1080/10635150600969864)
- Nilsson AN (2001) Dytiscidae (Coleoptera). World catalogue of insects 3: 1-395.
- Nilsson AN (2015) A World Catalogue of the Family Dytiscidae, or the Diving Beetles (Coleoptera, Adepaga). Version 1.1.2015 1: 1-298. URL: <http://www2.emg.umu.se/projects/biginst/andersn/>
- Ordish RG (1966) A systematic revision of the New Zealand water beetles (Coleoptera: Dytiscidae). Records of the Dominion Museum Wellington 5: 217-264.
- Riedel A, Sagata K, Suhardjono YR, Tänzler R, Balke M (2013a) Integrative taxonomy on the fast track - towards more sustainability in biodiversity research. Frontiers in Zoology 10 (1): 15. DOI: [10.1186/1742-9994-10-15](https://doi.org/10.1186/1742-9994-10-15)
- Riedel A, Sagata K, Surbakti S, Tänzler R, Balke M (2013b) One hundred and one new species of *Trigonopterus* weevils from New Guinea. ZooKeys 280: 1-150. DOI: [10.3897/zookeys.280.3906](https://doi.org/10.3897/zookeys.280.3906)
- Ruthensteiner B, Heß M (2008) Embedding 3D models of biological specimens in PDF publications. Microscopy research and technique 71 (11): 778-786. DOI: [10.1002/jemt.20618](https://doi.org/10.1002/jemt.20618)
- Sharp D (1882) On aquatic carnivorous Coleoptera or Dytiscidae. Scientific Transactions of the Royal Dublin Society 2 (2): 179-1003.



- Shaverdo H, Hendrich L, Balke M (2013) *Exocelina baliem* sp. n., the only known pond species of New Guinea *Exocelina* Broun, 1886 (Coleoptera, Dytiscidae, Copelatinae). ZooKeys 304: 83-99. DOI: [10.3897/zookeys.304.4852](https://doi.org/10.3897/zookeys.304.4852)
- Simonsen T, Kitching I (2014) Virtual dissections through micro-CT scanning: a method for non-destructive genitalia 'dissections' of valuable Lepidoptera material. Syst Entomol 39 (3): 606-618. DOI: [10.1111/syen.12067](https://doi.org/10.1111/syen.12067)
- Watts CHS, Humphreys WF (1999) Three new genera and five new species of Dytiscidae (Coleoptera) from underground waters in Australia. Records of the South Australian Museum 32: 121-142.
- Watts CHS, Leijs R (2005) Review of the epigeal species of Australian *Limbodessus* Guignot (Insecta: Coleoptera: Dytiscidae). Transactions of the Royal Society of South Australia 129 (1): 1-13.

## Supplementary material

### Suppl. material 1: *Limbodessus* Papua sequences fasta file

**Authors:** Michael balke

**Data type:** DNA sequences

**Brief description:** A fasta file containing the cox1 sequences newly generated for this project.

**Filename:** Limbodessus Papua sequences fasta file.pdf - [Download file](#) (20.77 kb)