

Rehabilitation following meniscal repair: a systematic review

Robert C Spang III,¹ Michael C Nasr,² Amin Mohamadi,² Joseph P DeAngelis,¹ Ara Nazarian,^{2,3} Arun J Ramappa¹

To cite: Spang III RC, Nasr MC, Mohamadi A, *et al.* Rehabilitation following meniscal repair: a systematic review. *BMJ Open Sport & Exercise Medicine* 2018;**4**:e000212. doi:10.1136/bmjsem-2016-000212

JPD, AN and AJR contributed equally.

JPD, AN and AJR are joint senior authors.

Accepted 9 February 2018

ABSTRACT

Objective To review existing biomechanical and clinical evidence regarding postoperative weight-bearing and range of motion restrictions for patients following meniscal repair surgery.

Methods and data sources Following the Preferred Reporting Items for Systematic Reviews and Meta-Analyses guideline, we searched MEDLINE using following search strategy: (((("Weight-Bearing/physiology"[Mesh]) OR "Range of Motion, Articular"[Mesh]) OR "Rehabilitation"[Mesh])) AND ("Menisci, Tibial"[Mesh]). Additional articles were derived from previous reviews. Eligible studies were published in English and reported a rehabilitation protocol following meniscal repair on human. We summarised rehabilitation protocols and patients' outcome among original studies.

Results Seventeen clinical studies were included in this systematic review. There was wide variation in rehabilitation protocols among clinical studies. Biomechanical evidence from small cadaveric studies suggests that higher degrees of knee flexion and weight-bearing may be safe following meniscal repair and may not compromise the repair. An accelerated protocol with immediate weight-bearing at tolerance and early motion to non-weight-bearing with immobilising up to 6 weeks postoperatively is reported. Accelerated rehabilitation protocols are not associated with higher failure rates following meniscal repair.

Conclusions There is a lack of consensus regarding the optimal postoperative protocol following meniscal repair. Small clinical studies support rehabilitation protocols that allow early motion. Additional studies are needed to better clarify the interplay between tear type, repair method and optimal rehabilitation protocol.

INTRODUCTION

The menisci reduce stress by increasing the contact area between the femur and tibia. They buffer against axial, rotational and shearing forces about the knee during motion.¹ The loss of meniscal tissue localises tibiofemoral contact and leads to progressive arthrosis^{2,3} and functional decline in the long term.⁴ To prevent these degenerative changes, meniscal repair has become more common.⁵ Kim *et al*⁶ documented a 25% increase in medial and lateral meniscal repairs between 1996 and 2006. Even though the majority of surgeries

What is already known?

- ▶ The menisci reduce stress by increasing the contact area between the femur and tibia.
- ▶ Meniscal repair is becoming a more appealing treatment for meniscal injuries.
- ▶ There is a wide variation between postoperative rehabilitation protocols following meniscal repair.

What are the new findings?

- ▶ There is no consensus regarding postoperative rehabilitation protocol for meniscal repair.
- ▶ The quality of existing evidence is low.
- ▶ An accelerated rehabilitation protocol may be safely implemented for appropriate patients.
- ▶ Further studies are needed to determine an optimal rehabilitation protocol.

on the meniscus remain meniscectomies, Abrams *et al*⁷ found that between 2005 and 2011 more isolated meniscal repairs were performed in the USA without an increase in the number of meniscectomies. Meniscal repairs may be performed more frequently because there has been a significant advance in surgical techniques and repair devices (figure 1A–E). Historically, the gold standard for meniscal repair has been the inside-out technique (figure 1C,D). Long flexible needles are used to pass sutures through the tissue under arthroscopic guidance.⁸ The sutures are then retrieved using a separate incision and are tied over the joint capsule. This technique may place neurovascular structures at risk and requires an additional incision.⁹ To avoid the morbidity associated with an inside-out repair, 'all-inside' arthroscopic techniques have been developed (figure 1A,B). These include anchor-based repairs and suture-based repairs. The most popular of these designs employ pre-tied sutures between non-absorbable anchors. The anchors are deployed when an introducer is passed through the meniscal tear and the joint capsule (figure 1E,F). In a study of porcine meniscal repairs,¹⁰ the inside-out



¹Carl J. Shapiro Department of Orthopaedic Surgery, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, Massachusetts, USA

²Center for Advanced Orthopaedic Studies, Carl J. Shapiro Department of Orthopaedic Surgery, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, Massachusetts, USA

³Department of Orthopaedic Surgery, Yerevan State Medical University, Yerevan, Armenia

Correspondence to

Dr Arun J Ramappa;
aramappa@bidmc.harvard.edu




Figure 1 (A,B) All-inside suture-based repair, (C,D) inside-out suture repair and (E, F) anchor-based repair techniques.

technique was significantly stronger compared with recent all-inside repair devices. A study on fresh-frozen human menisci¹¹ showed no difference. Although potentially less invasive than the inside-out technique, all-inside repairs can result in neurovascular injury, irritation from anchors and implant failure.¹²

Several factors may influence meniscal healing. The most important may be the meniscal blood supply. Scapinelli¹³ in 1968 and Arnoczky and Warren¹⁴ in 1982 described the limited peripheral blood supply to the outer one-third to one-quarter of the meniscus. From this finding, peripheral meniscal tears (ie, tears in the

'red-red' zone) are felt to have better healing potential (figure 2A).

The timing and type of meniscal tear may also impact healing. Acute, traumatic tears tend to have higher healing rates than chronic, atraumatic tears.¹⁵ Longitudinal tears are more amenable to repair due to their vertical orientation (figure 2B), whereas radial tears extending to the central relatively avascular 'white-white' zone are more challenging. Some surgeons have reported success when repairing bucket-handle tears extending to the white-white zone.¹⁶

Age is another topic for consideration. Preserving meniscal tissue is particularly important for the long-term health of young athletes, and younger patients may have a higher healing potential. In one study of 26 patients aged 17 years or younger, none required a repeat surgery at an average of 5 years of follow-up.¹⁷ In another report on two very young cases, meniscal repair for traumatic tears followed by limited weight-bearing rehabilitation resulted in a positive outcome.¹⁸ When age is not a factor, a meta-analysis investigating outcomes at least 5 years after meniscal repair showed a pooled failure rate of 23.1%.¹⁹

As biomechanical factors, postoperative range of motion (ROM) and weight-bearing status can impact meniscal healing after repair. The interplay between tear type and knee biomechanics can help define the most appropriate postoperative plan.

Restricting a patient's postoperative ROM intends to limit the risk of re-tear. Cadaveric studies have shown that femorotibial contact pressures increase with knee flexion.²⁰ If the ROM is restricted, the meniscal repair may be protected from increased mechanical stress.²¹ However, Richards *et al*²² investigated the effects of compressive loads in porcine longitudinal lateral meniscus repairs and found that weight-bearing reduced the meniscus and stabilised the repair. The highest compressive force occurred at full extension and the




Figure 2 (A) Outer red zones receive blood supply; (B) longitudinal tears have a higher likelihood of being vascularized.




Figure 3 (A) The pressure transducer 'P' was placed in the lateral meniscal cut and the knee was cycled into flexion and extension. (B) Intrameniscal pressures were reflected in neutral, internal, and external rotation.

lowest was at 90° of flexion. Beyond 100° of flexion, it increased steadily (figure 3). Higher pressures were seen with internal rotation of the tibia, suggesting torsional forces may be different than axial loads. Conversely, for radial tears, axial loading might displace, rather than reduce, the injury.²² Of note, cadaveric investigations are limited in their ability to recreate the biomechanics of the knee in vivo and are not able to predict how specific rehabilitation protocols impact a meniscal repair. For this reason, rehabilitation following a meniscal repair is particularly conservative in an effort to protect the repaired meniscus.

Postoperative rehabilitation aims to foster healing after meniscal repair and facilitate the patient's return to full function. Generally, these programmes are initially focused on protecting the repair while regaining ROM and gradually introducing progressive strengthening en-

route to a return to preinjury activity level. At present, there is a paucity of evidence to support one best practice and there is a high degree of variability among postoperative rehabilitation programmes. Considering the increased frequency and evolution of meniscal repairs, this review intends to summarise the best-available evidence and practices regarding the postoperative care and rehabilitation of patients undergoing a meniscal repair.

METHODS

Search strategy and data sources

Following the Preferred Reporting Items for Systematic Reviews and Meta-Analyses guideline, on 15 June 2017 we conducted an electronic search on MEDLINE with the following search strategy: (((("Weight-Bearing/physiology"[Mesh]) OR "Range of Motion, Articular"[Mesh]) OR "Rehabilitation"[Mesh])) AND ("Menisci, Tibial"[Mesh]). Additional sources included references of previous reviews.^{19 23 24}

A total of 453 studies were screened for eligibility. Only studies published in the English language in peer-reviewed journals were considered. Review papers, commentaries and studies on rehabilitation protocol following meniscectomy were excluded. Finally, 17 studies were included in the review (figure 4).

Bibliographic data, patients' characteristics, rehabilitation protocol and clinical outcome were recorded. Patient's outcome included meniscal healing, return to activities and clinical assessments based on the original studies. We did not confirm collected data by authors. The level of evidence for original studies is reported for each study.²⁵

Cochrane tool for evaluating risk of bias was used for assessing the methodological quality of the included studies.

RESULTS

Seventeen clinical studies including 798 patients were reviewed in this systematic review. There was wide variation in methodological quality of clinical studies. The majority of studies had considerable risk of bias (table 1).

A restricted rehabilitation protocol was used for 438 patients. An accelerated protocol with immediate weight-bearing at tolerance was used in 360 patients. Three studies compared restricted and accelerated protocol, which did not show any significant difference in complication rate or functional assessment (table 2).

Although a meta-analysis was not possible, it seems accelerated rehabilitation protocols are not associated with higher failure rates following meniscal repair.

DISCUSSION

Many rehabilitation programmes propose avoiding weight-bearing forces as an important goal in the immediate postoperative period to protect the repair from high compressive and shear forces. An MRI study of




Figure 4 Flow chart of the systematic review.

Table 1 Assessment of risk of bias using Cochrane tool

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias)	Blinding of outcome assessment (detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)
Morgan and Casscells ⁵⁴	–	–	–	–	+	+
Morgan <i>et al</i> ³¹	–	–	–	–	+	+
Barber ³⁸	–	–	–	–	+	+
Horibe <i>et al</i> ⁵⁵	–	–	–	–	?	?
Fritz <i>et al</i> ²⁹	–	–	–	–	+	+
Mariani <i>et al</i> ⁴⁰	–	–	–	–	+	+
Shelbourne <i>et al</i> ⁴¹	–	–	–	–	+	+
Barber and Click ⁵⁰	–	–	–	–	?	+
Mintzer <i>et al</i> ¹⁷	–	–	–	–	+	+
Bloome <i>et al</i> ^{†18}	–	–	–	–	+	+
Noyes and Barber-Westin ⁴⁶	–	–	–	–	+	+
O'Shea and Shelbourne ⁴⁷	–	–	–	–	?	+
Kocabey <i>et al</i> ⁴²	–	–	–	–	+	+
Bryant <i>et al</i> ⁴⁸	+ (Randomisation for method of repair)	+	?	?	?	+
Haklar <i>et al</i> ⁴⁵	–	–	–	–	+	+
Logan <i>et al</i> ⁴⁹	–	–	–	–	+	+
Lind <i>et al</i> ³⁵	+	+	–	–	?	+

–, high risk of bias; +, low risk of bias; ?, unclear risk of bias.

*Athletes under 17 years old.

†Very young children.

weight-bearing and non-weight-bearing knees found that the relative tibiofemoral movements of the loaded knee were similar to those in the unloaded knee. However, the medial femoral condyle moves approximately 4 mm anterior when the knee bends from full extension to 10° of flexion while bearing weight. In the unloaded knee, the position of the medial femoral condyle did not change from extension to flexion. Laterally, the femoral condyle rolls forward 13 mm from 110° to 60° of flexion and 1 mm from 60° to 0° in the unloaded knee.²⁶ In isolation, this pattern of motion suggests that non-weight-bearing knee flexion would be safe to 110° for medial meniscal repairs and to 60° for lateral meniscal repairs. However, this finding has not been validated clinically.

Becker *et al*²⁰ investigated changes in the menisiofemoral contact pressure after meniscal repair. Knees were loaded to approximately 50% of body weight, and menisiofemoral contact pressure was measured (Tekscan, Boston, Massachusetts, USA) while cycling the knee from extension to 90° of flexion. They found that menisiofemoral pressures increased in both compartments as the knee flexed and that meniscal repair had no impact. Their study did not investigate the effect of this pressure difference on the meniscal repair, nor did they include

torsional forces or higher impact loading. Ganley *et al*²⁷ sought to further investigate knee flexion and loading on meniscal healing in a cadaveric model. They produced full-thickness posteromedial meniscal tears in cadaveric knees and imbedded metal markers into the tear following repair. Using CT scans, the marker position was assessed at 30°, 60° and 90° of knee flexion after loading of 100 lbs to simulate partial weight-bearing. They determined that neither flexion angle, loading nor suture had a significant impact (figures 5 and 6). In this way, accelerated rehabilitation programme with partial weight-bearing may be appropriate. Torsional forces, higher degrees of flexion and loads larger than 100 lbs were not assessed.

Lin *et al*²⁸ sought to assess the effect of postoperative ROM following meniscal repair using a cadaveric model. They created a 2.5 cm posteromedial meniscal tear and repaired it with inside-out vertical mattress sutures (figure 7A). They measured the displacement at high degrees of flexion (90°, 110° and 135°) when loaded (figure 7B). Specimens were subjected to simulated open-chain flexion and extension with a load of 29 N applied to the hamstrings and 150 N to the quadriceps, exceeding the normal joint reactive force encountered

Table 2 Previously published rehabilitation protocols

Paper	No of patients (meniscal repairs)	WB restrictions	ROM restrictions	Other restrictions	Outcome	Level of evidence
Morgan and Casscells ⁵⁴	67 (70)	Immediate WBAT in extension 4 weeks	Full extension for 4 weeks	Pivoting not until 4 months	Excellent results in 69 (98%) repairs, 1 patient had second tear and 2 patients had surgical complications.	(Retrospective) case series; level IV evidence
Morgan <i>et al</i> ³¹	353 repairs, 74 had second-look arthroscopy	Immediate WBAT in full extension 4 weeks	Active 0°–60° after 1 week	No pivoting for 6 months	Asymptomatic healing occurred in 84% of patients at second-look arthroscopy.	Non-randomised cohort/ follow-up study; level III evidence
Barber ³⁸	95 (98) 56 (58) in standard protocol 39 (40) in accelerated protocol	Standard protocol: NWB for 12 weeks Accelerated protocol: Immediate WBAT	Standard protocol: immobilisation at flexion for 6 weeks Accelerated protocol: immediate unlimited ROM	Standard protocol: No pivoting for 6 months Accelerated protocol: Pivoting sports as soon as the patient desired	Standard protocol: 11/20 failure at second-look arthroscopy. Accelerated protocol: 4/10 failure at second-look arthroscopy.	Non-randomised cohort/ follow-up study; level III evidence
Horibe <i>et al</i> ⁵⁵	122 (132)	WBAT after 5–6 weeks	Immobilisation for 1–2 weeks	Vigorous not for 4–6 months	97 menisci (73%) had complete healing; 21 of which had new tear at second-look arthroscopy.	Non-randomised cohort/ follow-up study; level III evidence
Fritz <i>et al</i> ²⁹	1	Immediate WBAT with two crutches	Brace locked in extension for 6 weeks, 6–8 weeks unlocked for gait training ROM limited to 0°–90°	Return to full activity approximately in 1 year	Full ROM and no effusion with 4+/5 quadriceps strength on clinical examination, no progression of degenerative changes on X-ray.	Case report; level V evidence
Mariani <i>et al</i> ⁴⁰	22	Immediate WBAT	Immobilisation with brace locked at 0° during ambulation for 1 month, passive 0°–90° ROM from day 2 to 2 weeks	Progressive resistance exercise from 4 weeks, running and biking after 2 months; full return to sport after 6 months	17 (77%) patients showed 'good clinical' results. 3 (14%) showed signs of meniscal re-tear on MRI one of which had second surgery	Non-randomised cohort/ follow-up study; level III evidence
Shelbourne <i>et al</i> ⁴¹	61 17 in conventional protocol 39 in accelerated protocol	Conventional protocol: NWB until 6 weeks. Accelerated protocol: immediate WBAT	Conventional protocol: limited ROM until 6 weeks Accelerated protocol: immediate ROMAT	Conventional protocol: restricted activities Accelerated protocol: Bike and swim as tolerated 2–4 weeks, strength work	No significant difference between two protocols in Lysholm score, Noyes questionnaire score or self-evaluation score.	Non-randomised cohort/ follow-up study; level III evidence

Continued

Table 2 Continued

Paper	No of patients (meniscal repairs)	WB restrictions	ROM restrictions	Other restrictions	Outcome	Level of evidence
Barber and Click ⁵⁰	63 (65)	Immediate WBAT	Immediate unrestricted ROM; no braces were used	After adequate motion (0° to 120°), good strength and no effusion are achieved, return to all activities—including pivoting sports—is allowed	Second-look arthroscopy in 17 (26%) showed repair failure in 7 patients.	Non-randomised cohort/follow-up study; level III evidence
Mintzer <i>et al</i> ¹⁷	26 (29)	5 patients: NWB in a knee immobiliser for 4 weeks 21 patients: Allowed WBAT in a knee immobiliser for 4 weeks	Immobilisation for 4 weeks	NR	24 patients returned to their previous level of sports activity. The remaining two patients cited reasons other than surgery for limiting their sports activity	Non-randomised cohort/follow-up study; level III evidence
Bloome <i>et al</i> ¹⁸	2	Case 1: Partial WB in cast for 4 weeks. Case 2: Partial WB using crutches 2 weeks, WBAT 2–6 weeks	Case 1: Long-leg splint/cast for 4 weeks at full extension, then removable posterior splint for 2 weeks to use when ambulating Case 2: Long-leg cylinder cast until 6 weeks, then immobiliser for walking	NR	Case 1: Full return to activities at 7-month follow-up. Case 2: Normal gait at 3 months and full activities at 6 months.	Case report; level V evidence
Noyes and Barber-Westin ⁴⁶	29 (30)	Partial WB for 4 or 6 weeks	ROM progressed to 135° over 6 weeks Restriction of squatting or deep flexion beyond 125° for 4 months	No vigorous activities for 6 months	26 (87%) were asymptomatic at follow-up. Three repairs failed to heal, requiring partial meniscectomy, and one knee with tibiofemoral symptoms related to the repair was treated conservatively.	Non-randomised cohort/follow-up study; level III evidence
O'Shea and Shelbourne ⁴⁷	52 (55)	WBAT postoperative day 3	Immediate ROMAT	NR	At second-look arthroscopy showed 30 menisci (55%) appeared healed, 19 menisci (34%) appeared partially healed, and 6 menisci (11%) showed no healing.	Non-randomised cohort/follow-up study; level III evidence

Continued

Table 2 Continued

Paper	No of patients (meniscal repairs)	WB restrictions	ROM restrictions	Other restrictions	Outcome	Level of evidence
Kocabey <i>et al</i> ⁴²	52 (55)	Immediate WBAT	ROM 0°–125°	Return 3–5 months depending on tear type	Excellent results in all with combined ACL–meniscus repair ³² and in 96% (22 cases) with isolated meniscus repair.	Retrospective case series; level IV evidence
Bryant <i>et al</i> ⁴⁸	100	Protected WB for 3 weeks, then WBAT	Locked in extension for 3 weeks, then full ROM	No squatting, pivoting and twisting for a minimum of 6 months	Of 88 patients at follow-up, 22 (25%) patients had failed meniscal repairs.	Randomised controlled clinical trial; level I evidence
Haklar <i>et al</i> ⁴⁵	5	NWB 6–8 weeks	No squat beyond 120°	No running until 4 months, then return to normal activities	MRI showed that all five patients had fully healed meniscus.	Non-randomised cohort/follow-up study; level III evidence
Logan <i>et al</i> ⁴⁹	42 (45)	Protected WB for 6 weeks	90° flexion by 6 weeks	NR	34 (81%) patients returned to their main sport. There were 11 (24%) failures in meniscal repair.	Case series; level IV evidence
Lind <i>et al</i> ⁸⁵	60 32 in free protocol 28 restricted protocol	Free protocol: 2 weeks TDWB. Restricted protocol: TDWB for 6 weeks	Free protocol: ROM 0°–90°, no brace, then return to normal activities Restricted protocol: 6 weeks hinged brace, gradual increase ROM to 90°	Free protocol: Running at 8 weeks contact sports at 4 months Restricted protocol: 12 weeks, contact sports 6 months	Second-look arthroscopy showed failure of healing in 9 (28%) patients in free and 10 (36%) patients in the restricted rehabilitation groups. No difference in failure rate and no difference in functional outcome at 1–2 years.	Randomised controlled clinical trial; level I evidence

*Athletes 17 years old or younger.

†Very young children.

NR, not reported; NWB, non-weight-bearing; ROM, range of motion; ROMAT, range of motion as tolerated; TDWB, touch-down weight-bearing; WB, weight-bearing; WBAT, weight-bearing as tolerated.




Figure 5 Transverse section of a cadaver knee under 100 lbs of load at 60°. (A) Longitudinal and (B) transverse measurements are depicted between the markers.

during active knee flexion.²⁸ In accordance with the findings of Richards *et al*²² and Ganley *et al*,²⁷ they found that neither the meniscal tear nor the meniscal repair demonstrated significant gapping. Rather they compressed in the transverse plane when flexed from 90° to 135° while subjected to physiologic loads. They conclude that 'non-restrictive un-resisted open chain ROM protocols do not place undue stress on meniscal repairs'.

Early weight-bearing might enhance the mechanical environment promoting healing and allowing earlier functional recovery and return to sport.²² It has been shown that early weight-bearing as tolerated and limited ROM resulted in acceptable outcomes (ie, Lysholm score of 71.5) at 17 month follow-up.²⁹ While Becker's cadaveric findings may alleviate concern over iatrogenic cartilage damage from implants, the hypothetical danger of increased meniscofemoral pressures with knee flexion after meniscal repair has not been shown to impact clinical outcomes.²⁰

Some authors recommend immobilisation in full extension, reportedly due to the observation that peripheral posterior horn tears move away from the capsule in flexion and reduce in extension.^{18 30-32} This

recommendation stems from direct viewing of peripheral posterior horn tears using a 70° arthroscope and observing the reduction of these tears during passive knee extension. However, this observation and the subsequent practice of immobilising meniscal repairs in extension has not been shown to be beneficial clinically and may not be relevant to other tear types. Some investigators recommend immobilisation in various degrees of flexion,³²⁻³⁴ and others still advocate for limited early motion. Despite more aggressive protocols allowing for free ROM immediately postoperatively, 90° of flexion appears to be a comfortable restriction for surgeons.³⁵ Up to 85% of the load travels through the menisci with the knee in 90° of flexion, while less (50%) of the load passes through the meniscus in extension.³⁶

Additionally, meniscal dynamics using MRI three-dimensional reconstructions show that during knee flexion, the posterior excursion of the medial meniscus is 5.1 mm and the lateral meniscus is 11.2 mm.³⁷ However, a more recent study by Lin suggests that higher degrees of flexion may be safe.²⁸ There is no clinical evidence that limiting weight-bearing and/or knee flexion improves healing rates. Long-term outcome studies are lacking.




Figure 6 Transverse section of a cadaver knee without load at 60°. (A) Longitudinal and (B) transverse) measurements are depicted between the markers.




Figure 7 (A) Schematic of roentgen stereophotogrammetric analysis bead pair placement in relation to tear. Distances measured by vectors: a—absolute, b—transverse and c—vertical. (B) Changes in separation for each vector. Positive values indicate widening. Negative values indicate compression. MCL, medial collateral ligament region of posterior horn of medial meniscus; mid post, middle of posterior horn; post root, posterior root area of medial meniscus.

Accelerated rehabilitation protocols

A number of investigators have advocated for accelerated rehabilitation protocols (figure 8).^{38–41} In a prospective randomised trial, Lind *et al*²⁸ compared the impact of a ‘free rehabilitation’ regimen versus ‘restricted rehabilitation’.³⁵ Sixty patients underwent isolated repair of a vertical meniscal lesion using an all-inside technique. They were randomised by rehabilitation regimens. The ‘free’ group was allowed to range the knee 0°–90° immediately while maintaining the knee in touch-down weight-bearing for 2 weeks, and weight-bearing as tolerated thereafter. They were allowed to return to contact sports at 4 months. The ‘restricted’ group wore a hinged brace for 6 weeks and gradually increased their ROM to 90°. They were touch-down weight-bearing for 6 weeks, followed by eventual return to sport at 6 months. The authors found no difference in the healing rate. At second-look arthroscopy, there were 9 and 10 failures in




Figure 8 Meniscus repair success: standard versus accelerated. No difference in success rates exists between the standard accelerated rehabilitation groups.

the free and restricted rehabilitation groups, respectively. There was no difference in functional outcome scores at 2 years. From this experience, the authors concluded that free rehabilitation was safe without a higher failure rate.

Kocabay *et al*⁴² reported excellent results using rehabilitation guidelines specific to the tear's characteristics. For anterior-posterior longitudinal tears less than 3 cm, they promoted weight-bearing as tolerated without a brace. ROM progressed to 125° between 3 and 6 weeks. For tears greater than 3 cm, weight-bearing was allowed in a locked brace. ROM was limited to 0°–125° until 6 or 8 weeks. Return to sport was allowed after 3 months. For complex and radial tears, patients were required to wear a brace in which they were weight-bearing as tolerated ranging from 0° to 125° for 6 to 8 weeks. They returned to sport between 4 and 5 months.

Mariani *et al*⁴⁰ followed 22 patients who underwent an outside-in meniscal repair. They were allowed to bear weight immediately without ROM restrictions. On re-examination with an MRI at an average of 28 months after surgery, only 3 of 22 patients showed signs of re-tear with greater than 1 mm of gapping. Based on this experience, they advocated for more aggressive rehabilitation regimens.³⁶

There remains a concern regarding the safety of accelerated rehabilitation in the setting of a radial meniscal tear. Most studies investigating rehabilitation after meniscal repair have included patients with a longitudinal tear. However, since radial tears experience distraction forces and increased strain with axial loading, it is thought that a more conservative postoperative rehabilitation approach may be prudent in this setting.⁴³ Choi *et al*⁴⁴ and Haklar *et al*⁴⁵ reported on their experiences repairing isolated radial tears of the lateral meniscus. Choi *et al*⁴⁴ used a weight restriction protocol, whereas Haklar used a dual restriction protocol.

Overall, there is considerable variability in the rehabilitation following a meniscal repair. There is no clear consensus regarding the ideal programme (table 2). On the one hand, Noyes limited weight-bearing initially for 4–6 weeks, with ROM progressively advanced to 135° of flexion over 6 weeks.⁴⁶ At the other end of the spectrum, O'Shea and Shelbourne⁴⁷ published favourable results after unrestricted ROM with weight-bearing as tolerated beginning 3 days after surgery. Similarly, Bryant *et al*⁴⁸ allowed weight-bearing to tolerance with the knee locked in extension for 3 weeks, then WBAT with unlimited ROM thereafter. The impact of weight-bearing combined with twisting or pivoting movements on the repaired meniscus has not been adequately investigated. Furthermore, rehabilitation protocols with respect to meniscal repair with or without augmentation have not been evaluated.

Return to sport

The decision to repair a meniscus influences both the long-term health of the knee as well as the more immediate ability to return to activity. The postoperative treatment is an important consideration that should be discussed with

the patient when considering a meniscus repair. Meniscal preservation offers long-term benefits. However, because the recovery requires a longer period of immobilisation with restrictions and delays the return to sport, some athletes might not want to have a meniscal repair. In one study of 45 meniscal repairs in elite athletes, 81% returned to sports, with the vast majority back to their prior sporting level.⁴⁹ The mean return to sport was 5.6 months (range 3–8 months) for an isolated meniscal tear, compared with 11.8 months for ACL reconstruction with meniscal repair.

Meniscal repair in association with ACL reconstruction

Accelerated, or aggressive, rehabilitation is important following ACL reconstruction to improve ROM.⁴¹ Several studies have shown that accelerated rehabilitation is safe following ACL reconstruction with meniscal repair.^{40 50} In a series of 63 consecutive patients, 58 meniscal tears were repaired arthroscopically using an inside-out technique at the time of ACL reconstruction. Barber *et al*⁵⁰ promoted a rapid return to full function. Patients were aggressively rehabilitated to playing non-contact pivoting sports at 10–12 weeks, with unlimited activity using a derotational brace as early as 3–4 months once adequate motion (0°–120°), good strength and no effusion were achieved. With regards to meniscal repairs, the authors noted a lack of consensus regarding rehabilitation protocols and called restrictions into question.

Many surgeons do not restrict patients after meniscal repair in the setting of concurrent ACL reconstructions. It has been hypothesised that meniscal repairs benefit from an abundance of healing factors due to the intra-articular bleeding present during an ACL reconstruction. In a matched cohort study by Wasserstein *et al*,⁵¹ the patient cohort with concomitant ACL reconstruction was found to have a meniscal reoperation rate of 9.7% compared with 16.7% in the meniscus repair alone cohort. Conversely, in an ACL-deficient knee, meniscal repairs are prone to failure due to the persistent mechanical stress on the tissue.^{31 52 53}

CONCLUSION

Meniscal repair is an important procedure that aims to preserve tissue and prevent future arthrosis. While treatment may alleviate symptoms and allow for a timely return to activity, there is a lack of consensus regarding the optimal postoperative rehabilitation protocol. Moreover, there is scarce evidence supporting many current practices. Biomechanical evidence suggests that high degrees of knee flexion may be safe, but these data are limited to a few cadaveric studies. The impact of rotation and torsion forces has not been determined, but have implications for the return to sport and work. It is unclear whether larger joint forces associated with running or jumping threaten the meniscal repair.

An accelerated rehabilitation protocol may be safely implemented for appropriate patients, but it is unclear how the type of meniscal tear and the repair technique should affect the postoperative programme. Additional biomechanical studies are needed to better clarify the

interplay between tear type, repair method, knee loading, knee positioning and torsional forces. Clinical studies investigating these specific elements will help to optimise patient outcomes.

Contributors RCS, MCN, AM, JPD, AN and AJR developed the idea of the study. RCS, MCN and AM collected the data. JPD, AN and AJR approved the accuracy of the collected data. RCS, MCN and AM prepared first draft of the manuscript. JPD, AN and AJR provided clinical and biomechanical insight to the study. All authors approved final manuscript.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement This is a systematic review of previously published studies. All studies are available to the public.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2018. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- Kurosawa H, Fukubayashi T, Nakajima H. Load-bearing mode of the knee joint: physical behavior of the knee joint with or without menisci. *Clin Orthop Relat Res* 1980;149:283–90.
- Fairbank TJ. Knee joint changes after meniscectomy. *J Bone Joint Surg Br* 1948;30B:664–70.
- Seitz AM, Lubomierski A, Friemert B, et al. Effect of partial meniscectomy at the medial posterior horn on tibiofemoral contact mechanics and meniscal hoop strains in human knees. *J Orthop Res* 2012;30:934–42.
- Roos EM, Ostenberg A, Roos H, et al. Long-term outcome of meniscectomy: symptoms, function, and performance tests in patients with or without radiographic osteoarthritis compared to matched controls. *Osteoarthritis Cartilage* 2001;9:316–24.
- Stein T, Mehling AP, Welsch F, et al. Long-term outcome after arthroscopic meniscal repair versus arthroscopic partial meniscectomy for traumatic meniscal tears. *Am J Sports Med* 2010;38:1542–8.
- Kim S, Bosque J, Meehan JP, et al. Increase in outpatient knee arthroscopy in the United States: a comparison of National Surveys of Ambulatory Surgery, 1996 and 2006. *J Bone Joint Surg Am* 2011;93:994–1000.
- Abrams GD, Frank RM, Gupta AK, et al. Trends in meniscus repair and meniscectomy in the United States, 2005–2011. *Am J Sports Med* 2013;41:2333–9.
- Johnson D, Weiss B. Meniscal repair using the inside-out suture technique. *Sports Med Arthrosc Rev* 2012;20:68–76.
- Baena AE, Castilla BM, Fernandez JS, et al. Inside-out medial meniscus suture: an analysis of the risk of injury to the popliteal neurovascular bundle. *Arthroscopy* 2011;27:516–21.
- Rosso C, Kovtun K, Dow W, et al. Comparison of all-inside meniscal repair devices with matched inside-out suture repair. *Am J Sports Med* 2011;39:2634–9.
- Seitz A, Kasisari R, Claes L, et al. Forces acting on the anterior meniscotibial ligaments. *Knee Surg Sports Traumatol Arthrosc* 2012;20:1488–95.
- Grant JA, Wilde J, Miller BS, et al. Comparison of inside-out and all-inside techniques for the repair of isolated meniscal tears: a systematic review. *Am J Sports Med* 2012;40:459–68.
- Scapinelli R. Studies on the vasculature of the human knee joint. *Acta Anat* 1968;70:305–31.
- Arnoczky SP, Warren RF. Microvasculature of the human meniscus. *Am J Sports Med* 1982;10:90–5.
- Venkatachalam S, Godsiff SP, Harding ML. Review of the clinical results of arthroscopic meniscal repair. *Knee* 2001;8:129–33.
- Gallacher PD, Gilbert RE, Kanes G, et al. White on white meniscal tears to fix or not to fix? *Knee* 2010;17:270–3.
- Mintzer CM, Richmond JC, Taylor J. Meniscal repair in the young athlete. *Am J Sports Med* 1998;26:630–3.
- Bloome DM, Blevins FT, Paletta GA, et al. Meniscal repair in very young children. *Arthroscopy* 2000;16:545–9.
- Nepple JJ, Dunn WR, Wright RW. Meniscal repair outcomes at greater than five years: a systematic literature review and meta-analysis. *J Bone Joint Surg Am* 2012;94:2222–7.
- Becker R, Wirz D, Wolf C, et al. Measurement of meniscofemoral contact pressure after repair of bucket-handle tears with biodegradable implants. *Arch Orthop Trauma Surg* 2005;125:254–60.
- Anderson DR, Gershuni DH, Nakhostine M, et al. The effects of non-weight-bearing and limited motion on the tensile properties of the meniscus. *Arthroscopy* 1993;9:440–5.
- Richards DP, Barber FA, Herbert MA. Compressive loads in longitudinal lateral meniscus tears: a biomechanical study in porcine knees. *Arthroscopy* 2005;21:1452–6.
- VanderHave KL, Perkins C, Le M. Weightbearing versus nonweightbearing after meniscus repair. *Sports Health* 2015;7:399–402.
- Cavanaugh JT, Killian SE. Rehabilitation following meniscal repair. *Curr Rev Musculoskelet Med* 2012;5:46–58.
- Hurwitz SR, Slawson D, Shaughnessy A. Orthopaedic information mastery: applying evidence-based information tools to improve patient outcomes while saving orthopaedists' time. *J Bone Joint Surg Am* 2000;82:888–94.
- Hill PF, Vedi V, Williams A, et al. Tibiofemoral movement 2: the loaded and unloaded living knee studied by MRI. *J Bone Joint Surg Br* 2000;82:1196–8.
- Ganley T, Arnold C, McKernan D, et al. The impact of loading on deformation about posteromedial meniscal tears. *Orthopedics* 2000;23:597–601.
- Lin DL, Ruh SS, Jones HL, et al. Does high knee flexion cause separation of meniscal repairs? *Am J Sports Med* 2013;41:2143–50.
- Fritz JM, Irrgang JJ, Harner CD. Rehabilitation following allograft meniscal transplantation: a review of the literature and case study. *J Orthop Sports Phys Ther* 1996;24:98–106.
- Busenkell GL, Lee CS. Arthroscopic meniscal repair: a posterior cannulated technique. *Arthroscopy* 1992;8:247–53.
- Morgan CD, Wojtyk EM, Casscells CD, et al. Arthroscopic meniscal repair evaluated by second-look arthroscopy. *Am J Sports Med* 1991;19:632–8.
- Sommerlath K, Hamberg P. Healed meniscal tears in unstable knees. A long-term followup of seven years. *Am J Sports Med* 1989;17:161–3.
- Marzo J, Warren R, Arnoczky S, et al. Arthroscopic meniscal repair review of the outside-in technique. *Am J Knee Surg* 1991;4:164–71.
- Scott GA, Jolly BL, Henning CE. Combined posterior incision and arthroscopic intra-articular repair of the meniscus. An examination of factors affecting healing. *J Bone Joint Surg Am* 1986;68:847–61.
- Lind M, Nielsen T, Faunø P, et al. Free rehabilitation is safe after isolated meniscus repair: a prospective randomized trial comparing free with restricted rehabilitation regimens. *Am J Sports Med* 2013;41:2753–8.
- Ahmed AM. A pressure distribution transducer for in-vitro static measurements in synovial joints. *J Biomech Eng* 1983;105:309–14.
- Thompson WO, Thaete FL, Fu FH, et al. Tibial meniscal dynamics using three-dimensional reconstruction of magnetic resonance images. *Am J Sports Med* 1991;19:210–6.
- Barber FA. Accelerated rehabilitation for meniscus repairs. *Arthroscopy* 1994;10:206–10.
- Kozłowski EJ, Barcia AM, Tokish JM. Meniscus repair: the role of accelerated rehabilitation in return to sport. *Sports Med Arthrosc Rev* 2012;20:121–6.
- Mariani PP, Santori N, Adriani E, et al. Accelerated rehabilitation after arthroscopic meniscal repair: a clinical and magnetic resonance imaging evaluation. *Arthroscopy* 1996;12:680–6.
- Shelbourne KD, Patel DV, Adsit WS, et al. Rehabilitation after meniscal repair. *Clin Sports Med* 1996;15:595–612.
- Kocabay Y, Nyland J, Isbell WM, et al. Patient outcomes following T-Fix meniscal repair and a modifiable, progressive rehabilitation program, a retrospective study. *Arch Orthop Trauma Surg* 2004;124:592–6.
- Jones RS, Keene GC, Learmonth DJ, et al. Direct measurement of hoop strains in the intact and torn human medial meniscus. *Clin Biomech* 1996;11:295–300.
- Choi NH, Kim TH, Son KM, et al. Meniscal repair for radial tears of the midbody of the lateral meniscus. *Am J Sports Med* 2010;38:2472–6.

45. Haklar U, Kocaoglu B, Nalbantoglu U, *et al.* Arthroscopic repair of radial lateral meniscus [corrected] tear by double horizontal sutures with inside-outside technique. *Knee* 2008;15:355–9.
46. Noyes FR, Barber-Westin SD. Arthroscopic repair of meniscus tears extending into the avascular zone with or without anterior cruciate ligament reconstruction in patients 40 years of age and older. *Arthroscopy* 2000;16:822–9.
47. O'Shea JJ, Shelbourne KD. Repair of locked bucket-handle meniscal tears in knees with chronic anterior cruciate ligament deficiency. *Am J Sports Med* 2003;31:216–20.
48. Bryant D, Dill J, Litchfield R, *et al.* Effectiveness of bioabsorbable arrows compared with inside-out suturing for vertical, reparable meniscal lesions: a randomized clinical trial. *Am J Sports Med* 2007;35:889–96.
49. Logan M, Watts M, Owen J, *et al.* Meniscal repair in the elite athlete: results of 45 repairs with a minimum 5-year follow-up. *Am J Sports Med* 2009;37:1131–4.
50. Barber FA, Click SD. Meniscus repair rehabilitation with concurrent anterior cruciate reconstruction. *Arthroscopy* 1997;13:433–7.
51. Wasserstein D, Dwyer T, Gandhi R, *et al.* A matched-cohort population study of reoperation after meniscal repair with and without concomitant anterior cruciate ligament reconstruction. *Am J Sports Med* 2013;41:349–55.
52. DeHaven KE, Black KP, Griffiths HJ. Open meniscus repair. Technique and two to nine year results. *Am J Sports Med* 1989;17:788–95.
53. Hamberg P, Gillquist J, Lysholm J. Suture of new and old peripheral meniscus tears. *J Bone Joint Surg Am* 1983;65:193–7.
54. Morgan CD, Casscells SW. Arthroscopic meniscus repair: a safe approach to the posterior horns. *Arthroscopy* 1986;2:3–12.
55. Horibe S, Shino K, Nakata K, *et al.* Second-look arthroscopy after meniscal repair. Review of 132 menisci repaired by an arthroscopic inside-out technique. *J Bone Joint Surg Br* 1995;77:245–9.