

RESEARCH ARTICLE


Isolation and characterization of a N4-like lytic bacteriophage infecting *Vibrio splendidus*, a pathogen of fish and bivalves

Pantelis Katharios¹*, Panos G. Kalatzis^{1,2}, Constantina Kokkari¹, Elena Sarropoulou¹, Mathias Middelboe²

1 Institute of Marine Biology, Biotechnology and Aquaculture, Hellenic Centre for Marine Research, Crete, Greece, **2** Marine Biological Section, University of Copenhagen, Helsingør, Denmark

✉ These authors contributed equally to this work.

* katharios@hcmr.gr



OPEN ACCESS

Citation: Katharios P, Kalatzis PG, Kokkari C, Sarropoulou E, Middelboe M (2017) Isolation and characterization of a N4-like lytic bacteriophage infecting *Vibrio splendidus*, a pathogen of fish and bivalves. PLoS ONE 12(12): e0190083. <https://doi.org/10.1371/journal.pone.0190083>

Editor: Francisco Rodriguez-Valera, Universidad Miguel Hernandez de Elche, SPAIN

Received: September 4, 2017

Accepted: December 7, 2017

Published: December 28, 2017

Copyright: © 2017 Katharios et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files. The phage and its bacterial host Va5 have been deposited in the open collection of the Leibniz Institute - Deutsche Sammlung von Mikroorganismen und Zellkulturen (DSMZ) under the accession numbers DSM 104622 and DSM 104620, respectively. The annotated genome sequence for the phage pVa5 was deposited in GenBank database under the accession number KX889068.

Abstract

A novel virulent bacteriophage, vB_VspP_pVa5, infecting a strain of *Vibrio splendidus* was isolated from a sea-cage aquaculture farm in Greece, and characterized using microbiological methods and genomic analysis. Bacteriophage vB_VspP_pVa5 is a N4-like podovirus with an icosahedral head measuring 85 nm in length and a short non-contractile tail. The phage had a narrow host range infecting only the bacterial host, a latent period of 30 min and a burst size of 24 virions per infected bacterium. Its genome size was 78,145 bp and genomic analysis identified 107 densely-packed genes, 40 of which could be annotated. In addition to the very large virion encapsulated DNA-dependent RNA polymerase which is the signature of the N4-like genus, an interesting feature of the novel phage is the presence of a self-splicing group I intron in the thymidylate synthase gene. A tRNA^{Stop} interrupted by a ~2.5kb open reading frame—containing area was also identified. The absence of genes related to lysogeny along with the high efficacy observed during *in vitro* cell lysis trials, indicate that the vB_VspP_pVa5 is a potential candidate component in a bacteriophage cocktail suitable for the biological control of *V. splendidus* in aquaculture.

Introduction

Vibrio splendidus is a ubiquitous inhabitant of marine and brackish water with prodigious genotypic diversity that plays a major ecological part in the oceanic and coastal environments [1,2]. It has been associated with severe epizootics in many farmed aquatic animals including fishes [3–5], crustaceans [6], bivalves and echinoderms [7]. In bivalve aquaculture, *V. splendidus* is considered to be one of the most important bacterial pathogens [8–11] responsible for severe financial losses, while in fish culture it has been reported to cause significant mortalities mostly in turbot larvae [3,12]. Skin Ulceration Syndrome (SUS) caused by *V. splendidus* [7] is an important threat for the viability and profitability of the rapidly expanding industry of holothurian culture in China and Southeast Asia [13]. In fish and invertebrate larviculture where the use of antibiotics has specific drawbacks and therefore is not applicable, control of vibriosis

Funding: The study was supported by the Danish Council for Strategic Research (ProAqua project 12-132390), the Independent Research Fund Denmark (DFF – 7014-00080), and by the Greek National Strategic Reference Framework 2007–2013 (co-funded by European Social Fund and Greek National Funds) FISHPHAGE project 131, www.gsrt.gr.

Competing interests: The authors have declared that no competing interests exist.

is extremely challenging. In these cases, bacterial control is based on the enhancement of the innate immune system of the farmed animals [14] or on the use of probiotic bacteria [7,15].

Phages have been considered as a promising alternative to antibiotics since they present several benefits over chemotherapy for microbial control. These include their high host specificity indicating that they are harmless to the natural microbiota and their autonomous transfer between animals following initial administration [16]. Phage therapy could be an ideal option for microbial control in the fragile environments of the fish and invertebrate hatcheries [17].

A few phages infecting *V. splendidus* have previously been isolated and phage therapy against *V. splendidus* in sea cucumber (*Apostichopus japonicus*), hatcheries has recently produced encouraging results leading to increased survival of the phage-treated populations. However, the phages used in that study, PVS-1, PVS-2 and PVS-3 were only morphologically characterized [18]. Additionally, although the genome sequences of three more phage isolates infecting *V. splendidus* (Helene, Henriette and Martha) are available in GenBank, as yet no publications about them have been released.

Here we report the isolation and characterization of vB_VspP_pVa5 (pVa5), a lytic N4-like podovirus which infects a *V. splendidus* strain. To our knowledge, this is the first *V. splendidus* phage characterized both morphologically and genetically.

Materials and methods

Bacterial strains

The bacterial strain used as a host (VaAn) was isolated from mussels of a bio-fouled fish cage in Greece during a vibriosis incidence. The bacterial strain has previously been fully sequenced (Accession number: PRJNA349813) and identified as *Vibrio splendidus* by multilocus sequence analysis (MLSA) [19]. It is a moderate virulent strain able to cause mortality to fish as assessed in *in vivo* studies using cod, turbot and halibut larvae [19] and is resistant to sulfadiazine/trimethoprim and oxytetracycline. In addition to the host, ten *Vibrio splendidus* strains and four clinical *Vibriosis* associated with fish diseases from HCMR's collection were used in order to assess the host range of the phage. The strains used in this assay are presented in Table 1. Bacterial strains were maintained in microbeads (MicroBank) at -80°C and grown in sea water supplemented with 1% tryptone and 0.5% yeast extract at 25°C. Host range was assessed using the agar-overlay method [20].

Phage isolation

The phage was isolated from sea water near a cage aquaculture farm in central Greece in January 2015. Water samples (1 L) were supplemented with 1% tryptone (Difco) and 0.5% yeast extract (Difco), inoculated with VaAn and incubated at 25°C for 24h. Following filtration through 0.22 µm filters, 20 µL aliquots were tested for clearing zones on bacterial lawns of the host strain. Single phage plaques were detected following serial dilutions. Isolated plaques were picked and purified by re-plating five times to ensure clonal phage stock. The purified phage was propagated in high titer (10^{10} pfu mL⁻¹) and stored at 4°C. The phage and its bacterial host VaAn were deposited in the open collection of the Leibniz Institute—Deutsche Sammlung von Mikroorganismen und Zellkulturen (DSMZ) under the accession numbers DSM 104622 and DSM 104620, respectively.

Table 1. Host range of pVa5.

#	Strain	Bacterial species	Origin	Infectivity
1	VaAn	<i>V. splendidus</i>	Greece	+
2	3D3-3/pop17	<i>V. splendidus</i>	France	-
3	4D6-8/pop19	<i>V. splendidus</i>	France	-
4	3F1-17/pop25	<i>V. splendidus</i>	France	-
5	3F1-22/pop21	<i>V. splendidus</i>	France	-
6	3F1-45/pop20	<i>V. splendidus</i>	France	-
7	3H2-4/pop24	<i>V. splendidus</i>	France	-
8	3Z-15/pop23	<i>V. splendidus</i>	France	-
9	3Z-28/pop22	<i>V. splendidus</i>	France	-
10	3Z-31/pop16	<i>V. splendidus</i>	France	-
11	3Z-39/pop18	<i>V. splendidus</i>	France	-
12	DSMZ 2171	<i>V. alginolyticus</i>	Japan	-
13	DSMZ 19623	<i>V. harveyi</i>	USA	-
14	ATCC 19264	<i>V. anguillarum</i>	Sweden	-
15	ATCC 33509	<i>V. ordalii</i>	USA	-

<https://doi.org/10.1371/journal.pone.0190083.t001>

Phage morphology

Phage preparation in high titer (10^{10} pfu mL⁻¹) was negatively stained with 4% (w/v) uranyl acetate (pH 7.2) and observed with a JOEL JEM2100 transmission electron microscope (TEM) operated at 60 kV at the Electron Microscopy Laboratory of the University of Crete.

One-step growth analysis

The latency period and burst size were determined from the one-step growth curve, according to standard methodology. To summarize, 1 mL from a bacterial culture at the exponential phase was centrifuged and washed with saline buffer (0.9% NaCl), re-suspended in 1 ml medium and then infected with the phage at a MOI of 0.01. After 10 minutes' incubation, the mixture was diluted 25 times in the medium and at this time point and every 10 minutes thereafter, a sample was removed, and serial decimal dilutions were spotted on a previously prepared lawn of the host on agar plates. The number of infective phages in each sample was quantified from the plaques formed after an overnight incubation of the plates [17].

In vitro lysis

An *in vitro* lysis assay was performed in sterile 96-well plates using a TECAN microplate reader (Infinite PRO 200) equipped with temperature control. Specifically, twelve 200-μL wells of the plate were loaded with freshly-prepared cultures of the host organism. The plate was placed in the reader and incubated at 25°C with orbital shaking. Phage was added at three different multiplicities of infection (MOI 1, 10 and 100) in triplicates when the bacterial culture was at the exponential phase. Three wells were not infected and these served as control. The growth curve of the cultures was monitored in real-time over an 8-hour period, and optical density measurements at a wavelength of 560 nm (OD560) were recorded every 10 minutes.

DNA extraction and sequencing

The bacteriophage's genomic DNA was extracted according to the phenol-chloroform protocol [21] and stored in sterile Eppendorf tubes in -20°C until sequenced. The genome of the

pVa5 bacteriophage was sequenced using the Genome Sequencer GS Junior System (Roche Diagnostic). Shotgun sequencing was performed according to the manufacturer's instructions using 5 µg of the bacteriophage genomic DNA. Low quality sequences were trimmed off and shotgun reads were assembled using the De novo assembly of the GS Assembler software (Newbler).

Genome structure determination

The physical structure of the genome was determined by digesting the phage genome with restriction endonucleases. The enzyme ApaI (Thermo Scientific) was used in order to obtain the genomic restriction profile of the phage. The results were visualized on 1% agarose gel and the sizes of the DNA fragments were estimated in comparison with Lambda DNA/Hind III marker (Thermo Scientific).

Additionally, four sets of primers (Eurofins) were manually designed in order to determine by Polymerase Chain Reaction (PCR), whether the phage genome was circularly-permuted (S1 Fig). Two sets of primers specifically targeted two genomic areas which enclosed the first and the last bp of the obtained *de novo* assembled phage genome contig. The other two sets of primers were used as positive controls for the PCR, since they were targeting two random genomic areas of the phage.

Genomic analysis and phylogeny

Open reading frames (ORFs) were determined using RAST [22,23] and then manually curated using Geneious 9.1.5 based on three different start codons (ATG, GTG and TTG) and putative Shine Dalgarno sequences upstream the coding regions [24]. Genes and coding DNA sequences (CDSs) were predicted using a combined gene model from GeneMarkS [25] and Glimmer3 [26]. The predicted proteins were annotated by BLAST against the NCBI non-redundant database. Additional annotation was subsequently performed using Blast2Go [27] and InterProScan [28]. The presence of tRNAs was examined using both tRNAscan [29] and ARAGORN [30]. The annotated genome sequence for the phage pVa5 was deposited in GenBank database under the accession number KX889068.

Phylogenetic analysis was performed using two genetic markers: the DNA polymerase (DNAP I) and the virion-encapsulated RNA polymerase genes (vRNAP) of 30 available N4-like phages, including pVa5. The selected DNA sequences were aligned by ClustalX with default settings and the phylogenetic tree was constructed using Neighbor-Joining method with 1000 bootstraps through Geneious bioinformatic platform 9.1.5 [24]. The genomes of the N4-like phages which were classified in the same taxonomic group with pVa5 were compared to the novel phage using the progressive Mauve algorithm [31] in Geneious platform 9.1.5, in order to unravel its genomic architecture [24]. The determination of protein structure and RNA families was determined by Phyre² [32] and Rfam [33] online bioinformatic tools, respectively, in those cases where such analyses were necessary. The evaluation of the bacteriophage's lifestyle was performed through the combination of results obtained by Phyre² and BLAST. The presence of genes, such as integrase, fitness or virulence factors, which might indicate a lysogenic lifestyle for pVa5 was examined.

Results and discussion

Morphology, life cycle and phage efficacy

The phage produced small (pinhead size) clear plaques on the host lawn. Following TEM observations, it was classified to Podoviridae family. It had an icosahedral head measuring 85




Fig 1. Transmission electron micrograph of *Vibrio splendidus* phage demonstrating a Podoviridae morphology.

<https://doi.org/10.1371/journal.pone.0190083.g001>

nm in length and a short (25 nm) non-contractile tail (Fig 1). It was designated as vB_VspP_pVa5 according to the suggestions for bacterial viruses' nomenclature [34], while its abbreviation, pVa5, will be used in the manuscript.

One-step growth curve analysis (Fig 2) showed that pVa5 had a latent period of 30 min and a burst size of 24 virions per infected cell. N4-like bacteriophages demonstrate a great variability in both these life cycle parameters. Latency periods between 10–15 min to 6h and burst sizes between 10 to even 9000 virions per cell have already been reported in the literature. For




Fig 2. One-step growth curve of bacteriophage pVa5. The values are means \pm standard deviation of three replicates.

<https://doi.org/10.1371/journal.pone.0190083.g002>




Fig 3. *In vitro* lysis assay using bacteriophage pVa5 and the host bacterium *V. splendidus* VaAn at 3 different MOIs at 25°C. The values are means \pm standard deviation of 3 replicates.

<https://doi.org/10.1371/journal.pone.0190083.g003>

instance, in the case of phiAxp-3 a latency time of 80 min and burst size of 9000 virions per infected cell have been observed [35–37].

A spot test showed that the phage has a very limited lytic spectrum based on the 15 tested strains, since it could infect only the *V. splendidus* host strain VaAn and no other *Vibrio splendidus* strains, nor *Vibrio* species tested for phage susceptibility (Table 1).

In vitro lysis assay indicated that pVa5 could almost completely eliminate bacterial growth for 8 hours (Fig 3). The intense bactericidal activity that the phage demonstrated, renders it a very powerful candidate for *in vivo* phage therapy trials against *V. splendidus* strain VaAn. The inhibition of bacterial growth was equally marked with all three different applied MOIs, even at MOI: 1. The high efficiency of the phage makes it also practically convenient for use even in large volumes of water, such as fish and shellfish hatcheries, which would eventually be the ultimate goal. The regrowth of the bacterial population which takes place after almost 8h, even though it is very slow, could be considered as a constraint to phage therapy application. However, it has been adequately reported in the literature that bacteriophage resistance is associated with reduced virulence [38,39]. Hence, the exact effect of resistance development needs to be further evaluated through meticulously designed *in vivo* trials.

Genomic characterization

The genome of pVa5 had a size of 78,145 bp and a G+C content of 43.2% which is comparable to the host G+C content (44.3%). 107 genes were identified (S1 Table) of which 101 had ATG as starting codon, 4 GTG genes: 40, 45, 47 and 55) and 2 TTG (genes: 33 and 36). The coding region size of the genome was 71,258 bp (91.18% of the whole genome).

The digestion of the viral genome with ApaI produced four distinct DNA fragments, which clearly indicated that the genome contains three single restriction sites. ApaI recognizes and cleaves GGGCC[^]C genomic sites and the presence of three such sites were *in silico* confirmed in the phage genome. If the genome were circular, we should expect to find three distinct DNA fragments; however in our case, ApaI treatment yielded four distinct DNA fragments. Hence, it can be concluded that the physical structure of the pVa5 phage genome is linear and not circular. However, specific PCR using two sets of primers that match the start (bp 1) and end (bp 78,145) point of the physical phage genome, was able in both sets to amplify the




Fig 4. Gene map of pVa5. Genes with attributed function are noted in the figure. Yellow: CDS, Green: genes, White: introns, Dark green: putative tRNA.

<https://doi.org/10.1371/journal.pone.0190083.g004>

targeted genomic parts. In physically-linear genome phages, the packaging mechanism during replication may explain this result. Circularly permuted genomes that follow the formation of a linear concatemer during replication packaging do not have defined ends. Similarly, in our case, the replication process of pVa5 ensures the successful packaging of a complete 107-gene set (Fig 4) in each of the newly produced virions [40]; however, the DNA molecule is not always packed with the same start and end positions. A hydrolase CDS (gene 9) was also detected as soon as the obtained phage genome contig was *in silico* circularized, corroborating the circularly permuted propagation strategy of the pVa5. According to the convention followed for the circularly permuted phages, bp1 is set as the first nucleotide of the closest intergenic region immediately to the left of the terminase gene. In our case, the small terminase gene (gene 1) and large terminase gene (gene 4) were annotated as gp69 and gp68 since they resembled, respectively, the originally annotated small and large terminase genes of the N4 phage [41] by 61.7% and 71.2%.

None of the tRNA detection bioinformatic tools used were able to identify any tRNA genes, indicating that the phage is adapted to the host tRNA makeup for its protein synthesis [42]. However, after modifying the default parameters of the ARAGORN software (Allow introns 0–3000 bases: yes), an intervening ~2.5 kbp coding sequence (CDS) containing genomic area was present in a tRNA^{Stop} gene. Bacteriophages often carry tRNA genes for the regulation of translation-associated functions [43,44]. Even though intron-mediated tRNAs have been reported in both eukaryotic nuclei and Archaea [45], this finding is, to our knowledge, the first observed in a bacteriophage genome. This tRNA^{Stop} gene matches to the TAA stop codon and since TAA is the prevalent stop codon in pVa5 genome (72 out of 107 CDS), it could potentially contribute to regulatory procedures during viral infection. It has to be stated however that although the genomic structure of the tRNA gene matches to a tRNA^{Stop}, this identification might be an artifact. In order to be certain about its existence and its functionality, the specific mechanism of this gene’s function still needs to be elucidated.

Analysis of the pVa5 nucleotide sequence using the BLASTn algorithm of the NCBI Database [46] showed that the novel phage is unique as its closest similarity is with *Pseudoalteromonas* phage pYD6-A (E-value, 0.0; query coverage, 5%; identity, 78%) and the vibriophages VPB32 and VPB47 (E-value, 0.0; query coverage, 7%; identity, 76%) all of which are also N4-like podoviruses. Individual-gene BLAST search led to 34 genes with e-value > 0.01 and query coverage > 30% (S1 Table). The novelty of pVa5 is also supported by the fact that 29% of its total genes (31 out of 34), correspond to their homologues in pYD6-A, VPB32 and

VPB47. After mapping each of the pVa5 phage genes against the nr nucleotide collection of NCBI, no relationship with any potential virulence or fitness factor was documented. The structure of the proteins or hypothetical proteins did not either indicate any correlations with known toxins, corroborating to the lytic nature of the bacteriophage. After whole genome comparison on the amino acid level, pVa5 shared 24%, 22.7% and 14.4% identical sites with VBP47, VBP32 and pYD6-A, respectively. However, when the comparison was focused on the homologues, the pairwise alignment percentages ranged from at least 50% up to 90%. Our data suggest that pVa5 is no lysogenic like all other N4-like phages which have been recorded so far.

The key features characterizing the N4-like phages are the presence of a very large virion-encapsulated DNA-dependent RNA polymerase (vRNAP) responsible for the early gene transcription, 2 RNA polymerases and a single stranded DNA-binding protein [47,48]. The vRNAP of pVa5 (gene 25) is 3,345 aa in size, while the phage contains two more RNA polymerases (gene 79 and gene 90) 405 and 311 aa in size. The phage-encoded RNAPs of pVa5 are able, as in all N4-like phages, to establish the transcriptional independence of the virus [49]. Gene 28 was predicted to encode the single-stranded DNA binding (SSB) protein for pVa5. In the N4 phage, the SSB acts as an architectural transcription factor by providing an active promoter conformation for vRNAP binding [50].

The genomic organization of N4-like phages is considered to be highly conserved [42,51]. The genomes of N4-like phages are organized basically in two large clusters transcribed in opposite directions [52]; this type of organization is followed by pVa5, nevertheless with two exceptions, genes 14 and 15, which encode a peptidase M15A related to lysis and a phosphate starvation-inducible protein (PhoH) (possibly a phosphorus regulation gene). It has been demonstrated that many marine phages contain genes related to pho regulation [53] and although its function remains unclear, there has been speculation that the phages might use these genes to take over the phosphorus acquisition mechanism for their own benefit during phage DNA replication [54]. Similarly, ribonucleotide reductase (RNR), the enzyme responsible for the formation of deoxyribonucleotides from ribonucleotides, is also considered to be a tool used by the phage to sustain host metabolism promoting its own replication [55]. Bacteriophage pVa5 has three consecutive genes organized in a cluster (genes 33–35) encoding RNRs belonging to class I and class III which is not very common in podoviridae [56]. One of the most interesting findings in the genome of pVa5 is the presence of an intervening sequence containing an HNH endonuclease (gene 41) splicing the thymidylate synthase (*td*) gene (CDS 40.1 and 40.2). The concatenated amino acid sequence of the two thymidylate synthase CDS of pVa5 was aligned and mapped against their homologue *td* synthase genes from the bacteriophages VBP32, VBP47 and pYD6-A (S2 Fig). Apart from a 57 aa sequence that was completely missing from the *td* synthase gene of pVa5, their pairwise percentage identity was estimated at 82.8%, indicating a very high degree of conservation. Thymidylate synthase (*td*) is one of the most conserved enzymes among bacteriophage genomes [57] and partition of the gene by an intron into two separate genes has been observed on two other occasions: in the *E. coli* bacteriophage T4 [58] and in *Bacillus* bacteriophage b22 [59]. It has been suggested that *td* has a dual role related to catalysis and to the structure of phage baseplate, thus a differential expression that might be facilitated by the splicing could be the underlying reason [58]. Additionally, the *td* synthase intron is characterized as a group I intron (E-value: 1E-18, Accession: RF00028) in the Rfam database [33]. Group I introns are self-splicing introns that even though initially considered as molecular parasites, have been found to act in a co-evolution context which promotes mutualistic interactions with their host genes [60]. The first reported microbial group I intron was the intron that was also found in the *td* gene of the T4 bacteriophage [58]. Several group I introns found in bacteriophage genomes are being reported in the literature with an

increasing frequency [61,62]. Their self-splicing nature allows them to act as ribozymes catalyzing the expression of the gene in which they have intervened. Furthermore, the homing (HNH) endonuclease genes that are usually encoded by such introns, allow the introns to act also as mobile genetic elements and expand their distribution in intron-less genes through DNA-based recombination mobility [63,64]. This finding is also in accordance to the rules that should be followed by group I intron ribozymes [65]: 1) the pVa5 group I intron is located in a highly conserved gene (td gene); 2) it has a T preceding the insertion site and a G as last nucleotide of the insertion; 3) the size of the intron without the HNH endonuclease encoding area is 279 nt. However, it should be stated that HNH endonucleases are quite common, not only in the N4-like phages but generally in phage genomes since they constitute widespread components of phage DNA packaging machinery [66].

The lysis cassette of N4-like phages typically contains an endolysin, an N-acetylmuramoyl-L-alanine amidase, a holin and a peptidoglycan-degrading protein [52]. In pVa5 no lysis cluster could be identified and the only protein that could be related to lysis was a peptidase M15 encoded by gene 14. Similarly, a lysis cluster was also absent in the phages of *Vibrio parahaemolyticus* VBP32, VBP47 and *Pseudoalteromonas phage* pYD6-A [52]. Interestingly, these phages were evolutionarily closer to pVa5 as demonstrated by the phylogeny based on DNAP and vRNAP genes (Fig 5). Multiple alignment of the whole genomes of these phages using the progressive Mauve algorithm showed that pVa5 shares similar synteny characteristics with its taxonomic group phages (Fig 6). In addition to the closely related N4-like vibrio phages VBP47 and VBP32 which are identical and infect *Vibrio parahaemolyticus* strain RIMD2210633, the two other sequenced N4-like phages Ja-1 and VCO139 which infect *Vibrio cholerae* [67] were phylogenetically clustered in different but still vicinal branches. Especially in the RNAP-based phylogeny, all Vibriophages are clustered into the same taxonomic group which is then further divided into two distinct but closely related branches (Fig 5B).




Fig 5. Molecular phylogenetic analysis by Neighbor-joining method based on the Tamura-Nei model. The consensus tree which was produced by 1000 bootstraps is shown. The novel bacteriophage vB_VspP_pVa5 is classified by two different genetic markers: DNA Polymerase I (A) and virion-encapsulated RNA Polymerase (B) in a statistically robust taxonomic group which is highlighted by green areas and includes two *Vibrio* phages (NC020868 and HQ634194) and one *Pseudoalteromonas* phage (JF974296). The analysis involved 30 nucleotide sequences.

<https://doi.org/10.1371/journal.pone.0190083.g005>




Fig 6. Multiple alignment of the whole genomes of all N4-like Vibriophages using the progressive Mauve algorithm. Same coloured blocks indicate high synteny between genomes without genomic rearrangements. Genomic similarity is represented by the height of the bars, which correspond to the average level of conservation in that region of the genome sequence. Completely white regions represent fragments that were not aligned or contained sequence elements specific to a particular genome. The four bacteriophages included in the analysis are classified in the same taxonomic group by both DNAP and vRNAP gene markers.

<https://doi.org/10.1371/journal.pone.0190083.g006>

Conclusions

Bacteriophage vB_VspP_pVa5 is a novel virulent podovirus that infects *Vibrio splendidus*. Although it is a virulent phage based on the genomic analysis, it has a narrow host range and therefore is not suitable for a universal phage therapy scheme. On the other hand, it has interesting genomic features that significantly contribute to our current knowledge about N4-like phages and could therefore be exploited in phage infection studies in combination with the genomic information available for its bacterial host.

Supporting information

S1 Table. Annotation of the vB_VspP_pVa5 Vibriophage accompanied by the BLAST results of the genes. Genes with e-value > 0.01 and query coverage > 30% were omitted from the BLAST results in order to prevent a false interpretation of the results. (DOCX)

S1 Fig. Four sets of primers manually designed; sets 1 and 2 were able to amplify the genomic area between the first and last bp, whereas sets pos1 and pos2 served as positive controls in order to confirm the functionality of the PCR. The relative position of the primers as well as the genomic parts expected to be amplified, are also schematically depicted below. (PDF)

S2 Fig. Pairwise alignment and comparison of the amino acid sequences of the 4 thymidylate synthase genes from 4 N4-like phages classified in the same taxonomic group: *Vibrio* phage VBP47 (HQ634194), *Vibrio* phage VBP32 (NC_020868), *Pseudoalteromonas* phage pYD6-a (JF974296) and *Vibrio* phage vB_VspP_pVa5 (KX889068). (PDF)

Acknowledgments

We would like to thank Dr. F. Le Roux from Roscoff Marine Station for kindly providing the 10 *V. splendidus* strains that were used in the host range analysis, Dr. Manolis Mandalakis for

assisting in *in vitro* cell lysis assays, Prof. George Chalepakis and the staff of the Laboratory of Electron Microscopy of the University of Crete and Mrs. Margaret Eleftheriou for critical reading of the manuscript. The study was supported by the Danish Council for Strategic Research (ProAqua project 12–132390), the Independent Research Fund Denmark (DFF– 7014–00080), and by the Greek National Strategic Reference Framework 2007–2013 (co-funded by European Social Fund and Greek National Funds) FISHPHAGE project 131, www.gsrt.gr.

Author Contributions

Conceptualization: Pantelis Katharios, Panos G. Kalatzis.

Data curation: Pantelis Katharios.

Formal analysis: Pantelis Katharios, Panos G. Kalatzis, Constantina Kokkari, Elena Sarropoulou.

Investigation: Elena Sarropoulou.

Methodology: Pantelis Katharios, Constantina Kokkari, Elena Sarropoulou.

Project administration: Pantelis Katharios, Mathias Middelboe.

Resources: Pantelis Katharios.

Supervision: Pantelis Katharios.

Validation: Constantina Kokkari, Mathias Middelboe.

Writing – original draft: Pantelis Katharios, Panos G. Kalatzis, Constantina Kokkari, Elena Sarropoulou.

Writing – review & editing: Pantelis Katharios, Panos G. Kalatzis, Mathias Middelboe.

References

1. Le Roux F, Zouine M, Chakroun N, Binesse J, Saulnier D, Bouchier C, et al. Genome sequence of *Vibrio splendidus*: An abundant planktonic marine species with a large genotypic diversity. *Environ Microbiol.* 2009; 11: 1959–1970. <https://doi.org/10.1111/j.1462-2920.2009.01918.x> PMID: 19364337
2. Preheim SP, Timberlake S, Polz MF. Merging taxonomy with ecological population prediction in a case study of Vibrionaceae. *Appl Environ Microbiol.* 2011; 77: 7195–7206. <https://doi.org/10.1128/AEM.00665-11> PMID: 21873482
3. Gatesoupe F-J, Lambert C, Nicolas J-L. Pathogenicity of *Vibrio splendidus* strains associated with turbot larvae, *Scophthalmus maximus*. *J Appl Microbiol.* Wiley Online Library; 1999; 87: 757–763.
4. Jensen S, Samuelsen OB, Andersen K, Torkildsen L, Lambert C, Choquet G, et al. Characterization of strains of *Vibrio splendidus* and *V. tapetis* isolated from corkwing wrasse *Symphodus melops* suffering vibriosis. 2003;
5. Balcázar JL, Gallo-Bueno A, Planas M, Pintado J. Isolation of *Vibrio alginolyticus* and *Vibrio splendidus* from captive-bred seahorses with disease symptoms. *Antonie van Leeuwenhoek, Int J Gen Mol Microbiol.* 2010; 97: 207–210. <https://doi.org/10.1007/s10482-009-9398-4> PMID: 19921544
6. Baticados MCL, Lavilla-Pitogo CR, Cruz-Lacierda ER, de la Pena LD. Studies on the chemical control of luminous bacteria *Vibrio harveyi* and *V. splendidus* isolated from diseased *Penaeus monodon* larvae and rearing water. *Dis aquat Org.* 1990; 9: 133–139.
7. Zhao Y, Zhang W, Xu W, Mai K, Zhang Y, Liufu Z. Effects of potential probiotic *Bacillus subtilis* T13 on growth, immunity and disease resistance against *Vibrio splendidus* infection in juvenile sea cucumber *Apostichopus japonicus*. *Fish Shellfish Immunol.* 2012; <https://doi.org/10.1016/j.fsi.2012.01.027> PMID: 22342649
8. Gómez-León J, Villamil L, Lemos ML, Novoa B, Figueras A. Isolation of *Vibrio alginolyticus* and *Vibrio splendidus* from aquacultured carpet shell clam (*Ruditapes decussatus*) larvae associated with mass mortalities. *Appl Environ Microbiol. Am Soc Microbiol;* 2005; 71: 98–104.
9. Lacoste A, Jalabert F, Malham S, Cueff A, Gelebart F, Cordevant C, et al. A *Vibrio splendidus* strain is associated with summer mortality of juvenile oysters *Crassostrea gigas* in the Bay of Morlaix (North

- Brittany, France). *Dis Aquat Organ.* 2001; 46: 139–145. <https://doi.org/10.3354/dao046139> PMID: 11678229
10. Sugumar G, Nakai T, Hirata Y, Matsubara D, Muroga K. *Vibrio splendidus* biovar II as the causative agent of bacillary necrosis of Japanese oyster *Crassostrea gigas* larvae. *Dis Aquat Organ.* 1998; 33: 111–118. <https://doi.org/10.3354/dao033111> PMID: 9684317
 11. Dubert J, Barja JL, Romalde JL. New insights into pathogenic vibrios affecting bivalves in hatcheries: Present and future prospects. *Front Microbiol.* 2017; 8: 1–16.
 12. Thomson R, Macpherson HL, Riaza A, Birkbeck TH. *Vibrio splendidus* biotype 1 as a cause of mortalities in hatchery-reared larval turbot, *Scophthalmus maximus* (L.). *J Appl Microbiol.* Wiley Online Library; 2005; 99: 243–250.
 13. Conand C. Advances in sea cucumber aquaculture and management. Food & Agriculture Org.; 2004.
 14. Gu M, Ma H, Mai K, Zhang W, Bai N, Wang X. Effects of dietary β -glucan, mannan oligosaccharide and their combinations on growth performance, immunity and resistance against *Vibrio splendidus* of sea cucumber, *Apostichopus japonicus*. *Fish Shellfish Immunol.* Elsevier; 2011; 31: 303–309.
 15. Hjelm M, Bergh O, Riaza A, Nielsen J, Melchiorson J, Jensen S, et al. Selection and identification of autochthonous potential probiotic bacteria from turbot larvae (*Scophthalmus maximus*) rearing units. *Syst Appl Microbiol.* 2004; 27: 360–71. <https://doi.org/10.1078/0723-2020-00256> PMID: 15214642
 16. Nakai T, Park SC. Bacteriophage therapy of infectious diseases in aquaculture. *Res Microbiol.* 2002; 153: 13–18. [http://dx.doi.org/10.1016/S0923-2508\(01\)01280-3](http://dx.doi.org/10.1016/S0923-2508(01)01280-3) PMID: 11881893
 17. Kalatzis PG, Bastías R, Kokkari C, Katharios P. Isolation and Characterization of Two Lytic Bacteriophages, ϕ St2 and ϕ Grn1; Phage Therapy Application for Biological Control of *Vibrio alginolyticus* in Aquaculture Live Feeds. *PLoS One.* 2016; 1–18. <https://doi.org/10.1371/journal.pone.0151101> PMID: 26950336
 18. Li Z, Li X, Zhang J, Wang X, Wang L, Cao Z, et al. Use of phages to control *Vibrio splendidus* infection in the juvenile sea cucumber *Apostichopus japonicus*. *Fish Shellfish Immunol.* Elsevier Ltd; 2016; 54: 302–311. <https://doi.org/10.1016/j.fsi.2016.04.026> PMID: 27108378
 19. Rønneseth A, Castillo D, D'Alvise P, Tønnesen, Haugland G, Grotkjær T, et al. Comparative assessment of *Vibrio* virulence in marine fish larvae. *J Fish Dis.* 2017; 40: 1373–1385. <https://doi.org/10.1111/jfd.12612> PMID: 28160295
 20. Kutter E. Phage host range and efficiency of plating. *Methods Mol Biol.* 2009; 501: 141–9. https://doi.org/10.1007/978-1-60327-164-6_14 PMID: 19066818
 21. Higuera G, Bastías R, Tsertsvadze G, Romero J, Espejo RT. Recently discovered *Vibrio anguillarum* phages can protect against experimentally induced vibriosis in Atlantic salmon, *Salmo salar*. *Aquaculture.* Elsevier B.V.; 2013; 392–395: 128–133. <https://doi.org/10.1016/j.aquaculture.2013.02.013>
 22. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, et al. The RAST Server: rapid annotations using subsystems technology. *BMC Genomics.* 2008; 9: 75. <https://doi.org/10.1186/1471-2164-9-75> PMID: 18261238
 23. Overbeek R, Olson R, Pusch GD, Olsen GJ, Davis JJ, Disz T, et al. The SEED and the Rapid Annotation of microbial genomes using Subsystems Technology (RAST). *Nucleic Acids Res.* 2014; 42: 206–214. <https://doi.org/10.1093/nar/gkt1226> PMID: 24293654
 24. Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, et al. Geneious Basic: An integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics.* 2012; 28: 1647–1649. <https://doi.org/10.1093/bioinformatics/bts199> PMID: 22543367
 25. Besemer J, Lomsadze A, Borodovsky M. GeneMarkS: a self-training method for prediction of gene starts in microbial genomes. Implications for finding sequence motifs in regulatory regions. *Nucleic Acids Res.* 2001; 29: 2607–2618. PMID: 11410670
 26. Delcher AL, Bratke KA, Powers EC, Salzberg SL. Identifying bacterial genes and endosymbiont DNA with Glimmer. *Bioinformatics.* 2007; 23: 673–679. <https://doi.org/10.1093/bioinformatics/btm009> PMID: 17237039
 27. Conesa A, Götz S, García-Gómez JM, Terol J, Talón M, Robles M. Blast2GO: A universal tool for annotation, visualization and analysis in functional genomics research. *Bioinformatics.* 2005; 21: 3674–3676. <https://doi.org/10.1093/bioinformatics/bti610> PMID: 16081474
 28. Jones P, Binns D, Chang HY, Fraser M, Li W, McAnulla C, et al. InterProScan 5: Genome-scale protein function classification. *Bioinformatics.* 2014; 30: 1236–1240. <https://doi.org/10.1093/bioinformatics/btu031> PMID: 24451626
 29. Lowe TM, Chan PP. tRNAscan-SE On-line: integrating search and context for analysis of transfer RNA genes. *Nucleic Acids Res.* 2016; 44: W54–7. <https://doi.org/10.1093/nar/gkw413> PMID: 27174935
 30. Laslett D, Canback B. ARAGORN, a program to detect tRNA genes and tmRNA genes in nucleotide sequences. *Nucleic Acids Res.* 2004; 32: 11–16. <https://doi.org/10.1093/nar/gkh152> PMID: 14704338

31. Darling ACE, Mau B, Blattner FR, Perna NT. Mauve: Multiple Alignment of Conserved Genomic Sequence With Rearrangements. *Bioinformatics*. 2004; 19: 1394–1403. <https://doi.org/10.1101/gr.2289704> PMID: 15231754
32. Kelly LA, Mezulis S, Yates C, Wass M, Sternberg M. The Phyre2 web portal for protein modelling, prediction, and analysis. *Nat Protoc*. 2015; 10: 845–858. <https://doi.org/10.1038/nprot.2015.053> PMID: 25950237
33. Nawrocki EP, Burge SW, Bateman A, Daub J, Eberhardt RY, Eddy SR, et al. Rfam 12.0: Updates to the RNA families database. *Nucleic Acids Res*. 2015; 43: D130–D137. <https://doi.org/10.1093/nar/gku1063> PMID: 25392425
34. Kropinski AM, Prangishvili D, Lavigne R. Position paper: The creation of a rational scheme for the nomenclature of viruses of Bacteria and Archaea. *Environ Microbiol*. 2009; 11: 2775–2777. <https://doi.org/10.1111/j.1462-2920.2009.01970.x> PMID: 19519870
35. Chan JZM, Millard AD, Mann NH, Schfer H. Comparative genomics defines the core genome of the growing N4-like phage genus and identifies N4-like roseophage specific genes. *Front Microbiol*. 2014; 5: 1–14.
36. Li B, Zhang S, Long L, Huang S. Characterization and Complete Genome Sequences of Three N4-Like Roseobacter Phages Isolated from the South China Sea. *Curr Microbiol*. Springer US; 2016; 73: 409–418. <https://doi.org/10.1007/s00284-016-1071-3> PMID: 27270945
37. Ma Y, Li E, Qi Z, Li H, Wei X, Lin W, et al. Isolation and molecular characterisation of Achromobacter phage phiAxp-3, an N4-like bacteriophage. *Sci Rep*. Nature Publishing Group; 2016; 6: 24776. <https://doi.org/10.1038/srep24776> PMID: 27094846
38. Leon M, Bastias R. Virulence reduction in bacteriophage resistant bacteria. *Front Microbiol*. 2015; 6: 1–7.
39. Castillo D, Christiansen RH, Dalsgaard I, Madsen L, Middelboe M. Bacteriophage resistance mechanisms in the fish pathogen *Flavobacterium psychrophilum*: Linking genomic mutations to changes in bacterial virulence factors. *Appl Environ Microbiol*. 2015; 1.
40. Merrill BD, Ward AT, Grose JH, Hope S. Software-based analysis of bacteriophage genomes, physical ends, and packaging strategies. *BMC Genomics*. BMC Genomics; 2016; 17: 679. <https://doi.org/10.1186/s12864-016-3018-2> PMID: 27561606
41. Kulikov E, Kropinski AM, Golomidova A, Lingohr E, Govorun V, Serebryakova M, et al. Isolation and characterization of a novel indigenous intestinal N4-related coliphage vB_EcoP_G7C. *Virology*. Elsevier B.V.; 2012; 426: 93–99. <https://doi.org/10.1016/j.virol.2012.01.027> PMID: 22341309
42. Wittmann J, Dreiseikelmann B, Rohde M, Meier-Kolthoff JP, Bunk B, Rohde C. First genome sequences of Achromobacter phages reveal new members of the N4 family. *Virology*. 2014; 11: 14. <https://doi.org/10.1186/1743-422X-11-14> PMID: 24468270
43. Bailly-Bechet M, Vergassola M, Rocha E. Causes for the intriguing presence of tRNAs in phages. *Genome Res*. 2007; 17: 1486–1495. <https://doi.org/10.1101/gr.6649807> PMID: 17785533
44. Skliros D, Kalatzis PG, Katharios P, Flemetakis E. Comparative Functional Genomic Analysis of Two Vibrio Phages Reveals Complex Metabolic Interactions with the Host Cell. *Front Microbiol*. 2016; 7: 1–13.
45. Haugen P, Simon DM, Bhattacharya D. The natural history of group I introns. *Trends Genet*. 2005; 21: 111–119. <https://doi.org/10.1016/j.tig.2004.12.007> PMID: 15661357
46. Boratyn GM, Camacho C, Cooper PS, Coulouris G, Fong A, Ma N, et al. BLAST: a more efficient report with usability improvements. *Nucleic Acids Res*. 2013; 41: 29–33. <https://doi.org/10.1093/nar/gkt282> PMID: 23609542
47. Kazmierczak KM, Davydova EK, Mustaev AA, Rothman-Denes LB. The phage N4 virion RNA polymerase catalytic domain is related to single-subunit RNA polymerases. *EMBO J*. 2002; 21: 5815–5823. <https://doi.org/10.1093/emboj/cdf584> PMID: 12411499
48. Choi KH, McPartland J, Kaganman I, Bowman VD, Rothman-Denes LB, Rossmann MG. Insight into DNA and Protein Transport in Double-Stranded DNA Viruses: The Structure of Bacteriophage N4. *J Mol Biol*. 2008; <https://doi.org/10.1016/j.jmb.2008.02.059> PMID: 18374942
49. Lenneman BR, Rothman-Denes LB. Structural and biochemical investigation of bacteriophage N4-encoded RNA polymerases. *Biomolecules*. 2015; 5: 647–67. <https://doi.org/10.3390/biom5020647> PMID: 25924224
50. Lenneman BR, Rothman-Denes LB. Structural and biochemical investigation of bacteriophage N4-encoded RNA polymerases. *Biomolecules*. 2015; 5: 647–67. <https://doi.org/10.3390/biom5020647> PMID: 25924224
51. Wittmann J, Klumpp J, Moreno Switt AI, Yagubi A, Ackermann H-W, Wiedmann M, et al. Taxonomic reassessment of N4-like viruses using comparative genomics and proteomics suggests a new

- subfamily—“*Enquartavirinae*”. Arch Virol. Springer Vienna; 2015; 160: 3053–3062. <https://doi.org/10.1007/s00705-015-2609-6> PMID: 26395091
52. Wittmann J, Klumpp J, Moreno Switt AI, Yagubi A, Ackermann H-W, Wiedmann M, et al. Taxonomic reassessment of N4-like viruses using comparative genomics and proteomics suggests a new subfamily—“*Enquartavirinae*”. Arch Virol. Springer Vienna; 2015; 160: 3053–3062. <https://doi.org/10.1007/s00705-015-2609-6> PMID: 26395091
 53. Goldsmith DB, Crosti G, Dwivedi B, McDaniel LD, Varsani A, Suttle CA, et al. Development of phoH as a novel signature gene for assessing marine phage diversity. Appl Environ Microbiol. 2011; 77: 7730–7739. <https://doi.org/10.1128/AEM.05531-11> PMID: 21926220
 54. Zeng Q, Chisholm SW. Marine viruses exploit their host’s two-component regulatory system in response to resource limitation. Curr Biol. Elsevier Ltd; 2012; 22: 124–128. <https://doi.org/10.1016/j.cub.2011.11.055> PMID: 22244998
 55. Dwivedi B, Xue B, Lundin D, Edwards R a, Breitbart M. A bioinformatic analysis of ribonucleotide reductase genes in phage genomes and metagenomes. BMC Evol Biol. 2013; 13: 33. <https://doi.org/10.1186/1471-2148-13-33> PMID: 23391036
 56. Dwivedi B, Xue B, Lundin D, Edwards R a, Breitbart M. A bioinformatic analysis of ribonucleotide reductase genes in phage genomes and metagenomes. BMC Evol Biol. 2013; 13: 33. <https://doi.org/10.1186/1471-2148-13-33> PMID: 23391036
 57. Perry KM, Fauman EB, Finer-Moore JS, Montfort WR, Maley GF, Maley F, et al. Plastic adaptation toward mutations in proteins: Structural comparison of thymidylate synthases. Proteins Struct Funct Bioinforma. 1990; 8: 315–333. <https://doi.org/10.1002/prot.340080406> PMID: 2128651
 58. Chu FK, Maley GF, Maley F, Belfort M. Intervening sequence in the thymidylate synthase gene of bacteriophage T4. Proc Natl Acad Sci U S A. 1984; 81: 3049–53. <https://doi.org/10.1073/pnas.81.10.3049> PMID: 6328492
 59. Bechhofer DH, Hue KK, Shub DA. An intron in the thymidylate synthase gene of Bacillus bacteriophage b22: Evidence for independent evolution of a gene, its group I intron, and the intron open reading frame. Proc Natl Acad Sci U S A. 1994; 91: 11669–11673. PMID: 7972121
 60. Edgell DR, Chalamcharla VR, Belfort M. Learning to live together: mutualism between self-splicing introns and their hosts. BMC Biol. 2011; 9: 22. <https://doi.org/10.1186/1741-7007-9-22> PMID: 21481283
 61. Lavigne R, Vandersteegen K. Group I introns in Staphylococcus bacteriophages. Future Virol. 2013; 8: 997–1005. <https://doi.org/10.2217/fvl.13.84>
 62. Edgell DR, Gibb EA, Belfort M. Mobile DNA elements in T4 and related phages. Virol J. 2010; 7: 290. <https://doi.org/10.1186/1743-422X-7-290> PMID: 21029434
 63. Belfort M, Roberts RJ. Homing endonucleases: Keeping the house in order. Nucleic Acids Res. 1997; 25: 3379–3388. <https://doi.org/10.1093/nar/25.17.3379> PMID: 9254693
 64. Hausner G, Hafez M, Edgell DR. Bacterial group I introns: mobile RNA catalysts. Mob DNA. 2014; 5: 8. <https://doi.org/10.1186/1759-8753-5-8> PMID: 24612670
 65. Nielsen H. Group I Intron Ribozymes. In: Hartig JS, editor. Ribozymes: Methods and Protocols. Totowa, NJ: Humana Press; 2012. pp. 73–89. https://doi.org/10.1007/978-1-61779-545-9_6 PMID: 22315064
 66. Kala S, Cumby N, Sadowski PD, Hyder BZ, Kanelis V, Davidson AR, et al. HNH proteins are a widespread component of phage DNA packaging machines. Proc Natl Acad Sci U S A. 2014; 111: 6022–7. <https://doi.org/10.1073/pnas.1320952111> PMID: 24711378
 67. Fouts DE, Klumpp J, Bishop-Lilly KA, Rajavel M, Willner KM, Butani A, et al. Whole genome sequencing and comparative genomic analyses of two *Vibrio cholerae* O139 Bengal-specific Podoviruses to other N4-like phages reveal extensive genetic diversity. Virol J. 2013; 10: 165. <https://doi.org/10.1186/1743-422X-10-165> PMID: 23714204