



RESEARCH

Open Access

Estrogen receptor α (ER α) mediates 17 β -estradiol (E2)-activated expression of HBO1

Wen-zhong Wang^{1†}, Hai-ou Liu^{1,3†}, Yi-hong Wu¹, Yi Hong¹, Jun-wu Yang¹, Ye-heng Liu², Wei-bin Wu², Lei Zhou¹, Lin-lin Sun^{1,3}, Jie-jie Xu^{1*}, Xiao-jing Yun^{1*}, Jian-xin Gu²

Abstract

Background: HBO1 (histone acetyltransferase binding to ORC1) is a histone acetyltransferase (HAT) which could exert oncogenic function in breast cancer. However, the biological role and underlying mechanism of HBO1 in breast cancer remains largely unknown. In the current study, we aimed to investigate the role of HBO1 in breast cancer and uncover the underlying molecular mechanism.

Methods: Immunohistochemistry was applied to detect HBO1 protein expression in breast cancer specimens (n = 112). The expression of protein level was scored by integral optical density (IOD) for further statistical analyses using SPSS. Real-time PCR was used to simultaneously measure mRNA levels of HBO1. The HBO1 protein expression in breast cancer cells was confirmed by western blot.

Results: HBO1 was highly expressed in breast cancer tissues and significantly correlated with estrogen receptor α (ER α) (p < 0.001) and progesterone hormone (PR) (p = 0.002). HBO1 protein level also correlated positively with histology grade in ER α positive tumors (p = 0.016) rather than ER α negative tumors. 17 β -estradiol (E2) could upregulate HBO1 gene expression which was significantly inhibited by ICI 182,780 or ER α RNAi. E2-increased HBO1 protein expression was significantly suppressed by treatment with inhibitor of MEK1/2 (U0126) in T47 D and MCF-7 cells.

Conclusions: HBO1 was an important downstream molecule of ER α , and ERK1/2 signaling pathway may involved in the expression of HBO1 increased by E2.

Background

Breast cancer is a major public health issue, with more than one million new cases observed around the world in 2002 [1]. The pathogenesis of breast cancer is quite complex. Lifetime exposure to estrogen is reported to be associated with women's risk for breast cancer and the biological actions of estrogens are mediated primarily by ER α which belongs to the nuclear receptor superfamily, a family of ligand-regulated transcription factors [2-4]. ER α , which promotes cell growth, metastasis and also mediates resistance to apoptosis, plays a key role in progression of breast cancer [5,6].

HBO1 (histone acetyltransferase binding to ORC1), also named MYST2, belongs to the MYST family which is

characterized by a highly conserved C2HC zinc finger and a putative histone acetyltransferase domain. The role of HBO1 in cancer remains unclear, although its expression has been reported in testicular germ cell tumors, breast adenocarcinomas, and ovarian serous carcinomas [7]. Recent investigations have revealed that over-expression of HBO1 dramatically enhances the anchorage-independent growth of both MCF7 and SKBR3 breast cancer cells [8]. Furthermore, it also functions as a transcriptional coactivator for hormone receptors including ER α and PR [9], leading to consideration of this protein as a carcinogenic factor.

To elucidate the role of HBO1 in breast cancer, 112 breast cancer specimens were investigated using immunohistochemistry with an anti-HBO1 antibody. We found that HBO1 was notably increased in breast cancer. E2-upregulated HBO1 expression could be inhibited by ICI 182,780 or ER α RNAi in breast cancer cells.

* Correspondence: jxufdu@hotmail.com; xjyun@fudan.edu.cn

† Contributed equally

¹Department of Biochemistry and Molecular Biology, Shanghai Medical College of Fudan University, Shanghai, 200032, China

Full list of author information is available at the end of the article

Furthermore, we also showed that ERK1/2 signaling pathway was involved in the expression of HBO1 increased by E2.

Methods

Materials

Dulbecco's modified Eagle's medium (DMEM), phenol red-free DMEM (PR-free DMEM), U0126 and 17 β -estradiol (E2) were purchased from Sigma. Lipofectamine 2000, Trizol Reagent and fetal bovine serum (FBS) were purchased from Invitrogen. Charcoal-stripped fetal bovine serum (CS-FBS) was purchased from Biowest. PVDF membrane, leupeptin, aprotinin, phenylmethylsulfonyl fluoride (PMSF) and X-tremeGENE siRNA Transfection Reagent were purchased from Roche. RNA PCR Kit (AMV) Ver.3.0 was purchased from TaKaRa. Polinl-2-plus kit was a product of GBI. Anti-phospho-ERK1/2 (Thr202/Tyr204) and anti-ERK1/2 antibodies were purchased from Cell Signaling Technology. ER α siRNA, mouse monoclonal anti-ER α , anti-GAPDH, horseradish peroxidase (HRP)-conjugated goat anti-rabbit and HRP-conjugated goat anti-mouse IgG secondary antibodies were from Santa Cruz Biotechnology. Rabbit polyclonal anti-HBO1 antibody (Catalog No: 13751-1-AP) was purchased from Protein Tech Group, Inc. The enhanced chemiluminescence (ECL) assay kit was purchased from Tiangen. Dual-luciferase reporter assay system was bought from Promega. ICI 182,780 was purchased from Tocris Bioscience.

Cell culture and transfection

Human MCF-7, T47 D, MDA-MB-453 and MDA-MB-435 breast cancer cells were obtained from the Cell Bank of Type Culture Collection of Chinese Academy of Sciences. MCF-7 cells were cultured in DMEM supplemented with 10% fetal calf serum (FBS), 10 μ g/ml of insulin, 100 U/ml of penicillin and 50 μ g/ml of streptomycin. T47 D cells were maintained in DMEM supplemented with 10% FBS, 100 U/ml of penicillin and 50 μ g/ml streptomycin. SiRNA was transfected with X-tremeGENE siRNA Transfection Reagent according to the manufacturer's instructions.

Western blot analyses

Proteins were detected by western blot analysis as described previously [10]. The cells were lysed by lysis buffer, separated in 10% SDS-PAGE and then transferred to a PVDF membrane. The membrane was incubated with primary antibody followed by incubation with horseradish peroxidase-conjugated secondary antibody. Then the membrane was developed using the ECL detection system.

Tissue samples

112 primary breast cancer specimens were consecutively obtained from pathological archives of Huashan Hospital,

Fudan University from 2005 to 2008. There was no any bias for selection. Tissues were formalin-fixed and paraffin-embedded for histopathologic diagnosis and immunohistochemical study. The malignancy degree of tumor was scored according to the Scarff-Bloom-Richardson system.

Immunohistochemistry (IHC)

Serial sections (5 μ m thick) were mounted on glass slides coated with 10% polylysine. Sections were dewaxed in xylene and rehydrated in graded ethanol. Endogenous peroxidase activity was blocked by immersion in 0.3% methanolic peroxide for 30 minutes. Next, sections were microwaved in citrate buffer for antigen retrieval. Rabbit polyclonal anti-HBO1 (1:100 dilutions) was used as primary antibodies. Staining was completed with Polink-2-plus kit, in accordance with the manufacturer's instructions. Color reaction product was developed with diaminobenzidine. All sections were counterstained with hematoxylin. Two pathologists separately blinded to the clinical outcome of the patients evaluated all samples. The immunoreactivity intensity was evaluated as 0 (absent); 1 (weak nuclear staining); 2 (moderate nuclear staining of intermediate level); 3 (more coarse nuclear staining). Positive cells were quantified as a percentage of the total number of tumor cells with observation of 1000 cells in 5 high power field (\times 400), and assigned to one of five categories: 0: < 5%, 1: 5-25%, 2: 26-50%, 3: 51-75% and 4: > 75%. HBO1 expression was scored semi-quantitatively using the Remmele-score (immunoreactive score (IS) = score of percentage of positive tumor cells \times score of staining intensity). Then the slide of IS > 3 was classified as a positive case [11]. Reproducibility of the scoring method used by both observers was greater than 95%.

Total RNA extraction and real-time PCR

Total mRNA samples of T47 D and MCF-7 breast cancer cells were extracted using trizol reagent according to the manufacturer's instructions. One microgram of total RNA extracted from the cells was subjected to reverse transcription (RT). The RT and real-time PCR were performed by using TaKaRa RNA PCR Kit (AMV) Ver.3.0 and SYBR Premix Ex Taq II according to the manual respectively. Primers used for real-time PCR were as follows: HBO1-F 5'-GATGCCCACTGTATCATAACC-3' and HBO1-R 5'-TTCTTCCTGAGTTCAGCCACT-3'; GAPDH-F 5'-GGCTGAGAACGGGAAGCTTGTCAT-3' and GAPDH-R 5'-CAGCCTTCTCCATGGTGGTGAAGA-3'. Real-time PCR was performed using 7500 multicolor real-time PCR detection system (ABI) with the following cycling conditions: (i) 30s at 95°C and (ii) 40 cycles, with 1 cycle consisting of 5s at 95°C, 34s at

60°C. GAPDH was employed as an internal control under the same experimental conditions. Data were analyzed by using 7500 software (ABI). The values were obtained through normalizing to GAPDH copies.

Statistical analysis

Statistical data analyses were performed using SPSS 11.5 statistical software package. The relationship between protein levels and different clinical and pathological features were explored using cross tabulation and Pearson's χ^2 . P values less than 0.05 were selected.

Results

HBO1 protein level correlates positively with ER α

In order to explore the biological role of HBO1 in human breast cancer *in vivo*, we examined the expression of HBO1 in tumor samples of primary breast cancer (n = 112) by IHC analysis. We found that HBO1 which was predominantly detected in nucleus (Figure 1B-C) was

highly expressed in breast cancer tissues (Table 1) and significantly correlated with ER α (p < 0.001) and PR (p = 0.002) (Table 2). Further statistical analysis revealed HBO1 protein level correlated positively with histology grade in ER α positive tumors (p = 0.016) rather than ER α negative tumors (Table 2). For benign breast tissues, low HBO1 immunoreactivity was observed and epithelial cells displayed a minimal granular staining (Figure 1A). Moderately and poorly differentiated breast cancer tissues showed intense HBO1 staining (Figure 1B-C). To further study the relationship between HBO1 and ER α , we examined HBO1 expression level in several breast cancer cell lines by western blot, which showed that ER α positive breast cancer cell lines exhibited higher HBO1 protein than ER α negative breast cancer cell lines (Figure 1D).

E2 induces HBO1 expression in breast cancer cells

In order to further investigate the relationship between ER α and HBO1, we treated breast cancer cells with




Figure 1 Expression of HBO1 in human breast cancer. (A) Immunohistochemical staining for HBO1 in benign breast epithelial tissues (magnification: A, 400X). (B) Moderately differentiated breast cancer tissues (brown staining) (magnification: B, 400X). (C) Poorly differentiated breast cancer tissues (brown staining) (magnification: C, 400X). (D) HBO1 protein level in several breast cancer cell lines based on the western blot results.

Table 1 Clinical and pathological characteristics of patients

characteristic	value
Tumor grade (n[%])	
I	45 [40.2%]
II/III	67 [59.8%]
Ki67 status (n[%])	
negative	50 [44.6%]
positive	62 [55.4%]
Estrogen receptor α status (n[%])	
negative	39 [34.8%]
positive	73 [65.2%]
Progesterone receptor status (n[%])	
negative	60 [53.6%]
positive	52 [46.4%]
P53 status (n[%])	
Negative	62 [55.4%]
Positive	50 [44.6%]
HBO1 status (n[%])	
negative	48 [42.9%]
positive	64 [57.1%]
Histology Grade (n[%])	
G1	32 [28.6%]
G2/G3	80 [71.4%]

17 β -estradiol (E2). Quantitative real-time PCR was used to determine the effect of E2 on the mRNA level of HBO1. With the dose of E2 increasing from 10⁻⁹ to 10⁻⁷ M, the mRNA level of HBO1 gradually increased and reached a plateau at 10⁻⁸ M in T47 D cells (Figure 2A). Hereby, 10⁻⁸ M of E2 was applied for all subsequent experiments. Western blot was also performed to confirm that E2 could upregulate HBO1 protein expression in T47 D (Figure 2B) and MCF-7 cells (Figure 2C). Taken together, these results demonstrated that E2 increased expression of HBO1 at transcriptional level.

E2-induced HBO1 expression is inhibited by ICI 182,780 or ER α RNAi

To further study the role of ER α in HBO1 expression, ICI 182,780 was further applied in T47 D cells (Figure 3A) and MCF-7 cells (Figure 3B) to block the effect of E2. As shown in Figure 3A, E2 increased HBO1 protein expression (lane 2), which was significantly reduced by ICI 182,780 (lane 4). Similar results were obtained in MCF-7 cells (Figure 3B). ICI 182,780 was reported to act by binding ER α , causing disassociation of ER α associated proteins and resulting in impaired receptor dimerisation and increased receptor degradation [12,13]. As expected, ER α expression was decreased by ICI 182,780 (Figure 3A and 3B). These results indicated that

Table 2 Expression of HBO1 in relation to the clinical and pathological characteristics of patients



clinical feature	total	HBO1 expression		P-value
		negative	positive	
Tumor grade				
I	45	21	24	0.610
II/III	67	28	39	
Ki67				
negative	50	23	27	0.550
positive	62	25	37	
Estrogen receptor				
negative	39	26	13	< 0.001**
positive	73	22	51	
Progesterone receptor				
negative	60	34	26	0.002**
positive	52	14	38	
P53				
negative	62	23	39	0.170
positive	50	25	25	
Histology Grade				
G1	32	17	15	0.165
G2/G3	80	31	49	
among ER positive tumors				
Histology Grade				
G1	25	12	13	0.016*
G2/G3	48	10	38	
among ER negative tumors				
Histology Grade				
G1	8	5	3	0.779
G2/G3	31	21	10	

*P < 0.05 **P < 0.01

E2-induced HBO1 upregulation could be inhibited by ICI 182,780. In order to figure out the role of ER α in E2-induced HBO1 upregulation, ER α siRNA was transfected into T47 D and MCF7 cells to knockdown ER α . We observed that E2-induced HBO1 upregulation was inhibited by ER α siRNA (lane 4), indicating that HBO1 might be a downstream signaling molecule for ER α .

ERK1/2 signaling pathway was involved in the expression of HBO1 increased by E2

Using TRSEARCH software based on the sequence database, the promoter region of the HBO1 gene has no putative binding sites for ER α (data not shown). And previous studies have shown that steroidal estrogens are able to rapidly and transiently activate the ERK1/2 pathway [13], so we examined whether the ERK1/2 pathway was involved in the effect of E2 on HBO1 expression in breast cancer cells. As shown in Figure 4, E2-increased HBO1 protein expression was significantly suppressed




by treatment with inhibitor of MEK1/2 (U0126) in T47 D (Figure 4A) and MCF-7 (Figure 4B) cells as analyzed by western blot. These results indicated that ERK1/2 signaling pathway was involved in the E2-induced HBO1 upregulation in breast cancer cells.

Discussion

HBO1 is a potential oncogene which maps to17q21.3, a region where frequent allelic gains are found in breast cancers and this amplification is associated with a poor prognosis of clinical outcome [14-16]. Previous studies demonstrated over-expression of HBO1 dramatically enhanced the anchorage-independent growth of both MCF7 and SKBR3 breast cancer cells while depletion of HBO1 reduced the rate of DNA synthesis, the amount of MCM complex bound to chromatin, and progression through S phase. HBO1 has also been shown to enhance transcription mediated by steroid receptors including ER α and PR [9]. However, little is known about the role of HBO1 in breast cancer and the underlying molecular mechanism.

In this study, we first investigated the HBO1 protein expression in large numbers of tumor samples of



primary breast cancer (n = 112) by IHC analysis, and showed that HBO1 was highly expressed in breast cancer (Table 1) and positively correlated with ER α (p < 0.001) and PR (p = 0.002). Moreover, HBO1 protein level correlated positively with histology grade in ER positive tumors (p = 0.016) rather than ER α negative tumors through statistical analysis. As a coactivator of the replication licensing factor Cdt1 [17], HBO1 belongs to one component of the Replication Initiation Proteins known as prereplicative complex (pre-RC) proteins. Several pre-RC proteins are over-expressed in cancer and serve as good tumor markers. And some of them, such as Cdc6 and Cdt1, are elevated by E2 treatment in breast cancer. To determine whether HBO1 was also affected by E2, quantitative real-time PCR and western blot were performed. The results suggested HBO1 was elevated after E2 treatment. Further study demonstrated the E2-induced HBO1 upregulation could be inhibited by ICI 182,780 as well as ER α siRNA.

Estrogen receptor (ER) signaling is involved in many human diseases such as breast cancer [18], lung cancer [19,20] and cardiovascular diseases [21,22]. It was reported that E2 activated the signal-transducing ERK1/2 pathway, which were critical for cell proliferation [23-25], in human mammary cancer-derived cell lines, MCF-7 and T47 D [26-28]. We explored whether ERK1/2 signaling pathway was involved in the expression of HBO1 increased by E2. The MEK1/2 inhibitor U0126 significantly inhibited the expression of HBO1 in T47 D and MCF-7 cells, suggesting that E2 increased the expression of HBO1 through the ERK1/2 signaling pathway. Since previous studies have shown that progesterone receptor activates the Src/p21ras/ERK pathway via cross-talk with estrogen receptor in breast cancer [29], the positive correlation between HBO1 protein levels and PR which we obtained in statistical analysis was reasonable.

Estrogen exposure has been regarded as high risk factor of breast cancer. It has been reported that HBO1 strongly enhanced ER-mediated transcription [9], which indicated that HBO1 might play a role in the progress of breast cancer. In this study, we proved E2 could upregulate the mRNA and protein level of HBO1. Meanwhile, knock-down of ER α with siRNA significantly inhibited the upregulation of HBO1, indicating that cross-talking was happening between ER α and HBO1 in breast cancer, the biological functions of which need to be further studied.

Conclusion

Our data have demonstrated that the HBO1 protein levels correlated positively with ER α (p < 0.001) and PR (p = 0.002) expression in breast cancer. Further more, HBO1 protein levels correlated positively with histology grade in ER α positive tumors (p = 0.016). We also showed that the ERK1/2 signaling pathway was involved

in the expression of HBO1 increased by E2. These findings suggested a potential for targeting HBO1 as a novel means of breast cancer therapy as well as a potential diagnosis marker for ER α positive breast cancer.

Acknowledgements

This work is supported by grants from National Natural Science Funds: 30770426, 30930025, 30900266 and 31000348; State Key Project Specialized for Infectious Diseases: 2008ZX10002-015, 2008ZX10002-021, 2008ZX10001-02 and 2008ZX10004-014; National Basic Research Program of China (973 Program): 2010CB912100 (2010CB912104); National Fundamental Fund Project: J0730860; Shanghai Leading Academic Discipline Project: B110.

Author details

¹Department of Biochemistry and Molecular Biology, Shanghai Medical College of Fudan University, Shanghai, 200032, China. ²Institutes of Biomedical Science of Fudan University, Shanghai, 200032, China. ³Institute of Nautical Medicine, Nantong University, Nantong, 226001, China.

Authors' contributions

WWZ carried out the design of the study, performed IHC, real-time PCR, drafted the manuscript. LHO performed the western blot. WYH participated in SPSS Statistical Analysis. LYH participated in IHC and IOD scoring. WWB participated in real-time PCR and cell culture. ZL performed SPSS Statistical Analysis. HY and YJW participated in IHC. SLL participated in collection of breast cancer specimens. XJJ participated in the design of the study, drafted the figure. YXJ performed the design of the study, and helped drafting the manuscript. GJX performed the collection of breast cancer specimens. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 29 July 2010 Accepted: 1 November 2010

Published: 1 November 2010

References

1. Parkin DM, Bray F, Ferlay J, Pisani P: Global cancer statistics, 2002. *CA Cancer J Clin* 2005, **55**:74-108.
2. Liao DZ, Pantazis CG, Hou X, Li SA: Promotion of estrogen-induced mammary gland carcinogenesis by androgen in the male Noble rat: probable mediation by steroid receptors. *Carcinogenesis* 1998, **19**:2173-80.
3. Pettersson K, Gustafsson JA: Role of estrogen receptor beta in estrogen action. *Annu Rev Physiol* 2001, **63**:165-92.
4. Hilakivi-Clarke L, Cho E, Cabanes A, et al: Dietary modulation of pregnancy estrogen levels and breast cancer risk among female rat offspring. *Clin Cancer Res* 2002, **8**:3601-10.
5. Clarke R, Liu MC, Bouker KB, et al: Antiestrogen resistance in breast cancer and the role of estrogen receptor signaling. *Oncogene* 2003, **22**:7316-39.
6. O'Lone R, Frith MC, Karlsson EK, Hansen U: Genomic targets of nuclear estrogen receptors. *Mol Endocrinol* 2004, **18**:1859-75.
7. Iizuka M, Takahashi Y, Mizzen CA, et al: Histone acetyltransferase Hbo1: catalytic activity, cellular abundance, and links to primary cancers. *Gene* 2009, **436**:108-14.
8. Hu X, Stern HM, Ge L, et al: Genetic alterations and oncogenic pathways associated with breast cancer subtypes. *Mol Cancer Res* 2009, **7**:511-22.
9. Georgiakiaki M, Chabbert-Buffer N, Dasen B, et al: Ligand-controlled interaction of histone acetyltransferase binding to ORC-1 (HBO1) with the N-terminal transactivating domain of progesterone receptor induces steroid receptor coactivator 1-dependent coactivation of transcription. *Mol Endocrinol* 2006, **20**:2122-40.
10. Wang Y, Zong H, Chi Y, et al: Repression of estrogen receptor alpha by CDK11p58 through promoting its ubiquitin-proteasome degradation. *J Biochem* 2009, **145**:331-43.
11. Remmele W, Stegner HE: Recommendation for uniform definition of an immunoreactive score (IRS) for immunohistochemical estrogen receptor detection (ER-ICA) in breast cancer tissue. *Pathologie* 1987, **8**:138-40.
12. MacGregor JI, Jordan VC: Basic guide to the mechanisms of antiestrogen action. *Pharmacol Rev* 1998, **50**:151-96.

13. Wakeling AE: Similarities and distinctions in the mode of action of different classes of antioestrogens. *Endocr Relat Cancer* 2000, **7**:17-28.
14. Clark J, Edwards S, John M, *et al*: Identification of amplified and expressed genes in breast cancer by comparative hybridization onto microarrays of randomly selected cDNA clones. *Genes Chromosomes Cancer* 2002, **34**:104-14.
15. Hyman E, Kauraniemi P, Hautaniemi S, *et al*: Impact of DNA amplification on gene expression patterns in breast cancer. *Cancer Res* 2002, **62**:6240-5.
16. Pollack JR, Sorlie T, Perou CM, *et al*: Microarray analysis reveals a major direct role of DNA copy number alteration in the transcriptional program of human breast tumors. *Proc Natl Acad Sci USA* 2002, **99**:12963-8.
17. Miotto B, Struhl K: HBO1 histone acetylase is a coactivator of the replication licensing factor Cdt1. *Genes Dev* 2008, **22**:2633-8.
18. Yager JD, Davidson NE: Estrogen carcinogenesis in breast cancer. *N Engl J Med* 2006, **354**:270-82.
19. Stabile LP, Siegfried JM: Estrogen receptor pathways in lung cancer. *Curr Oncol Rep* 2004, **6**:259-67.
20. Marquez-Garban DC, Chen HW, Fishbein MC, Goodglick L, Pietras RJ: Estrogen receptor signaling pathways in human non-small cell lung cancer. *Steroids* 2007, **72**:135-43.
21. Huss JM, Kelly DP: Nuclear receptor signaling and cardiac energetics. *Circ Res* 2004, **95**:568-78.
22. Meyer MR, Haas E, Barton M: Gender differences of cardiovascular disease: new perspectives for estrogen receptor signaling. *Hypertension* 2006, **47**:1019-26.
23. Atanaskova N, Keshamouni VG, Krueger JS, Schwartz JA, Miller F, Reddy KB: MAP kinase/estrogen receptor cross-talk enhances estrogen-mediated signaling and tumor growth but does not confer tamoxifen resistance. *Oncogene* 2002, **21**:4000-8.
24. Martin LA, Farmer I, Johnston SR, Ali S, Dowsett M: Elevated ERK1/ERK2/estrogen receptor cross-talk enhances estrogen-mediated signaling during long-term estrogen deprivation. *Endocr Relat Cancer* 2005, **12**(Suppl 1):S75-84.
25. Santen RJ, Song RX, McPherson R, *et al*: The role of mitogen-activated protein (MAP) kinase in breast cancer. *J Steroid Biochem Mol Biol* 2002, **80**:239-56.
26. Migliaccio A, Pagano M, Auricchio F: Immediate and transient stimulation of protein tyrosine phosphorylation by estradiol in MCF-7 cells. *Oncogene* 1993, **8**:2183-91.
27. Auricchio A, di Domenico M, Castoria G, Bilancio A, Migliaccio A: Epidermal growth factor induces protein tyrosine phosphorylation and association of p190 with ras-GTP-ase activating protein in Caco-2 cells. *FEBS Lett* 1994, **353**:16-20.
28. Auricchio F, Migliaccio A, Castoria G, Di Domenico M, Bilancio A, Rotondi A: Protein tyrosine phosphorylation and estradiol action. *Ann N Y Acad Sci* 1996, **784**:149-72.
29. Migliaccio A, Piccolo D, Castoria G, *et al*: Activation of the Src/p21ras/Erk pathway by progesterone receptor via cross-talk with estrogen receptor. *EMBO J* 1998, **17**:2008-18.

doi:10.1186/1756-9966-29-140

Cite this article as: Wang *et al.*: Estrogen receptor α (ER α) mediates 17 β -estradiol (E2)-activated expression of HBO1. *Journal of Experimental & Clinical Cancer Research* 2010 **29**:140.

Submit your next manuscript to BioMed Central
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

