

Editorial

Open Access

## Towards a framework for the evolutionary genomics of Kinetoplastids: what kind of data and how much?

Alberto MR Dávila\*<sup>1</sup> and Julius Lukeš<sup>2</sup>

Address: <sup>1</sup>Lab. de Biologia Molecular de Tripanosomatídeos, Departamento de Bioquímica e Biologia Molecular, Instituto Oswaldo Cruz, Fiocruz, Av. Brasil 4365, Rio de Janeiro, RJ, Brasil. 21045-900 and <sup>2</sup>Institute of Parasitology, Czech Academy of Sciences and Faculty of Biology, University of South Bohemia, Branišovská 31, 37005 Ceske Budejovice, Czech Republic

Email: Alberto MR Dávila\* - [davila@fiocruz.br](mailto:davila@fiocruz.br); Julius Lukeš - [jula@paru.cas.cz](mailto:jula@paru.cas.cz)

\* Corresponding author

Published: 28 October 2003

Received: 30 July 2003

*Kinetoplastid Biology and Disease* 2003, **2**:16

Accepted: 28 October 2003

This article is available from: <http://www.kinetoplastids.com/content/2/1/16>

© 2003 Dávila and Lukeš; licensee BioMed Central Ltd. This is an Open Access article: verbatim copying and redistribution of this article are permitted in all media for any purpose, provided this notice is preserved along with the article's original URL.

### Abstract

The current status of kinetoplastids phylogeny and evolution is discussed in view of the recent progresses on genomics. Some ideas on a potential framework for the evolutionary genomics of kinetoplastids are presented.

### Editorial

In this journal, the concept of species are presented by Wendy Gibson [1] and Hooman Momen [2] while Michel Tibayrenc [3] reviews the evolution and variability of *Trypanosoma cruzi*. Veronique Hannaert and collaborators [4] provide a comprehensive review on the evolution of energy metabolism in Kinetoplastida and Austin Hughes and Helen Piontkivska [5] discuss the controversial issue of the monophyletic versus paraphyletic status of the genus *Trypanosoma*. The latter authors concluded that the markers used to date failed to provide a consistent answer about the origin of trypanosomes. The arguments for the use of more protein-based sequences and a larger number of species are admirable. However, some important questions remain: which genes and how many are needed? Which other species should be selected and how many should be explored?

Generally, a limited number of genes available for a large number of taxa is sufficient for most phylogenetic studies. However, in order to resolve the relationships among closely related species and strains, and to better understand their origin, the availability of a set of evolutionary informative genes would certainly be useful. The ribosomal (r)RNA genes (18S, 5S, SL) still dominate the field,

while only a few informative protein-coding genes have been described. The identification and choice of individual genes or gene families is sometimes difficult and less cost-effective when compared with high throughput sequencing initiatives [6]. This observation makes us to reflect about the need to discover new candidate genes for the study of kinetoplastid evolution. According to Podlipaev (2000, 2001) [7,8], from the 10 trypanosomatid genera [9], those (six) associated with insects have so far received little attention or have been "neglected" compared to species with an impact on human health. The same is true for all bodonids.

Still, Trypanosomatina flagellates are in the forefront among kinetoplastids, since several species (*T. cruzi*, *T. brucei* and *Leishmania major*) will have their genomes completely sequenced within the next few months. While the recent start of genome projects for *L. braziliensis*, *L. infantum*, *T. vivax* and *T. congolense* is very welcome, it is worth noticing that all the above mentioned species belong to only two (*Trypanosoma* and *Leishmania*) of the 20 kinetoplastid genera, showing the stark under-representation of the other genera in the sequencing initiative.

**Table 1: Survey of kinetoplastid sequences available in the Genbank.**

	Nucleotides	Proteins
<b><u>Trypanosomatina</u></b>		
<i>Trypanosoma</i>	139667	3695
<i>Leishmania</i>	29777	2945
<i>Crithidia</i>	163	206
<i>Herpetomas</i>	137	15
<i>Phytomonas</i>	112	12
<i>Leptomonas</i>	55	23
<i>Blastocrithidia</i>	12	7
<i>Wallaceina</i>	10	3
<i>Endotrypanum</i>	9	5
<b><u>Bodonina</u></b>		
<i>Bodo</i>	64	12
<i>Trypanoplasma</i>	22	24
<i>Cryptobia</i>	14	5
<i>Dimastigella</i>	9	1
<i>Rhynchomonas</i>	6	1
<i>Perkinsiella</i>	6	0
<i>Rhynchobodo</i>	3	0
<i>Cruzella</i>	3	0
<i>Ichthyobodo</i>	2	0
<i>Parabodo</i>	2	0
<i>Proccryptobia</i>	1	0

A casual search in the Entrez division of NCBI showed that although virtually omnipresent in nature, leptomonads and phytomonads have only dozens of entries, a mere trickle when compared with 139667 and 29777 entries for *Trypanosoma* and *Leishmania*, respectively (see Table 1). It is clear from the pattern of data available that efforts should be focused on obtaining more sequences from the "non-*Leishmania*" and "non-*Trypanosoma*" kinetoplastids. Centering on one "representative" or "model" species for each of the underrepresented genera *Bodo*, *Trypanoplasma* and *Cryptobia* would certainly be both worthwhile and rewarding.

Having a genome fully sequenced is an expensive initiative (although the cost per base is decreasing annually, rendering the whole genome sequencing efforts increasingly cost-effective), then centering the resources on EST (expressed sequence tags) projects to explore the transcriptome of some species would be certainly helpful for the comparative "transcriptomics" of kinetoplastids. It has been demonstrated that EST projects are relatively cheap when compared with whole genome shotgun strategies, with the advantage that a kinetoplastid EST project would provide with the coding sequences that would be of great help for evolutionary genomics. As an example, it would be particularly interesting to know more about the evolution and polymorphism of kinetoplastid-specific genes associated with metabolic pathways or even mark-

ers for typing. On the other hand, a potential disadvantage of the EST approach would be that one could not unequivocally evaluate the gene loss and genome structure because only a part of the genome would be surveyed. However, studies on gene and protein content, orthologs and paralogs would be perfectly allowed. Another advantage of having the transcriptome of other trypanosomatids sequenced would be the usefulness of such sequences for the annotation of the *T. cruzi*, *L. major* and *T. brucei* genomes. The sequencing consortia for these three species have been reporting a high number (sometimes >70 %) of orphan genes during the annotation process [10,11]. New data from other trypanosomatids may help to discover which of them are real "orphans" and/or species-specific genes, and which ones are genes confined to Kinetoplastida.

An important issue towards the exploration of the transcriptome of kinetoplastids outside Trypanosomatina is the choice of "model species". These "models" would be used to get insights into the evolution of bizarre structures and processes so far mostly described in *Trypanosoma* and less so in *Leishmania*. One proposal could be the study of the evolution of VSGs, as well as complex proteins involved in RNA editing or trans-splicing. A comparative approach has recently proved to be useful for our understanding of how the complex kinetoplast DNA network may have evolved [12]. The "models" should be members of major kinetoplastid clades in the current phylogenetic trees [13] that grow in reasonable densities in simple culture media, preferably axenic. *Bodo saltans* may be a good model, because it branches off on the border between bodonids and trypanosomatids, and is an ecologically important free-living species of potentially world-wide distribution. Its sole disadvantage is the growth with feeder bacteria. As far as we know, the only bodonid that grows in an axenic medium is *Trypanoplasma borreli*, a parasite of marine and freshwater fish, another potential candidate. The final proposed candidate is *Diplonema papillatum*, since it grows axenically in simple media to high densities. Moreover, it would be an excellent out-group for kinetoplastids, since it represents the most related group and likely shares with them numerous peculiarities.

Inferring from the recently unveiled diversity of extremely small eukaryotes [14], one would predict that a great diversity of kinetoplastids is still to be discovered. Perhaps an enthusiastic overestimate, some discuss the existence of a great trypanosomatid biodiversity in insects alone [15]. Recent surveys have found new species in ocean depths [16,17] or unexpected niches [18]. One of the approaches used for the discovery of these new eukaryote taxa is culture-independent-survey by PCR (ciPCR), which has also helped to double the number of prokaryotic

phyla [19]. This approach would be particularly interesting in a search for difficult-to-cultivate Kinetoplastida. For such a survey, a number of known conserved markers/genes would be needed for the design of PCR-based assays and, unfortunately, that is not possible with the limited data currently available. There may be some requirements for choosing the markers since according to Charlebois and collaborators [20] protein-coding regions are usually less universal, more difficult to PCR-amplify and often shorter and less information-rich than the rRNA genes. Possible discrepancies between the rRNA- and protein-based trees can be avoided by choosing informative and unbiased genes and by constructing trees on the basis of concatenated alignments. Whole-genome trees are becoming popular given the availability of a large number of bacterial genomes in the databases, however, for the near future, it is not going to be an option for non-pathogenic kinetoplastids and in general for eukaryotes because of the size of their genomes. New technologies for sequencing large genomes and cheaper costs per base could, in the future, facilitate that approach even for protozoans.

It seems that with the available genomics approaches, there are now more chances to map the real diversity of kinetoplastids in nature, and explore their fascinating biology, evolution and "roots".

### Acknowledgements

We are very grateful to Wendy Gibson, Sergei Podlipaev and Hooman Momen for their comments on the manuscript, however, all mistakes and wrong interpretations remain ours.

### References

- Gibson W: **Species concepts for trypanosomes: from morphological to molecular definitions?** *Kinetoplastid Biology and Disease* 2003, **2(10)**..
- Momen H: **Concepts of species in trypanosomatids.** *Kinetoplastid Biology and Disease* 2003, **2(13)**..
- Tibayrenc M: **Genetic subdivisions within *Trypanosoma cruzi* (Discrete Typing Units) and their relevance for molecular epidemiology and experimental evolution.** *Kinetoplastid Biology and Disease* 2003, **2(12)**..
- Hannaert V, Bringaud F, Opperdoes FR and Michels PAM: **Evolution of energy metabolism and its compartmentation in Kinetoplastida.** *Kinetoplastid Biology and Disease* 2003, **2(11)**..
- Hughes AL and Piontkivska H: **Molecular Phylogenetics of Trypanosomatidae: Contrasting Results from 18S rRNA and Protein Phylogenies.** *Kinetoplastid Biology and Disease* 2003, **2(15)**..
- Dávila AMR, Majiwa PAO, Grisard EC, Aksoy S and Melville SE: **Comparative genomics to uncover the secrets of tsetse and livestock-infective trypanosomes.** *Trends Parasitol* 2003, **19(10)**:436-439.
- Podlipaev S: **The more insect trypanosomatids under study the more diverse Trypanosomatidae appears.** *Int J Parasitol* 2001, **31(5-6)**:648-52.
- Podlipaev S: **Insect trypanosomatids: the need to know more.** *Mem Inst Oswaldo Cruz* 2000, **95(4)**:517-22.
- Vickerman K: **Order Kinetoplastida Honigberg, 1963.** In: *An Illustrated Guide to the Protozoa Volume 2*. Secondth edition. Edited by: Lee JJ, Leedale GF, Bradbury P. Society of Protozoologists, USA; 2000:1159-1180.
- El-Sayed NM and Donelson JE: **A survey of the *Trypanosoma brucei rhodesiense* genome using shotgun sequencing.** *Mol Biochem Parasitol* 1997, **84(2)**:167-78.
- Porcel BM, Tran AN, Tammi M, Nyarady Z, Rydaker M, Urmenyi TP, Rondinelli E, Pettersson U, Andersson B and Aslund L: **Gene survey of the pathogenic protozoan *Trypanosoma cruzi*.** *Genome Res* 2000, **10(8)**:1103-7.
- Lukeš J, Guilbride DL, Votýpka J, Zíková A, Benne R and Englund PT: **Kinetoplast DNA network: evolution of an improbable structure.** *Eukaryot Cell* 2002, **1(4)**:495-502.
- Simpson AGB, Lukeš J and Roger AJ: **The evolutionary history of kinetoplasts and their kinetoplasts.** *Mol Biol Evol* 2002, **19**:2071-2083.
- Massana R, Guillou L, Diez B and Pedros-Alio C: **Unveiling the organisms behind novel eukaryotic ribosomal DNA sequences from the ocean.** *Appl Environ Microbiol* 2002, **68(9)**:4554-8.
- Stevens JR: **One million insects – a lot of parasites?** *Trends Parasitol* 2001, **17**:119-120.
- Atkins MS, Teske AP and Anderson OR: **A survey of flagellate diversity at four deep-sea hydrothermal vents in the Eastern Pacific Ocean using structural and molecular approaches.** *J Eukaryot Microbiol* 2000, **47(4)**:400-11.
- Lopez-Garcia P, Philippe H, Gail F and Moreira D: **Autochthonous eukaryotic diversity in hydrothermal sediment and experimental microcolonizers at the Mid-Atlantic Ridge.** *Proc Natl Acad Sci U S A* 2003, **100(2)**:697-702.
- Dyková I, Fiala I, Lom J and Lukeš J: ***Perkinsiella amoebae*-like endosymbionts of *Neoparamoeba* spp, relatives of the kinetoplastid *Ichthyobodo*.** *Europ J Protistol* 2003, **39**:37-52.
- DeLong EF and Pace NR: **Environmental diversity of bacteria and archaea.** *Syst Biol* 2001, **50(4)**:470-8.
- Charlebois RL, Beiko RG and Ragan MA: **Microbial phylogenomics: Branching out.** *Nature* **421(6920)**:217. 2003 Jan 16

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
http://www.biomedcentral.com/info/publishing\_adv.asp

