

in the future and inform prospective theoretical and experimental research in this field.

FUNDING

This work was supported by the Basque Government through the ELKARTEK-2016 program.

Conflict of interest statement. None declared.

Michel Armand
Centre for Cooperative Research on
Alternative Energies,
Basque Research and Technology Alliance,
Spain
E-mail: marmand@cicenergigune.com

2. Kang B and Ceder G. *Nature* 2009; **458**: 190–3.
3. Armand M and Touzain P. *Mater Sci Eng* 1977; **31**: 319–29.
4. Wang Z, Wang D and Zou Z *et al.* *Natl Sci Rev* 2020; **7**: 1768–75.

REFERENCES

1. Armand M and Tarascon JM. *Nature* 2008; **451**: 652–7.

National Science Review
8: nwa185, 2021
doi: 10.1093/nsr/nwa185
Advance access publication 2 September 2020

CHEMISTRY

Tackling the challenge of controlling the spin with electric field

Roberta Sessoli 

Electric (E) fields are commonly employed to control magnetization in some classes of materials such as multiferroics. However, the extreme confinement of the electric field, for instance under the tip of a scanning tunnel microscope, makes it an ideal tool to manipulate individual spins, generating interest in spins as potential quantum bits. Spins are in fact intrinsic quantum systems that can be manipulated easily by means of pulses of electromagnetic microwaves, as commonly done in magnetic resonance spectroscopies. Although the coupling usually involves the magnetic field of microwaves, the first experiments with E -field pulses date back to the 1970s when Mims investigated the symmetry of the lanthanide ion coordination environment [1]. A linear effect is characteristic of an acentric environment. Renewed interest in spin-electric coupling is motivated by recent advances in scanning probe spectroscopies able to perform single atom CW [2] and pulsed [3] EPR spectroscopy. These exceptional tools make possible the optimization of spin-electric coupling in the qubit and exploration of the potential for gate control. In this respect, ensemble measurements (diluted single crystals or frozen solutions) provide an immense playground. Excluding rare exceptions, spin-electric coupling is enhanced by spin-orbit coupling. Lanthanide-based materials

with the orbitally unquenched contribution of 4f electrons are promising candidates. Liu Z *et al.* recently coordinated a pulsed EPR study on Ce^{3+} ions (f^1 , $S = 1/2$ and $J = 5/2$ but with a well isolated ground doublet) diluted in a polar diamagnetic crystal of yttrium aluminum garnet (YAG), as shown in Fig. 1 [4]. The standard $\pi/2 - t - \pi$ Hahn sequence generating an echo was modified by insertion of an E -field pulse. The Hahn echo

intensity was found to depend on both E -pulse intensity and length, demonstrating that the E -field is controlling the spin system. The effect, often quantified as a frequency shift per V m^{-1} , is about one order of magnitude larger than that detected in molecular qubits based on 3d ions. In the same study, the authors used the E -pulses to control evolution of the spin system through a phase gate and to implement the Deutsch–Jozsa




Figure 1. Diluted Ce^{3+} ions (f^1 , $J = 5/2$) substituting diamagnetic Y^{3+} ions in the YAG lattice are operated as spin qubits using mw pulses, as commonly done in pulsed electron paramagnetic resonance spectroscopy (color code: Ce, green; Y, cyan; Al, grey; O, red). Liu Z *et al.* [4] have added an electric field pulse and demonstrated that it acts as a phase gate modifying the spin echo signal. The large spin-orbit coupling of Ce^{3+} ion is a key ingredient for coherent electric field control of the spin, while the rigid garnet lattice warrants long spin coherence also at moderate temperatures.

algorithm, a well-known example of a deterministic quantum algorithm which is exponentially faster than its classical analogue.

However, an E -driven phase gate is usable only if a large number of operations can be made before the coherence is lost. Efficient operation requires strong spin-electric coupling but the same mechanisms that originate the coupling (spin-orbit interaction and atomic displacements) are also responsible for magnetic relaxation and loss of coherence. The Ce^{3+} in YAG has good coherence lifetime, 15 μs at 10 K, thanks to its rigid lattice and low nuclear spin concentration. Interestingly, a contemporaneous investigation by Liu J *et al.* highlighted even stronger spin-electric coupling in a holmium(III) polyoxometalate [5]. The strength of the effect has been rationalized considering the

coupling with a vibration mode of the Ho coordination sphere. Shorter coherence lifetimes were observed, although these were enhanced when protected at clock transitions.

In conclusion, recent advances in E -field coherent control of spin qubits suggest that single spin qubit operation is viable and particularly promising for discrete molecular systems. However, optimization of coherence time and spin-electric coupling poses antithetic conditions and remains an open challenge for chemists and physicists working in this fascinating field.

Conflict of interest statement. None declared.

Roberta Sessoli 
Dipartimento di Chimica 'Ugo Schiff' & INSTM RU,
Università degli Studi di Firenze, Italy
E-mail: roberta.sesso@unifi.it

REFERENCES

1. Mims WB. *The Linear Electric Field Effect in Paramagnetic Resonance*. Oxford: Clarendon Press, 1976.
2. Baumann S, Paul W and Choi T *et al.* *Science* 2015; **350**: 417–20.
3. Yang K, Paul W and Phark SH *et al.* *Science* 2019; **366**: 509–12.
4. Liu Z, Wang YX and Fang YH *et al.* *Natl Sci Rev* 2020; **7**: 1557–63.
5. Liu J, Mrozek J and Duan Y *et al.* arXiv: 2005.01029.

National Science Review
8: nwa267, 2021
doi: 10.1093/nsr/nwa267
Advance access publication 4 December 2020

MATERIALS SCIENCE

A promising route to neuromorphic vision

Leon Chua

The human vision system has a very strong capability in perceiving surroundings. Such visual perception starts in the retina, which receives and preprocesses the visual input in the form of light, and ends up with high-level processing in the visual cortex of the brain (Fig. 1a). As such, we can understand what the visual inputs represent while consuming little energy. It has been a long-sought dream for human beings to build a powerful and energy-efficient intelligent vision system that has a superior ability, similar to the human brain. Computer vision, as a similar model to the human brain, aims at viewing, processing and understanding images in the same way as human beings, and has become one major technological advancement in building intelligent machines [1]. However, the mainstream technology for computer vision is based on algorithms running on a von Neumann architecture computer, and cannot

emulate the hierarchical organizations and biological functions of the human vision system. In particular, traditional computer vision in conjunction with the conventional charge-coupled device (CCD) and complementary metal-oxide-semiconductor (CMOS) image sensors suffers from challenges in latency

and power consumption as a high volume of redundant visual information is sensed and then has to be processed.

A recent work [2] on the development of the neuromorphic visual system attempted to replicate the human's capability of visual perception and represents a significant advancement towards the




Figure 1. Human vision system (a) and neuromorphic vision system (b) composed of crossbar array and van der Waals heterostructure materials.