



Back to the Origin: *bla*_{OXA-204} and *bla*_{NDM-1} Genes in *Shewanella* spp. from a Tunisian River

Sana Ncir,^{a,b} Agnese Lupo,^a Antoine Drapeau,^a Pierre Châtre,^a Meriem Souguir,^{a,b} Sana Azaiez,^b Jean-Yves Madec,^a Wejdene Mansour,^b  Marisa Haenni^a

^aUnité Antibiorésistance et Virulence Bactériennes, ANSES - Université de Lyon, Lyon, France

^bLaboratoire de Biophysique métabolique et Pharmacologie Appliquée (LR12ES02), Faculté de Médecine Ibn Al Jazzar Sousse, Sousse, Tunisia

KEYWORDS carbapenem, *Shewanella xiamenensis*, *Shewanella bicestrii*, Tunisia, NDM-1, OXA-204, IS91, PER-1

The environmental bacteria *Xanthomonas* spp. and *Shewanella* spp. are progenitors of the *bla*_{NDM} and *bla*_{OXA-48-like} carbapenemase genes, respectively (1, 2). The most likely hypothesis is that these genes have been mobilized in the environment on mobile genetic elements, and subsequently transferred to Gram-negative bacteria (GNB) of clinical importance. The pathogens (primarily *Escherichia coli* and *Klebsiella pneumoniae*) that acquired these carbapenemase genes then emerged as a major One Health issue and are currently reported not only in clinical settings but also in animals and the environment (3). In January 2019, two carbapenem-resistant (CP-R) *Shewanella* spp. were isolated from fresh water sampled at two different locations along the El Hammam Oued river in Sousse, Tunisia, as part of a pilot study on multidrug-resistant GNB in aquatic environments. This river collects urban treated wastewater but no hospital effluents. The two isolates resistant to ertapenem (MIC ≥32 mg/L) were whole-genome sequenced using short-read (NovaSeq-6000, Illumina) and long-read (MinION, Oxford Nanopore) technologies (all genomic sequences were deposited in DDBJ/EMBL/GenBank under the accession number [PRJNA830355](https://www.ncbi.nlm.nih.gov/nuclink/PRJNA830355)). Genomes were assembled with Unicycler and analyzed using tools of the Center for Genomic Epidemiology (<https://www.genomicepidemiology.org/>).

S. xiamenensis 8A possessed a 4,959,567 bp chromosome and one nontypeable 216,160 bp plasmid (p8A) showing poor or fragmented homologies with published sequences. Except for *bla*_{OXA-538}, all resistance genes—conferring resistances to beta-lactams, aminoglycosides, sulfonamides, trimethoprim, rifampicin and erythromycin—were located on p8A (Fig. 1A). Specifically, p8A carried the *bla*_{PER-1} extended-spectrum beta-lactamase (ESBL) and the *bla*_{OXA-204} carbapenemase genes. *bla*_{OXA-204} has already been reported in one *S. xiamenensis* strain from water in Portugal (4), flanked by the peptidase C15 and *lysR* gene as intrinsically found on the *Shewanella* spp. chromosome (5). Here, the *bla*_{OXA-204} gene was surrounded by an *ISEcp1* insertion sequence and a truncated *lysR* gene, as reported in the *K. pneumoniae* KP49 isolated from a Tunisian patient ([KP027886](https://pubmed.ncbi.nlm.nih.gov/27886/)) (6). This indicates that *bla*_{OXA-204} in p8A is not intrinsic but has transited through another bacterial host before being reintegrated in *S. xiamenensis* p8A via a transposition event.

Shewanella bicestrii 10B possessed a 4,717,299 bp chromosome carrying a *bla*_{OXA-48-like} and the *aadA2* resistance genes. The strain additionally presented a nontypeable 160,879 bp plasmid (p10B) sharing conserved regions with the pSA70-3 and pKC-Na-R1 plasmids, respectively, isolated from *Shewanella putrefaciens* and *Shewanella algae* (Fig. 1B) (7). p10B presented genes conferring resistance to beta-lactams, aminoglycosides, sulfonamides, trimethoprim, chloramphenicol, quinolones, and rifampicin, but also to mercury and biocides. CP-R was conferred by the *bla*_{NDM-1} gene; *bla*_{NDM-1}, together with the *bla*_{OXA-10}, *sul1*, and *aadA1* genes, was surrounded by two IS91-like that might have mediated its insertion/excision (8). A similar genetic organization was observed in pSA70-3 with only the left

Editor Daria Van Tyne, University of Pittsburgh School of Medicine

Copyright © 2022 Ncir et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Marisa Haenni, marisa.haenni@anses.fr.

The authors declare no conflict of interest.

Published 20 September 2022




FIG 1 Schematic representation of the p8A plasmid (A). Comparative schematic representation of the p10B, pSA70-3 (CP078040), and pKC-Na-R1 (CP033574) plasmids. Genetic elements were analyzed by BLASTn and visualized using Easyfig 2.2.5_win. The window size used for BLASTn analysis was 2,000 bp.

IS91-like missing (7), while the element was more fragmented in pKC-Na-R1, where only the *sul1*-IS91-like and *aadA1*-*bla*_{OXA-10} genes were conserved upstream the *bla*_{VEB-1} ESBL gene.

Environmental bacteria are progenitors of carbapenemases (1, 2), but reacquisition of these genes by environmental bacteria is rarely reported. This study showed that the *bla*_{OXA-204} and *bla*_{NDM-1} genes, which are highly prevalent in Tunisian hospital settings and effluents (9, 10), can spread back to environmental bacteria. In both cases, resistance genes moved to a *Shewanella* spp. nontypeable plasmid after IS-mediated acquisition, a mechanism that might be more efficient than plasmid transfer for bacterial genera that are not closely related.

ACKNOWLEDGMENT

This work was funded by the “PHC Utique” program of the French Ministry of Foreign Affairs, the French Ministry of higher education, research, and innovation and the Tunisian Ministry of higher education and scientific research under the CMCU project number 21G0807. This work was also supported by internal funding of the French Agency for Food, Environmental and Occupational Health & Safety (ANSES).

REFERENCES

- Poirel L, Potron A, Nordmann P. 2012. OXA-48-like carbapenemases: the phantom menace. *J Antimicrob Chemother* 67:1597–1606. <https://doi.org/10.1093/jac/dks121>.
- Sekizuka T, Matsui M, Yamane K, Takeuchi F, Ohnishi M, Hishinuma A, Arakawa Y, Kuroda M. 2011. Complete sequencing of the *bla*_{NDM-1}-positive IncA/C plasmid from *Escherichia coli* ST38 isolate suggests a possible origin from plant pathogens. *PLoS One* 6:e25334. <https://doi.org/10.1371/journal.pone.0025334>.
- Dandachi I, Chaddad A, Hanna J, Matta J, Daoud Z. 2019. Understanding the epidemiology of multi-drug resistant gram-negative bacilli in the middle east using a One Health approach. *Front Microbiol* 10:1941. <https://doi.org/10.3389/fmicb.2019.01941>.
- Tacao M, Correia A, Henriques I. 2013. Environmental *Shewanella xiamensis* strains that carry *bla*_{OXA-48} or *bla*_{OXA-204} genes: additional proof for *bla*_{OXA-48}-like gene origin. *Antimicrob Agents Chemother* 57:6399–6400. <https://doi.org/10.1128/AAC.00771-13>.
- Pitout JDD, Peirano G, Kock MM, Strydom K-A, Matsumura Y. 2019. The global ascendancy of OXA-48-type carbapenemases. *Clin Microbiol Rev* 33:e00102-19. <https://doi.org/10.1128/CMR.00102-19>.
- Charfi K, Mansour W, Khalifa AB, Mastouri M, Aouni M, Mammeri H. 2015. Emergence of OXA-204 beta-lactamase in Tunisia. *Diagn Microbiol Infect Dis* 82:314–317. <https://doi.org/10.1016/j.diagmicrobio.2015.04.003>.
- D'Souza AW, Potter RF, Wallace M, Shupe A, Patel S, Sun X, Gul D, Kwon JH, Andleeb S, Burnham C-AD, Dantas G. 2019. Spatiotemporal

- dynamics of multidrug resistant bacteria on intensive care unit surfaces. *Nat Commun* 10:4569. <https://doi.org/10.1038/s41467-019-12563-1>.
8. Chen C-J, Wu T-L, Lu P-L, Chen Y-T, Fung C-P, Chuang Y-C, Lin J-C, Siu LK. 2014. Closely related NDM-1-encoding plasmids from *Escherichia coli* and *Klebsiella pneumoniae* in Taiwan. *PLoS One* 9:e104899. <https://doi.org/10.1371/journal.pone.0104899>.
 9. Dziri O, Dziri R, Ali El Salabi A, Chouchani C. 2020. Carbapenemase producing gram-negative bacteria in Tunisia: history of thirteen years of challenge. *Infect Drug Resist* 13:4177–4191. <https://doi.org/10.2147/IDR.S259562>.
 10. Nasri E, Subirats J, Sánchez-Melsió A, Mansour HB, Borrego CM, Balcázar JL. 2017. Abundance of carbapenemase genes (*bla_{KPC}*, *bla_{NDM}* and *bla_{OXA-48}*) in wastewater effluents from Tunisian hospitals. *Environ Pollut* 229:371–374. <https://doi.org/10.1016/j.envpol.2017.05.095>.