



Vibrio neptunius Produces Piscibactin and Amphibactin and Both Siderophores Contribute Significantly to Virulence for Clams

Fabián Galvis¹, Lucía Ageitos², Jaime Rodríguez², Carlos Jiménez², Juan L. Barja¹, Manuel L. Lemos^{1*} and Miguel Balado^{1*}

¹ Departamento de Microbiología y Parasitología, Instituto de Acuicultura y Facultad de Biología-CIBUS, Universidade de Santiago de Compostela, Santiago de Compostela, Spain, ² Centro de Investigaciones Científicas Avanzadas (CICA) e Departamento de Química, Facultad de Ciencias, Universidade da Coruña, A Coruña, Spain

OPEN ACCESS

Edited by:

Cristian Oliver,
Austral University of Chile, Chile

Reviewed by:

Javier Santander,
Memorial University of Newfoundland,
Canada

Jorge Alberto Giron,
University of Puebla, Mexico

*Correspondence:

Manuel L. Lemos
manuel.lemos@usc.es
Miguel Balado
miguel.balado@usc.es

Specialty section:

This article was submitted to
Molecular Bacterial Pathogenesis,
a section of the journal
Frontiers in Cellular and
Infection Microbiology

Received: 30 July 2021

Accepted: 07 October 2021

Published: 25 October 2021

Citation:

Galvis F, Ageitos L, Rodríguez J, Jiménez C, Barja JL, Lemos ML and Balado M (2021) *Vibrio neptunius* Produces Piscibactin and Amphibactin and Both Siderophores Contribute Significantly to Virulence for Clams. *Front. Cell. Infect. Microbiol.* 11:750567. doi: 10.3389/fcimb.2021.750567

Vibrio neptunius is an inhabitant of mollusc microbiota and an opportunistic pathogen causing disease outbreaks in marine bivalve mollusc species including oysters and clams. Virulence of mollusc pathogenic vibrios is mainly associated with the production of extracellular products. However, siderophore production is a common feature in pathogenic marine bacteria but its role in fitness and virulence of mollusc pathogens remains unknown. We previously found that *V. neptunius* produces amphibactin, one of the most abundant siderophores in marine microbes. In this work, synthesis of the siderophore piscibactin was identified as the second siderophore produced by *V. neptunius*. Single and double mutants in biosynthetic genes of each siderophore system, piscibactin and amphibactin, were constructed in *V. neptunius* and their role in growth ability and virulence was characterized. Although the High Pathogenicity Island encoding piscibactin is a major virulence factor in vibrios pathogenic for fish, the *V. neptunius* wild type did not cause mortality in turbot. The results showed that amphibactin contributes more than piscibactin to bacterial fitness *in vitro*. However, infection challenges showed that each siderophore system contributes equally to virulence for molluscs. The *V. neptunius* strain unable to produce any siderophore was severely impaired to cause vibriosis in clams. Although the inactivation of one of the two siderophore systems (either amphibactin or piscibactin) significantly reduced virulence compared to the wild type strain, the ability to produce both siderophores simultaneously maximised the degree of virulence. Evaluation of the gene expression pattern of each siderophore system showed that they are simultaneously expressed when *V. neptunius* is cultivated under low iron availability *in vitro* and *ex vivo*. Finally, the analysis of the distribution of siderophore systems in genomes of *Vibrio* spp. pathogenic for molluscs showed that the gene clusters encoding amphibactin and piscibactin are widespread in the Coralliilyticus clade. Thus, siderophore production would constitute a key virulence factor for bivalve molluscs pathogenic vibrios.

Keywords: Coralliilyticus, *Vibrio neptunius*, bivalve molluscs pathogens, virulence factors, siderophores, piscibactin, amphibactin, aquaculture

INTRODUCTION

Bacteria belonging to the genus *Vibrio* (Vibrios) are ubiquitously distributed in the marine environment and are also a dominant fraction of bivalve microbiota (Vezzulli et al., 2018). Mollusc hemolymph is a critical site for the host immune response (Potgieter et al., 2015). Bivalve hemocytes can kill vibrios by phagocytosis and production of reactive oxygen species, highly reactive nitric oxide, antimicrobial peptides and hydrolytic enzymes (Destoumieux-Garzón et al., 2020). Interestingly, some Vibrios are part of the resident microbiota of bivalves as they persist in the hemolymph in the absence of an environmental source of population (Potgieter et al., 2015; Lokmer et al., 2016; Vezzulli et al., 2018; Zhang et al., 2018). The microbiota benefits the host as it boosts the immune system, and promotes reproduction, nutrition and defence mechanisms (Engel et al., 2002; McFall-Ngai et al., 2013; Cahill et al., 2016; Utermann et al., 2018). Nonetheless, under unfavourable conditions some bacteria are responsible for disease outbreaks. Among them, vibriosis is a serious epizootic disease caused by some *Vibrio* spp. that has become the most important limiting factor of the intensive fish and shellfish mariculture industry worldwide (Paillard et al., 2004; Toranzo et al., 2005; Travers et al., 2015; Dubert et al., 2017).

Vibrio species belonging to the Coralliilyticus and Orientalis clades are among the best-known species of bivalve pathogens (Dubert et al., 2017). They include *V. neptunius*, a marine bacterium that was isolated from marine water samples and animals such as turbot larvae (*Scophthalmus maximus*), rotifers (*Brachiomus plicatilis*) and larval stages of cephalopods (*Octopus vulgaris*) (Thompson et al., 2003; García-Amado et al., 2011; Farto et al., 2019). Notably, this bacterium is a relevant pathogen of aquacultured marine invertebrates, including artemia, oysters and clams (Prado et al., 2005; Kesarcodi-Watson et al., 2009a; Kesarcodi-Watson et al., 2009b; Romalde et al., 2014; Dubert et al., 2017). *V. neptunius* rapidly invades bivalve larvae tissues by entering them through the filtration feeding process. Its virulence is commonly associated with the production of thermolabile extracellular products with cytotoxic activity for fish and homeothermic animal cell lines (Dubert et al., 2016). Remarkably, vibriolysin-like protease VnpA and collagenase ColA were recently characterized as relevant virulence factors of *V. neptunius* since their production is required for full virulence in oyster larvae (*Ostrea edulis*) (Galvis et al., 2021).

It is well established that the production and utilization of siderophores is a key virulence factor for most vertebrate pathogenic bacteria (Kramer et al., 2020). However, although siderophore production would be a common feature of *Vibrio* pathogens affecting bivalves (Gómez-León et al., 2005; Mechri et al., 2017), its role in bacterial fitness and/or virulence is understudied in this type of bacterial pathogens. Our previous works demonstrated that *V. neptunius* produces a set of 9 amphibactin forms to overcome low-iron conditions. The gene cluster involved in its production and utilization was identified (genes *absABDEF* and *abtABCDE*) and the amphibactin outer

membrane transporter gene *abtA* was characterized. It is noteworthy that *abtA* was regularly found in mollusc microbiota including some of the most devastating pathogens such as *V. coralliilyticus* and *V. tubiashii*. Interestingly, a *V. neptunius* Δ *absE* mutant (impaired to synthesize amphibactins) showed a weak but not null siderophore activity in cell free supernatants, which could imply the production of a second siderophore. Besides, the putative role of amphibactins in virulence for bivalves was not yet evaluated (Galvis et al., 2020).

In the present work the siderophore piscibactin was identified as the second siderophore produced by *V. neptunius*. Single and double *V. neptunius* mutants in biosynthetic genes of each siderophore system, piscibactin and amphibactin, were constructed and their role in virulence determined. In addition, the gene expression patterns of each siderophore were studied *in vitro* and *ex vivo*. The results showed that both siderophore systems, amphibactin and piscibactin, are expressed during infection, playing a key role in the virulence of *V. neptunius* for clams.

MATERIALS AND METHODS

Bacterial Strains, Plasmids, and Media

The bacterial strains and plasmids used, as well as those derived from this study, are listed in **Table 1**. *V. neptunius* and *V. anguillarum* strains were grown at 25°C in Tryptic Soy Agar and Broth (Pronadisa, Madrid, Spain) supplemented with 1% NaCl (TSA-1 and TSB-1, respectively), as well as in M9 minimal medium supplemented with 0.2% Casamino Acids (Difco) (CM9) (Lemos et al., 1988). *Escherichia coli* strains were grown at 37°C in Luria-Bertani (LB) medium (Pronadisa) or LB supplemented with the appropriate antibiotics. Ampicillin sodium salt was used at 100 µg/mL, kanamycin 50 µg/mL and gentamycin 15 µg/mL (final concentrations).

DNA Manipulations and Bioinformatics Tools

Total genomic DNA from *V. neptunius* PP-145.98 was purified with the InstaGene™ Matrix (BioRad, Hercules, California, CA, USA). PCR reactions were all carried out with Taq polymerase NZYTaQ (Nzytech, Lisboa, Portugal) according to manufacturer protocol in a T-Gradient Thermal Cycler (Biometra, Göttingen, Germany). The extraction of DNA from agarose gels and purification of plasmid DNA were carried out using NucleoSpin Gel and a PCR clean-up kit (Macherey-Nagel, Düren, Germany) and a GeneJET Plasmid Miniprep Kit (Thermo-Fisher, Waltham, MA, USA).

The genome of *V. neptunius* PP-145.98 strain (accession number JAFHLB000000000) was screened, using antiSMASH 5.0 (Blin et al., 2019), for the presence of siderophore-related sequences. The NCBI services (<http://ncbi.nlm.nih.gov>) were used to consult the DNA and protein sequence databases with BLAST algorithm. Prediction of protein domains was carried out by using the Pfam protein families database (Finn et al., 2014).

TABLE 1 | Strains and plasmids used in this study.

Strain or plasmid	Relevant characteristic(s)	Reference or source
<i>V. neptunius</i>		
PP-145.98	Wild type strain, isolated from <i>Ruditapes philippinarum</i> (larvae), Ap ^r	
FG109	PP-145.98 <i>absF</i> defective mutant, Ap ^r	This study
FG113	PP-145.98 <i>irp2</i> defective mutant, Ap ^r	This study
FG115	PP-145.98 <i>absF</i> and <i>irp2</i> defective mutant, Ap ^r	This study
<i>V. anguillarum</i>		
RV22	Wild-type serotype O2 strain isolated from diseased turbot (Spain)	(Lemos et al., 1988)
MB14	RV22 <i>vabF</i> defective mutant	(Balado et al., 2006)
MB67	RV22 <i>vabD</i> defective mutant	(Balado et al., 2018)
ML210	RV22 <i>vabD</i> and <i>frpA</i> defective mutant	submitted manuscript
<i>E. coli</i>		
DH5 α	SupE4 Δ lacU169 (Φ 80 <i>lacZ</i> Δ M15) <i>hsdR17 recA1 endA1 gyrA96 thi-1 relA1</i>	Laboratory stock
S17-1- λ pir	Tp ^r Sm ^r <i>recA, thi, pro, hsdR-M+RP4: 2- Tc : Mu:Km Tn7 λpir</i>	(Herrero et al., 1990)
Plasmids		
pWKS30	Low-copy-number cloning vector, Ap ^r	(Wang and Kushner, 1991)
pNidKan	Suicide vector derived from pCVD442, Km ^r	(Mouriño et al., 2004)
pHRP309	Low-copy lacZ reporter plasmid, mob Gm ^r	(Parales and Harwood, 1993)
pFG154	<i>proC</i> promoter (<i>PproC</i>) fused to promoterless <i>lacZ</i> gene in pHRP309, Gm ^r	This study
pFG156	<i>entD</i> promoter (<i>PentD</i>) fused to promoterless <i>lacZ</i> gene in pHRP309, Gm ^r	This study
pFG180	<i>abtA</i> promoter (<i>PabtA</i>) fused to promoterless <i>lacZ</i> gene in pHRP309, Gm ^r	This study
pFG166	<i>absE</i> promoter (<i>PabsE</i>) fused to promoterless <i>lacZ</i> gene in pHRP309, Gm ^r	This study
pFG175	<i>abtA</i> (reverse) promoter (<i>PentD</i> -Reverse) fused to promoterless <i>lacZ</i> gene in pHRP309, Gm ^r	This study
pFG176	<i>araC1</i> promoter (<i>ParaC1</i>) fused to promoterless <i>lacZ</i> gene in pHRP309, Gm ^r	This study
pFG188	<i>frpA</i> promoter (<i>PfrpA</i>) fused to promoterless <i>lacZ</i> gene in pHRP309, Gm ^r	This study

The sequences of housekeeping genes *ftsZ*, *gyrB*, *mreB*, *pyrH* y *recA* of bivalve pathogenic vibrios were downloaded from the NCBI database and used for multilocus sequence analysis (MLSA). Sequences of each gene were concatenated into a single 3880 bp sequence and aligned using the MUSCLE. The final phylogenetic tree was constructed based on concatenated sequences of the five housekeeping genes by maximum-likelihood (GTR+G model). Sequence alignments and phylogenetic tree were performed using MEGAX (v. 10.2.2) (Kumar et al., 2018).

Construction of *absF* and *irp2* Mutants by Allelic Exchange

In-frame deletions of *absF* and *irp2* in *V. neptunius* PP-145.98 were constructed by using PCR amplification of two fragments of each gene and flanking regions that, when ligated together, would result in an in-frame (nonpolar) deletion. The oligonucleotides used to amplify the upstream and downstream ends of each gene are shown in **Table S1**. Once deleted alleles were constructed by sequential cloning of the PCR products into pWKS30 plasmid, they were liberated by digestion with *NotI* and *ApaI* and cloned into the suicide vector pCAR109 (Mouriño et al., 2004). The resulting plasmids (**Table 1**) were mated from *E. coli* S17-1- λ pir into *V. neptunius* PP-145.98 and transconjugants with the plasmid integrated in the chromosome by homologous recombination, were selected on TSA-1 containing 50 mg/mL of kanamycin (resistance conferred by pNidKan) and 100 mg/mL of ampicillin (antibiotic to select *V. neptunius* PP-145.98). A second recombination event was obtained by selecting for sucrose (10%) resistance. This process led to the generation of the *V. neptunius* PP-145.98 single mutants Δ *absF* and Δ *irp2*, and the

double mutant Δ *absF* Δ *irp2*, named FG109, FG113 and FG115, respectively.

Growth Promotion and Siderophore Production Assays

Growth measurement of *V. neptunius* PP-145.98 strains was performed using 96-well microtiter plates. Each well contained 200 μ L of CM9 medium (Lemos et al., 1988) supplemented with FeCl₃ at 10 μ M (PROBUS) or with the iron chelators ethylenediamine-di(o-hydroxyphenyl-acetic acid) (EDDA) at 5 μ M or 2,2'-dipyridyl (dipyridyl) (Sigma) at 50 μ M or 30 μ M. Each well was inoculated with a 1:50 dilution of an overnight culture of the strain to be tested in TSB-1 at OD₆₀₀ = 0.5. The plates were incubated at 25°C with shaking at 150 rpm. After 18 h of incubation, growth (OD₆₀₀) was recorded in an iMACK Microplate reader (Bio-Rad). Bacterial cultures in CM9 with 30 μ M 2,2'-dipyridyl and an OD₆₀₀ \approx 0.6 (after 6 h of incubation) were used to measure siderophore production using the chrome azurol-S (CAS) liquid assay (Schwyn and Neilands, 1987). Equal volumes of each cell free supernatant and CAS reagent were mixed and absorbance at 630 nm (A₆₃₀) was measured in a UV-VIS spectrophotometer (Hitachi) after 15 min of incubation at room temperature.

Cross-Feeding Assays

The biological activities of the supernatants produced by the parental and mutant strains were determined by cross-feeding experiments. We tested the ability of culture supernatants from *V. neptunius* PP-145.98 mutants defective in piscibactin and amphibactin synthesis to cross-feed different indicator strains defective in the synthesis and/or transport of piscibactin.

To test whether *V. neptunius* wild type or derivative mutants produce piscibactin, a cross-feeding assay was conducted using two *V. anguillarum* mutants derived from RV22 strain that lack siderophore synthesis: RV22 Δ *avabD*, a single mutant (strain MB67) that does not produce siderophores and that it is able to use piscibactin as iron source since it carries the piscibactin outer membrane transporter FrpA; and RV22 Δ *avabD* Δ *frpA*, a double mutant that does not use piscibactin since it has inactivated the piscibactin transporter FrpA. Indicator strains were inoculated into CM9 plates as follows: 0.5 mL of an overnight culture in TSB-1 at an OD₆₀₀ = 0.5 were mixed with 5 mL of CM9 medium containing 0.5% agarose and 2,2'-dipyridyl 100 μ M, a concentration close to the Minimal Inhibitory Concentration (MIC) and at which growth halos can be easily visualized (Balado et al., 2006). The *V. neptunius* strains to be tested for piscibactin production were cultured in TSA-1 plates and the cells were harvested with a sterile loop and placed on the surface of the plates inoculated with the *V. anguillarum* strains. A piscibactin producer *V. anguillarum* strain (RV22 Δ *avabF*) was used as control. The presence of growth halos of the *V. anguillarum* indicator strains around cells of *V. neptunius* after overnight incubation at 25°C was indicative of piscibactin production.

RNA Purification and RT-PCR

The organization of amphibactin genes into operon(s) was tested by reverse transcription PCR. *V. neptunius* PP-145.98 was grown until exponential phase (ca. OD₆₀₀ = 1) in 10 mL CM9 medium containing 2.5 μ M EDDA. Cells were pelleted by centrifugation at 10,000 \times g for 2 min and total RNA was isolated with Trizol (Invitrogen) following the manufacturer's recommendations. RT-PCR was performed with 1 μ g RNA pre-treated with RQ1 RNase-Free DNase (Promega) by using the M-MLV reverse transcriptase (Invitrogen). A primer located at the 3'-end of *absA* gene was used to obtain a cDNA that was used as template for subsequent PCR reactions targeted in *abtE* (PCR2) and in the region between *abtC* and *abtD* (PCR1), *absE* and *absF* (PCR3) and between *abtA* and *absB* (PCR4) (Table S1). A negative control reaction was performed with total RNA treated with DNase without M-MLV reverse transcriptase to confirm the lack of genomic DNA contamination in each reaction mixture. The PCR positive control reaction was done using 100 ng of genomic DNA as template.

lacZ Transcriptional Fusions and β -Galactosidase Assays

The DNA fragments corresponding to the amphibactin and piscibactin promoter regions of *V. neptunius* PP-145.98 were amplified by PCR. The PCR products spanned from the first nucleotides of the coding sequence to ca. 700-900 bp upstream sequence. These putative promoter regions were fused to a promoterless *lacZ* gene in the low-copy-number reporter plasmid pHRP309 (Parales and Harwood, 1993). Transcriptional fusions of genes *abtC* and *abtA* for amphibactin, and *araC1* and *frpA* for piscibactin were constructed. The resulting transcriptional fusion constructs, *abtC::lacZ* (pFG156), *abtA::lacZ* (pFG180), *araC1::lacZ* (pFG176) and *frpA::lacZ* (pFG188), were mobilized

from *E. coli* S17-1- λ pir into *V. neptunius* PP-145.98 by conjugation. *V. neptunius* PP-145.98 derivatives carrying the transcriptional fusions were grown in CM9 at 25°C under conditions of iron availability (CM9 with 10 μ M Fe₂(SO₄)₃) and iron deficiency (CM9 with 50 μ M 2,2'-dipyridyl). For the *ex vivo* assay *V. neptunius* PP-145.98 derivatives carrying the promoter-*lacZ* fusions were grown at 25°C in the hemolymph of commercial mussels (*Mitylus galloprovincialis*) under normal or excess iron conditions. Hemolymph was collected from approximately 30 mussels by puncture of the adductor anterior muscle, then centrifuged at 4,000 rpm for 10 min and filtered through a nitrocellulose membrane with a pore size of 0.25 μ m. The β -galactosidase (LacZ) activity in CM9 cultures and hemolymph were measured by the method of Miller (Miller, 1992). Results showed are means of three independent experiments, each one measured in triplicate. Statistical significance of differences was determined using *t*-test. *P*-values were considered significant when *P* was <0.05.

Virulence Assays

Experimental infections in healthy clams (*Ruditapes philippinarum*) were performed with the wild-type and mutants of *V. neptunius* PP-145.98. For acclimatization the clams (mean size of 10 mm) were kept for 2 days in a seawater tank at 20°C with continuous aeration. The wild-type strain and the mutants Δ *absF*, Δ *irp2* and Δ *absF* Δ *irp2* of *V. neptunius* PP-145.98 were cultured in CM9 medium with 2,2'-dipyridyl 20 μ M for 12 h at 25°C to achieve an OD₆₀₀ = 0.6. The infection was carried out by bath as follows: groups of 50 clams were introduced in 50 mL of seawater containing the bacterial strain to be tested at a final concentration of 10⁶ cfu/mL, and incubated for 24 h. The clams were then washed twice with abundant seawater and placed in containers with 200 mL of seawater with aeration. The assay was performed by duplicate and a group of clams under the same conditions but without bacterial inoculation was used as negative control. Pathogenicity was evaluated after 96 h and was expressed as a percentage of survival. Mortalities were recorded daily, and the statistical significance of differences in percentage of survival for the different *V. neptunius* strains was determined using the Kaplan–Meier method with the Mantel–Cox log-rank test using SPSS (version 20; IBM SPSS Inc., Chicago, IL, USA). *P*-values were considered significant when *P* was <0.05, <0.01 and <0.001.

Virulence assays on fish were performed using turbot (*Scophthalmus maximus*) fingerlings weighting 5 g on average. The fish were divided into groups of 10 animals. All groups, one per strain tested and a control group, were maintained in 50 L seawater tanks at 18°C with continuous aeration and water recirculation. The wild-type strain and mutants were grown in CM9 medium for 12 h at 25°C to reach an OD₆₀₀ = 0.6. The inoculum used was a 10-fold dilution of this suspension in saline solution (0.85% NaCl in distilled H₂O). Fish were inoculated intraperitoneally (ip) with 100 μ L of bacterial suspensions. A control group was inoculated with 100 μ L of saline solution. Mortalities were recorded daily for 15 days after injection. All animal experimentation protocols used in this study were reviewed and approved by the Animal Ethics Committee of the University of Santiago de Compostela (protocol id. 15004/14/003).

Piscibactin Detection by Mass Spectrometry


The presence of piscibactin was studied following the SPE-HLB/HPLC-MS methodology previously described (Souto et al., 2012; Balado et al., 2018). Briefly, *V. neptunius* $\Delta absF$ was grown in CM9 medium supplemented with 30 μM 2,2'-dipyridyl at 25°C under continuous shaking (150 rpm) for 24 h. Once achieved an $\text{OD}_{600} = 1$, the bacterial culture was centrifuged at $10,000 \times g$ for 10 min (Beckman J-21 High Speed Centrifuge) and filtrated through a 0.45- μm pore size membrane. 800 mL of the resultant cell-free supernatant were treated with 17 mg of FeCl_3 , incubated at 4°C for 12 h, and concentrated under reduced pressure to 300 mL. Half of this volume was fractionated using an Oasis[®] hydrophilic lipophilic balance (HLB) cartridge (Waters) (35 cm^3 , 6 g), previously conditioned with 60 mL of acetonitrile (ACN, solvent B) and deionized water (H_2O , solvent A), in three batches of 75 mL. Each batch was fractionated with 30 mL of the following solvent mixtures: 1:0, 7:3, 1:1, 3:7, and 0:1 of A:B (v/v) obtaining the fractions VN $\Delta absFH1$ -5 respectively. VN $\Delta absFH3$ was analysed by HPLC/HRMS in a HPLC Acela (Thermo) coupled to a PDA and HRMS (LQT-Orbitrap Discovery) detector in full positive ion using the Atlantis dC18 (100 x 4.6 mm, 5 μm) column (Waters) and the following method (solvent A: H_2O , solvent B:

ACN): 35 min from 10% to 100% of B, 5 min isocratic at 100% of B and 10 min from 100 to 10% of B at 1 mL min^{-1} . The results showed the presence of ferri-piscibactin in the chromatographic peak with a $R_t=10.69$ min displaying the $[\text{M}+\text{H}]^+$ adduct at m/z 507.0032 (calcd. for $\text{C}_{19}\text{H}_{21}\text{N}_3\text{O}_4\text{S}_3\text{Fe}^+$ m/z 507.0038) and the absorbance maxima at 227, 256, 307, and 388 nm in its UV spectrum.

RESULTS

V. neptunius Genome Harbors a Version of the High Pathogenicity Island (irp-HPI) Encoding Piscibactin Biosynthesis and Transport

In silico analysis of *V. neptunius* PP-145.98 genome sequence (NZ_JAFHLB010000000) led to the identification of a gene cluster with high homology to *irp* genes previously identified in *P. damsela* subsp. *piscicida* and several *Vibrios*. These genes are harbored in the High Pathogenicity Island named *irp*-HPI and confer the ability to produce and use the siderophore piscibactin (Osorio et al., 2015) (Figure 1). The genomic island *irp*-HPI of *V. neptunius* (*irp*-HPI_{Vnep}) is a DNA region of approximately 33 kb that is located in the chromosome II






FIGURE 3 | Growth levels under low and high-iron availability and siderophore production (CAS assay) of *V. neptunius* PP-145.98 wild type and its derivative mutants $\Delta absF$, $\Delta irp2$ and $\Delta irp2\Delta absF$. Asterisks denote statistically significant differences with the wild type strain (**P* < 0.05 and ****P* < 0.001).

mutant were studied following the SPE-HLB/HPLC-HRMS methodology previously described by our research group (Souto et al., 2012). Cell-free supernatant was treated with FeCl₃, to obtain the stable ferri-siderophores, and fractionated

using an HLB cartridge. HPLC/HRMS analysis of the CAS-positive fraction VN $\Delta absFH3$, eluted with 1:1 of ACN: H₂O, confirmed the presence of ferri-piscibactin in the chromatographic peak at *R*_t=10.69 min (**Figures 5A, B**) by




FIGURE 4 | Cross feeding assay to test the production of piscibactin by *V. neptunius* 145.98 wild type strain and its derivative $\Delta absF$ and $\Delta irp2$ single mutants and $\Delta irp2\Delta absF$ double mutant. Indicator strains were inoculated within the CM9 plates containing 100 μ M of 2,2'-dipyridyl. Tested strains were placed as a loop full of bacterial biomass on the agar surface. The presence of a growth halo shows that the indicator strains can use the siderophores produced by the tested strains to overcome the iron limitation.




FIGURE 5 | HPLC/HRMS analysis for the detection of ferri-piscibactin in the fraction VN Δ absFH3, from the *V. neptunius* Δ absF mutant, eluted with 1:1 H₂O:ACN from a HLB cartridge. **(A)** Total Ion Current (TIC) chromatogram of VN Δ absFH3 in which ferri-piscibactin was detected at $R_t = 10.69$ min, highlighted in blue. **(B)** HPLC-DAD chromatogram of VN Δ absFH3 in which ferri-piscibactin was detected at $R_t = 10.69$ min, highlighted in blue. **(C)** UV spectrum of the peak at $R_t = 10.69$ min, where ferri-piscibactin was detected, showing the absorption maxima at 207, 256, 307 and 388 nm. **(D)** (+)-HRESIMS of the peak at $R_t = 10.69$ min, where ferri-piscibactin was identified, displaying: m/z 507.0032 (calcd. for C₁₉H₂₁N₃O₄S₃Fe⁺, m/z 507.0038), expanded region of MS in the range m/z 501-514, showing the presence of the characteristic Fe isotopic distribution, and structure of ferri-piscibactin.

comparison of its MS and UV spectra with the previously described data (Souto et al., 2012). Specifically, the mass spectrum (MS) of this peak showed the $[M + H]^+$ adduct at m/z 507.0032 (calcd. for C₁₉H₂₁N₃O₄S₃Fe⁺, m/z 507.0038) along with the isotopic distribution of Fe ($M_r = 54, 56, 57, 58$; ratio 0.6:9.2:0.2:0.02) (**Figure 5D**); while the UV spectrum displayed the absorbance maxima at 227, 256, 307, and 388 nm (**Figure 5C**).

All these results demonstrate that each one of the *V. neptunius* single mutants (Δ irp2 or Δ absF) produce one unique siderophore when cultivated under iron-restricted conditions. Thus, while *V. neptunius* Δ irp2 mutant produces only amphibactin and Δ absF produces only piscibactin, the Δ irp2 Δ absF double mutant does not produce any of these siderophores.

Piscibactin and Amphibactin Significantly Contribute to *V. neptunius* Virulence for Clams

To study the role of piscibactin and/or amphibactin production in virulence of the bivalve mollusc pathogen *V. neptunius*, experimental infection challenges were performed. For this purpose, groups of 50 clams larvae were inoculated with the *V. neptunius* wild type or with one of the mutants. A 100% mortality was obtained in the group inoculated with *V. neptunius* wild-type strain 4 days after the challenge (**Figure 6A**) and no mortality was observed in the non-challenged control group. Interestingly, mortality registered after inoculation with Δ absF or Δ irp2 single mutants decreased up to 75-80% (**Figure 6A**), which represents statistically significant differences with respect to the




FIGURE 6 | Survival curves after infection challenge in clams (*Ruditapes philippinarum*) (A) and turbot (B) with *V. neptunius* wild-type strain and its derivatives $\Delta absF$, $\Delta irp2$ and $\Delta irp2\Delta absF$ mutants. Asterisks denote statistically significant differences between strains (* $P < 0.05$; ** $P < 0.01$ and *** $P < 0.001$).

group inoculated with the wild type strain. Most notably, mortality was strongly reduced in the clams group inoculated with the $\Delta absF\Delta irp2$ double mutant, where almost 80% of larvae survived to the challenge (Figure 6A). To test the possible virulence of *V. neptunius* wild type for fish, an experimental infection was done also in fish, using turbot fingerlings. In this case, there was not registered any mortality (Figure 6B), which showed that *V. neptunius* PP-145.98 wild type strain is pathogenic only for bivalve molluscs.

Amphibactin Gene Cluster Transcriptional Organization


Although amphibactin is one of the most abundant siderophores in the seawater (Boiteau et al., 2016; Gauglitz et al., 2021), transcriptional organization of amphibactin genes was not studied to date. To define the transcriptional organization of amphibactin genes, a series of reverse transcriptase PCR reactions (RT-PCRs) were performed. Results are shown in Figure 7. Since all amphibactin genes are encoded in the same DNA strand (Figure 7A), a reverse transcriptase reaction was done by using a primer targeted on *absA*, the last gene of the cluster (Figure 7A). Positive amplification was found in a PCR targeted into the *abtC* gene using previously obtained cDNA as template. This result clearly shows that amphibactin genes are transcribed into a polycistronic mRNA spanning from *abtC* to *absA* genes. Consequently, they must be transcribed from a divergent promoter located in the *entD-abtC* intergenic region (Figure 7B). Nevertheless, this result does not rule out the possibility that additional promoters exist within the gene cluster driving independent transcription of some genes. *In silico* analysis of the intergenic sequences of the amphibactin cluster suggests the existence of two putative FUR box motifs: one located 113 bp upstream of *abtC* start codon (GCAAACCATTTTCATTTGC) and another one located 83 bp upstream of *abtA* (*absF-abtA* intergenic region) (GATAACCATTTATCATTAGC) (Figure 7A). *AbtA* was previously characterized as the TonB-dependent outer membrane transporter required for ferri-amphibactin internalization (Galvis et al., 2020). These Fur box motifs showed an identity of 58 and 79%, respectively, to the FUR

box motif consensus sequence of *E. coli* (GATAATGATA ATCATTATCATTATC) (Ochsner and Vasil, 1996; Payne et al., 2016). Thus, the existence of a promoter upstream of *abtA* controlling the expression of amphibactin transporter will be further studied.

Both Siderophore Systems, Amphibactin and Piscibactin, Are Significantly Expressed When *V. neptunius* Is Cultivated Under Iron Deficiency

To evaluate the transcription levels of both amphibactin and piscibactin siderophore systems, the promoter regions of each siderophore gene cluster were cloned into the promoterless plasmid pHRP309 upstream of the *lacZ* gene. Resulting plasmids were mobilized into *V. neptunius* PP-145.98 wild type strain and the transcription levels of each promoter were evaluated *in vitro* and *ex vivo* (Figure 8A). To identify amphibactin promoters, fusions of the putative promoters *abtA*, *abtC*, and the sequence upstream of *absE* were evaluated by measuring β -galactosidase activity. These regions were named *PabtA*, *PabtC* and *PabsE*, respectively (Figure 8A). On the other hand, since *ParaC1* and *PfrpA* were previously characterized as the promoters that control piscibactin expression in *V. anguillarum* (Balado et al., 2018), *lacZ* fusions of the *V. neptunius* sequences upstream of *frpA* and *araC1* (*PfrpA* and *ParaC1*) were also obtained. The *PabtA* promoter cloned in reverse orientation was used as negative control and its transcriptional levels were almost undetectable under all the growth conditions tested (data not shown).

Piscibactin promoters (*ParaC1* and *PfrpA*) as well as amphibactin promoters *PabtC* and *PabtA* were assayed under low-iron availability *in vitro* (CM9 with 50 μ M dipyrindyl). The results show that all these promoters displayed high β -galactosidase activities (Figure 8A). However, the sequence upstream of *absE* showed an almost undetectable β -galactosidase activity under all the conditions tested (Figure 8A). Thus, the presence of an independent promoter upstream of the biosynthetic gene *absE* was discarded. It is noteworthy that while piscibactin promoter *PfrpA* achieved an activity of ca.



22,000 U under low iron availability, the activity of *PabtC*, *PabtA* and *ParaC1* were 50% lower (ca. 10,000 U). These results would suggest that, under iron deficiency, the expression of piscibactin genes would be predominant. As expected, the addition of 10 μ M FeCl_3 to the culture medium greatly reduced (>40%) the transcriptional activity of amphibactin and piscibactin promoters, indicating that the two siderophore systems are significantly less expressed when iron availability increase (**Figure 8A**). The transcriptional activity of the constitutive promoter *proC* was independent of iron concentration.

To assess the expression of the siderophore systems during bacteria-bivalve interaction, an *ex vivo* assay was performed by the incubation of *V. neptunius lacZ* fusion carrier strains in mussel hemolymph (*Mytilus galloprovincialis*). Interestingly, the expression pattern registered *ex vivo* (**Figure 8B**) showed some differences with that observed *in vitro* (**Figure 8A**). Specifically, the transcriptional activity of the amphibactin promoter *PabtC* was 2-fold higher when incubated in hemolymph, achieving ca. 25,000 U. Thus, while the piscibactin promoter *PfprA* showed *in vivo* a notable higher expression than amphibactin promoter *PabtC*, there were no statistically significant differences between them *ex vivo*. Addition of 10 μ M FeCl_3 to the hemolymph also significantly reduced transcriptional activity of both siderophore systems, which suggests that iron limitation must be a signal that bacteria detect when entering the host (**Figure 8B**).

Piscibactin and Amphibactin Are Present in Most Pathogenic *Vibrio* Species of the Coralliilyticus Clade

To evaluate the distribution of amphibactin and piscibactin siderophore systems in species of *Vibrio* with importance in bivalve aquaculture, a total of 234 genomes of *Vibrio* species belonging to the Coralliilyticus, Anguillarum, Harveyi, Orientalis, Pecteneccida and Splendidus clades were tested *in silico* for the presence of complete amphibactin and piscibactin gene clusters. These species were selected since they are commonly reported as bacteria associated with mortality events in larvae and spats of bivalves hatcheries and their virulence for bivalves is well-established (Beaz-Hidalgo et al., 2010; Dubert et al., 2017). The results (**Figure 9** and **Table S3**) showed that most *Vibrio* genomes belonging to the Coralliilyticus clade harbor complete gene clusters encoding amphibactin and piscibactin siderophore systems. All *V. neptunius*, *V. ostreicida* and *V. coralliilyticus* genomes available in GenBank harbor close homologues of amphibactin genes, sharing identical structure and an average identity of 99%, 67% and 84%, respectively, with those found in *V. neptunius* PP-145.98 strain. The genomic island *irp*-HPI is present in all *V. neptunius* and *V. ostreicida* genomes sequenced and also in most *V. coralliilyticus* (18 of the 29). These results altogether suggest that most *Vibrio* species that belong to the Coralliilyticus




FIGURE 8 | Transcriptional activity (β -galactosidase units) of piscibactin (*ParaC1* and *PfrpA*) and amphibactin (*PabtC* and *PabtA*) promoters carried out using *lacZ* fusions. Expression assay was performed *in vitro* by cultivating *V. neptunius* PP-145.98 in CM9 minimal medium supplemented with FeCl₃ 10 μ M (iron excess) or 2,2'-dipyridyl 50 μ M (iron deficiency) (A) and *ex vivo* by cultivating bacteria in mussel hemolymph (B). *PabtA* promoter cloned in reverse orientation upstream of the *lacZ* gene was used as a negative control. A *lacZ* fusion with the constitutive promoter *proC* (*PproC*) was used as control. Asterisks indicate statistically significant differences between bars ($P < 0.05$); ns, no statistically significant differences. Error bars denote standard deviations.

clade harbor close homologues of both amphibactin and piscibactin siderophore systems. In addition, the amphibactin system is also widespread among vibrios of the Orientalis clade (*V. bivalvicida*, *V. europaeus* and *V. tubiashii*). Finally, among vibrios of the Splendidus clade, 9.5% of *V. splendidus* and 25% of *V. tasmaniensis* genomes also harbor amphibactin genes.

DISCUSSION

Bacterial virulence is context-dependent and mainly depends on the interaction among diverse virulence factors functioning at the same time or sequentially when bacteria enter the host (Won and Park, 2008). The main virulence factors characterized to date in bacteria pathogenic for marine bivalve molluscs are those related to motility, chemotaxis and, in a major extent, to extracellular products (de O Santos et al., 2011; Kimes et al.,

2012; Ushijima et al., 2014; Ushijima et al., 2018). In mollusc tissues, iron is mainly associated to ferritins, whose role includes transport, detoxification and iron storage (Webb et al., 1991). Hemolymph bacterial inhabitants include both mutualistic and pathogenic bacteria that must overcome hemolymph antibacterial activity and compete for the nutrients present in the host sources (Vezzulli et al., 2018).

Although the *V. neptunius* pathogenicity is not yet well known, some studies made with *V. coralliilyticus* could shed some light to the *V. neptunius* virulence factors, since these are two close-related bacteria (Thompson et al., 2003; Sawabe et al., 2013; Dubert et al., 2017). *V. coralliilyticus* pathogenicity depends on the coordinated expression of multiple virulence factors such as flagella, quorum sensing (QS), T6SS, and extracellular products including hemolysins and proteases (Jeffries, 1982; Austin et al., 2005; de O Santos et al., 2011; Kimes et al., 2012; Richards et al., 2015; Guillemette et al., 2020). It has been suggested that siderophore




FIGURE 9 | Distribution of amphibactin and piscibactin siderophore systems in *Vibrio* species responsible for outbreaks in bivalves hatcheries. Presence (+) or absence (-) of complete piscibactin/amphibactin gene cluster determined by Blast search in WGS database indexed in the NCBI database using the nucleotide sequences of *V. neptunius* PP-145.98 as query. The ratio (*) indicates the number of genomes of a *Vibrio* sp. harboring close homologues with respect to the total number of genomes analyzed of that species. Phylogenetic tree was constructed by maximum-likelihood (GTR+G model) with the concatenated sequences of the five housekeeping genes *ftsZ*, *gyrB*, *mreB*, *pyrH* and *recA*.

production should play a main role in *Vibrio* infection disease affecting bivalve molluscs (Soto-Rodríguez et al., 2003; Gómez-León et al., 2005; Mechri et al., 2017; Zhang et al., 2020). However, this hypothesis was based on the high frequency of siderophore producer strains found in mollusc microbiota and not in functional analysis (Desriac et al., 2014; Hernández-Robles et al., 2016; Leite et al., 2017).

The present work shows that *V. neptunius* PP-145.98 genome harbours a close homologue of the piscibactin system (*irp* genes), which is part of the highly pathogenicity island *irp*-HPI (Osorio et al., 2006; Balado et al., 2018). This genomic island harbours the genes that encode most functions required for piscibactin biosynthesis (*irp123459*) and uptake (*frpABC*), but lacks a gene homolog to *entD*, which encodes an 4'-phosphopantetheinyl transferase (PPTases) (Osorio et al., 2015; Balado et al., 2018). PPTases are required to activate the non-ribosomal peptide synthetases (NRPSs) assembly lines enabling siderophore synthesis (Orikasa et al., 2006). This function must be supplied by one of the two PPTases found in *V. neptunius* genome (loci WP_206370543.1 and WP_206368753.1). Genetic and chemical data proved that *irp* genes of *V. neptunius* PP-145.98 are expressed and that piscibactin is being synthesized. Since our previous works demonstrated that *V. neptunius* PP-145.98 also produces the siderophore amphibactin (Galvis et al., 2020), *V. neptunius* PP-145.98 would produce two different siderophores, amphibactin and piscibactin, simultaneously. This is not a rare characteristic, since the same feature has been reported in other bacteria such as *Pseudomonas aeruginosa* (Cornelis and

Dingemans, 2013) or *Escherichia coli* (Valdebenito et al., 2006), or in the fish pathogens *V. anguillarum* (Balado et al., 2018) and *Aeromonas salmonicida* (Balado et al., 2015). The synthesis of siderophores demands a high energetic cost to the bacterium (Cornelis et al., 2011). Thus, production of redundant siderophore systems by a cell is deleterious (Cordero et al., 2012). However, in some cases, switching between siderophore systems would enhance niche flexibility (Sandy and Butler, 2009; Dumas et al., 2013; Zhang et al., 2019) and pathogenesis (Garcia et al., 2011; Balado et al., 2018).

Inactivation of both siderophore systems, piscibactin and amphibactin, greatly reduced growth capacity of *V. neptunius* under low-iron conditions. Interestingly, amphibactin and piscibactin are siderophores with different chemical characteristics and different affinity for Fe(III). Amphibactin is an amphiphilic siderophore produced by the addition of a variable fatty acid moiety to a head group that coordinates a strong affinity for iron. Thus, amphibactin is mainly associated to the cell membrane, which would minimize the diffusive loss of siderophore in the marine environment (Martinez et al., 2003; Boiteau et al., 2016). On the other hand, piscibactin is a labile yersiniabactin-like siderophore with low affinity for iron (Souto et al., 2012). Congruently, the *V. neptunius* $\Delta irp2$ single mutant, that only produces amphibactin, showed higher growth under strong iron restriction than the piscibactin-only producer *V. neptunius* $\Delta abtF$ mutant. The affinity of a siderophore for iron does not predict their impact in virulence, e.g. highly virulent *V. anguillarum* strains produce piscibactin and also vanchrobactin,

a siderophore with higher affinity for iron, but piscibactin production is the most relevant for virulence (Balado et al., 2018).

Infection challenges showed that inactivation of siderophore synthesis strongly reduced the mortality caused by *V. neptunius* in clams. Thus, our results proved that siderophore production constitutes a key virulence factor of pathogenic *Vibrios* in molluscs. Interestingly, inactivation of one of the two siderophore systems (either amphibactin or piscibactin) significantly reduced virulence compared to the wild type strain. This finding showed that the ability to produce both siderophores maximises virulence of *V. neptunius*. In this concern, piscibactin a yersiniabactin-like siderophore, belongs to a type of siderophores that could have other roles during infection besides iron uptake, since they could promote bacterial colonization, dissemination and resistance against phagocytosis (Chaturvedi et al., 2012; Bobrov et al., 2014; Holden and Bachman, 2015; Koh and Henderson, 2015). Thus, piscibactin and amphibactin could have specialized role(s) in the survival of *V. neptunius* within the host and the environment. However, the fact that *V. neptunius* single mutants deficient in each one of these siderophores have showed indistinguishable virulence properties, suggests that both siderophore systems contribute equally to *V. neptunius* virulence.

Although amphibactin is one of the most abundant siderophores in seawater (Boiteau et al., 2016; Gauglitz et al., 2021), transcriptional organization of amphibactin genes was not studied to date. The results demonstrated that amphibactin and piscibactin siderophore systems are significantly expressed when *V. neptunius* is cultivated under low-iron availability *in vitro*. More notably, both siderophore systems are also simultaneously expressed at similar levels when *V. neptunius* is cultivated in bivalve hemolymph (*ex vivo*). The co-expression of siderophores with different chemical properties could contribute to enhance bacterial fitness during host-pathogen interactions (Dumas et al., 2013; Lages et al., 2019). Thus, these results confirm that the high virulence for clams showed by *V. neptunius* wild-type strain must depend on the simultaneous production of piscibactin and amphibactin.

Austin and col. (2005) showed that the *V. neptunius* LMG 20536 and LMG 20610 strains are pathogenic to rainbow trout causing up to 100% mortality at a dose of 10^6 cells per fish inoculated intraperitoneally. The present work showed that, although PP-145.98 strain harbors the Highly Pathogenicity Island *irp*-HPI and produces the siderophore piscibactin, wild type strain does not cause mortality in turbot (*Scophthalmus maximus*). Interestingly, Austin and col. (2005) also showed that while extracellular products (ECPs) of *V. neptunius* LMG 20610 were highly toxic for fish, ECPs of *V. neptunius* LMG 20536 were almost non toxic (Austin et al., 2005). These findings greatly suggest the existence of a high heterogeneity among *V. neptunius* strains and are in agreement with previous works showing that the pathogenicity mechanisms necessary for infecting fish or shellfish are different (Won and Park, 2008; Zhang et al., 2020).

In a previous work it was found that a high proportion of the *Vibrio*-like isolates from mussel hemolymph were PCR positive for a target in amphibactin outer membrane transporter *abtA*

(Galvis et al., 2020). Those *abtA*-positive isolates were mainly non-pathogenic vibrios belonging to the Splendidus clade (Galvis et al., 2020). The *in silico* analysis of 234 genome sequences of well characterized mollusc pathogens belonging to the clades Coralliilyticus, Anguillarum, Harveyi, Orientalis, Pectenecida and Splendidus showed that amphibactin is widespread not only in the Splendidus clade, but also in Orientalis and Coralliilyticus clades. However, *irp*-HPI element encoding piscibactin seems to be limited to the Coralliilyticus clade, since it is present in all *V. neptunius* and *V. ostreicida* genomes and in most *V. coralliilyticus* genomes available. Thus, simultaneous production of both siderophores would be a widespread character in the well-recognized mollusc pathogens *V. coralliilyticus*, *V. neptunius* and *V. ostreicida*. According to the available data, siderophores could have a dual function since they can be key drivers of the microbial community structure and would be key virulence factors that enable infection in an animal host (Vezzulli et al., 2008; Cordero et al., 2012; Kramer et al., 2020; Gauglitz et al., 2021). Rubio-Portillo and col. (2020) showed that siderophore production has a central role in competition among bacteria of mollusc microbiota, enhancing the ability of *Vibrio* coral pathogens such as *V. coralliilyticus* and *V. mediterranei* to invade the host and cause tissue necrosis.

It is known that some bacterial diseases emerged after rapid spread of selectively favoured virulence factors by horizontal gene transfer (HGT) (Bruto et al., 2017; Le Roux and Blokesch, 2018). Molluscs are a major group of marine animals that play key roles in marine ecosystems including water clarification and providing an habitat for other organisms (Schatte Olivier et al., 2020). In addition, molluscs serve as alternative/reservoir hosts for pathogenic bacteria, including human pathogens (Destoumieux-Garzón et al., 2020). Although mollusc microbiota itself would not be a primary source of pathogens for higher animals, the high prevalence of siderophore systems in this niche would imply that the microbiota could serve as test beds (reservoir) of virulence factors like the High-Pathogenicity Island *irp*-HPI encoding the piscibactin system, which is a proved virulence factor in some fish pathogenic bacteria (Osorio et al., 2015; Balado et al., 2018).

In conclusion, synthesis of the siderophores amphibactin and piscibactin, which are widespread in several different *Vibrio* species pathogenic for marine bivalve molluscs, would constitute key virulence factors for disease outbreaks in molluscs aquaculture systems.

DATA AVAILABILITY STATEMENT

Publicly available datasets were analyzed in this study. This data can be found here: NCBI, accession number: JAFHLB000000000.

ETHICS STATEMENT

The animal study was reviewed and approved by Bioethics Committee of the University of Santiago de Compostela.

AUTHOR CONTRIBUTIONS

FG and LA performed the lab experiments. MB, JR, and CJ analyzed the data. FG, LA, CJ, and MB wrote the first draft of the manuscript. MB and ML corrected the draft and built the final version of the manuscript. All authors conceived and designed the study, contributed to manuscript revision, and read and approved the submitted version.

FUNDING

This work was supported by grants AGL2017-86183-R (AEI/FEDER, EU), RTI2018-093634-B-C21/C22 (AEI/FEDER, EU) and PID2019-103891RJ-100 (AEI) from the State Agency for Research (AEI) of Spain. AGL2017-86183-R and RTI2018-093634-B-C21/C22 were co-funded by the FEDER Programme from the European Union. Work in University of Santiago de Compostela was also supported by grants GRC2018/018 and 2021-CP112, and in University of A Coruña by grant GRC2018/039 from Xunta de Galicia (Spain). FG was financed with a fellowship ‘Programa de formación de recurso humano de alto nivel doctorado en el exterior’ granted by Colciencias and the

government of Norte de Santander, Colombia. LA was financed with a fellowship (ED481A-2019/081) from Xunta de Galicia (Spain), co-financed by ESF (European Social Fund).

ACKNOWLEDGMENTS

The authors acknowledge Proameixa Fernández S.L. and IGafa (Instituto Galego de Formación en Acuicultura) from Galicia, Spain, for providing the seed and larvae of molluscs for the pathogenesis assays. FG also acknowledges a fellowship from ‘Programa de formación de recurso humano de alto nivel doctorado en el exterior’ granted by Colciencias and the government of Norte de Santander, Colombia; and ‘Pasaporte a la Ciencia’ granted by Ictex, Colombia.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fcimb.2021.750567/full#supplementary-material>

REFERENCES

- Austin, B., Austin, D., Sutherland, R., Thompson, F., and Swings, J. (2005). Pathogenicity of Vibrios to Rainbow Trout (*Oncorhynchus mykiss*, Walbaum) and *Artemia nauplii*. *Environ. Microbiol.* 7, 1488–1495. doi: 10.1111/j.1462-2920.2005.00847.x
- Balado, M., Lages, M. A., Fuentes-Monteverde, J. C., Martínez-Matamoros, D., Rodríguez, J., Jiménez, C., et al (2018). The Siderophore Piscibactin is a Relevant Virulence Factor for *Vibrio Anguillarum* Favored at Low Temperatures. *Front. Microbiol.* 9, 1766. doi: 10.3389/fmicb.2018.01766
- Balado, M., Osorio, C. R., and Lemos, M. L. (2006). A Gene Cluster Involved in the Biosynthesis of Vanchrobactin, a Chromosome-Encoded Siderophore Produced by *Vibrio anguillarum*. *Microbiology* 152, 3517–3528. doi: 10.1099/mic.0.29298-0
- Balado, M., Souto, A., Vences, A., Careaga, V. P., Valderrama, K., Segade, Y., et al (2015). Two Catechol Siderophores, Acinetobactin and Amonabactin, Are Simultaneously Produced by *Aeromonas salmonicida subsp. salmonicida* Sharing Part of the Biosynthetic Pathway. *ACS Chem. Biol.* 10, 2850–2860. doi: 10.1021/acschembio.5b00624
- Beaz-Hidalgo, R., Balboa, S., Romalde, J. L., and Figueras, M. J. (2010). Diversity and Pathogenicity of *Vibrio* Species in Cultured Bivalve Molluscs. *Environ. Microbiol. Rep.* 2, 34–43. doi: 10.1111/j.1758-2229.2010.00135.x
- Blin, K., Shaw, S., Steinke, K., Villebro, R., Ziemert, N., Lee, S. Y., et al (2019). antiSMASH 5.0: Updates to the Secondary Metabolite Genome Mining Pipeline. *Nucleic Acids Res.* 47, W81–W87. doi: 10.1093/nar/gkz310
- Bobrov, A. G., Kirillina, O., Fetherston, J. D., Miller, M. C., Burlison, J. A., and Perry, R. D. (2014). The *Yersinia Pestis* Siderophore, Yersiniabactin, and the ZnuABC System Both Contribute to Zinc Acquisition and the Development of Lethal Septicaemic Plague in Mice. *Mol. Microbiol.* 93, 759–775. doi: 10.1111/mmi.12693
- Boiteau, R. M., Mende, D. R., Hawco, N. J., McIlvin, M. R., Fitzsimmons, J. N., Saito, M. A., et al (2016). Siderophore-Based Microbial Adaptations to Iron Scarcity Across the Eastern Pacific Ocean. *Proc. Natl. Acad. Sci. U. S. A.* 113, 14237–14242. doi: 10.1073/pnas.1608594113
- Bruto, M., James, A., Petton, B., Labreuche, Y., Chenivesse, S., Alunno-Bruscia, M., et al (2017). *Vibrio Crassostreae*, a Benign Oyster Colonizer Turned Into a Pathogen After Plasmid Acquisition. *ISME J.* 11, 1043–1052. doi: 10.1038/ismej.2016.162
- Cahill, P. L., Fidler, A. E., Hopkins, G. A., and Wood, S. A. (2016). Geographically Conserved Microbiomes of Four Temperate Water Tunicates. *Environ. Microbiol. Rep.* 8, 470–478. doi: 10.1111/1758-2229.12391
- Chaturvedi, K. S., Hung, C. S., Crowley, J. R., Stapleton, A. E., and Henderson, J. P. (2012). The Siderophore Yersiniabactin Binds Copper to Protect Pathogens During Infection. *Nat. Chem. Biol.* 8, 731–736. doi: 10.1038/nchembio.1020
- Cordero, O. X., Ventouras, L.-A., DeLong, E. F., and Polz, M. F. (2012). Public Good Dynamics Drive Evolution of Iron Acquisition Strategies in Natural Bacterioplankton Populations. *Proc. Natl. Acad. Sci. U. S. A.* 109, 20059–20064. doi: 10.1073/pnas.1213344109
- Cornelis, P., and Dingemans, J. (2013). *Pseudomonas Aeruginosa* Adapts its Iron Uptake Strategies in Function of the Type of Infections. *Front. Cell. Infect. Microbiol.* 3, 75. doi: 10.3389/fcimb.2013.00075
- Cornelis, P., Wei, Q., Andrews, S. C., and Vinckx, T. (2011). Iron Homeostasis and Management of Oxidative Stress Response in Bacteria. *Metallomics* 3, 540. doi: 10.1039/c1mt00022e
- de O Santos, E., Alves, N., Dias, G. M., Mazotto, A. M., Vermelho, A., Vora, G. J., et al (2011). Genomic and Proteomic Analyses of the Coral Pathogen *Vibrio coralliilyticus* Reveal a Diverse Virulence Repertoire. *ISME J.* 5, 1471–1483. doi: 10.1038/ismej.2011.19
- Desriac, F., Le Chevalier, P., Brillet, B., Leguerinel, I., Thuillier, B., Paillard, C., et al (2014). Exploring the Hologenome Concept in Marine Bivalvia: Haemolymph Microbiota as a Pertinent Source of Probiotics for Aquaculture. *FEMS Microbiol. Lett.* 350, 107–116. doi: 10.1111/1574-6968.12308
- Destoumieux-Garzon, D., Canesi, L., Oyanedel, D., Travers, M. A., Charrière, G. M., Pruzzo, C., et al (2020). *Vibrio*-bivalve Interactions in Health and Disease. *Environ. Microbiol.* 22, 4323–4341. doi: 10.1111/1462-2920.15055
- Dubert, J., Barja, J. L., and Romalde, J. L. (2017). New Insights Into Pathogenic Vibrios Affecting Bivalves in Hatcheries: Present and Future Prospects. *Front. Microbiol.* 8, 762. doi: 10.3389/fmicb.2017.00762
- Dubert, J., Nelson, D. R., Spinard, E. J., Kessner, L., Gomez-Chiarri, M., Costa, F., et al (2016). Following the Infection Process of Vibrios in Manila Clam (*Ruditapes philippinarum*) Larvae Through GFP-Tagged Pathogenic *Vibrio* Species. *J. Invertebr. Pathol.* 133, 27–33. doi: 10.1016/j.jip.2015.11.008
- Dumas, Z., Ross-Gillespie, A., and Kummerli, R. (2013). Switching Between Apparently Redundant Iron-Uptake Mechanisms Benefits Bacteria in Changeable Environments. *Proc. Biol. Sci.* 280, 20131055. doi: 10.1098/rspb.2013.1055

- Engel, S., Jensen, P. R., and Fenical, W. (2002). Chemical Ecology of Marine Microbial Defense. *J. Chem. Ecol.* 28, 1971–1985. doi: 10.1023/A:1020793726898
- Farto, R., Fichi, G., Gestal, C., Pascual, S., and Nieto, T. P. (2019). “Bacteria-Affecting Cephalopods,” in *Handbook of Pathogens and Diseases in Cephalopods* (Cham: Springer International Publishing), 127–142. doi: 10.1007/978-3-030-11330-8_8
- Finn, R. D., Bateman, A., Clements, J., Coggill, P., Eberhardt, R. Y., Eddy, S. R., et al (2014). Pfam: The Protein Families Database. *Nucleic Acids Res.* 42, D222–D230. doi: 10.1093/nar/gkt1223
- Galvis, F., Ageitos, L., Martínez-Matamoros, D., Barja, J. L., Rodríguez, J., Lemos, M. L., et al (2020). The Marine Bivalve Mollusks Pathogen *Vibrio neptunius* Produces the Siderophore Amphibactin, Which Is Widespread in Molluscs Microbiota. *Environ. Microbiol.* 22, 5467–5482. doi: 10.1111/1462-2920.15312
- Galvis, F., Barja, J. L., Lemos, M. L., and Balado, M. (2021). The Vibriolysin-Like Protease VnpA and the Collagenase ColA Are Required for Full Virulence of the Bivalve Mollusks Pathogen *Vibrio neptunius*. *Antibiotics* 10, 391. doi: 10.3390/antibiotics10040391
- García-Amado, M. A., Bozo-Hurtado, L., Astor, Y., Suárez, P., and Chistoserdov, A. (2011). Denaturing Gradient Gel Electrophoresis Analyses of the Vertical Distribution and Diversity of *Vibrio* Spp. Populations in the Cariaco Basin. *FEMS Microbiol. Ecol.* 77, 347–356. doi: 10.1111/j.1574-6941.2011.01116.x
- García, E. C., Brumbaugh, A. R., and Mobley, H. L. T. (2011). Redundancy and Specificity of *Escherichia Coli* Iron Acquisition Systems During Urinary Tract Infection. *Infect. Immun.* 79, 1225–1235. doi: 10.1128/IAI.01222-10
- Gauglitz, J. M., Boiteau, R. M., McLean, C., Babcock-Adams, L., McIlvin, M. R., Moran, D. M., et al (2021). Dynamic Proteome Response of a Marine *Vibrio* to a Gradient of Iron and Ferrioxamine Bioavailability. *Mar. Chem.* 229, 103913. doi: 10.1016/j.marchem.2020.103913
- Gómez-León, J., Villamil, L., Lemos, M. L., Novoa, B., and Figueras, A. (2005). Isolation of *Vibrio alginolyticus* and *Vibrio splendidus* From Aquacultured Carpet Shell Clam (*Ruditapes decussatus*) Larvae Associated With Mass Mortalities. *Appl. Environ. Microbiol.* 71, 98–104. doi: 10.1128/AEM.71.1.98-104.2005
- Guillemette, R., Ushijima, B., Jalan, M., Häse, C. C., and Azam, F. (2020). Insight Into the Resilience and Susceptibility of Marine Bacteria to T6SS Attack by *Vibrio cholerae* and *Vibrio coralliilyticus*. *PLoS One* 15 (1), e0227864. doi: 10.1371/journal.pone.0227864
- Hernández-Robles, M. F., Álvarez-Contreras, A. K., Juárez-García, P., Natividad-Bonifacio, I., Curiel-Quesada, E., Vázquez-Salinas, C., et al (2016). Virulence Factors and Antimicrobial Resistance in Environmental Strains of *Vibrio alginolyticus*. *Int. Microbiol.* 19, 191–198. doi: 10.2436/20.1501.01.277
- Herrero, M., de Lorenzo, V., and Timmis, K. N. (1990). Transposon Vectors Containing Non-Antibiotic Resistance Selection Markers for Cloning and Stable Chromosomal Insertion of Foreign Genes in Gram-Negative Bacteria. *J. Bacteriol.* 172, 6557–6567. doi: 10.1128/jb.172.11.6557-6567.1990
- Holden, V. I., and Bachman, M. A. (2015). Diverging Roles of Bacterial Siderophores During Infection. *Metallomics* 7, 986–995. doi: 10.1039/c4mt00333k
- Jeffries, V. E. (1982). Three *Vibrio* Strains Pathogenic to Larvae of *Crassostrea Gigas* and *Ostrea Edulis*. *Aquaculture* 29, 201–226. doi: 10.1016/0044-8486(82)90136-3
- Kesarcodi-Watson, A., Kaspar, H., Lategan, M., and Gibson, L. (2009a). Challenge of New Zealand Greenshell™ Mussel *Perna canaliculus* Larvae Using Two *Vibrio* Pathogens: A Hatchery Study. *Dis. Aquat. Organ.* 86, 15–20. doi: 10.3354/dao02100
- Kesarcodi-Watson, A., Kaspar, H., Lategan, M. J., and Gibson, L. (2009b). Two Pathogens of Greenshell™ Mussel Larvae, *Perna canaliculus*: *Vibrio splendidus* and a *V. coralliilyticus/neptunius* -Like Isolate. *J. Fish Dis.* 32, 499–507. doi: 10.1111/j.1365-2761.2009.01006.x
- Kimes, N. E., Grim, C. J., Johnson, W. R., Hasan, N. A., Tall, B. D., Kothary, M. H., et al (2012). Temperature Regulation of Virulence Factors in the Pathogen *Vibrio coralliilyticus*. *ISME J.* 6, 835–846. doi: 10.1038/ismej.2011.154
- Koh, E.-I., and Henderson, J. P. (2015). Microbial Copper-Binding Siderophores at the Host-Pathogen Interface. *J. Biol. Chem.* 290, 18967–18974. doi: 10.1074/jbc.R115.644328
- Kramer, J., Özkaya, Ö., and Kümmerli, R. (2020). Bacterial Siderophores in Community and Host Interactions. *Nat. Rev. Microbiol.* 18, 152–163. doi: 10.1038/s41579-019-0284-4
- Kumar, S., Stecher, G., Li, M., Knyaz, C., and Tamura, K. (2018). MEGA X: Molecular Evolutionary Genetics Analysis Across Computing Platforms. *Mol. Biol. Evol.* 35, 1547–1549. doi: 10.1093/molbev/msy096
- Lages, M. A., Balado, M., and Lemos, M. L. (2019). The Expression of Virulence Factors in *Vibrio Anguillarum* Is Dually Regulated by Iron Levels and Temperature. *Front. Microbiol.* 10, 2335. doi: 10.3389/FMICB.2019.02335
- Lambalot, R. H., Gehring, A. M., Flugel, R. S., Zuber, P., LaCelle, M., Marahiel, M. A., et al (1996). A New Enzyme Superfamily - the Phosphopantetheinyl Transferases. *Chem. Biol.* 3, 923–936. doi: 10.1016/s1074-5521(96)90181-7
- Leite, L., Jude-Lemeilleur, F., Raymond, N., Henriques, I., Garabetian, F., and Alves, A. (2017). Phylogenetic Diversity and Functional Characterization of the Manila Clam Microbiota: A Culture-Based Approach. *Environ. Sci. Pollut. Res.* 24, 21721–21732. doi: 10.1007/s11356-017-9838-z
- Lemos, M. L., Salinas, P., Toranzo, A. E., Barja, J. L., and Crosa, J. H. (1988). Chromosome-Mediated Iron Uptake System in Pathogenic Strains of *Vibrio anguillarum*. *J. Bacteriol.* 170, 1920–1925. doi: 10.1128/jb.170.4.1920-1925.1988
- Le Roux, F., and Blokesch, M. (2018). Eco-Evolutionary Dynamics Linked to Horizontal Gene Transfer in Vibrios. *Annu. Rev. Microbiol.* 72, 89–110. doi: 10.1146/annurev-micro-090817-062148
- Liu, Q., Ma, Y., Zhou, L., and Zhang, Y. (2005). Gene Cloning, Expression and Functional Characterization of a Phosphopantetheinyl Transferase From *Vibrio anguillarum* Serotype O1. *Arch. Microbiol.* 183, 37–44. doi: 10.1007/s00203-004-0745-6
- Lokmer, A., Goedknecht, M. A., Thielges, D. W., Fiorentino, D., Kuenzel, S., Baines, J. F., et al (2016). Spatial and Temporal Dynamics of Pacific Oyster Hemolymph Microbiota Across Multiple Scales. *Front. Microbiol.* 7, 1367. doi: 10.3389/fmicb.2016.01367
- Martinez, J. S., Carter-franklin, J. N., Mann, E. L., Martin, J. D., Haygood, M. G., and Butler, A. (2003). Structure and Membrane Affinity of a Suite of Amphiphilic Siderophores Produced by a Marine Bacterium. *Proc. Natl. Acad. Sci.* 100, 3754–3759. doi: 10.1073/pnas.0637444100
- McFall-Ngai, M., Hadfield, M. G., Bosch, T. C. G., Carey, H. V., Domazet-Lošo, T., Douglas, A. E., et al (2013). Animals in a Bacterial World, a New Imperative for the Life Sciences. *Proc. Natl. Acad. Sci.* 110, 3229–3236. doi: 10.1073/pnas.1218525110
- McMillan, D. G. G., Velasquez, I., Nunn, B. L., Goodlett, D. R., Hunter, K. A., Lamont, I., et al (2010). Acquisition of Iron by Alkaliphilic *Bacillus* species. *Appl. Environ. Microbiol.* 76, 6955–6961. doi: 10.1128/AEM.01393-10
- Mechri, B., Monastiri, A., Medhioub, A., Medhioub, M. N., and Aouni, M. (2017). Molecular Characterization and Phylogenetic Analysis of Highly Pathogenic *Vibrio alginolyticus* Strains Isolated During Mortality Outbreaks in Cultured *Ruditapes decussatus* Juvenile. *Microb. Pathog.* 111, 487–496. doi: 10.1016/J.MICPATH.2017.09.020
- Miller, J. H. (1992). *A Short Course in Bacterial Genetics* (Plainview, N.Y: Cold Spring Harbor Laboratory Press).
- Mourinho, S., Osorio, C. R., and Lemos, M. L. (2004). Characterization of Heme Uptake Cluster Genes in the Fish Pathogen *Vibrio Anguillarum*. *J. Bacteriol.* 186, 6159–6167. doi: 10.1128/JB.186.18.6159-6167.2004
- Ochsner, U. A., and Vasil, M. L. (1996). Gene Repression by the Ferric Uptake Regulator in *Pseudomonas Aeruginosa*: Cycle Selection of Iron-Regulated Genes. *Proc. Natl. Acad. Sci. U. S. A.* 93, 4409–4414. doi: 10.1073/pnas.93.9.4409
- Orikasa, Y., Nishida, T., Hase, A., Watanabe, K., Morita, N., and Okuyama, H. (2006). A Phosphopantetheinyl Transferase Gene Essential for Biosynthesis of N - 3 Polyunsaturated Fatty Acids From *Moritella marina* Strain MP-1. *FEBS Lett.* 580, 4423–4429. doi: 10.1016/j.febslet.2006.07.008
- Osorio, C. R., Juiz-Rio, S., and Lemos, M. L. (2006). A Siderophore Biosynthesis Gene Cluster From the Fish Pathogen *Photobacterium Damselae* Subsp. *Piscicida* Is Structurally and Functionally Related to the *Yersinia* High-Pathogenicity Island. *Microbiology* 152, 3327–3341. doi: 10.1099/mic.0.29190-0
- Osorio, C. R., Rivas, A. J., Balado, M., Fuentes-Monteverde, J. C., Rodríguez, J., Jiménez, C., et al (2015). A Transmissible Plasmid-Borne Pathogenicity Island Confers Piscibactin Biosynthesis in the Fish Pathogen *Photobacterium damsela* Subsp. *Piscicida*. *Appl. Environ. Microbiol.* 81, 5867–5879. doi: 10.1128/AEM.01580-15
- Paillard, C., Le Roux, F., and Borrego, J. J. (2004). Bacterial Disease in Marine Bivalves, a Review of Recent Studies: Trends and Evolution. *Aquat. Living Resour.* 17, 477–498. doi: 10.1051/alr:2004054

- Parales, R. E., and Harwood, C. S. (1993). Construction and Use of a New Broad-Host-Range *lacZ* Transcriptional Fusion Vector, pHRP309, for Gram-Bacteria. *Gene* 133, 23–30. doi: 10.1016/0378-1119(93)90220-W
- Payne, S. M., Mey, A. R., and Wyckoff, E. E. (2016). *Vibrio* Iron Transport: Evolutionary Adaptation to Life in Multiple Environments. *Microbiol. Mol. Biol. Rev.* 80, 69–90. doi: 10.1128/MMBR.00046-15
- Potgieter, M., Bester, J., Kell, D. B., and Pretorius, E. (2015). The Dormant Blood Microbiome in Chronic, Inflammatory Diseases. *FEMS Microbiol. Rev.* 39, 567–591. doi: 10.1093/femsre/fuv013
- Prado, S., Romalde, J., Montes, J., and Barja, J. (2005). Pathogenic Bacteria Isolated From Disease Outbreaks in Shellfish Hatcheries. First Description of *Vibrio neptunius* as an Oyster Pathogen. *Dis. Aquat. Organ.* 67, 209–215. doi: 10.3354/dao067209
- Richards, G. P., Watson, M. A., Needleman, D. S., Church, K. M., and Häse, C. C. (2015). Mortalities of Eastern and Pacific Oyster Larvae Caused by the Pathogens *Vibrio Coralliilyticus* and *Vibrio Tubiashii*. *Appl. Environ. Microbiol.* 81, 292–297. doi: 10.1128/AEM.02930-14
- Romalde, J. L., Dieguez, A. L., Lasa, A., and Balboa, S. (2014). New *Vibrio* Species Associated to Molluscan Microbiota: A Review. *Front. Microbiol.* 4, 413. doi: 10.3389/fmicb.2013.00413
- Rubio-Portillo, E., Martin-Cuadrado, A. B., Caraballo-Rodríguez, A. M., Rohwer, F., Dorrestein, P. C., and Antón, J. (2020). Virulence as a Side Effect of Interspecies Interaction in *Vibrio* Coral Pathogens. *mbio* 11, 201–221. doi: 10.1128/mBio.00201-20
- Sandy, M., and Butler, A. (2009). Microbial Iron Acquisition: Marine and Terrestrial Siderophores. *Chem. Rev.* 109, 4580–4595. doi: 10.1021/cr9002787
- Sawabe, T., Ogura, Y., Matsumura, Y., Feng, G., Amin, A. R., Mino, S., et al (2013). Updating the *Vibrio* Clades Defined by Multilocus Sequence Phylogeny: Proposal of Eight New Clades, and the Description of *Vibrio tritonius* Sp. Nov. *Front. Microbiol.* 4, 414. doi: 10.3389/fmicb.2013.00414
- Schatte Olivier, A., Jones, L., Vay, L., Christie, M., Wilson, J., and Malham, S. K. (2020). A Global Review of the Ecosystem Services Provided by Bivalve Aquaculture. *Rev. Aquacult.* 12, 3–25. doi: 10.1111/raq.12301
- Schwyn, B., and Neilands, J. B. (1987). Universal Chemical Assay for the Detection and Determination of Siderophores. *Anal. Biochem.* 160, 47–56. doi: 10.1016/0003-2697(87)90612-9
- Soto-Rodríguez, S., Roque, A., Lizarraga-Partida, M., Guerra-Flores, A., and Gomez-Gil, B. (2003). Virulence of Luminous *Vibrios* to *Artemia franciscana* Nauplii. *Dis. Aquat. Organ.* 53, 231–240. doi: 10.3354/dao053231
- Souto, A., Montaos, M. A. M. A., Rivas, A. J. A. J., Balado, M., Osorio, C. R., Rodríguez, J., et al (2012). Structure and Biosynthetic Assembly of Piscibactin, a Siderophore From *Photobacterium damsela* Subsp. *Piscicida*, Predicted From Genome Analysis. *Eur. J. Org. Chem.* 2012, 5693–5700. doi: 10.1002/ejoc.201200818
- Thompson, F. L., Li, Y., Gomez-Gil, B., Thompson, C. C., Hoste, B., Vandemeulebroecke, K., et al (2003). *Vibrio Neptunius* Sp. Nov., *Vibrio brasiliensis* Sp. Nov. And *Vibrio xuii* Sp. Nov., Isolated From the Marine Aquaculture Environment (Bivalves, Fish, Rotifers and Shrimps). *Int. J. Syst. Evol. Microbiol.* 53, 245–252. doi: 10.1099/ijs.0.02447-0
- Toranzo, A. E., Magariños, B., and Romalde, J. L. (2005). A Review of the Main Bacterial Fish Diseases in Mariculture Systems. *Aquaculture* 246, 37–61. doi: 10.1016/j.aquaculture.2005.01.002
- Travers, M.-A., Boettcher Miller, K., Roque, A., and Friedman, C. S. (2015). Bacterial Diseases in Marine Bivalves. *J. Invertebr. Pathol.* 131, 11–31. doi: 10.1016/j.jip.2015.07.010
- Ushijima, B., Richards, G. P., Watson, M. A., Schubiger, C. B., and Häse, C. C. (2018). Factors Affecting Infection of Corals and Larval Oysters by *Vibrio Coralliilyticus*. *PLoS One* 13, e0199475. doi: 10.1371/journal.pone.0199475
- Ushijima, B., Videau, P., Burger, A. H., Shore-Maggio, A., Runyon, C. M., Sudek, M., et al (2014). *Vibrio Coralliilyticus* Strain OCN008 Is an Etiological Agent of Acute Montipora White Syndrome. *Appl. Environ. Microbiol.* 80, 2102–2109. doi: 10.1128/AEM.03463-13
- Utermann, C., Parrot, D., Breusing, C., Stuckas, H., Staufenberger, T., Blümel, M., et al (2018). Combined Genotyping, Microbial Diversity and Metabolite Profiling Studies on Farmed *Mytilus* Spp. From Kiel Fjord. *Sci. Rep.* 8, 7983. doi: 10.1038/s41598-018-26177-y
- Valdebenito, M., Crumbliss, A. L., Winkelmann, G., and Hantke, K. (2006). Environmental Factors Influence the Production of Enterobactin, Salmochelin, Aerobactin, and Yersiniabactin in *Escherichia coli* Strain Nissle 1917. *Int. J. Med. Microbiol.* 296, 513–520. doi: 10.1016/j.ijmm.2006.06.003
- Vezzulli, L., Guzmán, C. A., Colwell, R. R., and Pruzzo, C. (2008). Dual Role Colonization Factors Connecting *Vibrio cholerae*'s Lifestyles in Human and Aquatic Environments Open New Perspectives for Combating Infectious Diseases. *Curr. Opin. Biotechnol.* 19, 254–259. doi: 10.1016/j.copbio.2008.04.002
- Vezzulli, L., Stagnaro, L., Grande, C., Tassistro, G., Canesi, L., and Pruzzo, C. (2018). Comparative 16S rDNA Gene-Based Microbiota Profiles of the Pacific Oyster (*Crassostrea Gigas*) and the Mediterranean Mussel (*Mytilus Galloprovincialis*) From a Shellfish Farm (Ligurian Sea, Italy). *Microb. Ecol.* 75, 495–504. doi: 10.1007/s00248-017-1051-6
- Wang, R. F., and Kushner, S. R. (1991). Construction of Versatile Low-Copy-Number Vectors for Cloning, Sequencing and Gene Expression in *Escherichia Coli*. *Gene* 100, 195–199. doi: 10.1016/0378-1119(91)90366-J
- Webb, J., St-Pierre, T. G., and Macey, D. J. (1991). “Iron Biomineralization in Invertebrates,” in *Iron Biominerals* (Boston, MA: Springer US), 193–220. doi: 10.1007/978-1-4615-3810-3_14
- Won, K. M., and Park, S. (2008). Pathogenicity of *Vibrio Harveyi* to Cultured Marine Fishes in Korea. *Aquaculture* 285, 8–13. doi: 10.1016/j.aquaculture.2008.08.013
- Zhang, X., Baars, O., and Morel, F. M. M. (2019). Genetic, Structural, and Functional Diversity of Low and High-Affinity Siderophores in Strains of Nitrogen Fixing *Azotobacter Chroococcum*. *Metallomics* 11, 201–212. doi: 10.1039/C8MT00236C
- Zhang, X.-H., He, X., and Austin, B. (2020). *Vibrio Harveyi*: A Serious Pathogen of Fish and Invertebrates in Mariculture. *Mar. Life Sci. Technol.* 2, 231–245. doi: 10.1007/s42995-020-00037-z
- Zhang, X. X., Sun, Z., Zhang, X. X., Zhang, M., and Li, S. (2018). Hemolymph Microbiomes of Three Aquatic Invertebrates as Revealed by a New Cell Extraction Method. *Appl. Environ. Microbiol.* 84, e02824–e02817. doi: 10.1128/AEM.02824-17

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2021 Galvis, Ageitos, Rodríguez, Jiménez, Barja, Lemos and Balado. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.