

Research Article

Sex Differences in the Renal Vascular Responses of AT₁ and Mas Receptors in Two-Kidney-One-Clip Hypertension

Zahra Pezeshki ^{1,2} and Mehdi Nematbakhsh ^{1,2,3}

¹Water and Electrolytes Research Center, Isfahan University of Medical Sciences, Isfahan, Iran

²Department of Physiology, Isfahan University of Medical Sciences, Isfahan, Iran

³Isfahan^{MN} Institute of Basic and Applied Sciences Research, Isfahan, Iran

Correspondence should be addressed to Mehdi Nematbakhsh; nematbakhsh@med.mui.ac.ir

Received 1 October 2020; Revised 5 January 2021; Accepted 20 January 2021; Published 20 February 2021

Academic Editor: Fulvio Morello

Copyright © 2021 Zahra Pezeshki and Mehdi Nematbakhsh. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Background. The prevalence and severity of hypertension, as well as the activity of the systemic and local renin angiotensin systems (RASs), are gender related, with more symptoms in males than in females. However, the underlying mechanisms are not well understood. In this study, we investigated sex differences in renal vascular responses to angiotensin II (Ang II) administration with and without Ang II type 1 and Mas receptor (AT₁R and MasR) antagonists (losartan and A779) in the 2K1C rat model of renovascular hypertension. **Methods.** Male and female 2K1C rats were divided into 8 experimental groups (4 of each sex) treated with vehicle, losartan, A779, or A779+losartan. Responses of mean arterial pressure (MAP), renal blood flow (RBF), and renal vascular resistance (RVR) to Ang II were determined. **Results.** In both sexes, the basal MAP, RBF, and RVR were not significantly different between the four groups during the control period. The Ang II administration decreased RBF and increased RVR in a dose-related manner in both sexes pretreated with vehicle or A779 ($P_{\text{dose}} < 0.0001$), but in vehicle pretreated groups, RBF and RVR responses were different between male and female rats ($P_{\text{group}} < 0.05$). AT₁R blockade increased RBF and decreased RVR responses to Ang II, and no difference between the sexes was detected. Coblockades of AT₁R and MasR receptors increased RBF response to Ang II significantly in males alone but not in females ($P_{\text{group}} = 0.04$). **Conclusion.** The impact of Ang II on RBF and RVR responses seems to be gender related with a greater effect on males, and this sex difference abolishes by Mas receptor blockade. However, the paradoxical role of dual losartan and A779 may provide the different receptor interaction in RAS between male and female rats.

1. Introduction

Hypertension is a complex condition which induces abnormalities such as left ventricular hypertrophy, carotid atherosclerosis, and renal dysfunction [1–3], and it is the leading modifiable risk factor for death [2, 4]. Males have a higher prevalence of hypertension compared to age-matched premenopausal females [5, 6]. Hypertensive nephropathy also is gender dependent [7, 8]. Females are relatively resistant to renal and cardiovascular abnormalities than males. This has been attributed in part to physiologic sex-based differences in the function of the renin angiotensin system (RAS) [6, 9–11]. Despite these apparent gender differences in human hypertension,

similar treatment guidelines are applied in both males and females [8].

Activation of systemic and local RAS via angiotensin II (Ang II) is an important contributing mechanism in multiple forms of hypertension [12–14]. Ang II type 1 receptor antagonists, as well as angiotensin-converting enzyme (ACE) and renin inhibitors, are widely used for treatment of hypertension [12–15]. Therapeutic strategies targeting the RAS have differential effects, in males and females, on the complications of hypertension, including renal disease [12, 14, 16, 17]. The mechanisms that underlie these differences are incompletely defined.

To improve management of hypertension, a better understanding of sex and gender differences in the function of

the RAS, and in particular the renal responses to Ang II, is essential. In females, Ang II type 2 receptor (AT₂R) and Mas receptor (MasR) are more abundant, while Ang II type 1 receptors (AT₁R) are less abundant, in females than males [10, 11, 18]. Ang II has low affinity for the MasR, but interactions between the MasR and AT₁R may alter the Ang II activity [19, 20]. However, these interactions have not been well documented in renovascular hypertension. In the current study, we tested the hypothesis that the relative roles of renal AT₁R and MasR differ in males and females in two-kidney-one-clip (2K1C) renin-dependent hypertension. To test this hypothesis, we examined the effects of blockade of the AT₁R (with losartan) and the MasR (with A779) on the renal response to Ang II in male and female rats with 2K1C hypertension.

2. Materials and Methods

2.1. Animals. The experiments were conducted on 64 (30 males and 34 females) in total age-matched male and female Wistar rats. All animals were housed in cages under controlled conditions (the room temperature of 23–25°C with 12 h light/12 h dark cycle) and had free access to water and rat chow. The experimental procedures were approved by the Ethics Committee of Isfahan Medical Sciences University (ethical code #: IR.MUI.REC.1397.3.461).

2.2. Surgical Procedures and Measurements

2.2.1. Induction of 2K1C Hypertension. Fifty- to sixty-day-old male and female rats were anesthetized with chloral hydrate (450 mg/kg, i.p.) (Merck, Germany) and xylazine (10 mg/kg, i.p.). The animals were subjected to right kidney clipping with a U-shaped silver clip (0.2 mm internal diameter). The right abdominal wall was shaved, and the skin and tissue of the right flank incised to expose the right kidney. The renal artery of the right kidney was isolated, and a clip was placed close to the aorta. Then, the kidney was carefully pushed back into the retroperitoneal cavity and the abdominal wall and skin were sutured. Povidone-iodine 10% was used for skin disinfection before and after surgery. Following the surgery, all rats were monitored until recovery and then individually housed for 24 h in a recovery cage and allowed access to standard rat chow and water ad libitum for 28 days post clipping.

2.2.2. Sham-Operated Rats. We also assigned a total of 15 rats (7 males and 8 females) as sham-operated groups which were subjected to surgical operation process without renal clipping. The data from these sham groups were used to compare the blood pressures from 2K1C groups to verify hypertension in 2K1C rats.

2.2.3. Catheterization and Measurements. Twenty-eight days after clipping, the rats (male: 276 ± 6, female: 184 ± 3 g) were anesthetized with urethane (1.7 g/kg i.p., Merck, Germany). Initially, the animals were tracheostomized to facilitate spontaneous ventilation. Then, the left carotid artery was catheterized for measurement of systolic (SYS)

and mean arterial pressure (MAP). In addition, the left jugular vein was isolated, and the polyethylene catheter was implanted into the vein for drug infusion. Also, the left femoral artery was catheterized to determine renal perfusion pressure (RPP). The arterial pressures were measured by pressure transducers and bridge amplifiers connected to the PowerLab system (AD Instruments, Australia). An incision was made in the subabdominal area to insert a polyethylene catheter in the bladder to facilitate urine flow. Then, with the animals lying on the right side on a flat table, an incision was made in left flank using an electrical surgical cutter and the left kidney was carefully excised. To measure RBF, the renal artery was isolated followed by placing an ultrasonic flow probe (0.7 mm in diameter) (Transonic MA0.7PSB, flow probe, USA) directly around the renal artery. The probe was connected to a flow meter (T402, Transonic Systems Inc., Ithaca, NY 14850 USA). An adjustable occluder also was placed around the aorta between the renal and the mesenteric arteries to regulate RPP during Ang II infusion. MAP, RPP, and RBF were recorded continuously during the experiment. After surgical procedure, the animals were monitored for about 30 minutes to achieve the steady state condition. At the end of this period, the animal's SYS, MAP, RPP, RBF, and RVR (calculated by RPP/RBF ratio) were initially recorded as baseline data (control phase).

2.2.4. Experimental Protocol

(1) Vehicle/Antagonist Infusion. Male rats receive saline (the vehicle group, group 1), AT₁R blocker; losartan (the losartan group, group 2), MasR antagonist; A779 (Bachem Bioscience Inc., King of Prussia, PA, USA) (the A779 group, group 3); and the combination of A779 and losartan (the A779+losartan group, group 4). The female rats also had the same protocol as male rats of groups 1–4. Losartan and A779 were dissolved in 0.9% w/v saline. Losartan and A779 either alone or together were administered in bolus doses of 5 mg·kg⁻¹ and 50 μg·kg⁻¹, respectively, in relative groups, which then were followed by continuous infusions of 5 mg·kg⁻¹·h⁻¹ losartan and 50 μg·kg⁻¹·h⁻¹ A779 with a microsyringe infusion pump (New Era Pump System Inc., Farmingdale, NY, USA) during the experiment. Vehicle-treated groups followed a similar protocol but received the saline vehicle instead of antagonists during the experiment. Subsequently, 30 minutes after vehicle/antagonist infusion, the parameters were measured and considered as data to determine the vehicle/antagonist effects (antagonist phase). However, the vehicle or antagonists were continuously infused until end Ang II administration (end of experiment).

(2) Ang II Infusion. The vascular responses to graded Ang II (Sigma, St Louis MI, USA) infusion were measured in all experimental groups, while the antagonists or vehicle injection continued during the Ang II infusion. Each dose of Ang II (0, 100, 300, or 1000 ng/kg/min) was administered for 15 min, and the last 3-minute measurements in each dose were obtained. During Ang II administration, the aortic diameter was adjusted by an aortic occluder to control RPP at constant level.




FIGURE 1: The protocol of experimental study (see Section 2.2.4).

Finally, the rats were killed humanly, and the right and left kidney were rapidly removed and weighed. The protocol of the experimental study is demonstrated in (Figure 1)

2.2.5. Statistical Analysis. Data are expressed as mean \pm standard error of mean (SEM) and were analyzed using SPSS software (version 20). The baseline data were compared using the one-way analysis of variance (ANOVA) followed by the least significant difference (LSD) test. The paired-sample *T*-test was applied to compare the left and right kidney weights. The responses to antagonists or vehicle are reported as percentage (%) change from the baseline values prior to administration of the antagonists. Responses to Ang II are reported as % change from the average values after administration of the antagonists or its vehicle, but prior to administration of Ang II. These data were then subjected to ANOVA for repeated measures, with the between-subject factor “group” (P_{group}), the within-subject factor “time” (P_{time}), and their interaction ($P_{\text{time}\&\text{group}}$). $P \leq 0.05$ was considered statistically significant.

3. Results

3.1. Verification of 2K1C Implementation. The verification of 2K1C hypertension was determined on the day of experiment after direct blood pressure measurement. Two exclusion criteria were assigned to verify hypertension by 2K1C. The animals with SYS blood pressure lower than 145 mmHg and the rats with infarcted (partially or totally) kidney tissue were excluded from the study. Accordingly, the success rate of the accepted 2K1C hypertensive model was 32% in male and 31% in female rats. Therefore, a total of 64 (30 males and 34 females) 2K1C rats were included in this study. The SYS pressure in male 2K1C rats was 157 ± 2.2 mmHg and in female 2K1C rats was 148.8 ± 2.1 mmHg which was significantly different $P = 0.023$. These data revealed less elevation of SYS pressure in female 2K1C rats compared with male 2K1C rats. The SYS blood pressure in male and female 2K1C groups was also compared with the sham-operated group (Figure 2). Significant differences were detected between 2K1C and sham-operated groups ($P < 0.05$). The comparison between right and left kidney weight per 100 gr body weight indicated that the weight of the clipped kidney (right kidney) was lower than that of an unclipped one in both sexes, while no significant difference between right and left kidney weight was observed in male and female sham-operated groups (Table 1). These data confirmed the induction of hypertension in the clip cased hypertension.

3.2. Baseline Data (Control Phase). The baseline data within each sex showed that there was no significant difference in MAP, RPP, RBF, and RVR between the groups prior to administration of antagonists or vehicle (Table 2).

3.3. Hemodynamics Responses to Antagonist Infusion (Antagonist Phase). The results of the antagonist phase indicated the reduction of MAP, RPP, and RVR percentage change in male and female rats treated with losartan alone and A779 plus losartan about 40–60 percent from the baseline. These findings confirmed the important role of Ang II action through AT_1R in maintaining arterial pressure and causing vasoconstriction. On the other hand, A779 had little effect, and it does not appear to be important in basal control of hemodynamics parameters. However, no significant difference was detected between male and female rats in MAP, RPP, RBF, and RVR percentage change by vehicle or antagonists (Figure 3).

3.4. Hemodynamics Response to Ang II Administration. Ang II infusion increased MAP percentage change in male and female groups, and no differences were detected between the sexes (Figure 4). However, the raw values of MAP in the equilibrium time (base), 30 min after antagonist administration (treat), and graded angiotensin II infusion are tabulated in Table 3. As previously described, RPP was fixed by aortic occluder during Ang II infusion. So, no difference was expected in RPP percentage change response to Ang II between the male and female groups. Administration of losartan alone increased RBF percentage change response to Ang II infusion in male and female rats. For example, Ang II with dose of 1000 ng/kg/min increased RBF about 19.3% in male and 13% in female rats, while coadministration of A779 and losartan increased RBF response to Ang II (1000 ng/kg/min) 17.2% in male rats and decreased 3.3% in female rats (Figure 4). The Ang II infusion decreased RBF and increased RVR percentage change in a dose-related manner in both sexes, and these responses were significantly different between male and female rats (Figure 4). That means the greater RBF and RVR responses to Ang II were detected in male rats received vehicle instead antagonist when compared with female rats. However, this sex difference in RBF and RVR responses to Ang II was eliminated by AT_1R and MasR antagonists’ infusion. Administration of losartan alone decreased RVR response to Ang II infusion in male and female rats. For example, Ang II with a dose of 1000 ng/kg/min decreased RVR 17% in male and 12% in female rats,




FIGURE 2: Systolic pressure (SYS, mmHg) in sham-operated and 2K1C groups. The data are reported as mean \pm standard error of mean. *Significant difference from sham-operated groups. The statistical P values were obtained using one-way ANOVA followed by the LSD post hoc test.

TABLE 1: The male and female data for body weight before clipping operation (BW1) and 28 days after 2K1C implementation (BW2).

	Groups	BW1 (g)	BW2 (g)	P	RKW (g)/100 gBW	LKW (g)/100 gBW	P
Male	Sham operated ($n=7$)	193 \pm 7.2	277 \pm 5.8	0 < 0.01	0.37 \pm 0.03	0.36 \pm 0.01	0.64
	Vehicle ($n=9$)	192 \pm 7.1	272 \pm 7.3	0 < 0.0001	0.37 \pm 0.01	0.47 \pm 0.02	0 < 0.001
	Losartan ($n=8$)	214 \pm 9.2	280 \pm 14.5	0 < 0.0001	0.29 \pm 0.03	0.46 \pm 0.04	0.05
	A779 ($n=7$)	212 \pm 7.1	256 \pm 12.9	0.01	0.28 \pm 0.04	0.48 \pm 0.03	0.02
	A779 + losartan ($n=6$)	225 \pm 8.7	298 \pm 7.1	0 < 0.0001	0.31 \pm 0.01	0.36 \pm 0.02	0.02
Female	Sham operated ($n=8$)	151 \pm 3.6	195 \pm 5.9	0 < 0.0001	0.37 \pm 0.02	0.35 \pm 0.01	0.37
	Vehicle ($n=10$)	149 \pm 2.1	191 \pm 3.7	0 < 0.0001	0.32 \pm 0.02	0.42 \pm 0.02	0.02
	Losartan ($n=7$)	152 \pm 2.7	182 \pm 3.5	0 < 0.0001	0.36 \pm 0.02	0.43 \pm 0.02	0.05
	A779 ($n=9$)	145 \pm 7.5	187 \pm 7.1	0 < 0.0001	0.33 \pm 0.01	0.39 \pm 0.01	0.02
	A779 + losartan ($n=8$)	143 \pm 2.2	173 \pm 3.4	0 < 0.0001	0.27 \pm 0.03	0.42 \pm 0.03	0.03

The right kidney weight per 100 gram BW (RKWg/100gBW) and left kidney weight per 100 gram BW (LKWg/100gBW). The data are reported as mean \pm standard error of mean. The statistical P values were obtained using the paired-sample T -test.

TABLE 2: The baseline hemodynamics data.

	Groups	MAP (mmHg)	RPP (mmHg)	RBF (ml/min)	RVR (mmHg·min/ml)
Male	Vehicle ($n=9$)	120.4 \pm 6.4	109.1 \pm 10.6	2.9 \pm 0.2	41.9 \pm 7.5
	Losartan ($n=8$)	124.2 \pm 3.3	115.4 \pm 3.5	3.1 \pm 0.3	41.2 \pm 5.1
	A779 ($n=7$)	112.7 \pm 6.6	100.1 \pm 7.3	2.6 \pm 0.3	41.8 \pm 5.3
	A779 + losartan ($n=6$)	124.3 \pm 6.8	114.1 \pm 8.97	2.2 \pm 0.3	57.3 \pm 9.9
	$P_{(ANOVA)}$	0.51	0.60	0.22	0.39
Female	Vehicle ($n=10$)	114.8 \pm 6.4	105.2 \pm 5.5	2.4 \pm 0.3	48.5 \pm 6.1
	Losartan ($n=7$)	116.2 \pm 8.6	108.9 \pm 9.3	2.3 \pm 0.3	47.1 \pm 5.5
	A779 ($n=9$)	114.5 \pm 3.5	108.0 \pm 3.6	2.4 \pm 0.3	50.0 \pm 5.5
	A779 + losartan ($n=8$)	113.2 \pm 6.6	101.6 \pm 6.6	1.8 \pm 0.1	59.7 \pm 5.8
	$P_{(ANOVA)}$	0.99	0.85	0.35	0.48

MAP, RPP, RBF, and RVR stand for mean arterial pressure, renal perfusion pressure, renal blood flow, and renal vascular resistance, respectively. The data were reported as mean \pm standard error of mean. The statistical P values were obtained using one-way ANOVA followed by the LSD post hoc test.

while coadministration of A779 + losartan decreased RVR response to Ang II (1000 ng/kg/min) 12.8% in male rats and increased 2.7% in female rats (Figure 4).

4. Discussion

Our major findings showed that in vehicle-treated rats, the renal vasoconstriction was greater in males than in females, confirming sex differences in the renal response to Ang II. Losartan abolished the renal vasoconstriction in response to Ang II, and in both sexes, renal vasodilatation was seen with no sex difference. Sex differences in the renal vasoconstrictor response to Ang II also were not seen after MasR blockade. This provides some level of evidence to support a role of

MasR in underlying the sex difference in the response to Ang II. The most striking sex difference in the response to Ang II was after combined blockade of AT₁R and MasR. Such finding highlights the role of gender and receptor interaction between male and female rats in the 2K1C model.

Our data support the fewer RBF and higher RVR responses to Ang II in female rats. It seems that these findings are related to hypertension and sex which affect RAS components. The RAS components such as renin, ACE, angiotensinogen, AT₁R, AT₂R, MasR, and AT₄R are expressed in the kidney [14, 21–23], and different distributions of receptors in the kidneys and other organs have been observed in both sexes [24]. Hypertension itself changes distribution and regulation of vascular and renal




FIGURE 3: The effect of antagonist on hemodynamics parameters. MAP, RPP, RBF, and RVR stand for mean arterial pressure, renal perfusion pressure, renal blood flow, and renal vascular resistance in male M and female F rats. Data are presented as mean \pm SEM and were analyzed using repeated-measures ANOVA with the factors group (sex), treat (antagonist), and their interaction.

TABLE 3: Mean arterial pressure (MAP) in the equilibrium time (base), 30 min after antagonist administration (treat), and graded angiotensin II infusion.

Treatment	Group gender	Mean arterial pressure (mmHg)				Angiotensin II (ng/kg/min)		
		Antagonists	Base	Treat	30	100	300	1000
Vehicle	Male		120 \pm 6.4	121 \pm 5.0	124 \pm 4.5	129 \pm 4.8	134 \pm 5.6	150 \pm 5.2
	Female		114 \pm 6.3	112 \pm 4.8	113 \pm 4.5	116 \pm 4.3	120 \pm 4.0	135 \pm 4.5
			$P_{time} = 0.8$	$P_{group} = 0.3$		$P_{time} < 0.0001$	$P_{group} = 0.04$	
			$P_{time\&group} = 0.4$		$P_{time\&group} = 0.5$			
Losartan	Male		124 \pm 3.3	76 \pm 4.7	77 \pm 5.4	81 \pm 6.3	87 \pm 6.5	91 \pm 8.2
	Female		116 \pm 8.6	72 \pm 5.7	71 \pm 6.4	72 \pm 7.4	79 \pm 8.9	83 \pm 7.0
			$P_{time} < 0.0001$	$P_{group} = 0.3$		$P_{time} = 0.004$	$P_{group} = 0.4$	
			$P_{time\&group} = 0.7$		$P_{time\&group} = 0.9$			
A779	Male		113 \pm 6.6	102 \pm 8.1	108 \pm 7.7	122 \pm 7.3	131 \pm 7.1	144 \pm 6.3
	Female		114 \pm 3.5	102 \pm 5.4	106 \pm 6.1	113 \pm 6.0	124 \pm 4.4	140 \pm 5.1
			$P_{time} = 0.001$	$P_{group} = 0.9$		$P_{time} < 0.0001$	$P_{group} = 0.5$	
			$P_{time\&group} = 0.8$		$P_{time\&group} = 0.5$			
A779 + losartan	Male		124 \pm 6.8	65 \pm 2.5	68 \pm 2.8	74 \pm 3.5	81 \pm 5.2	87 \pm 3.5
	Female		113 \pm 6.6	64 \pm 6.0	70 \pm 4.0	76 \pm 4.9	78 \pm 6.3	85 \pm 5.8
			$P_{time} < 0.0001$	$P_{group} = 0.5$		$P_{time} < 0.0001$	$P_{group} = 0.9$	
			$P_{time\&group} = 0.1$		$P_{time\&group} = 0.5$			

Data are presented as mean \pm SEM and were analyzed using repeated-measures ANOVA with the factors group (sex), time (treated with antagonist or dose of Ang II), and their interaction.




FIGURE 4: The effect of graded Ang II infusion on hemodynamics parameters. MAP, RPP, RBF, and RVR stand for mean arterial pressure, renal perfusion pressure, renal blood flow, and renal vascular resistance in male M and female F rats. Data are presented as mean \pm SEM and were analyzed using repeated-measures ANOVA with the factors group (sex), time (dose of Ang II), and their interaction.

RAS component which may be caused different systemic and renal responses [23, 25–27]. In the current model of hypertension, 2K1C-induced Ang II-dependent hypertension, it may change the systemic and local renal RAS distribution and function [28–30]. Ang II-dependent hypertension was gender related in kidney transplantations subjects [31]. Renal ACE2 mRNA expression level in males is less than that in females, and treatment with Ang II has enhanced ACE2 mRNA expression in females 2.5 folds greater than that in males [32]. 2K1C hypertension reduced vascular and renal AT₁R [26, 27]. Also, a lower ratio of AT₁R/AT₂R is found in female rats than in male rats spontaneously hypertensive rats (SHRs) [18]. Ang II infusion decreased AT₁R and increased AT₂R in the renal cortex of male SHRs without change in AT₁R and AT₂R expression in female SHRs [33]. The enhancement of ACE/Ang II/AT₁R pathways is observed in males, and the balance is shifted towards the ACE2/Ang 1–7/MasR and AT₂R pathways in females [11, 24, 34]. Augmentation of the vasodilatory arm of the RAS in females contributes to the gender differences in response to Ang II.

AT₁R antagonist, losartan, eliminated the vasoconstriction response to Ang II. The renal AT₁R distribution and function in males is more than that in females [31]. Thus, the

disappearance of the sex difference in vasoconstriction response to Ang II may be due to the elimination of the greater AT₁R response to Ang II in male rats.

The other finding indicates that no differences were detected in RBF and RVR responses to Ang II infusion between males and females in condition of MasR blockade male rats have a greater blood pressure response to Ang II than female rats and A779 eliminated gender difference response to Ang II [33]. Coadministration of A779 and losartan in female rats had shown more RBF reduction compared to male rats. MasR participates in vasorelaxation and blood pressure reduction via Ang 1–7 [22, 35]. MasR is not activated by Ang II but indirectly involved in Ang II action through an interaction with the AT₁R. Coexpression of MasR and AT₁R inhibits the actions of Ang II and acts as a physiological antagonist for AT₁R [19]. MasR distribution and function is also gender dependent with a greater distribution and function in female rats than in male rats [11, 36, 37]. In addition, a study indicated the enhancement of MasR protein expression in carotid of 2K1C hypertensive rats [35], while the renal Mas mRNA was not different between SHRs and normotensive Wistar rats [38]. Ang II infusion enhances MasR protein expression in female SHRs with no change in male SHRs [33]. Also there was not gender

difference in the intratubular MasR expression 3 weeks after 2K1C while 5-week flowing clipping operation intratubular MasR expression was elevated in female rats compared with male 2K1C rats [39], and renal MasR was decreased in male 2K1C rats [40]. Ang 1–7 via MasR and Ang II via AT₁R interact with each other on rat renal mesangial cells [41]. Women plasma Ang 1–7 is higher compared with men [42]. Renal cortical Ang 1–7 levels are higher in female SHR compared with male SHR [33]. Therefore, MasR blocking in the A779 groups eliminated vasodilatory effect of MasR and may remove the inhibitory effect of the MasR on AT₁R and induce a greater response, especially in females, which have a greater distribution of the MasR.

5. Conclusion

The fewer RBF and RVR responses to Ang II in female rats may be related to more ACE2/Ang 1–7/MasR and AT₂R pathway activities in female rats than in male 2K1C rats. Also, enhancement of Ang II due to 2K1C may affect distribution of different receptors and transporters in male and female rats and change the expected systemic and local RAS function. Moreover, the stronger RBF response to Ang II in A779 and A779 + losartan 2K1C female treated groups is related to greater distribution and function of ACE2/Ang 1–7/MasR arm of the RAS and fewer activity of ACE/Ang II/AT₁R arm of the RAS in female 2K1C rats. Therefore, rather than blocking the ACE/Ang II/AT₁R axis, activating the ACE2/Ang 1–7/MasR axis of the RAS may be more effective in improving the renal and systemic side effects of Ang II-dependent hypertension in female rats but not in male rats.

Data Availability

The data used to support the findings of this study are available from the corresponding author on request.

Conflicts of Interest

The authors declare no conflicts of interest.

Acknowledgments

This research was supported by Isfahan University of Medical Sciences (grant no. 397461). The authors thank Dr. Roger Evan for his excellent and helpful discussion.

References

- [1] European Society of Hypertension-European Society of Cardiology Guidelines, “European Society of Hypertension-European Society of Cardiology guidelines for the management of arterial hypertension,” *Journal of Hypertension*, vol. 21, no. 6, pp. 1011–1053, 2003.
- [2] G. Leoncini, F. Viazzi, G. Storage, G. Deferrari, and R. Pontremoli, “Blood pressure variability and multiple organ damage in primary hypertension,” *Journal of Human Hypertension*, vol. 27, no. 11, pp. 663–670, 2013.
- [3] F. Viazzi, D. Parodi, G. Leoncini et al., “Serum uric acid and target organ damage in primary hypertension,” *Hypertension*, vol. 45, no. 5, pp. 991–996, 2005.
- [4] E. Pringle, C. Phillips, L. Thijs et al., “Systolic blood pressure variability as a risk factor for stroke and cardiovascular mortality in the elderly hypertensive population,” *Journal of Hypertension*, vol. 21, no. 12, pp. 2251–2257, 2003.
- [5] E. E. Gillis and J. C. Sullivan, “Sex differences in hypertension,” *Hypertension*, vol. 68, no. 6, pp. 1322–1327, 2016.
- [6] D. Mozaffarian, E. J. Benjamin, A. S. Go et al., “Heart disease and stroke statistics-2016 update: a report from the American Heart Association,” *Circulation*, vol. 133, no. 4, pp. e38–360, 2016.
- [7] S. R. Silbiger and J. Neugarten, “The role of gender in the progression of renal disease,” *Advances in Renal Replacement Therapy*, vol. 10, no. 1, pp. 3–14, 2003.
- [8] P. A. James, S. Oparil, B. L. Carter et al., “2014 evidence-based guideline for the management of high blood pressure in adults,” *JAMA*, vol. 311, no. 5, pp. 507–520, 2014.
- [9] G. Baiardi, M. Macova, I. Armando, H. Ando, D. Tyurmin, and J. M. Saavedra, “Estrogen upregulates renal angiotensin II AT1 and AT2 receptors in the rat,” *Regulatory Peptides*, vol. 124, no. 1–3, pp. 7–17, 2005.
- [10] L. M. Hilliard, M. Nematbakhsh, M. M. Kett et al., “Gender differences in pressure-natriuresis and renal autoregulation,” *Hypertension*, vol. 57, no. 2, pp. 275–282, 2011.
- [11] L. M. Hilliard, A. K. Sampson, R. D. Brown, and K. M. Denton, “The “his and hers” of the renin-angiotensin system,” *Current Hypertension Reports*, vol. 15, no. 1, pp. 71–79, 2013.
- [12] R. M. Carey, P. K. Whelton, and Committee AHAHGW, “Prevention, detection, evaluation, and management of high blood pressure in adults: synopsis of the 2017 American College of Cardiology/American Heart Association hypertension guideline,” *Annals of Internal Medicine*, vol. 168, no. 5, pp. 351–358, 2018.
- [13] A. S. Bombardieri and R. Toto, “Dual blockade of the renin-angiotensin-aldosterone system: beyond the ACE inhibitor and angiotensin-II receptor blocker combination,” *American Journal of Hypertension*, vol. 22, no. 10, pp. 1032–1040, 2009.
- [14] X. C. Li, D. Zhu, X. Zheng, J. Zhang, and J. L. Zhuo, “Intratubular and intracellular renin-angiotensin system in the kidney: a unifying perspective in blood pressure control,” *Clinical Science*, vol. 132, no. 13, pp. 1383–1401, 2018.
- [15] U. P. Jorde, P. V. Ennezat, J. Lisker et al., “Maximally recommended doses of angiotensin-converting enzyme (ACE) inhibitors do not completely prevent ACE-mediated formation of angiotensin II in chronic heart failure,” *Circulation*, vol. 101, no. 8, pp. 844–846, 2000.
- [16] M. Hudson, E. Rahme, H. Behloul, R. Sheppard, and L. Pilote, “Sex differences in the effectiveness of angiotensin receptor blockers and angiotensin converting enzyme inhibitors in patients with congestive heart failure—a population study,” *European Journal of Heart Failure*, vol. 9, no. 6-7, pp. 602–609, 2007.
- [17] C. Falconnet, M. Bochud, P. Bovet, M. Maillard, and M. Burnier, “Gender difference in the response to an angiotensin-converting enzyme inhibitor and a diuretic in hypertensive patients of African descent,” *Journal of Hypertension*, vol. 22, no. 6, pp. 1213–1220, 2004.
- [18] M. Silvaantonialli, R. C. Tostes, L. Fernandes et al., “A lower ratio of AT1/AT2 receptors of angiotensin II is found in female than in male spontaneously hypertensive rats,” *Cardiovascular Research*, vol. 62, no. 3, pp. 587–593, 2004.

- [19] E. Kostenis, G. Milligan, A. Christopoulos et al., "G-protein-coupled receptor Mas is a physiological antagonist of the angiotensin II type 1 receptor," *Circulation*, vol. 111, no. 14, pp. 1806–1813, 2005.
- [20] J. O'Neill, V. Healy, and E. J. Johns, "Intrarenal Mas and AT1 receptors play a role in mediating the excretory actions of renal interstitial angiotensin-(1-7) infusion in anaesthetized rats," *Experimental Physiology*, vol. 102, no. 12, pp. 1700–1715, 2017.
- [21] T. Matsusaka, F. Niimura, A. Shimizu et al., "Liver angiotensinogen is the primary source of renal angiotensin II," *Journal of the American Society of Nephrology*, vol. 23, no. 7, pp. 1181–1189, 2012.
- [22] S. N. Patel, Q. Ali, P. Samuel, U. M. Steckelings, and T. Hussain, "Angiotensin II type 2 receptor and receptor Mas are colocalized and functionally interdependent in obese Zucker rat kidney," *Hypertension*, vol. 70, no. 4, pp. 831–838, 2017.
- [23] L. G. Navar, L. M. Harrison-Bernard, A. Nishiyama, and H. Kobori, "Regulation of intrarenal angiotensin II in hypertension," *Hypertension*, vol. 39, no. 2, pp. 316–322, 2002.
- [24] A. K. Sampson, K. M. Moritz, and K. M. Denton, "Postnatal ontogeny of angiotensin receptors and ACE2 in male and female rats," *Gender Medicine*, vol. 9, no. 1, pp. 21–32, 2012.
- [25] R. Romero-Nava, J. E. Rodriguez, A. A. Reséndiz-Albor et al., "Changes in protein and gene expression of angiotensin II receptors (AT1 and AT2) in aorta of diabetic and hypertensive rats," *Clinical and Experimental Hypertension*, vol. 38, no. 1, pp. 56–62, 2016.
- [26] Z.-Q. Wang, L. J. Millatt, N. T. Heiderstadt, H. M. Siragy, R. A. Johns, and R. M. Carey, "Differential regulation of renal angiotensin subtype AT1A and AT2 receptor protein in rats with angiotensin-dependent hypertension," *Hypertension*, vol. 33, no. 1, pp. 96–101, 1999.
- [27] F. Amiri and R. Garcia, "Renal angiotensin II receptor regulation in two-kidney, one clip hypertensive rats," *Hypertension*, vol. 30, no. 3, pp. 337–344, 1997.
- [28] M. C. Prieto, R. A. González-Villalobos, F. T. Botros et al., "Reciprocal changes in renal ACE/ANG II and ACE2/ANG 1-7 are associated with enhanced collecting duct renin in Goldblatt hypertensive rats," *American Journal of Physiology-Renal Physiology*, vol. 300, no. 3, pp. F749–F755, 2011.
- [29] W. Shao, K. Miyata, A. Katsurada et al., "Increased angiotensinogen expression, urinary angiotensinogen excretion, and tissue injury in nonclipped kidneys of two-kidney, one-clip hypertensive rats," *American Journal of Physiology-Renal Physiology*, vol. 311, no. 2, pp. F278–F290, 2016.
- [30] L. C. Matavelli, J. Huang, and H. M. Siragy, "Angiotensin AT2 receptor stimulation inhibits early renal inflammation in renovascular hypertension," *Hypertension*, vol. 57, no. 2, pp. 308–313, 2011.
- [31] L. Wang, X. Wang, H. Y. Qu et al., "Role of kidneys in sex differences in angiotensin II-induced hypertension," *Hypertension*, vol. 70, no. 6, pp. 1219–1227, 2017.
- [32] A. K. Sampson, K. M. Moritz, E. S. Jones, R. L. Flower, R. E. Widdop, and K. M. Denton, "Enhanced angiotensin II type 2 receptor mechanisms mediate decreases in arterial pressure attributable to chronic low-dose angiotensin II in female rats," *Hypertension*, vol. 52, no. 4, pp. 666–671, 2008.
- [33] J. C. Sullivan, K. Bhatia, T. Yamamoto, and A. A. Elmarakby, "Angiotensin (1-7) receptor antagonism equalizes angiotensin II-induced hypertension in male and female spontaneously hypertensive rats," *Hypertension*, vol. 56, no. 4, pp. 658–666, 2010.
- [34] M. A. Zimmerman and J. C. Sullivan, "Hypertension: what's sex got to do with it?" *Physiology*, vol. 28, no. 4, pp. 234–244, 2013.
- [35] V. C. Olivon, R. D. Aires, L. B. Santiago, L. Z. N. Ramalho, S. F. Cortes, and V. S. Lemos, "Mas receptor overexpression increased Ang-(1-7) relaxation response in renovascular hypertensive rat carotid," *Peptides*, vol. 71, pp. 250–258, 2015.
- [36] T. Safari, M. Nematbakhsh, L. M. Hilliard, R. G. Evans, and K. M. Denton, "Sex differences in the renal vascular response to angiotensin II involves the Mas receptor," *Acta Physiologica*, vol. 206, no. 2, pp. 150–156, 2012.
- [37] M. Maleki and M. Nematbakhsh, "Mas receptor antagonist (A799) alters the renal hemodynamics responses to angiotensin II administration after renal moderate ischemia/reperfusion in rats: gender related differences," *Research in Pharmaceutical Sciences*, vol. 14, no. 1, pp. 12–19, 2019.
- [38] Z. Tan, J. Wu, and H. Ma, "Regulation of angiotensin-converting enzyme 2 and Mas receptor by Ang-(1-7) in heart and kidney of spontaneously hypertensive rats," *Journal of the Renin-Angiotensin-Aldosterone System*, vol. 12, no. 4, pp. 413–419, 2011.
- [39] S. H. Lee, Y. H. Lee, S. W. Jung et al., "Sex-related differences in the intratubular renin-angiotensin system in two-kidney, one-clip hypertensive rats," *American Journal of Physiology-Renal Physiology*, vol. 317, no. 3, pp. F670–F682, 2019.
- [40] Y. G. Kim, S. H. Lee, S.-Y. Kim et al., "Sequential activation of the intrarenal renin-angiotensin system in the progression of hypertensive nephropathy in Goldblatt rats," *American Journal of Physiology-Renal Physiology*, vol. 311, no. 1, pp. F195–F206, 2016.
- [41] H. Xue, L. Zhou, P. Yuan et al., "Counteraction between angiotensin II and angiotensin-(1-7) via activating angiotensin type I and Mas receptor on rat renal mesangial cells," *Regulatory Peptides*, vol. 177, no. 1-3, pp. 12–20, 2012.
- [42] J. C. Sullivan, P. Rodriguez-Miguel, M. A. Zimmerman, and R. A. Harris, "Differences in angiotensin (1-7) between men and women," *American Journal of Physiology-Heart and Circulatory Physiology*, vol. 308, no. 9, pp. H1171–H1176, 2015.