

Disclosure: No significant relationships.

Keywords: Psychoneuroimmunology; mindfulness; Selfmanagement; Autopoiesis

EPV0488

Efficacy of IV immunoglobulins on depressive symptoms and self-injury: A case report

A. Silva, P. Politi, N. Brondino and M. Olivola*

Department Of Brain And Behavioral Sciences, University of Pavia, Pavia, Italy

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.2021

Introduction: Some studies in literature highlight the correlation between immune-mediated inflammatory processes and psychiatric pathologies. However, there are few studies about the efficacy of IV immunoglobulins in psychiatric features (1). (1) ZUNSZAIN, Patricia A.; HEPGUL, Nilay; PARIANTE, Carmine M. Inflammation and depression. In: Behavioral neurobiology of depression and its treatment. Springer, Berlin, Heidelberg, 2012. p. 135-151.

Objectives: Case report: a 39 year patient diagnosed with borderline personality disorder and myasthenia was hospitalized for self-injury ideation, acting out and depressive episode treated with acid valproic, aripiprazole, gabapentin; flare-up of myasthenia that needed treatment.


Methods: Clinical and test evaluation was performed in three stages: before (t0), immediately after (t1) and 3 weeks after (t2) the administration of the IgEV without other treatment modifications. We have used: - Inventory of Statements About Self-Injury (ISAS) - Barrat Impulsiveness Scale, Version 11 (BIS-11) - Hamilton Anxiety Rating Scale (HAM-A) - Montgomery-Asberg Depression Rating Scale (MADRS) - Alexian Brothers Urge to Self-Injure Scale (ABUSI)

Results: The patient has a score of 79 at BIS-11. She used to have a huge number of acting aout as we see on ISAS (Fig.1).

Figure 1

ISAS		
Self-Injury	Life time	t0-t2
Cutting	100	0
Biting	200	0
Carving	10	0
Pulling Hair	10	0
Severe Scratching	100	0
Banging or Hitting Self	250	0
Interfering/Would Healing	50	0
Rubbing Skin Againts Rough Surface	2	0
Sticking Self/Needles	5	0
Swallowing Dangerous Substance	300	0

Figure 2



Conclusions: We observed a reduction in non-suicidal self-injurious ideation, the suspension of acting-out, a complete remission of depressive symptoms with mild persistence of anxious symptoms immediately after the administration of immunoglobulins, and the remission continue until one month after the administration (Fig.2).

Disclosure: No significant relationships.

Keywords: acting-out; immunoglobulins; Borderline; Depression

EPV0489

Prospects for immunotherapy of depression based on cell technologies

E. Markova* and M. Knyazheva

Neuroimmunology Lab, State Research Institute of Fundamental and Clinical Immunology, Novosibirsk, Russian Federation

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.2022