


# Isolated Neutropenia/Benign Ethnic Neutropenia: A Common Clinical and Laboratory Finding in Southern and Western Saudi Arabia

This article was published in the following Dove Press journal:  
*International Journal of General Medicine*

Zuhier A Awan<sup>1,2</sup>  
Saeed M Al Amoudi<sup>2</sup>  
Muhammad Saboor<sup>3,4</sup>  
Husain Y Alkhalidy<sup>5,6</sup> 

<sup>1</sup>Department of Clinical Biochemistry, Faculty of Medicine, King Abdulaziz University, Jeddah, Saudi Arabia;

<sup>2</sup>Department of Clinical Pathology, Al-Borg Medical Laboratories, Jeddah, Saudi Arabia; <sup>3</sup>Department of Medical Laboratory Technology, Faculty of Applied Medical Sciences, Jazan University, Jazan, Saudi Arabia; <sup>4</sup>Medical Research Centre (MRC), Jazan University, Jazan, Saudi Arabia; <sup>5</sup>Department of Internal Medicine, College of Medicine, King Khalid University, Abha, Saudi Arabia; <sup>6</sup>Research Center for Advanced Materials Science, King Khalid University, Abha, Saudi Arabia

**Objective:** Isolated mild neutropenia is a common clinical problem in some ethnicities including Arabs and Middle Eastern population. The current study aims to authenticate the prevalence of isolated neutropenia in Southern and Southwestern Saudi Arabia, explore the effect of altitude or regional differences and to suggest a new reference range for neutrophil count.

**Methods:** In this retrospective cross-sectional study, laboratory results of a commercial laboratory were screened over a period of 5 years (2016–2020) in seven different cities of different altitudes in South and southwestern Saudi Arabia. Participants' laboratory investigations were reviewed and excluded for any abnormal complete blood count, renal profile, liver profile, lipid profile, thyroid function test, fasting blood glucose, or HbA1c findings. Descriptive analysis and 95th percentile range were calculated using standard statistical methods.

**Results:** A total of 91,880 complete blood count results were included in the final analysis. Isolated neutropenia was common laboratory finding, with a prevalence ranging from 11% to 23%. The 2.5th percentile of the neutrophil count was lower than currently utilized  $1.5 \times 10^9/L$  in all studied seven cities.

**Conclusion:** Mild to moderate neutropenia is common in Southern and Southwestern Saudi Arabia. Benign ethnic neutropenia (BEN) likely explains this high prevalence. Since BEN has no clinical significance, the reference range for normal neutrophil counts needs to be adjusted to reflect the effect of BEN.

**Keywords:** isolated neutropenia, benign ethnic neutropenia, reference intervals, Saudi Arabia

Correspondence: Husain Y Alkhalidy  
Department of Internal Medicine, College of Medicine, Research Center for Advanced Materials Science, King Khalid University, P.O.Box 9004, Abha, 61413, Saudi Arabia  
Tel +966599915567  
Email halkhalidy@kku.edu.sa

## Introduction

Neutrophils are the most abundant leukocytes in the peripheral blood. They play a pivotal role in the innate immune system and are key regulators of the adaptive immune system.<sup>1,2</sup> The absolute neutrophil count (ANC) normally ranges from  $1.8$  to  $7.5 \times 10^9/L$ .<sup>3</sup> Marked reduction in the peripheral neutrophil count is regarded as a serious clinical problem that needs to be addressed. Peripheral neutrophils constitute only a minute fraction of the total body neutrophils that are mainly in the bone marrow. The number of neutrophils in the peripheral circulation may be affected by multiple factors, including stress, smoking, pregnancy, and drugs. Altitude may also be associated with lower neutrophil counts.<sup>4,5</sup> Reduced ANC is defined as a count less than  $1.5 \times 10^9/L$  and is known as neutropenia. In severe cases, neutropenia predisposes an individual to infections or even life-threatening

consequences. Neutropenia is categorized into three subtypes based on the ANC: mild ( $1.0\text{--}1.5 \times 10^9/\text{L}$ ), moderate ( $0.5\text{--}1.0 \times 10^9/\text{L}$ ), and severe ( $<0.5 \times 10^9/\text{L}$ ).

Decreased production or increased destruction of neutrophils may cause neutropenia.<sup>3</sup> Severe congenital neutropenia, as the name suggests, is an inherited condition that leads to life-threatening infections in the first year of life. This condition has also a propensity to progress to acute leukemia.<sup>6</sup> Patients whom absolute neutrophil remains low due to decreased production are more prone to develop infections as compared to conditions associated with increased peripheral destruction or margination of neutrophils. Moreover, as first reported in 1941 by Forbes et al, low neutrophil count ( $<1.5 \times 10^9/\text{L}$ ) in certain ethnicities with no predisposition to infections indicates no clinical disadvantage in the affected individuals. This condition is known as benign ethnic neutropenia (BEN).<sup>7,8</sup>

In ethnicities with reduced circulating neutrophil counts, the bone marrow cellularity, morphology, number of granulocyte colony-forming units, and maturation of myeloid cells remain normal. However, slight reduction in the number of progenitors has been observed in some cases of BEN.<sup>7</sup> Additionally, a slight increase in ANC has been reported in individuals with BEN after exercise.<sup>9</sup> Subjects with BEN show a circadian variation of the neutrophil count similar to that observed in subjects without BEN.<sup>10</sup>

BEN is most common in Africans, Afro-Americans, Ethiopian Jews, and Yemenite Jews.<sup>8,11–13</sup> We previously showed that a higher prevalence of isolated neutropenia was observed in two cities (Jeddah and Abha) in Saudi Arabia.<sup>14</sup> Saudi Arabia, representing the larger part of the Arabian peninsula, is inhabited by mainly Arab tribes of Semitic origin. In this study, we aimed to confirm our previous findings through a large sample size from seven different cities at different altitudes (Jeddah, Taif, Al Baha, Abha, Khamis Mushait, Najran, and Gizan) in Southern and Southwestern Saudi Arabia. We also aimed to explore any potential association of neutropenia prevalence with altitude or any regional differences.

## Methods

This study was designed and conducted in accordance with the Declaration of Helsinki principles. The study protocol and study subject consent waiver, due to use of completely de-identified data with no associated risk to the participants, were approved by the ethical and research committee of King Khalid University, Abha, Saudi Arabia. (ECM#2020-

0905) – (HAPO-06-B-001). Neither personal identification nor bioinformation of study subjects were collected or disclosed. In this retrospective study, results of a commercial laboratory (Al-Borg Laboratories) were screened over a period of 5 years (2016–2020) in seven different cities in Saudi Arabia. Laboratory investigations included complete blood count (CBC), renal profile, liver profile, lipid profile, thyroid function test, fasting blood glucose measurement, and HbA1c measurement. Screened participants were either customers who do lab tests for health check-ups or follow up of chronic health conditions. Data were collected from the laboratories located in Jeddah (sea level), Taif (1898 meters above sea level), Al Baha (2000 meters), Abha (2270 meters), Khamis Mushait (1700 meters), Najran (1293 meters), and Gizan (sea level). Exclusion lab criteria were applied to exclude any pre-existing health conditions. Subjects selected were all having normal CBC, biochemical, kidney, liver, sugar, lipid and thyroid profiles and thus were assumed healthy and proper to estimate the neutrophil count normal range.

## Inclusion and Exclusion Criteria

Male and female patients aged 13–60 years were included. To estimate the white blood cell count (WBC) reference, the following exclusion criteria were applied: any abnormalities, unless mild, in the renal profile (serum blood urea nitrogen, creatinine), liver profile (ALT, AST, total bilirubin, and direct bilirubin), lipid profile (cholesterol, HDL, LDL, and triglyceride), thyroid function test (TSH, T4), and diabetic profile (fasting blood sugar and HbA1c).

For CBC, individuals with hemoglobin (Hb) concentration of 12–19 g/dl, platelet count of  $140\text{--}450 \times 10^9/\text{L}$ , and WBC count of less than  $11.5 \times 10^9/\text{L}$  were included. For low WBC count, the exclusion was applied only if other blood cell types were abnormal, eg, concurrent anemia or thrombocytopenia. Since BEN is usually mild to moderate, analyses were performed with both inclusion and exclusion of neutrophil counts less than  $0.5 \times 10^9/\text{L}$ .

## Blood Collection and Laboratory Analysis

To ensure no inter laboratory variations in the obtained results, the different branches follow organizational standard operating protocols of blood sample collection for CBC. Blood samples were collected into liquid K3 EDTA tubes and processed within 2 hours of collection. Sysmex automated analyzers (Sysmex Corporation, Kobe, Japan) were utilized for WBC enumeration and differential count.

## Statistical Analysis

The participants' data were de-identified for any personal information before importing to an Excel datasheet. After extraction, data were revised, coded, and transferred into the statistical software IBM SPSS Statistics for Windows, version 22 (IBM Corp., Armonk, N.Y., USA). Normal distribution assessment was applied for all scale parameters. Data for both sex and age were stratified into two groups: young adults (13–40 years old) and middle-aged adults (41–60 years old). Descriptive analyses using the mean and 95% confidence interval, median, range, standard deviation (SD), and percentiles (2.5th, 25th, 75th, and 97.5th) were utilized to assess the normal population range. Student's *t*-tests were used to compare continuous variables between different sex and age groups. A *p*-value <0.05 was considered statistically significant.

Since plasma volume is known to be reduced with altitude, we explored any association between neutrophil count and hematocrit considering that lower hematocrit is associated with higher plasma volume and vice versa.

## Results

For the present study, 91,880 CBC results were included in the final analysis ([supplement 1](#)). [Table 1](#) summarizes the frequency of neutropenia in the studied population, while [Table 2](#) depicts the statistical analysis of the mean, SEM, median, SD, range, and percentiles (2.5th, 25th, 75th, and 97.5th). Comparison between groups based on sex and age group is depicted in [Table 3](#).

Mild neutropenia was the most common type of neutropenia among the studied subjects of all cities. The city of Al Baha showed slightly lower frequency in comparison with

**Table 1** Frequency of Neutropenia in the Studied Populations

		Jeddah (Sea Level)	Taif (1898 m)	Al Baha (2000 m)	Abha (2270 m)	Khamis Mushait (1700 m)	Najran (1293m)	Gizan (Sea Level)
Total number analyzed	Female	29,738	3486	1867	4689	6554	2188	2885
	Male	24,001	2765	1359	3315	4807	1840	2386
	Overall	53,739	6251	3226	8004	11,361	4028	5271
Mild neutropenia ( $1.0-1.5 \times 10^9/L$ )	Female	3544	287	249	631	748	365	464
	Male	2744	200	190	310	564	302	412
	Overall	6288 (12%)	487 (8%)	439 (14%)	941 (12%)	1312 (12%)	667 (17%)	876 (17%)
Moderate neutropenia ( $0.5-1 \times 10^9/L$ )	Female	961	93	81	235	252	135	139
	Male	712	69	55	156	198	119	106
	Overall	1673 (3%)	162 (3%)	136 (4%)	391 (5%)	450 (4%)	254 (6%)	245 (5%)
Severe neutropenia ( $<0.5 \times 10^9/L$ )	Female	15	7	7	8	5	3	3
	Male	13	2	8	7	3	5	2
	Overall	28 (0.05%)	9 (0.14%)	15 (0.46%)	15 (0.19%)	8 (0.07%)	8 (0.20%)	5 (0.10%)

**Table 2** Descriptive Analysis of Distribution of Neutrophils Count ( $\times 10^9/L$ ) in the Studied Populations

City	Number	Mean	SEM	Median	SD	Min	Max	2.5th Percentile	25th Percentile	75th Percentile	97.5th Percentile
Jeddah (sea level)	53,739	3.18	0.01	3.05	1.35	0.50	8.98	1.05	2.14	4.03	6.19
Taif (1898 m)	6251	3.15	0.02	3.02	1.35	0.50	8.55	1.00	2.11	4.01	6.15
Al Baha (2000 m)	3226	2.74	0.02	2.55	1.29	0.50	8.65	0.90	1.73	3.55	5.61
Abha (2270 m)	8004	2.76	0.01	2.55	1.32	0.50	8.77	0.87	1.73	3.58	5.87
Khamis Mushait (1700 m)	11,361	2.90	0.01	2.74	1.34	0.51	8.83	0.91	1.84	3.74	5.93
Najran (1293m)	4028	2.56	0.02	2.40	1.22	0.52	7.99	0.83	1.57	3.34	5.39
Gizan (Sea level)	5271	2.43	0.02	2.16	1.16	0.50	8.47	0.89	1.59	3.05	5.42

**Notes:** Total numbers indicate all subjects analyzed after applying the exclusion criteria. Neutrophils <0.5 were excluded from this analysis with no change in distribution.  
**Abbreviations:** SEM, standard error of the mean; SD, standard deviation of the mean.

**Table 3** Mean ( $\pm$ Standard Deviation) of the Neutrophil Count with Respect to Gender and Age Group Distribution

Age Group	(Years)	13–40		41–60	
		M	F	M	F
Jeddah (sea level)		3.18 $\pm$ 1.36	3.22 $\pm$ 1.36	3.12 $\pm$ 1.32	3.17 $\pm$ 1.34
P values		0.34*		0.09**	
Taif (1898 m)		3.10 $\pm$ 1.37	3.20 $\pm$ 1.35	3.19 $\pm$ 1.33	3.05 $\pm$ 1.36
P values		0.73*		0.41**	
Al Baha (2000 m)		2.741 $\pm$ 0.29	2.77 $\pm$ 1.27	2.76 $\pm$ 1.30	2.69 $\pm$ 1.32
P values		0.1*		0.13**	
Abha (2270 m)		2.76 $\pm$ 1.26	2.74 $\pm$ 1.34	2.75 $\pm$ 1.31	2.81 $\pm$ 1.35
P values		0.8*		0.91**	
Khamis Mushait (1700 m)		2.90 $\pm$ 1.35	2.94 $\pm$ 1.36	2.85 $\pm$ 1.29	2.87 $\pm$ 1.33
P values		0.001*		0.01**	
Najran (1293m)		2.52 $\pm$ 1.19	2.62 $\pm$ 1.26	2.48 $\pm$ 1.14	2.61 $\pm$ 1.28
P values		0.71*		0.34**	
Gizan (sea level)		2.39 $\pm$ 1.17	2.49 $\pm$ 1.20	2.36 $\pm$ 1.11	2.44 $\pm$ 1.15
P values		0.001*		0.01**	

**Notes:** 5% trimmed mean reported. Independent Student's *t*-test was used. \*Comparison for gender. \*\*Comparison for age group. While some comparison is statistically significant, the difference between the mean neutrophil counts is not clinically significant.

others, while the highest prevalence was observed in Najran and Gizan (17%). A high prevalence of moderate neutropenia was observed in Abha (5%), Najran (5%), and Gizan (6%). Severe neutropenia was rare among all types, with only 88 cases out of 91,880 (Table 1). Pearson correlation between neutrophil count and hematocrit was not significant.

The lowest value of ANC for the 2.5th percentile was found in Najran ( $0.83 \times 10^9/L$ ), while the highest was found in Jeddah ( $1.05 \times 10^9/L$ ). Additionally, the 2.5th percentile ANC value of the study participants in the remaining cities was between the ANC values found in Jeddah and Najran ( $0.83$ – $1.05 \times 10^9/L$ ). The highest prevalence of isolated neutropenia was found in Najran (23% with mild and moderate forms). Gizan had a prevalence of 22%, making

it the second most prevalent city for neutropenia, while in Taif, Jeddah, Khamis Mushait, Abha, and Al Baha, the frequency of neutropenia (combined mild and moderate) was found to be 11%, 15%, 16%, 17%, and 18%, respectively. Severe neutropenia was the rarest, with only 88 cases out of 91,880 (0.096%).

## Discussion

The present study was conducted primarily to authenticate our previous findings of a high prevalence of isolated neutropenia and explore any geographical association by including other cities of different altitudes and a large dataset of participants. Our aims also include using a larger multi-cities sample to determine a modified reference range suitable to our population. After excluding abnormal tests results, 91,880 CBCs were included for determining the frequency of neutropenia in the study population. The current study validates our previous findings in which the prevalence of neutropenia was found to be 18%.<sup>14</sup> The findings of this study are consistent with those of Gari et al<sup>12</sup> who also reported a high prevalence (20%) of neutropenia in Jeddah. Several other studies have shown a variable frequency of neutropenia among different populations.<sup>11,15–17</sup> Moreover, the findings of this study indicated that the 2.5th percentile of neutrophil count was lower than what is utilized now as lower end of the reference range;  $1.5 \times 10^9/L$ , in all seven cities. This finding is most likely explained by racial effect. We postulate, as explained below, that this prevalence is most likely explained by Benign ethnic neutropenia.

BEN has a positive association with Duffy-null phenotype.<sup>18</sup> Atypical chemokine receptor-1 (*ACKR1*) gene, also known as the Duffy antigen receptor for chemokines (DARC), encodes the Duffy antigens ( $Fy^a$  and  $Fy^b$ ). A single nucleotide polymorphism (SNP) at codon 42 (G125A, rs12075) results in two types of encoded proteins,  $Fy^a$  and  $Fy^b$ .  $Fy^a$  antigen includes glycine, while  $Fy^b$  contains aspartic acid.<sup>19</sup> The Duffy-null phenotype ( $Fy^{a-b-}$ ), also symbolized as erythroid silent (ES), occurs due to another SNP, rs2814778 at position 46 (–46 T > C). The Duffy antigen expression is inhibited in individuals with homozygous C/C polymorphisms.<sup>20</sup> Individuals with rs2814778 suffer from an abnormal phenotype of neutrophils with enhanced egression into the tissues from the blood.<sup>21</sup> *Plasmodium vivax* enters the red cells through DARC. Duffy-null phenotypes therefore may provide a natural resistance to *P. vivax*. Consequently, this protective role of Duffy-null phenotypes in malaria

endemic regions has undergone positive selection leading to its high prevalence in some ethnicities.<sup>22</sup> Although molecular analysis for rs2814778 of Duffy-null phenotypes was not conducted in this study to correlate the high frequency of this phenotype with BEN, the prevalence of Duffy-null phenotypes in Saudi Arabia reported in various studies could be linked with BEN. Duffy-null phenotype (Fy<sup>a-b-</sup>) at a frequency of 62% to 78% has been reported in Saudi Arabia populations.<sup>23,24</sup> Therefore, the isolated neutropenia reported in this current study may be Duffy-null phenotype-induced BEN.

Arabs are among the ethnicities where higher prevalence of BEN has been reported. A previous study has shown that the prevalence of benign neutropenia was high in Saudi participants as compared with none Saudis.<sup>12</sup> This finding suggests that BEN is more common in certain ethnicities. Herein this point also needs to be mentioned that consanguineous marriages are common in Saudi Arabia that could be attributed to the high prevalence of BEN in Saudis.<sup>25</sup> Thirdly, as Saudi Arabia is in close proximity to the African countries where the prevalence of BEN is up to 50%, the migration of African to this region could be considered a contributing factor in the high prevalence of BEN in Saudi population. We previously noticed a higher prevalence of isolated neutropenia at a high altitude in comparison with sea level.<sup>14</sup> In this study, the prevalence of isolated neutropenia in Taif city (1898 m above sea level) was not different from that in the coastal Jeddah city in the current study. Furthermore, in this study, we did not find any association between altitude levels and hematocrit (as a surrogate for decreased plasma volumes). Nonetheless, owing to many confounding factors, including different prevalence of BEN, the effect of altitude on the neutrophil count should not be dismissed without further research.

The ANC, usually above  $1.5 \times 10^9/L$ , is used as an inclusion criterion for clinical trials.<sup>26</sup> This can potentially lead to exclusion of otherwise healthy individuals with physiologically lower neutrophil counts.<sup>27</sup> The same criterion is also used to grade chemotherapy-associated toxicity, which helps in dose adjustment of crucial treatments. Dose intensity of chemotherapy is important, and dose reduction or delay is associated with increased mortality.<sup>28,29</sup> In high-prevalence BEN populations, low baseline ANC should not be considered a factor for altering the treatment strategies in certain clinical conditions. For instance, use of clozapine and deferiprone also induces neutropenia. Administration of these drugs can commence if the ANC is above  $0.5 \times 10^9/L$ , in the absence of any other comorbid condition. In patients with

malignancies, if the ANC ranges between  $0.5 \times 10^9/L$  and  $1.0 \times 10^9/L$ , chemotherapy may be administered safely.<sup>30</sup>

In clinical practice, individuals with  $ANC < 1.5 \times 10^9/L$  usually undergo unwarranted laboratory and clinical investigations without any significant findings. Although the differential diagnoses of neutropenia are wide, BEN can be determined based on the appropriate clinical scenario. BEN is usually an incidental finding of isolated mild neutropenia with no other hematological abnormalities. It is not associated with increased risks of infections.<sup>31</sup> Previous studies have indicated the chronicity of the condition, and follow-up usually indicates the stability and benign nature of the condition.<sup>32</sup> Clinical history can also suggest the presence of other causes of neutropenia. For instance, severe congenital neutropenia is associated with severe neutropenia and increased risk of infections. In cyclic neutropenia, recurrent infections, especially oral lesions and low neutrophil counts, at 21-day intervals are common. Meanwhile, in BEN, recurrent infections are infrequent, while ANC usually are only mildly low. Moreover, since molecular analysis for rs2814778 of Duffy-null phenotypes is not available in clinical laboratories, the Duffy phenotype could be utilized as a marker for identifying individuals with BEN.<sup>33</sup> In current clinical practice, the diagnosis of BEN depends on the exclusion of other causes of neutropenia.

This study has the advantage of large sample size and the availability of extended lab tests to exclude many health conditions. Nonetheless, it has some limitations. It was a retrospective analysis with few demographics and health-related data. Some micronutrient deficiencies that may affect neutrophil counts such as B12 and folate deficiencies were not assessed.

## Conclusion

Isolated neutropenia is common in the southern and southwestern Saudi Arabia. BEN, extensively reported in Middle Eastern countries, is likely to explain many cases of mild isolated neutropenia. However, the effects of other factors, including geography, altitude, and autoimmune diseases, should be explored. In light of the above results, we suggest that the 2.5th percentile of ANC for Saudi Arabians could be set at  $1.0 \times 10^9/L$ . Additionally, ANC of  $> 0.8 \times 10^9/L$ , when isolated, chronic and stable with benign medical history, should not be considered alarming in the absence of fever, pyogenic infection, or inflammation.

## Data Sharing Statement

Data used in this research are available through corresponding author to readers upon request.

## Acknowledgment

The authors extend their appreciation to the research center for advanced materials (RCAMS) at King Khalid University, Abha, Saudi Arabia for supporting this work through the research group program under grant number (RCAMS/KKU/0020/20).

## Disclosure

All authors declare that there are no conflicts of interest in this work.

## References

- Nauseef WM. Neutrophils, from cradle to grave and beyond [Internet]. *Immunol Rev*. 2016;273:5–10. doi:10.1111/imr.12463
- Kobayashi Y. Neutrophil biology: an update [Internet]. *EXCLI J*. 2015;14:220–227. doi:10.17179/excli2015-102
- Hoffbrand AVMP. The white cells 1: granulocytes, monocytes and their benign disorders. In: Hoffbrand AVMP, editor. *Hoffbrand's Essential Hematology*. 7th ed. Wiley-Blackwell; 2016:87–101.
- Peterson RF, Peterson WG. The differential count at high altitudes. *J Lab Clin Med*. 1935;20(7):723–726.
- Siqués P, Brito J, León-Velarde F, et al. Hematological and lipid profile changes in sea-level natives after exposure to 3550-m altitude for 8 months. *High Alt Med Biol*. 2007;8(4):286–295. doi:10.1089/ham.2007.8405
- Xia J, Miller CA, Baty J, et al. Somatic mutations and clonal hematopoiesis in congenital neutropenia. *Blood*. 2018;131:408–416. doi:10.1182/blood-2017-08-801985
- Atallah-Yunes SA, Ready A, Newburger PE. Benign ethnic neutropenia. *Blood Rev*. 2019;37:100586. doi:10.1016/j.blre.2019.06.003
- Lakhota R, Aggarwal A, Link ME, Rodgers GP, Hsieh MM. Natural history of benign ethnic neutropenia in individuals of African ancestry. *Blood Cells Mol Dis*. 2019;77:12–16. doi:10.1016/j.bcmd.2019.01.009
- Phillips D, Rezvani K, Bain BJ. Exercise induced mobilisation of the marginated granulocyte pool in the investigation of ethnic neutropenia. *J Clin Pathol*. 2000;53(6):448–483. doi:10.1136/jcp.53.6.481
- Souto Filho JTD, Portugal RD, Nucci M. Effect of circadian variation on neutrophil mobilization to the peripheral blood in benign constitutional neutropenia. *Exp Hematol*. 2019;69:22–26. doi:10.1016/j.exphem.2018.10.007
- Denic S, Narchi H, Al Mekaini LA, Al-Hammadi S, Al Jabri ON, Souid AK. Prevalence of neutropenia in children by nationality. *BMC Hematol*. 2016;16(1):1–7. doi:10.1186/s12878-016-0054-8
- Gari M, Dakhkhni M, Gari A, et al. Incidence and potential causative factors associated with chronic benign neutropenia in the Kingdom of Saudi Arabia. *BMC Proc*. 2015;9(2):1–7. doi:10.1186/1753-6561-9-S2-S1
- Ortiz MV, Meier ER, Hsieh MM. Identification and clinical characterization of children with benign ethnic neutropenia. *J Pediatr Hematol Oncol*. 2016;38(3):e140–3. doi:10.1097/MPH.0000000000000528
- Alkhalidy H, Awan ZA, Abouzaid AA, et al. The prevalence of isolated neutropenia at high altitude in Southern Saudi Arabia: does altitude affect leucocyte count? *Int J Gen Med*. 2020;13:1373–1379. doi:10.2147/IJGM.S284133
- Wonodi I, Oduguwa T, Amoo I, et al. Benign ethnic neutropenia in a sample of Nigerian healthy controls and clozapine-treated schizophrenia patients with DARC null variant. *Schizophr Bull*. 2017;43(suppl\_1):S67. doi:10.1093/schbul/sbx021.178
- Goswami R, Kakkar N, John MJ. Lower total leukocyte and neutrophil counts in healthy young Africans from Uganda. *Indian J Hematol Blood Transfus*. 2018;34(2):278–281. doi:10.1007/s12288-017-0866-0
- Merz LE, Achebe M. When non-whiteness becomes a condition. *Blood*. 2020.
- Rappoport N, Simon AJ, Lev A, et al. Correlation between “ACKR1/DARC null” polymorphism and benign neutropenia in Yemenite Jews [Internet]. *Br J Haematol*. 2015;170:892–895. doi:10.1111/bjh.13345
- Rappoport N, Simon AJ, Amariglio N, Rechavi G. The Duffy antigen receptor for chemokines, ACKR1,– ‘Jeanne DARC’ of benign neutropenia. *Br J Haematol*. 2019;184(4):497–507. doi:10.1111/bjh.15730
- Fragiadaki I, Papadakis S, Sevastaki G, et al. Increased frequency of the single nucleotide polymorphism of the DARC/ACKR1 gene associated with ethnic neutropenia in a cohort of European patients with chronic idiopathic neutropenia. *Am J Hematol*. 2020;95(7):E163–6. doi:10.1002/ajh.25813
- Palmblad J, Höglund P. Ethnic benign neutropenia: A phenomenon finds an explanation. *Pediatr Blood Cancer*. 2018;65(12):e27361. doi:10.1002/pbc.27361
- Dinardo CL, Kerbaay MN, Santos TC, et al. Duffy null genotype or Fy(a-b-) phenotype are more accurate than self-declared race for diagnosing benign ethnic neutropenia in Brazilian population [Internet]. *Int J Lab Hematol*. 2017;39:e144–6. doi:10.1111/ijlh.12712
- Fahad Almsned LA. Frequency of the ABO and Duffy Phenotypes among Saudi Population in Eastern Provence. *J Med Sci Clin Res*. 2017;05(03):19295–19298. doi:10.18535/jmscr/v5i3.154
- Owaidah AY, Naffaa NM, Alumran A, Alzahrani F. Phenotype frequencies of major blood group systems (Rh, kell, kidd, duffy, mns, p, lewis, and lutheran) among blood donors in the eastern region of Saudi Arabia. *J Blood Med*. 2020;11:59–65. doi:10.2147/JBM.S236834
- El-Hazmi MAF, Al-Swailem AR, Warsy AS, Al-Swailem AM, Sulaimani R, Al-Meshari AA. Consanguinity among the Saudi Arabian population. *J Med Genet*. 1995;32:623–626. doi:10.1136/jmg.32.8.623
- Jin S, Pazdur R, Sridhara R. Re-evaluating eligibility criteria for oncology clinical trials: analysis of investigational new drug applications in 2015. *J Clin Oncol*. 2017;35(33):3745–3752. doi:10.1200/JCO.2017.73.4186
- Vastola ME, Yang DD, Muralidhar V, et al. Laboratory eligibility criteria as potential barriers to participation by black men in prostate cancer clinical trials [Internet]. *JAMA Oncol*. 2018;4:413–414. doi:10.1001/jamaoncol.2017.4658
- Denduluri N, Lyman GH, Wang Y, et al. Chemotherapy dose intensity and overall survival among patients with advanced breast or ovarian cancer. *Clin Breast Cancer*. 2018;18(5):380–386. doi:10.1016/j.clbc.2018.02.003
- Crawford J, Denduluri N, Patt D, et al. Relative dose intensity of first-line chemotherapy and overall survival in patients with advanced non-small-cell lung cancer. *Support Care Cancer*. 2020;28(2):925–932. doi:10.1007/s00520-019-04875-1
- Hsieh MM, Tisdale JF, Rodgers GP, Young NS, Trimble EL, Little RF. Neutrophil count in African Americans: lowering the target cutoff to initiate or resume chemotherapy? *J Clin Oncol*. 2010;28:1633–1637. doi:10.1200/JCO.2009.24.3881
- Legge SE, Christensen RH, Petersen L, et al. The Duffy-null genotype and risk of infection. *Hum Mol Genet*. 2020;1–9. doi:10.1093/hmg/ddz215

32. Donadieu J, Beaupain B, Fenneteau O, Bellanné-Chantelot C. Congenital neutropenia in the era of genomics: classification, diagnosis, and natural history [Internet]. *Br J Haematol*. 2017;557–574. doi:10.1111/bjh.14887
33. Barreto MESF, Lipay ME, Santos LD, et al. Duffy phenotyping and FY\*B-67T/C genotyping as screening test for benign constitutional neutropenia. *Hematol Transfus Cell Ther*. 2020. doi:10.1016/j.htct.2020.08.015

### International Journal of General Medicine

Dovepress

### Publish your work in this journal

The International Journal of General Medicine is an international, peer-reviewed open-access journal that focuses on general and internal medicine, pathogenesis, epidemiology, diagnosis, monitoring and treatment protocols. The journal is characterized by the rapid reporting of reviews, original research and clinical studies

across all disease areas. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/international-journal-of-general-medicine-journal>