

SCIENTIFIC REPORTS



OPEN

Maximum temperature accounts for annual soil CO₂ efflux in temperate forests of Northern China

Received: 11 January 2015

Accepted: 16 June 2015

Published: 16 July 2015

Zhiyong Zhou^{1,2}, Meili Xu^{1,2}, Fengfeng Kang¹ & Osbert Jianxin Sun^{1,2}

It will help understand the representation legality of soil temperature to explore the correlations of soil respiration with variant properties of soil temperature. Soil temperature at 10 cm depth was hourly logged through twelve months. Basing on the measured soil temperature, soil respiration at different temporal scales were calculated using empirical functions for temperate forests. On monthly scale, soil respiration significantly correlated with maximum, minimum, mean and accumulated effective soil temperatures. Annual soil respiration varied from 409 g C m⁻² in coniferous forest to 570 g C m⁻² in mixed forest and to 692 g C m⁻² in broadleaved forest, and was markedly explained by mean soil temperatures of the warmest day, July and summer, separately. These three soil temperatures reflected the maximum values on diurnal, monthly and annual scales. In accordance with their higher temperatures, summer soil respiration accounted for 51% of annual soil respiration across forest types, and broadleaved forest also had higher soil organic carbon content (SOC) and soil microbial biomass carbon content (SMBC), but a lower contribution of SMBC to SOC. This added proof to the findings that maximum soil temperature may accelerate the transformation of SOC to CO₂-C via stimulating activities of soil microorganisms.

Global surface temperature is predicted to keep increasing by about 4°C until the end of this century, which will particularly accelerate the carbon cycles of terrestrial ecosystems^{1,2}. Vice versa, increased CO₂ concentration of atmosphere will inevitably lead to a higher temperature increase in future scenarios of climate change. This apparently demonstrates a tight correlation between soil CO₂ efflux and temperature. Moreover, their robust relationship virtually depends on the properties of soil temperature and other biophysical factors under diverse environmental conditions.

The CO₂ emitted from soil surface (i.e. soil respiration) derives from a composite process, normally classified into autotrophic respiration by the activities of plant root, and heterotrophic respiration via the microbial decomposition of soil organic matter^{3,4}. Its magnitude is conjointly determined by many biotic and abiotic factors, including gross primary production, plant photosynthesis, plant roots, soil microbial biomass and activity, soil organic matter content, soil water content, and soil temperature, etc^{3–5}. Furthermore, the interactions among these environmental variables deepen the complexity of the mechanism underlying soil CO₂ efflux. For example, soil respiration was decoupled from soil temperature when soil moisture content was below the threshold value⁶. In addition, precipitation pulse events could abruptly increase soil respiration rate in short period⁷. It is indicated that empirical models integrating more variables could provide a more accurate estimation of soil CO₂ efflux than that with a single ecological variable^{4,8}. Although the effects of other variables on soil respiration have received much attention, soil temperature and soil moisture content are still the two predominant variables extensively used to

¹Ministry of Education Key Laboratory for Silviculture and Conservation, Beijing Forestry University, Beijing, 100083, China. ²The Institute of Forestry and Climate Change Research, Beijing Forestry University, Beijing, 100083, China. Correspondence and requests for materials should be addressed to Z.Z. (email: zhiyong@bjfu.edu.cn)

explain the variance of soil respiration. In principle, temperature is the most fundamental parameter, around which many models have been built, including the well-known Arrhenius function⁹. The effects of other environmental factors could be reduced by modifying the empirical models through clarifying the magnitude of the temperature dependence of soil respiration.

Accumulative soil CO₂ efflux on a long-term scale is a vital component of ecosystem respiration, and in turn determines the carbon balance between terrestrial ecosystems and the atmosphere. The emergence of automatically continuous measurement system of soil respiration offers an opportunity to evaluate this carbon balance⁷, however, annual soil respiration is still calculated via empirical functions using continuously (i.e., hourly or daily) monitored data of soil temperature¹⁰. This is because soil respiration is a temperature dependent process, and these two time series can fit well a number of positive mathematical relationships^{2,11}. Yet, these empirical functions have been founded basing on the steady state assumption of the temporal correlation between soil respiration and temperature with time. In essence, temporal variation is an inherent property of soil temperature. Although it has been extensively accepted that the seasonal variation of soil respiration is induced by seasonal changes in soil temperature, no much attention has been paid on the effects of maximum or minimum soil temperature on soil CO₂ efflux.

Soil temperature normally exhibits great heterogeneity and fluctuating values over different periods from hourly to diurnal, seasonal, and annual, especially in temperate regions, which mainly accounts for the temporal variations of soil respiration⁷. This is the reason that long-term soil respiration can be upscale estimated using daily, monthly, and mean annual temperature (MAT)¹². However, the surrogate efficacy of average soil temperature has been questioned by a number of studies^{2,4}. First of all, the temporal correlation of soil respiration with temperature has not been found to be significant on every day of the year⁴. Additionally, Bond-Lamberty and Thomson² have demonstrated the vital importance of temperature anomaly in determining the amplitude of soil respiration. These indicate that soil temperature is a more flexible biophysical factor and can reflect the changeable characteristics of natural circumstance. Particularly, the role of extreme soil temperature (i.e. maximum and minimum) still remains unclear in regulating soil respiration. In order to save the enormous efforts in measuring the soil CO₂ efflux at a long time period, it is worthy to make certain what property of soil temperature can be the suitable surrogate of soil respiration for forest ecosystems.

As the climate has been undergoing a warming scenario, we hypothesized that it was the temperature increase of warm season that would lead to a higher soil CO₂ efflux of temperate forests. To evaluate the impacts of the warmest soil temperatures at diurnal, monthly and seasonal scales on soil respiration, we measured instantaneous rate of soil respiration and coupled soil temperature periodically, and monitored hourly soil temperature for a whole year at nine forest plots within a catchment of temperate forests in northern China. In this study, our main objectives were to identify (1) the correlation of soil respiration with soil temperatures on variant temporal scales, (2) the manipulation of maximum temperature on accumulative soil CO₂ efflux, for temperate forests of northern China.

Results

Accumulative soil CO₂ efflux. Annual soil respiration markedly varied from 409 g C m⁻² in coniferous forest to 570 g C m⁻² in mixed forest and to 692 g C m⁻² in broadleaved forest ($P=0.004$, $n=3$). Significant difference in monthly soil respiration was also detected among these three forest types through warm seasons from September to October in 2012 and from April to August in 2013. The highest monthly soil respiration turned out in July and was 75.24 g C m⁻², 139.69 g C m⁻², and 116.64 g C m⁻², and the soil respiration in January was lowest at 14.03 g C m⁻², 13.77 g C m⁻², and 13.32 g C m⁻², respectively in coniferous, broadleaved, and mixed forests. Generally, soil CO₂ efflux during summer (Jun, Jul, Aug) accounted for 50.73% of annual soil respiration for temperate forests. Winter (Dec, Jan, Feb) soil respiration was only about 8.12% of total annual soil CO₂ efflux. Soil CO₂ efflux in spring (Mar, Apr, May) and autumn (Sep, Oct, Nov) equally took over 20% or so of the total CO₂ efflux during the study year (Fig. 1).

The effects of different forest types on ecological variables. Forest type greatly influenced basal soil respiration (R_{10}), SOC, soil total nitrogen content (TN), soil pH, and litter carbon stock in northern China. R_{10} was significantly higher in broadleaved and mixed forests than in coniferous forest ($P=0.015$, $n=3$). Additionally, broadleaved forest also had the highest SOC and TN contents among these three forest types. SOC and TN were 83.62% and 93.10%, 94.82% and 90.48% higher in broadleaved forest than in coniferous and mixed forests, separately. Soil pH of broadleaved forest was significant and 13.02% larger than that of mixed forest. However, litter carbon stock was highest in coniferous forest, and was 80.85% and 94.54% higher than in broadleaved and mixed forests. No significant differences were found in C:N ratio, SMBC, mean annual temperature (MAT), and Q_{10} among these three forest types (Table 1).

Seasonal variation in soil temperature. Distinct seasonal changes in mean, maximum and minimum soil temperature were detected in these three forest types, with the highest values in July and the lowest values in January, during the study period. On the monthly scale, maximum soil temperature varied from -0.52°C to 17.27°C , from -0.72°C to 19.60°C , and from -0.73°C to 18.79°C , and minimum soil temperature ranged from -3.04°C to 14.91°C , from -4.08°C to 15.61°C , and from -6.17°C to 15.71°C , and mean soil temperature varied from -1.71°C to 15.89°C , from -3.03°C to 16.99°C ,




Figure 1. Seasonal variation of soil respiration during September 2012 and August 2013 among three types of temperate forest. Different lowercase letters above bars mean significant difference ($P < 0.05$, $n = 3$) among these three forest types.

Forest type	SOC (g/kg)	TN (g/kg)	C : N	SMBC (mg/kg)	pH	Soil bulk density (g/cm ³)	Litter C stock (g C/m ²)	MAT (°C)	R ₁₀ (μmol CO ₂ m ⁻² s ⁻¹)	Q ₁₀
Coniferous forest	18.43 a	1.45 a	12.64 a	828.76 a	6.05 a	1.10 a	649.22 b	6.10 a	1.25 a	2.82 a
Broadleaved forest	33.86 b	2.80 b	11.99 a	868.55 a	6.51 b	0.95 a	359.04 a	7.03 a	1.85 b	3.17 a
Mixed forest	17.38 a	1.47 a	11.75 a	536.59 a	5.76 a	1.04 a	333.78 a	6.41 a	1.62 b	3.36 a

Table 1. Biophysical and chemical variables of three forest types in northern China. Different lowercase letters in the same column mean significant difference ($P < 0.05$, $n = 3$) among these three forest types.

and from -3.32°C to 16.98°C , respectively in coniferous forest, broadleaved forest, and mixed forest (Fig. 2a,b,d).

However, there was no general changing trend of soil temperature range among these forest types. The largest value of soil temperature range happened over April, the lowest was over February. The temperature range over the warmest month of July kept the second lowest value (Fig. 2c).

Correlation of warmest temperature with annual soil respiration. In northern China, July and summer are the warmest month and season with highest temperatures. Mean July soil temperature increased from 15.59°C to 17.92°C , mean summer soil temperature changed from 14.39°C to 16.59°C , and mean soil temperature on the warmest day varied from 16.94°C to 22.03°C , during the study period for these three forest types. These three properties of soil temperature extremely affected accumulative soil CO₂ efflux over an annual scale for temperate forests. Annual soil respiration significantly linearly increased with mean soil temperatures on the warmest day, in July and summer during the study period in temperate forests (Fig. 3a–c). In particular, no significant correlation was found between annual soil CO₂ efflux and mean annual soil temperature for these three forest types ($R^2 = 0.43$, $P = 0.06$, $n = 9$; original data not shown).

The variation of soil CO₂ efflux with soil temperature at monthly scale. The changing trend of soil CO₂ efflux was in a good parallel with that of soil temperature on a monthly scale for temperate forests in northern China. Monthly soil respiration was positively and exponentially correlated with the maximum soil temperature, the minimum soil temperature, and mean soil temperature during the corresponding period for all three types of temperate forest (Fig. 4a–f, 5a,c,e). Concurrently, an extremely significant and positive correlation existed between accumulative soil CO₂ efflux and accumulated effective soil temperature on a monthly scale in the studied forest communities. Most of the variation in monthly soil respiration could be ascribed to the increase in the accumulated effective soil temperature in all three types of temperate forest (Fig. 5b,d,f).




Figure 2. Temporal changes in maximum soil temperature (a), minimum soil temperature (b), soil temperature range (c), and mean soil temperature (d) during September 2012 and August 2013 among three types of temperate forest.

Discussion

Soil CO₂ efflux has received a great deal of attention since the earlier studies^{4,13}, and its importance has been extensively accepted in determining the role of a forest ecosystem as a carbon sink/source¹⁴. The carbon sequestration of a forest ecosystem largely depends on the difference between net primary production and heterotrophic soil CO₂ efflux. The CO₂ efflux through heterotrophic decomposition constitutes at least 50% of total soil respiration in northern temperate forests¹⁵.

Generally speaking, the dynamic balance between soil organic carbon input and output was virtually determined by the accumulative soil CO₂ efflux on a normally annual scale. In the study region of northern China, annual soil respiration varied from 409 g C m⁻² yr⁻¹ in coniferous forest to 692 g C m⁻² yr⁻¹ in broadleaved forest, which was comparable to those of similar forest types in the same climate zone^{10,16} and fell right within the range of six temperate forests in northeast China¹⁷. The larger accumulative soil CO₂ efflux of broadleaved forest was mainly due to its higher soil temperature over different temporal scales, as temperature increase in the warmest period was indicated by this study results to lead to a greater soil respiration.

Soil respiration of temperate forests displayed distinct seasonal variation with nearly 51% CO₂ emitted in summer and only 8.00% CO₂ respired in winter. The proportion of winter soil CO₂ efflux to total annual soil respiration agreed well with 4.92–7.83% of Wang *et al.*¹⁸. This was mainly attributed to the robust low value of winter soil temperature and the thickness and duration of snow cover¹⁹. Although winter temperature was indicated to have a vital impact on plant phenology²⁰, we speculated that future increase in winter temperature would have limited impact on annual soil CO₂ efflux in temperate forests of northern China, because of the lowest proportion of winter soil respiration.

Seasonal changes were clearly observed in maximum soil temperature, minimum soil temperature, mean soil temperature, soil temperature range, and accumulated effective soil temperature during the study period of twelve months for the three types of temperate forest in northern China. Generally, broadleaved forest had a comparatively higher value in these properties of soil temperature. This was because the broadleaved forest mainly consisted of *Q. wutaishanica*, which normally distributed along the south direction slope of the investigated mountain. On the northfacing slope stood the coniferous forest with a lower soil temperature.

In addition, soil CO₂ efflux on a monthly scale in three forest types was largely and significantly explained by the changes in maximum, minimum, mean, and accumulated effective soil temperatures at the corresponding temporal scales. It was evident that the temporal variation of soil respiration greatly




Figure 3. Correlations of soil CO₂ efflux during September 2012 and August 2013 with mean soil temperature of the warmest day (a), mean July soil temperature (b), and mean summer soil temperature (c) for temperate forests.

depended on the seasonal changing trend of soil temperature in temperate forests. Furthermore, soil CO₂ efflux on an annual scale was well interpreted respectively by mean soil temperatures on the warmest day, in the warmest month and season for these three temperate forests. This suggested that soil temperature in the warmest period was the most appropriate predictor in modeling the annual soil respiration in temperate forests of northern China. This finding slightly differed from the results of Bahn *et al.*¹² and Litton *et al.*²¹, who pointed out that total annual CO₂ emission could be predicted by the soil respiration rate at mean annual temperature or increase with mean annual temperature. In this study, no significant correlation was ever detected between mean annual soil temperature and annual soil CO₂ efflux for these three temperate forests. Mean annual value may obscure the effects of soil temperature fluctuations at diurnal or seasonal scales on instantaneous rate of soil respiration, especially in temperate climatic zone.




Figure 4. Correlations of monthly soil respiration with maximum and minimum soil temperature during September 2012 and August 2013 for coniferous forest (a,b), broadleaved forest (c,d) and mixed forest (e,f).

Furthermore, water-limitation was perhaps another contributor to the decoupling between mean annual soil temperature and annual soil respiration in the studied temperate forests².

Overall, the surrogate legality of soil temperature also relied on the period length over which accumulative soil CO₂ efflux was calculated. For temperate forests in northern China, appropriate predicting factors varied from maximum and minimum soil temperatures, mean soil temperature, and accumulated effective soil temperature for the estimation of monthly soil respiration to mean July soil temperature, mean summer soil temperature, and soil temperature of the warmest day for the calculation of annual soil CO₂ efflux. In these different properties of soil temperature, only the soil temperature in the warmest periods was a most suitable variable to predict the accumulative soil CO₂ efflux at both monthly and annual scales.

Soil respiration derived from two major biological processes, autotrophic respiration by the growth and maintenance of plant roots, and heterotrophic respiration via the microbial decomposition of carbohydrates from live roots, litter, and old soil organic matter^{3,4}. Both of these two fluxes were greatly affected by higher temperature. In warm season, plant has higher physiological activity, e.g. plant growth and photosynthesis, which simultaneously induced a larger proportion of autotrophic respiration²²; soil microbial activity was also motivated by warm soil temperature, which speeded the turnover rate of soil organic carbon and produced more heterotrophic CO₂ efflux²³. In this study, broadleaved forest had a significant higher SOC content than both coniferous and mixed forests ($P = 0.026$, $n = 3$). Soil microbial biomass carbon was comparatively higher in broadleaved forest than in coniferous and mixed forests, although the difference did not reach the significant level ($P = 0.40$, $n = 3$). This still agreed well with the higher soil temperature in broadleaved forest. The broadleaved forest with the higher soil temperature and soil organic matter could provide much substrate for microbial organism to decompose and release a larger amount of CO₂-C at both monthly and annual scale in the studied temperate forests. Another




Figure 5. Correlations of monthly soil respiration with mean monthly soil temperature and monthly accumulated effective soil temperature during September 2012 and August 2013 for coniferous forest (a,b), broadleaved forest (c,d) and mixed forest (e,f).

reason for the higher accumulative soil respiration in broadleaved forest was the slightly greater activity of soil microbial organism represented by the higher soil microbial biomass carbon content.

Otherwise, the microbial biomass carbon content was 25.52 mg g SOC⁻¹ and much lower in broadleaved forest than 32.11 mg g SOC⁻¹ and 50.17 mg g SOC⁻¹ in mixed and coniferous forests, respectively. The contribution of SMBC to SOC showed a contrary changing trend compared to soil temperature among these three forest types. This would help explain why the forest plot with higher soil temperature released much more CO₂-C, because the contribution of SMBC to SOC reflected changes in property of natural ecosystems and demonstrated the status of soil carbon accumulation²⁴. In essence, increased temperature might accelerate the transformation of soil organic matter into CO₂-C by soil microbes. A warming experiment indicated that soil microbial metabolic activity was increased 66% by warming, and new and recent fixed carbon were respired 38% larger than under control temperature conditions²⁵. More soil labile carbon was perhaps lost through the motivated heterotrophic respiration, which eventually slowed down the accumulation of soil organic carbon. Therefore, this kind of activation of heterotrophic respiration by higher temperature may be the underlying mechanism for maximum temperature accounting for the annual soil respiration in temperate forests of northern China.

Material and Methods

Study sites and experiment layout. This study was conducted at the Taiyueshan Long-Term Forest Ecosystem Research Station (36°04' N, 112°06' E; elevation during 600–2,600 m a.s.l.), which situates about 190 km southwest of Taiyuan in Shanxi province of China. Experimental site is about 3 km west to the station in a classic catchment named after Xiaoshegou. This area has four distinct seasons, i.e. spring from March to May, summer from June to August, autumn from September to November, and winter from December to February. However, most of precipitation occurs between June and August

Forest type	Tree height (m)	Diameter at breast height (cm)	Plant density (hm ⁻²)	Elevation range (m.a.s.l)	Forest age (yr)	Slope gradient
Coniferous forest	10.94	16.50	1,808	1,690	70	17°
Broadleaved forest	7.16	14.35	1492	1,653	70	19°
Mixed forest	7.56	10.41	4033	1,710	70	22°

Table 2. Basic characteristics of the studied temperate forests in northern China.

with a mean annual precipitation of 600 mm, and the mean annual temperature (MAT) is about 11°C, with 26°C in the warmest period of July and -23°C in the coldest month of January. The hill height at this study region averages 600 m from the baseline at 1200 m a.s.l. The edaphic characteristic of this hill slope belongs to Eutric Cambisols (FAO classification) or Cinnamon soil (Chinese Classification) with a mean soil depth between 30 cm and 50 cm^{26,27}. The overstory vegetation consists mainly of *Pinus tabulaeformis*, *Quercus wutaishanica*, *Betula dahurica*, *Larix gmelinii* var. *principis-rupprechtii*, *Tilia mongolica*. The understory shrub community mainly consists of *Corylus mandshurica*, *Corylus heterophylla*, *Acer ginnala*, *Lespedeza bicolor*, *Philadelphus incanus*, *Rosa bella*, *Lonicera chrysantha*. Major herbaceous species includes *Carex lanceolata*, *Spodiopogon sibiricus*, *Rubia chinensis*, *Thalictrum petaloideum*, *Melica pappiana*.

Along three ridges within this catchment, three forest types, i.e. coniferous, broadleaved and mixed forests, were investigated to complete this study. Simultaneously, three forest plots at 20 × 20 m were constructed for each forest type. It was needed to be specified that the six plots of broadleaved and mixed forests were ever used in another research of Zhou *et al.*²⁸, who mainly focused on basal rates of soil respiration and their determinants. Three coniferous forest plots were originally used to explore the relationship of soil respiration with soil temperature properties, although they had been built and investigated in 2008 and 2009. These nine forest plots distributed along the catchment topography with coniferous forest plots on northfacing slope, broadleaved forest plots on southfacing slope, and mixed forest plots on top slope. Basic characteristics of these three types of forest community were investigated in late August 2009 as described in Zhou *et al.*²⁸ (Table 2).

Soil respiration measurement. Nine soil collars were permanently inserted into the soil at each forest plot, and soil respiration was periodically measured during the growing season in both 2008 and 2009, using a Li – Cor infrared gas analyzer (LI – 8100, Li – Cor Inc., Lincoln, NE, U.S.A.), equipped with a portable chamber. Concurrently, soil temperature at 10 cm depth close to each soil collar was monitored using a thermocouple probe attached to LI-8100 system. The details about these measurements could be seen in Zhou *et al.*²⁸, and soil respiration rate at 10°C (R_{10}) and temperature sensitivity (Q_{10} , soil respiration change with a proportional change of 10°C in soil temperature) were calculated for each forest plot via the Arrhenius function using instantaneous soil respiration rate and the coupled soil temperature²⁸.

Soil temperature at 10 cm depth on each forest plot was hourly and automatically logged from September 1st 2012 to August 31st 2013 using a temperature monitor (HOBO U22-001, Onset, U.S.A.), which was buried 10 cm in the top soil through one year period. As R_{10} and Q_{10} were indicated to be two comparatively constant parameters for a specific ecosystem²⁸, then we were able to estimate the soil respiration rate at a given hour with the logged temperature complying with the empirical function (1). No change was proposed to happen in soil temperature over an hour, so did the soil respiration rate. Therefore, the accumulative soil respiration on daily, monthly, or annual scale could be calculated following the function (2).

$$R_{si} = R_{10} \times Q_{10}^{(T_{si}-10)/10} \quad (1)$$

$$R = \sum R_{si} \times 3,600 \times 12 \times 10^{-6} = \sum R_{10} \times Q_{10}^{(T_{si}-10)/10} \times 3,600 \times 12 \times 10^{-6} \quad (2)$$

where R_{si} ($\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$) represents estimated soil respiration rate at aliquot i^{th} hour, T_{si} (°C) is the soil temperature at 10 cm depth automatically logged at aliquot i^{th} hour, R_{10} ($\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$) and Q_{10} are the calculated parameters of soil respiration using field measurement data of instantaneous soil respiration rate and coupled soil temperature, R (g C m^{-2}) refers to the accumulative CO_2 efflux on daily, monthly, or annual scale, 3,600 means 3,600 seconds per hour, 12 means 12 gram carbon per mole CO_2 , and $1 \mu\text{mol}$ equals 10^{-6} mole. Mean values of R_{10} and Q_{10} used in functions (1) and (2) can be seen in Table 1 for each forest type.

In order to explore the effects of soil temperature properties on soil respiration, we calculated the maximum and the minimum soil temperature, i.e. the soil temperature respectively at the warmest and coldest time point during a month or a year, soil temperature range equaled the maximum temperature minus the minimum temperature during each month, and the effective soil temperature equaled mean daily temperature minus 5°C.

Analyses of soil biochemical properties. Five soil cores were sampled on each forest plot in late August of 2013 using a cylindrical driller of 4 cm in diameter and 20 cm in depth. Sampled soils were put in polyethylene bags and shipped immediately to the laboratory in a box cooled by ice. Prior to nutrient analysis, soils were sieved with a screen of 2 cm, and were evenly separated into two parts after any visible debris was manually picked out. One subsample was air-dried to constant weight and ground to pass through a 0.2 mm sieve before it was used to determine soil organic carbon (SOC) content via the standard Mebius method²⁹. Soil total nitrogen content was determined using the Kjeldahl digestion procedure with a Tector Kjeltex system 1026 Distilling unit³⁰. Another subsample was stored in refrigerator at 4 °C, and was analyzed for soil microbial biomass carbon (SMBC) through the chloroform fumigation extraction method no longer than seven days since it was collected³¹. Moist soil (equal 15 g oven-dried soil) was extracted with 60 ml of 0.5 M/L K₂SO₄ and was shaking for half an hour in a polyethylene bottle with elastic plug. Another 15 g soil (dry weight basis) was firstly fumigated with chloroform in a glass jar with CO₂-free air, and was incubated for 24 hours at 25 °C in a dark microcosm with air moisture content of 60%. The fumigated soil was extracted in the same way after the incubation was finished. The total amount of microbial biomass carbon was determined by the difference between K₂SO₄ – extracted carbon in fumigated and non-fumigated soil using K_{ec} factor of 0.38. Simultaneously, soil bulk density was determined using a cylindrical soil driller of 100 cm³ and oven-dried at 110 °C to constant weight in laboratory. Soil pH was measured in deionized H₂O via Sartorius AG (PB-10, Sartorius, Germany).

Data analysis. All biophysical variables including soil respiration, soil temperature, and soil resource content, etc. were averaged across forest plots for each forest type when they were analyzed in this article. Analysis of variance (ANOVA) at $\alpha = 0.05$ significance level was used to test the effect of forest type on biophysical variables. All data analyses were performed using the software of SPSS 15.0. Figures of this paper were completed via the software of Sigmaplot in version 10.0.

References

1. Van Vuuren, D. P. *et al.* Temperature increase of 21st century mitigation scenarios. *PNAS*. **105**, 15258–15262 (2008).
2. Bond-Lamberty, B. & Thomson, A. Temperature-associated increases in the global soil respiration record. *Nature*. **464**, 579–582 (2010).
3. Kuzyakov, Y. & Gavrichkova, O. Time lag between photosynthesis and carbon dioxide efflux from soil: a review of mechanisms and controls. *Global Change Biol.* **16**, 3386–3406 (2010).
4. Vargas, R., Carbone, M.S., Reichstein, M. & Baldocchi, D. D. Frontiers and challenges in soil respiration research: from measurements to model-data integration. *Biogeochemistry*. **102**, 1–13 (2011).
5. Vargas, R. *et al.* Looking deeper into the soil: biophysical controls and seasonal lags of soil CO₂ production and efflux. *Ecol. Appl.* **20**, 1569–1582 (2010).
6. Vincent, G., Shahriari, A. R., Lucot, E., Badot, P. M. & Epron, D. Spatial and seasonal variations in soil respiration in a temperate deciduous forest with fluctuating water table. *Soil Biol. Biochem.* **38**, 2527–2535 (2006).
7. Vargas, R., Detto, M., Baldocchi, D. D. & Allen, M. F. Multiscale analysis of temporal variability of soil CO₂ production as influenced by weather and vegetation. *Global Change Biol.* **16**, 1589–1605 (2010).
8. Tang, X. L. *et al.* Dependence of soil respiration on soil temperature and soil moisture in successional forests in Southern China. *J. Integr. Plant Biol.* **48**, 564–663 (2006).
9. Arrhenius, S. Über die Reaktionsgeschwindigkeit bei der Inversion von Rohrzucker durch Säuren. *Zeitschrift für Physik Chemie*. **4**, 226–248 (1889).
10. Zhou, Z. *et al.* Predicting soil respiration using carbon stock in roots, litter and soil organic matter in forests of Loess Plateau in China. *Soil Biol. Biochem.* **57**, 135–143 (2013).
11. Fang, C. & Moncrieff, J. B. The dependence of soil CO₂ efflux on temperature. *Soil Biol. Biochem.* **33**, 155–165 (2001).
12. Bahn, M. *et al.* Soil respiration at mean annual temperature predicts annual total across vegetation types and biomes. *Biogeosciences*. **7**, 2147–2157 (2010).
13. Lundegårdh, H. Carbon dioxide evolution of soil and crop growth. *Soil Sci.* **23**, 417–453 (1927).
14. Etzold, S. *et al.* The carbon balance of two contrasting mountain forest ecosystems in Switzerland: similar annual trends, but seasonal differences. *Ecosystems*. **14**, 1289–1309 (2011).
15. Levy-Varon, J. H., Schuster, W. S. F. & Griffin, K. L. The autotrophic contribution to soil respiration in a northern temperate deciduous forest and its response to stand disturbance. *Oecologia*. **169**, 211–220 (2012).
16. Luan, J., Liu, S., Wang, J., Zhu, X. & Shi, Z. Rhizospheric and heterotrophic respiration of a warm-temperate oak chronosequence in China. *Soil Biol. Biochem.* **43**, 503–512 (2011).
17. Wang, C. K. & Yang, J. Y. Rhizospheric and heterotrophic components of soil respiration in six Chinese temperate forests. *Global Change Biol.* **13**, 123–131 (2007).
18. Wang, W., Peng, S., Wang, T. & Fang, J. Winter soil CO₂ efflux and its contribution to annual soil respiration in different ecosystems of a forest-steppe ecotone, north China. *Soil Biol. Biochem.* **42**, 451–458 (2010).
19. Elberling, B. Annual soil CO₂ effluxes in the high Arctic: the role of snow thickness and vegetation type. *Soil Biol. Biochem.* **39**, 646–654 (2007).
20. Clark, J., Melillo, J., Mohan, J. & Salk, C. The seasonal timing of warming that controls onset of the growing season. *Global Change Biol.* **20**, 1136–1145 (2014).
21. Litton, C. M., Giardina, C. P., Albano, J. K., Long, M. S. & Asner, G. P. The magnitude and variability of soil-surface CO₂ efflux increase with mean annual temperature in Hawaiian tropical montane wet forests. *Soil Biol. Biochem.* **43**, 2315–2323 (2011).
22. Bahn, M., Janssens, I. A., Reichstein, M., Smith, P. & Trumbore, S. E. Soil respiration across scales: towards an integration of patterns and processes. *New Phytol.* **186**, 292–296 (2010).
23. Billings, S. A. & Ford Ballantyne, I. V. How interactions between microbial resource demands, soil organic matter stoichiometry, and substrate reactivity determine the direction and magnitude of soil respiratory response to warming. *Global Change Biol.* **19**, 90–102 (2013).
24. Insam, H. & Domsch, K. H. Relationship between soil organic carbon and microbial biomass on chronosequences of reclamation sites. *Micro. Ecol.* **15**, 177–188 (1988).
25. Streit, K. *et al.* Soil warming alters microbial substrate use in alpine soils. *Global Change Biol.* **20**, 1327–1338 (2014).

26. Ma, Q. Y. Determination of the biomass of individual trees in stands of *Pinus Tabulaeformis* plantation in north China. *J. Beijing Forestry University* **4**, 1–18 (1983). (in Chinese with English abstract)
27. Guo, J. T., Wang, W. X., Cao, X. F., Wang, Z. Q. & Lu, B. L. A study on forest soil groups of Taiyue mountain. *J. Beijing Forestry University* **14**, 134–142 (1992). (in Chinese with English abstract)
28. Zhou, Z., Guo, C. & Meng, H. Temperature sensitivity and basal rate of soil respiration and their determinants in temperate forests of north China. *PLoS ONE*. **8**, e81793 (2013).
29. Nelson, D. W. & Sommers, L. E. in Total carbon, organic carbon, and organic matter, Methods of soil analysis. (ed. Miller, R. H. & Keeney, D. R.) 101–129 (Madison, 1982).
30. Gallaher, R. N., Weldon, C.O. & Boswell, F. C. A semiautomated procedure for total nitrogen in plant and soil samples. *Soil Sci. Soc. Am. J.* **40**, 887–889 (1976).
31. Vance, E. D., Brookes, P. C. & Jenkinson, D. S. An extraction method for measuring soil microbial biomass C. *Soil Biol. Biochem.* **19**, 703–707 (1987).

Acknowledgements

This research was performed with funding from the Special Research Program for Public-Welfare Forestry of the State Forestry Administration of China (Grant No. 201104008) and from the National Foundation of Natural Science of China (Grant No. 41003029). We would like to thank Chao Guo for his assistance in field data collection. We were also grateful to the Taiyueshan Long-Term Forest Ecosystem Research Station for the permit of carrying out field investigation on its experiment sites.

Author Contributions

Z.Z, F.K. and O.J.S. designed the experiment and established the forest plots together; M.X. helped analyze the soil samples and the original datasets; Z.Z. analyzed the experimental data and mainly wrote the manuscript. All authors reviewed the manuscript.

Additional Information

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Zhou, Z. *et al.* Maximum temperature accounts for annual soil CO₂ efflux in temperate forests of Northern China. *Sci. Rep.* **5**, 12142; doi: 10.1038/srep12142 (2015).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>