



MEETING ABSTRACT

Open Access

P01-008 – FMF genotype-phenotype correlations in Germany

M Jeske¹, P Lohse², T Kallinich³, T Berger⁴, C Rietschel⁵, D Holzinger⁶, C Kamlah⁷, P Lankisch⁸, R Berendes⁹, G Dückers¹⁰, G Horneff¹¹, E Lilienthal¹², JP Haas¹³, A Giese¹⁴, F Dressler¹⁵, J Berrang¹⁶, C Pütter¹⁷, L Braunewell¹, U Neudorf^{1*}, T Niehues¹⁰, E Lainka¹

From 7th Congress of International Society of Systemic Auto-Inflammatory Diseases (ISSAID) Lausanne, Switzerland. 22-26 May 2013

Introduction

Familial Mediterranean fever (FMF) is one the most common autoinflammatory disease (AID). Pathogenomic relevant mutations in the MEFV gene show autosomal recessive inheritance, but co-dominant mutations have been described.

Objectives

We aimed to evaluate correlations between ethnic origin, phenotype and genotype for FMF patients in the German AID-Net-registry.

Methods

We used two common scoring systems modified for children (Mor et al., Pras et al.) to assess disease severity in 243 FMF patients of the AID-Net-registry. For the four most frequent mutations, we tested for a correlation of the genotype with the phenotype, mean CRP and ethnic origin, respectively. Furthermore, we evaluated the applicability of the two severity scores for children.

Results

Among the 243 patients, we detected a total of 433 pyrin mutations and 22 different sequence variants, including one new mutation (p.Gly488Asp). The four most frequent alterations were p.Met694Val (55%, n=238), p.Met680Ile (12%, n=52), p.Val726Ala (10%, n=44) and p.Glu148Gln (8%, n=34). Ethnic origin could be determined in 224 cases; the prevailing ancestry was Turkish (83%, n=185), 8% (n=18) were Lebanese. P.Met694Val in homozygous form (n=74; 30.5%) was correlated with a more severe disease activity, based on the score by Mor, as well as with

a higher mean CRP (74 mg/l, n=60, 31 mg/l, n=59) compared to patients without this mutation (p=0.01 and p<0.01, respectively). The score suggested by Pras did not yield a significant genotype-phenotype correlation; indeed, the two scoring systems were inconsistent with each other (κ <0.07). Although a typical distribution of mutations in different ethnic populations was obvious, this trend was not statistically significant, probably due to the divergent number of cases.

Conclusion

The homozygous p.Met694Val substitution was associated with a more severe disease activity. There was no origin-genotype correlation in this FMF population. The well-known severity scores for children (Mor, Pras) are inconsistent.

Disclosure of interest

M. Jeske: None Declared, P. Lohse: None Declared, T. Kallinich: None Declared, T. Berger: None Declared, C. Rietschel: None Declared, D. Holzinger : None Declared, C. Kamlah: None Declared, P. Lankisch: None Declared, R. Berendes: None Declared, G. Dückers: None Declared, G. Horneff Consultant for: financial support for clinical trials from Abbott, Pfizer and Roche, E. Lilienthal: None Declared, J. Haas Consultant for: Pfizer, Roche and Novartis, A. Giese: None Declared, F. Dressler: None Declared, J. Berrang: None Declared, C. Pütter: None Declared, L. Braunewell: None Declared, U. Neudorf: None Declared, T. Niehues: None Declared, E. Lainka: None Declared.

Authors' details

¹Children's Hospital, University Duisburg-Essen, Essen, Germany. ²Labor Blessing und Partner, Singen, Germany. ³Charité Berlin, Children's hospital,

¹Children's Hospital, University Duisburg-Essen, Essen, Germany
Full list of author information is available at the end of the article

Berlin, Germany. ⁴Children's Hospital, University Witten/Herdecke, Datteln, Germany. ⁵Department of Pediatric Rheumatology, Clementine Hospital, Frankfurt, Germany. ⁶Children's Hospital, University Münster, Münster, Germany. ⁷Children's Hospital, UKE Hamburg, Hamburg, Germany. ⁸Pediatric Rheumatology, University Düsseldorf, Düsseldorf, Germany. ⁹Pediatric Rheumatology, St. Marien's Children's Hospital, Landshut, Germany. ¹⁰Pediatric Immunology and Rheumatology, HELIOS Children's hospital, Krefeld, Germany. ¹¹Centre of Pediatric Rheumatology, Asklepios Clinic, St. Augustin, Germany. ¹²Children's Hospital, Ruhr-University, Bochum, Germany. ¹³German Center for Pediatric Rheumatology, Children's Hospital, Garmisch-Partenkirchen, Germany. ¹⁴Marienhospital Herne, Ruhr-University Bochum, Herne, Germany. ¹⁵Pediatric Rheumatology Clinic, Hannover Medical School, Hannover, Germany. ¹⁶Pediatric Rheumatology, Westfaelisches Kinderzentrum, Dortmund, Germany. ¹⁷Institute for Medical Informatics, Biometry and Epidemiology, University Hospital, Essen, Germany.

Published: 8 November 2013

doi:10.1186/1546-0096-11-S1-A12

Cite this article as: Jeske *et al.*: P01-008 – FMF genotype-phenotype correlations in Germany. *Pediatric Rheumatology* 2013 **11**(Suppl 1):A12.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

