

RESEARCH

Open Access

Characterization of proteome alterations in *Phanerochaete chrysosporium* in response to lead exposure

Volkan Yıldırım¹, Servet Özcan², Dörte Becher³, Knut Büttner³, Michael Hecker³, Gülay Özcengiz^{1*}

Abstract

Background: Total soluble proteome alterations of white rot fungus *Phanerochaete chrysosporium* in response to different doses (25, 50 and 100 μ M) of Pb (II) were characterized by 2DE in combination with MALDI-TOF-MS.

Results: Dose-dependent molecular response to Pb (II) involved a total of 14 up-regulated and 21 down-regulated proteins. The induction of an isoform of glyceraldehyde 3-phosphate dehydrogenase, alcohol dehydrogenase class V, mRNA splicing factor, ATP-dependent RNA helicase, thioredoxin reductase and actin required a Pb (II) dose of at least 50 μ M. Analysis of the proteome dynamics of mid-exponential phase cells of *P. chrysosporium* subjected to 50 μ M lead at exposure time intervals of 1, 2, 4 and 8 h, identified a total of 23 proteins in increased and 67 proteins in decreased amount. Overall, the newly induced/strongly up-regulated proteins involved in (i) amelioration of lipid peroxidation products, (ii) defense against oxidative damage and redox metabolism, (iii) transcription, recombination and DNA repair (iv) a yet unknown function represented by a putative protein.

Conclusion: The present study implicated the particular role of the elements of DNA repair, post-transcriptional regulation and heterotrimeric G protein signaling in response to Pb (II) stress as shown for the first time for a basidiomycete.

Background

Heavy metal pollution is a major environmental concern due to its toxic effects through the food chain and its high persistence in the environment [1]. Lead (Pb) is one of the most abundant toxic metal; mining and smelting activities, lead containing paints, paper and pulp, gasoline and explosives as well as the disposal of municipal sewage sludge enriched with Pb being the main sources of pollution [2]. Possible mechanisms involved in metal-induced oxidative stress were extensively reviewed by Ercal *et al.* [3]. As a redox-inactive metal, Pb is known to deplete cells' major antioxidants, thiol-containing antioxidants and enzymes in particular.

The unique oxidative enzyme system of white-rot basidiomycetes is directly involved in complete lignin mineralization and degradation of various xenobiotic compounds as well as dyes [4,5]. *Phanerochaete*

chrysosporium is one of the best studied white-rot fungi shown to be very promising for treatment of phenolic effluents from pulp and paper, coal conversion, textile and olive oil industries. This organism is also very effective in biosorbing heavy metal ions from dilute solutions [6-8], with an equilibrium adsorptive capacity order of lead (II) > chromium (III) > copper (II) = cadmium (II) > nickel (II) [9]. Recently, *P. chrysosporium* was successfully employed for bioremediation of lead-contaminated soil [10,11]. It is well-known that tolerance to different metals varies greatly among microorganisms. When the above-mentioned heavy metals were compared for their effects on growth of *P. chrysosporium*, lead was the best tolerated metal in that the concentrations up to 100 μ M did not interfere with growth in liquid cultures (unpublished). Comparatively, only 5 μ M of this metal inhibits the growth of *S. cerevisiae* by approximately 30% [12].

The concentration level of lead found in wastewaters varies greatly depending on the type of wastewater which is ranging between 0.04 to 0.05 ppm in urban wastewater [13,14], 0.06 to 2.60 ppm in industrial

* Correspondence: ozcengiz@metu.edu.tr

¹Department of Biological Sciences, Middle East Technical University, Ankara, Turkey

Full list of author information is available at the end of the article

wastewater [15], 1.27 ppm in metal finishing waste water in particular [16], and 20 to 100 ppm in waste biogas residual slurry [17]. Temporal variability of these concentrations as caused, for example, by rain events or regular daily fluctuations as seen in wastewater treatment plant effluents is to be noted [18].

While the effects of toxic metals on cell physiology can be studied at the level of individual proteins, proteomics has allowed the responses to be studied on a much wider scale [19,20]. Regarding eukaryotic microorganisms, heavy metal stress proteomics have been published for *S. cerevisiae* [21,22], *Schizosaccharomyces pombe* [23] and *Chlamydomonas reinhardtii* [24]. Our group reported the first reference proteome map of *P. chrysosporium* along with an analysis of cadmium and copper response in this organism [25]. A total of 80 Cd-up-regulated and 74 Cu-up-regulated protein spots were detected and identified, 34 being common to the stress caused by both metals. Thus the aim of the present study is to determine the dynamic response of *P. chrysosporium* proteome to subtoxic levels of lead.

Results and Discussion

To investigate the proteome response of *P. chrysosporium* to Pb (II) exposure we used 2D electrophoresis followed by MALDI-TOF analysis. The cells were exposed either three levels of Pb (25, 50 and 100 μ M) for 40 h or 50 μ M of Pb after 40 h incubation for 1, 2, 4 and 8 h. This kind of setup helped us to observe proteome response of *P. chrysosporium* to acute and chronic exposure of Pb (II). The highest lead concentration employed in this study (100 μ M; ca 20 ppm) is about 8 fold higher than its upper level reported for industrial wastewaters [15].

When subjected to different levels of Pb (II), a total of 14 up-regulated and 21 down-regulated protein spots were identified. (Table 1 and 2). Among the up-regulated proteins, the proteins common to all three dosages included isocitrate dehydrogenase alpha subunit, UDP-glucose pyrophosphorylase, septin family protein (P-loop GTPase), F0F1-type ATP synthase, alpha subunit, GTPase Ran/TC4/GSP1, short-chain acyl-CoA dehydrogenase, polyadenylate-binding protein (RRM superfamily) and G protein beta subunit-like protein. On the other hand, the induction of an isoform of glyceraldehyde 3-phosphate dehydrogenase, alcohol dehydrogenase class V, mRNA splicing factor, ATP-dependent RNA helicase, thioredoxin reductase and actin required a Pb (II) dose of at least 50 μ M.

Among the up-regulated proteins, 6 were newly induced. One such protein was mitochondrial F0F1-type ATP synthase, alpha subunit (Figure 1a) which pointed to galvanized energy metabolism under stress. The other one was **actin** (Figure 1b) constituting a cytoskeleton

that plays a pivotal role in many eukaryotic signaling pathways. Actin-induced hyperactivation of the Ras signaling pathway was demonstrated to lead to apoptosis in *S. cerevisiae* [26]. A septin family protein (P-loop GTPase) was also newly induced in response to Pb (II) exposure (Figure 1b). Septin family of cytoskeletal proteins with GTPase activity are involved in many processes including membrane dynamics, vesicle trafficking, apoptosis and infection [27]. Ras GTPase [GTPase Ran/TC4/GSP1 (nuclear protein transport pathway; small G protein superfamily)] was found as a Cd responsive protein in our previous study [25], providing clue for the existence of Ras signaling pathway in *P. chrysosporium*, a pathway accelerating programmed cell death in *C. albicans* under harsh environmental stress [28]. In the present work, Ras GTPase was detected as a newly expressed protein upon Pb exposure (Figure 1c), thus providing another evidence for its role in heavy metal stress response.

RNA-binding proteins (RBPs) shuttle between cellular compartments either constitutively or in response to stress and regulate localization, translation, or turnover of mRNAs [29]. Post-transcriptional regulation can also occur through stabilization of mRNAs by specific RBPs in response to certain stimuli. The process of bulk export of mRNAs from nucleus to cytoplasm is highly conserved across eukaryotes. The export-competent mRNP consists of mRNAs and a dozen nucleocytoplasmic shuttling nuclear proteins, including RNA export factors, poly(A)-binding proteins, DEAD-box protein 5 and nucleoporins in yeast [30]. The RNA recognition motif (RRM) domain is by far the most abundant type of eukaryotic RNA-binding motif and besides mRNA binding, RRM domains involve in diverse protein-protein interactions. RBPs have been shown to translocate to the cytoplasm in response to stress. For example, A18 hnRNP, as induced by UV radiation, targets stress-activated transcripts and stimulates translation, thereby increasing survival after genotoxic stress [31]. In the present study, the identified RBPs were a newly-induced polyadenylate-binding protein (RRM superfamily) (Figure 1d) as well as two up-regulated proteins, namely splicing factor RNPS1 (Figure 1e) and ATP-dependent RNA helicase (Figure 1f). Splicing factor RNPS1 is a versatile splicing regulator for a wide variety of alternatively spliced genes and regulates alternative splicing both negatively and positively through interaction with associated factors in vivo [32]. Although their exact mechanism of function remains unclear, several ATP-dependent RNA helicases of the DEAD-box family have been described to be involved in transcription, pre-mRNA splicing, ribosome biogenesis, nuclear export, resolution of inhibitory mRNA secondary structures and translation initiation, RNA degradation and even organelle

Table 1 Dose-dependent upregulated protein spots in response to Pb (II)

| KOG Class | Protein ID | Putative Function | Treatment/ Control Ratio | | | Subcellular Locations* | Multiple Spots |
|---------------------------------------------------------------------|------------|--------------------------------------------------------------------------------------|-----------------------------|------------|-------------|------------------------|----------------|
| | | | 25 μ M | 50 μ M | 100 μ M | | |
| Amino acid transport and metabolism | 139320a | Isocitrate dehydrogenase, alpha subunit | 5,14 | 5,02 | 4,06 | Mit | 2 |
| Carbohydrate transport and metabolism | 134115 | UDP-glucose pyrophosphorylase | 3,50 | 3,05 | 5,22 | Cyto_nuc | - |
| | 135471 | Septin family protein (P-loop GTPase) | New | New | New | M | - |
| | 132198d | Glyceraldehyde 3-phosphate dehydrogenase | - | New | New | C | 4 |
| Cytoskeleton | 139298a | Actin and related proteins | 1,38 | 11,01 | 1,43 | Cysk | 4 |
| Energy production and conversion | 137299d | F0F1-type ATP synthase, alpha subunit | New | New | New | Mit | 4 |
| General function prediction only | 134635 | mRNA splicing factor | 2,29 | 4,62 | 2,26 | C | - |
| Intracellular trafficking, secretion, and vesicular transport | 123314 | GTPase Ran/TC4/GSP1 (nuclear protein transport pathway), small G protein superfamily | New | New | New | C | - |
| Lipid transport and metabolism | 1819 | Short-chain acyl-CoA dehydrogenase | 3,60 | 3,70 | 4,30 | C | - |
| Posttranslational modification, protein turnover, chaperones | 8527 | Thioredoxin reductase | 2,54 | 3,49 | 4,46 | Mit | - |
| RNA processing and modification | 123005a | Polyadenylate-binding protein (RRM superfamily) | New | New | New | Cysk | 2 |
| | 126823 | ATP-dependent RNA helicase | 0,97 | 7,89 | 1,86 | Nuc | - |
| Secondary metabolites biosynthesis, transport and catabolism | 4796c | Alcohol dehydrogenase, class V | - | New | New | C | 4 |
| Signal transduction mechanisms | 10373 | G protein beta subunit-like protein | 3,10 | 3,30 | 4,60 | C | - |

*C; cytoplasmic, Ext; extracellular, Mit; mitochondrial, Nuc; Nuclear, Cysk; cytoskeleton, Cyto_nuc; cytoplasmic_nuclear.

gene expression [33]. This enzyme was shown to be involved in adaptive response to oxidative stress in *Clostridium perfringens* [34], heat shock response in *Aspergillus fumigatus* [35] and various kinds of stress in many plants, including salt response in barley [36], pathogen infection and oxidative stress in transgenic *Arabidopsis* [37] and salt stress in the halophyte *Apocynum venetum* [38]. Taken together, up-regulation of above-mentioned proteins in *P. chrysosporium* in response to Pb stress strongly suggests that these proteins might act in concert to mediate transcriptional and post-transcriptional regulations in a direction to overcome Pb toxicity.

An isoform of glyceraldehyde-3-phosphate dehydrogenase (GAPDH) (Figure 1g) was also among newly induced proteins in response to Pb. This extremely abundant glycolytic enzyme has multiple and unrelated functions. GAPDH expression was shown to increase in various organisms during apoptosis induced by a variety of stress factors. Potential role of its nuclear translocation in apoptosis and oxidative stress was proposed to be related with its activity as a DNA repair enzyme or as a nuclear carrier for pro-apoptotic molecules [39]. Different isoforms of GAPDH responding differently to H₂O₂ stress have been shown in *S. pombe* [40] and in budding yeast [41,42]. In *S. pombe*, peroxide stress signals are transmitted from the Mak2/3 sensor kinases to the Mpr1 histidine-containing phosphotransfer (HPT) protein and finally to the Mcs4 response regulator,

which activates a MAP kinase cascade. Morigasaki *et al.* [43] recently showed that GAPDH plays an essential role in the phosphorelay signaling by physically associating with the Mcs4 response regulator and stress-responsive MAP kinase kinase kinases (MAPKKKs), where its redox-sensitive cysteine residue which is transiently oxidized in response to H₂O₂ stress may enhance its association. In *Arabidopsis thaliana*, the steady-state mRNA level of the cytosolic GAPDH increased when plants were transferred from normal growth condition to heat-shock, anaerobiosis, or increased sucrose supply [44] and the enzyme was recently shown to suppress heat shock-induced H₂O₂ production and cell death [45].

The induction of redox enzyme thioredoxin reductase was quite expected given its role in the regeneration of reduced thioredoxin (Figure 2a). UDP-glucose is not only a necessary metabolite for cell wall biogenesis, but it is involved in the synthesis of the carbohydrate moiety of glycolipids and glycoproteins [46]. UDP-glucose pyrophosphorylase (Figure 2b) which is under the control of stationary phase transcription factor SigmaB in *B. subtilis* [47] was shown as a novel salt stress-responsive protein in rice [48]. G-protein-linked pathways evolved to allow responses to extracellular agonists in eukaryotic cells include those for nutrient sensing, pheromone response and mating, and pathogenesis in fungi [49], however except for the induction of Cd-induced G-protein β subunit in fission

Table 2 Dose-dependent downregulated protein spots in response to Pb (II)

| KOG Class | Protein ID | Putative Function | Treatment/ Control Ratio | | | Subcellular location* | Multiple spots |
|---------------------------------------------------------------------|----------------|----------------------------------------------------------------------|-----------------------------|----------|-----------|--------------------------|-------------------|
| | | | 25 uM | 50 uM | 100 uM | | |
| Amino acid transport and metabolism | 138721a | Glutamate/leucine/phenylalanine/valine dehydrogenases | 0,50 | 0,22 | 1,32 | C | 2 |
| | 139320b | Isocitrate dehydrogenase, alpha subunit | 0,12 | 0,02 | 0,01 | Mit | 3 |
| | 139663&138887a | Methionine synthase II (cobalamin-independent)-/-aconitate hydratase | 0,24 | 0,09 | 0,08 | C | 4 |
| | 139663&138887b | Methionine synthase II (cobalamin-independent)-/-aconitate hydratase | 0,30 | 0,06 | 0,15 | C | 4 |
| | 139663&138887c | Methionine synthase II (cobalamin-independent)-/-aconitate hydratase | 0,14 | 0,00 | 0,05 | C | 4 |
| | 139663&138887d | Methionine synthase II (cobalamin-independent)-/-aconitate hydratase | 0,26 | 0,18 | 0,06 | C | 4 |
| | 139663d | Methionine synthase II (cobalamin-independent) | 0,15 | 0,17 | 0,66 | C | 6 |
| | 139663e | Methionine synthase II (cobalamin-independent) | 0,20 | 0,20 | 0,60 | C | 6 |
| | 139663f | Methionine synthase II (cobalamin-independent) | 0,07 | 0,02 | 0,01 | C | 6 |
| Cytoskeleton | 139298c | Actin and related proteins | 0,59 | 0,16 | 1,15 | Cysk | 4 |
| Energy production and conversion | 1056 | Zinc-binding oxidoreductase | 0,14 | 0,09 | 0,25 | C | - |
| | 123932 | Fumarate reductase, flavoprotein subunit | 0,14 | 0,07 | 0,34 | C | - |
| Lipid transport and metabolism | 10015 | Acetyl-CoA acetyltransferase | 0,66 | 0,24 | 0,88 | C | 2 |
| NA | 8290 | hypothetical protein | 0,40 | 0,22 | 0,28 | C | - |
| | 140431b | hypothetical protein | 0,04 | 0,03 | 0,01 | C | 4 |
| | 140431c | hypothetical protein | 0,19 | 0,03 | 0,07 | C | 4 |
| | 140431d | hypothetical protein | 0,52 | 0,22 | 0,11 | C | 4 |
| Posttranslational modification, protein turnover, chaperones | 122440 | Molecular chaperones HSP70/HSC70, HSP70 superfamily | 0,53 | 0,09 | 0,84 | C | 6 |
| | 131983b | Molecular chaperones GRP78/BiP/KAR2, HSP70 superfamily | 0,71 | 0,11 | 0,19 | Ext | 2 |
| Secondary metabolites biosynthesis, transport and catabolism | 8565 | Hydroxysteroid 17-beta dehydrogenase 11 | 0,45 | 0,29 | 1,24 | Mit | - |
| Signal transduction mechanisms | 10895 | Glycosylphosphatidylinositol-specific phospholipase C | 0,11 | 0,30 | 1,19 | Cyto_ nucl | - |

*C; cytoplasmic, Ext; extracellular, Mit; mitochondrial, Nuc; Nuclear, Cysk; cytoskeleton, Cyto_nucl; cytoplasmic_nuclear.

yeast [50], there is scarcity of literature reports on the roles of these pathways in stress response. On the other hand, evidence is accumulating for heterotrimeric G protein signaling in stress-associated physiological processes in plants. In mature leaves, G proteins transmit signals to molecules, including small GTPases, ion channels, and phospholipases which are the effectors in the responses to various stress conditions, including pathogens, ozone treatment and water deficit [51]. *Arabidopsis thaliana* with null mutation in the gene encoding β subunit of heterotrimeric G protein was more sensitive to O₃ damage than wild-type plants [52]. The newly induced G protein beta subunit-like protein (Figure 2c) demonstrated in our work provided evidence for heterotrimeric G protein signaling under Pb(II) stress in *P. chrysosporium*.

Zn-containing alcohol dehydrogenase (Class V), with accession number 4796, gives the highest homology to *Cryptococcus neoformans* mannitol-1-P-dehydrogenase, a zinc-containing long chain alcohol/polyol dehydrogenase accumulating mannitol as an intracellular osmolyte and stress protectant [53]. Members of this family metabolize a wide variety of substrates, including ethanol, retinol, other aliphatic alcohols, hydroxysteroids, and lipid peroxidation products. The enzyme seemed to undertake one of the major functions to counteract with Cu toxicity [25] and Pb toxicity, as revealed by the present study (Figure 2d).

As to the down-regulated proteins, the most drastic effect was the decrease in abundance of certain isoforms of cobalamin-independent methionine synthase. Protein glutathionylation has been increasingly recognized as an




Figure 1 Dose-dependent upregulated protein spots.

important mode of regulation in eukaryotes and glutathionylation of key proteins involved in protein synthesis leads to inhibition of translation. MetE is so far one of the few proteins in bacteria known to be the most sensitive to oxidative damage. When stressed by an

oxidant, glutathionylation of the active site of MetE protects the enzyme from permanent oxidative damage as shown in *E. coli* [54]. Thus, by turning off MetE in the face of oxidative stress, protein synthesis can be slowed or stopped, freeing cellular resources to be used




Figure 2 Dose-dependent upregulated protein spots.

elsewhere. Also, diminished expression of genes encoding enzymes in methionine and cysteine biosynthesis can be interpreted as a mechanism of securing sulfur for production of proteins involved in reductive detoxification, such as thioredoxin and glutathione [55].

Analysis of the proteome dynamics of mid-exponential phase cells of *P. chrysosporium* subjected acute lead exposure identified a total of 88 differentially expressed protein spots, 23 up-regulated and 67 down-regulated

(Table 3 and 4). Regarding the up-regulated ones, seven proteins, namely aldehyde dehydrogenase, alcohol dehydrogenase class V, 60S acidic ribosomal protein P0 and particular isoforms of a putative protein, glyoxylate as well as certain isoforms of NAD-dependent malate dehydrogenase and glyceraldehyde-3-phosphate dehydrogenase were newly-induced upon lead exposure. For 17 out of a total of 23 up-regulated ones, the most significant increase was detected after 1 h exposure.

Table 3 Time-dependent upregulated protein spots in response to Pb

| KOG class | Protein ID | Putative Function | Treatment/Control ratio | | | | Subcellular location* | Multiple spots |
|---------------------------------------------------------------------|------------|---------------------------------------------------------------------------------------------------|-------------------------|------|-------|-------|-----------------------|----------------|
| | | | 1h | 2h | 4h | 8h | | |
| Amino acid transport and metabolism | 130118a | Glutamine synthetase | 1,50 | 3,61 | 3,88 | 1,08 | C | 4 |
| | 139663 | Methionine synthase II (cobalamin-independent) | 3,51 | 0,58 | 2,29 | 1,58 | C | 3 |
| Carbohydrate transport and metabolism | 137211 | Dihydroxyacetone kinase/glycerone kinase | 9,99 | 3,50 | 3,37 | 3,93 | C | - |
| | 132198c | Glyceraldehyde 3-phosphate dehydrogenase | 7,40 | 7,00 | 5,20 | 4,80 | C | 5 |
| | 132198d | Glyceraldehyde-3-phosphate dehydrogenase | new | new | new | new | C | 5 |
| | 132198e | Glyceraldehyde 3-phosphate dehydrogenase | 1,29 | 4,90 | 4,71 | 6,09 | C | 5 |
| Energy production and conversion | 133289 | Aldehyde dehydrogenase | 4,36 | 3,09 | 2,30 | 3,34 | C | - |
| | 134389 | NADH-dependent flavin oxidoreductase/12-oxophytodienoate reductase | 4,74 | 0,82 | 1,29 | 1,33 | C | - |
| | 138693 | Aldehyde dehydrogenase | new | new | new | new | C | - |
| | 123932a | Fumarate reductase, flavoprotein subunit | 3,54 | 2,39 | 1,37 | 1,87 | C | 2 |
| | 129245a | NAD-dependent malate dehydrogenase | new | new | new | - | Mit | 5 |
| | 133757c | Glyoxylate/hydroxypyruvate reductase (D-isomer-specific 2-hydroxy acid dehydrogenase superfamily) | new | new | new | new | C | 3 |
| | 135576b | Inorganic pyrophosphatase/Nucleosome remodeling factor, subunit NURF38 | 1,24 | 1,34 | 8,02 | 0,84 | Nuc | 4 |
| General function prediction only | 10307 | 1,4-benzoquinone reductase-like; Trp repressor binding protein-like/protoplast-secreted protein | 0,44 | 0,87 | 13,14 | 4,72 | C | - |
| | 132851 | Glyoxylase | 6,16 | 1,99 | 4,02 | 2,30 | C | - |
| NA | 140431a | Putative Protein | - | new | new | new | C | 4 |
| | 140431b | Putative Protein | 19,30 | 9,80 | 19,40 | 22,90 | C | 4 |
| | 140431c | Putative Protein | 4,29 | 2,2 | 0,75 | 2,3 | C | 4 |
| Posttranslational modification, protein turnover, chaperones | 1827 | 20S proteasome, regulatory subunit alpha type PSMA6/SCL1 | 4,60 | 2,20 | 2,10 | 2,70 | Mit | 2 |
| RNA processing and modification | 8607 | Splicing factor RNPS1, SR protein superfamily | 3,32 | 3,12 | 0,95 | 3,5 | Cyt_Nuc | - |
| Secondary metabolites biosynthesis, transport and catabolism | 4796d | Alcohol dehydrogenase, class V | new | new | new | new | C | 4 |
| | 127894 | Zinc-containing alcohol dehydrogenase superfamily | 5,21 | 3,22 | 5,32 | 6,55 | C | - |
| Translation, ribosomal structure and biogenesis | 130203 | 60S acidic ribosomal protein P0 | new | new | new | new | C | - |

*C; cytoplasmic, Ext; extracellular, Mit; mitochondrial, Nuc; Nuclear, Cysk; cytoskeleton, Cyto_nuc; cytoplasmic_nuclear.

Among the up-regulated spots identified, “energy production and conversion” proteins constituted the major KOG class as was found by Garcia-Leiro [56] in an analysis of oxidative stress response of *Kluyveromyces lactis*.

Although Pb (II) is not a redox active metal ion, it is known to cause oxidative stress resulting in increased production of reactive oxygen species (ROS) inducing lipid peroxidation [57,58]. There were several newly induced/strongly up-regulated proteins involved in amelioration of lipid peroxidation products, namely zinc-containing alcohol dehydrogenase, glyceraldehyde-3-phosphate dehydrogenase, glyoxylate/hydroxypyruvate reductase (Figure 3a)

and two different aldehyde dehydrogenases (ALDH; Figure 3c) detected after 1 h exposure. NAD-dependent malate dehydrogenase (MDH; Figure 3b), on the other hand, is a very well known up-regulated component of different stress conditions in various organisms. It provides protection against oxidative damage caused by Zn in *E. coli* through the action of oxaloacetate [59]. In the fission yeast, MDH was one of the component core environmental stress response (CESR) as determined by transcriptional profiling, proteomic and metabolomic analysis [40,50]. The other components of CESR included zinc-binding dehydrogenases, serine/threonine protein kinase,

Table 4 Time-dependent downregulated protein spots in response to Pb

| KOG Class | Protein ID | Putative Function | Treatment/control ratio | | | | Subcellular Location* | Multiple spots |
|---------------------------------------------------------------------|------------|---------------------------------------------------------------------------------------------------|-------------------------|------|------|------|-----------------------|----------------|
| | | | 1h | 2h | 4h | 8h | | |
| Amino acid transport and metabolism | 137931 | Glycine/serine hydroxymethyltransferase | 0,56 | 0,27 | 0,19 | 0,42 | C | 3 |
| | 1542 | 3-isopropylmalate dehydrogenase | 0,39 | 0,42 | 0,26 | 0,39 | C | - |
| | 10011 | 3-isopropylmalate dehydratase (aconitase superfamily) | 0,15 | 0,55 | 0,14 | 0,18 | Cysk | - |
| | 125842 | Lysine-ketoglutarate reductase/saccharopine dehydrogenase | 0,43 | 0,17 | 0,08 | 0,33 | C | - |
| | 131837 | Oxoprolinase | 0,56 | 0,32 | 0,31 | 0,24 | C | - |
| | 134775 | Glutamine amidotransferase/cyclase | 0,53 | 0,67 | 0,59 | 0,26 | C | - |
| | 138721 | Glutamate/leucine/phenylalanine/valine dehydrogenases | 0,84 | 0,35 | 0,23 | 0,34 | C | 2 |
| | 139663 | Methionine synthase II (cobalamin-independent) | 0,25 | 0,36 | 0,05 | 0,17 | C | 5 |
| Carbohydrate transport and metabolism | 8743 | Inositol monophosphatase | 0,12 | 0,04 | 0,06 | 0,39 | C | - |
| | 10433 | Transketolase | 0,92 | 0,89 | 0,89 | 0,18 | C | - |
| | 133884 | Glycolipid transfer protein | 0,04 | 1,12 | 0,39 | 0,27 | C | - |
| | 137623 | Phosphoglucomutase | 1,54 | 0,35 | 0,17 | 1,07 | C | - |
| | 122435c | 3-phosphoglycerate kinase | 0,95 | 0,31 | 0,24 | 1,59 | C | 3 |
| | 3052a | Mannose-6-phosphate isomerase, type II | 0,44 | 0,54 | 0,60 | 0,29 | Cysk | 2 |
| | 3052b | Mannose-6-phosphate isomerase, type II | 0,91 | 0,44 | 0,18 | 0,32 | Cysk | 2 |
| Coenzyme transport and metabolism | 10308 | S-adenosylhomocysteine hydrolase | 0,67 | 0,64 | 0,30 | 0,56 | C | - |
| Cytoskeleton | 139298 | Actin and related proteins | 0,30 | 0,37 | 0,07 | 0,11 | Cysk | 4 |
| Defense mechanism | 10742 | N-6 Adenine-specific DNA methylase | 1,05 | 0,6 | 0,32 | 0,19 | Mit | - |
| Energy production and conversion | 912 | Kynurenine 3-monooxygenase and related flavoprotein monooxygenases | 0,78 | 0,41 | 0,19 | 0,20 | Ext | - |
| | 1350 | Aldehyde dehydrogenase | 0,45 | 0,34 | 0,36 | 0,29 | C | - |
| | 132162 | NAD-dependent malate dehydrogenase | 0,00 | 0,37 | 0,55 | 0,51 | C | - |
| | 132918 | Sulfide:quinone oxidoreductase/flavo-binding protein | 0,85 | 0,49 | 0,22 | 0,55 | Mit | - |
| | 134368 | Vacuolar H ⁺ -ATPase V1 sector, subunit E | 0,29 | 0,67 | 0,32 | 1,98 | Nuc | - |
| | 135659 | NADP ⁺ -dependent malic enzyme | 1,13 | 0,72 | 0,18 | 1,18 | Mit | - |
| | 140211 | Glyoxylate/hydroxypyruvate reductase (D-isomer-specific 2-hydroxy acid dehydrogenase superfamily) | 1,51 | 0,51 | 0,29 | 0,33 | C | - |
| | 123932b | Fumarate reductase, flavoprotein subunit | 1,21 | 0,85 | 0,60 | 0,10 | C | 2 |
| | 131257a | NADH-ubiquinone oxidoreductase, NDUFS1/75 kDa subunit | 0,91 | 0,88 | 0,42 | 0,21 | C | 3 |
| | 131879a | Dihydroliipoamide dehydrogenase | 0,62 | 0,52 | 0,26 | 0,80 | C | 2 |
| | 133924a | Aldehyde dehydrogenase | 0,65 | 0,49 | 0,26 | 0,61 | C | 2 |
| | 133924b | Aldehyde dehydrogenase | 0,31 | 0,48 | 0,28 | 0,55 | C | 2 |
| | 563a | Pyruvate dehydrogenase E1, alpha subunit | 1,33 | 0,28 | 0,27 | 0,28 | Mit | 2 |
| General function prediction only | 3442 | Predicted NAD-dependent oxidoreductase | 0,30 | 0,22 | 0,54 | 0,50 | C | - |
| | 132767b | Serine/threonine protein kinase, active site | 0,55 | 0,85 | 0,55 | 0,27 | Mit | 2 |
| Lipid transport and metabolism | 511 | Enoyl-CoA hydratase | 1,04 | 0,74 | 0,14 | 0,96 | C | - |
| | 10355b | Mevalonate pyrophosphate decarboxylase | 0,73 | 0,36 | 0,24 | 0,48 | Mit | 2 |
| Posttranslational modification, protein turnover, chaperones | 361 | 26S proteasome regulatory complex, ATPase RPT3 | 0,25 | 0,17 | 0,06 | 0,05 | Nuc | - |
| | 1324 | Molecular co-chaperone STI1 | 1,02 | 0,68 | 0,01 | 1,11 | C | - |
| | 1846 | Chaperonin complex component, TCP-1 beta subunit (CCT2) | 0,79 | 0,85 | 0,46 | 0,28 | C | - |
| | 5061 | Dipeptidyl aminopeptidase | 1,06 | 1,07 | 0,99 | 0,29 | Ext | 2 |
| | 8527 | Thioredoxin reductase | 0,27 | 0,45 | 0,22 | 0,31 | Mit | - |

Table 4 Time-dependent downregulated protein spots in response to Pb (Continued)

| | | | | | | | | |
|---------------------------------------------------------------------|--------|-----------------------------------------------------------------------------------------------------------|------|------|------|------|-----|---|
| | 130274 | Glutathione peroxidase | 0,33 | 0,77 | 1,10 | 0,19 | C | - |
| | 131571 | Protein disulfide isomerase (prolyl 4-hydroxylase beta subunit) | 0,91 | 0,08 | 0,74 | 0,06 | Ext | - |
| | 133185 | 26S proteasome regulatory complex, ATPase RPT1 | 0,31 | 0,34 | 0,07 | 0,04 | Nuc | - |
| | 133717 | 20S proteasome, regulatory subunit alpha type PSMA4/PRE9 | 0,30 | 0,62 | 0,82 | 0,73 | Mit | - |
| | 134073 | Chaperonin complex component, TCP-1 zeta subunit (CCT6) | 0,70 | 0,42 | 0,14 | 0,10 | C | - |
| | 139500 | Multifunctional chaperone (14-3-3 family) | 0,27 | 0,38 | 0,20 | 0,47 | Nuc | - |
| | 10340a | HSP70(putative ortholog to <i>S. cerevisiae</i> Heat shock protein homolog SSE1 (Chaperone protein MSI3)) | 0,66 | 0,54 | 0,06 | 0,64 | C | 2 |
| | 10340b | HSP70(putative ortholog to <i>S. cerevisiae</i> Heat shock protein homolog SSE1 (Chaperone protein MSI3)) | 0,34 | 0,15 | 0,20 | 0,26 | C | 2 |
| Secondary metabolites biosynthesis, transport and catabolism | 133231 | Predicted dehydrogenase | 0,26 | 0,53 | 0,16 | 0,10 | C | - |
| | 4796b | Alcohol dehydrogenase, class V | 0,61 | 0,31 | 0,17 | 0,07 | C | 4 |
| Translation, ribosomal structure and biogenesis | 3216 | Prolyl-tRNA synthetase | 0,80 | 0,60 | 0,51 | 0,12 | C | - |
| | 6570 | Mitochondrial translation elongation factor Tu | 0,61 | 0,59 | 0,16 | 0,76 | Mit | - |
| | 10819 | Elongation factor 2 | 1,27 | 0,65 | 0,15 | 0,14 | C | - |

*C; cytoplasmic, Ext; extracellular, Mit; mitochondrial, Nuc; Nuclear, Cysk; cytoskeleton, Cyto_nucl; cytoplasmic_nuclear.

G-protein beta subunit, quinone oxidoreductase, flavin oxidoreductase, protein with RNA recognition motif and short chain dehydrogenase, all were found to be up-regulated in *P. chrysosporium* cells in our former [25] and present study. Hot pepper transcriptome profiling under cold stress [60] proteomic analysis of Cd response in marine alga *Nannochloropsis oculata* [61] DNA microarray and quantitative RT-PCR analyses in *Corynebacterium glutamicum* under oxygen deprivation [62], analysis of a metabolic network in *Pseudomonas fluorescens* exposed to oxidative stress [63] and transcriptomic analysis of Al stress in roots of *Arabidopsis thaliana* [64] were among other studies consistently detecting MDH involvement. As discussed by the latter authors, the burst of ROS generated by Al had to enhance the generation of NADPH to maintain a high ratio of reduced antioxidants.

In the present study, two different ALDHs with the protein IDs of 133289 and 138693 were found to display a biphasic up-regulation (at 1 h and 8 h) upon Pb treatment (Figure 3c). ALDH superfamily enzymes and their pathophysiological significance was recently reviewed by Marchitti *et al.* [65]. Induction of ALDH in response to oxidative stress was demonstrated also in bacteria, e.g. *Pseudomonas aeruginosa* [66] and yeast [67]. As to the plants, overexpression of ALDH3 genes in *A. thaliana* confers tolerance to various abiotic stress conditions and protects plants against lipid peroxidation [68,69].

The expression of remarkably many genes encoding proteins and enzymes involved in defense from oxidative damage and redox metabolism is stimulated under a range of stress conditions. Examples of such genes

include catalase, thioredoxin, glutaredoxin, genes encoding enzymes possibly involved in detoxification such as glyoxylase and dihydroxyacetone kinase, and numerous oxidoreductases that may be involved in metabolism of oxidized biomolecules or in adjusting redox metabolism to provide sufficient NADPH for detoxification. Dihydroxyacetone kinase/glycerone kinase (DAK) was induced in *P. chrysosporium* 1 h after Pb administration (Figure 3d). Its relation with salt stress [70], heat stress [71], osmotic stress [72], oxidative stress [73], starvation [74] and cadmium stress [21] was well documented. Glyoxylase system detoxifying glyoxal, methylglyoxal and other physiological alpha-oxoaldehydes formed by lipid peroxidation is a component of stress response, as shown for yeasts [75,76], various plants [77-79], a parasitic nematode [80] and in a basidiomycete, as shown by our study (Figure 3e).

Inorganic phosphatase constitutes NURF-38 subunit of the ATP-dependent Nucleosome Remodeling Factor complex (NURF) which was initially identified in *Drosophila* and then in yeast and vertebrates [81,82]. The complex, when targeted onto chromatin, affects major DNA-dependent processes including transcription, DNA repair and recombination. Thus, our finding that inorganic pyrophosphatase/nucleosome remodeling factor NURF38 is induced by a factor of 8 upon 4 h lead exposure pointed to its role in responding to metal toxicity in *P. chrysosporium* (Figure 3f). The 60S acidic ribosomal protein P0 was one of the newly-induced component of Pb (II)-stressed proteome. This protein plays an essential role by docking and forming the tip of the




Figure 3 Time-dependent upregulated protein spots.

whole eukaryotic ribosomal “stalk” complex [83]. Unlike the typical ribosomal proteins, P0 appears to have multiple functions in the cell. By overexpressing *Drosophila* P0 in *E. coli*, Yacoub *et al.* [84] demonstrated that P0 contained 5' APE activity distinct from the 3' AP lyase

activity associated with *Drosophila* rpS3 and also exhibited nuclease activity against both double and single-stranded DNA. The authors also reported that the protein is located in both nucleus and ribosomes. Recently, this protein was shown to be upregulated in

Ras-transformed NIH3T3 cells [85] and Jurkat cells during heat stress-induced apoptosis [86]. Additionally, when the *S. pombe* cells challenged with oxidative stress, 60S ribosomal protein P0 was among differentially-expressed ones though its induction ratio was only 1.6 [40] (Figure 3f).

Being consistent with our former report on Cu and Cd response of the organism [25], the downregulated proteins included some redox enzymes and certain molecular chaperons like HSP70, protein disulfide isomerase and chaperonin complex components.

Bacteria and fungi are among the first components of the biota in ecosystems affected by toxic pollutants including heavy metals. Their relatively small size and simplicity make them particularly attractive models for environmental proteomics [18]. The data obtained from the analysis of proteomes of such organisms help to gain insight into underlying mechanisms of toxicity which is of great value from the basic sciences point-view. Besides, the subtle changes detected in the level of individual proteins in response to environmental stressors lead to the discovery of biomarkers of exposure and also provide an opportunity to genetically engineer such microbes to express higher levels of specific proteins (e. g. DNA repair proteins), thereby conferring higher tolerance to increasing concentrations of heavy metals and potentiate bioremediation [87,88]. Our studies are under way to obtain complete coverage of the lead-induced stress proteome of *P. chrysosporium*, hence identifying all possible ecotoxicological biomarkers as well as targets for improved lead bioaccumulation.

The apoptotic machinery in fungi was recently reviewed by Sharon *et al.* [89]. Among fungi, apoptosis has only been studied in detail in *S. cerevisiae*. Filamentous fungal species are poorly analyzed for functions of the apoptosis-related genes and although homologs of some apoptotic genes could be identified in fungal genomes analyzed, to date only a few genes have been functionally analyzed. A better understanding of fungal apoptotic networks for identification of paralogs and putative homologs of apoptosis as compared to those of high eukaryotes is required which is possible through findings from more fungal species. As reported by Lorin *et al.* [90] for the filamentous fungus *Podospora anserina*, compensatory induction of an alternative oxidase (AOX) provides a decreased production of ROS and a striking increase in lifespan. In the present research, the elements like actin known to induce hyperactivation of the Ras signaling pathway, septin family protein (P-loop GTPase) functioning in many processes including apoptosis, Ras GTPase of acceleration of programmed cell death are identified for the first time in a multicellular fungus and expected to contribute to the knowledge on stress-induced apoptosis in fungi. When exposed to lead

stress, the elements of stress tolerance and initiation of apoptotic cell death counteracted while the cells of *P. chrysosporium* kept on growing. On the other hand, our findings on induction/upregulation of certain catabolic enzymes and mitochondrial F0F1-type ATP synthase might provide support for an intrinsic and mitochondrial nature of the apoptotic response in fungi and fit to the generally accepted view that apoptosis requires energy as it is a highly regulated process involving a number of ATP-dependent steps [91].

Conclusion

The present study draws attention particularly to the up-regulated elements of apoptosis, DNA repair, post-transcriptional regulation and heterotrimeric G protein signaling as shown for the first time for a metal-stressed basidiomycete. DNA-binding response regulator(s) mediating stress response in *P. chrysosporium*, like Yap1p TF and Yap2p TF of *S. cerevisiae* [92], Sty1p-activated Atf1p and Pap1p TF of the fission yeast [50,93] and Cap1p TF of *Candida albicans* [76] remains to be identified through further analysis.

Materials and methods

Culture conditions

P. chrysosporium (ATTC 24725) spores were separated from Sabaroud Dextrose agar slant surfaces by scraping, homogenized and suspended in sterile distilled water. A spore suspension was prepared to contain 2.5×10^6 spores.mL⁻¹ at an absorbance of 0.5 at 650 nm using a Shimadzu UV-1208 spectrophotometer. The suspension was then transferred into a 250 mL Erlenmeyer flasks each containing 150 ml of the growth medium described by Prouty [94] which was composed, in g.L⁻¹, of glucose, 10; KH₂PO₄, 2; MgSO₄, 0.5; CaCl, 0.1; NH₄Cl, 0.12 and thiamine, 0.001 and adjusted to a pH of 4.5. The cultures were incubated for 40 h at 200 rpm in a rotary shaker at 35°C. When the growth was terminated at 40th h, the cultures were still in exponential growth.

For proteomic characterization of the lead response, the cells of *P. chrysosporium* were grown in minimal media containing different levels of lead (25, 50 and 100 µM Pb(NO₃)₂, respectively) for 40 h till they reached mid-exponential phase and they were harvested by filtration, washed twice with distilled water and stored at -20° C for the extraction of proteins. As a parallel setup, *P. chrysosporium* cells were grown on minimal media for 40 h till they reached mid-exponential phase and then subjected to 50 µM lead for 1, 2, 4 and 8 h to investigate the effect of duration of temporal Pb (II) exposure on *P. chrysosporium* proteome. For both cases, a control sample, without lead exposure, was harvested at 40th h of the growth.

Protein Extraction

After crushing the cells with liquid nitrogen, TCA-acetone extraction was performed as in Damerval *et al.* [95]. After breaking a 500 mg of harvested mycelium in liquid nitrogen, 5 mL of 10% trichloroacetic acid in acetone containing 0.07% β -mercaptoethanol was added and vortexed, then incubated at -20°C for 45 min and centrifuged at 15 000 *g* for 15 min. The supernatant was decanted and the pellet was resuspended in 5 mL of acetone containing 0.07% β -mercaptoethanol which was then incubated at -20°C for 1 h (mixed every 15 min intervals by vortexing) and recentrifuged. The supernatant was discarded, the remaining pellet was vacuum-dried and stored as a powder at -20°C . The modified Bradford assay [96] was used to determine protein concentrations.

2DE

2D gels of the harvested cells were run in duplicates for control and each treatment. Isoelectric focusing was performed in 18 cm linear IPG-strips (pH range 3-10, Biorad, Hercules, CA, USA). IPG strips were passively rehydrated by applying 300 μl of rehydration buffer containing 8 M urea, 2 M thiourea, 1% w/v CHAPS, 20 mM DTT and 0.5% v/v ampholyte 3-10 with 300 μg protein sample for 16 h. Isoelectric focusing was performed with the Protean IEF Cell unit (Biorad, Hercules, CA, USA) employing a total of 80 000 Vh. After consecutive equilibration of the gels in solutions containing DTT and iodoacetamide as suggested by Görg *et al.* [97], the separation in the second dimension was done in polyacrylamide gels of 12.5% T and 2.6% C on the Biorad Protean Xii electrophoresis system (Biorad, Hercules, CA, USA) by applying 2 W per gel. Gels were stained with colloidal Coomassie blue [98].

Image analysis

Coomassie stained gels were digitized by using an HP scanner. Spot pattern analyses were accomplished by using the 2D image analysis software Delta2D version 3.3 (Decodon, Germany). Of the proteins found differentially expressed in Pb (II)-exposed cells, only those showing at least 3 fold difference in abundance were selected and subjected to MALDI-TOF analysis. To correct the quantitative variability, the spot volumes were normalized as a percentage of the total volume in all of the spots in the gel. SDs of the spot intensities from the two replicates were in the range of 20%.

Protein identification

The identifications were accomplished by mass spectrometry according to established protocols. Briefly, protein spots were excised from stained 2D gels, destained and digested with trypsin (Promega, Madison, WI, USA) and

for extraction of peptides, the gel pieces were covered with 60 μl 0.1% trifluoroacetic acid in 50% CH_3CN and incubated for 30min at 40°C . Peptide solutions were mixed with an equal volume of saturated α -cyano-3-hydroxycinnamic acid solution in 50% acetonitrile-0.1% trifluoroacetic acid (v/v) and applied to a sample plate for MALDI-TOF-MS. Mass analyses were carried out on the Proteome-Analyzer 4700 (Applied Biosystems, Foster City, CA, USA). The three most abundant peptides in each MS spectrum were chosen for MS/MS experiment. The resulting sequence data were included for the database search to increase the reliability of protein identification. Mass accuracy was usually in the range between 10 and 30 ppm.

Database searches

Amino acid sequences for *P. chrysosporium* proteins were obtained from organism's genome project [Joint Genome Institutes (JGI)] web site [99]. PMF and MS/MS data was searched in the *P. chrysosporium* data with the aid of MASCOT software [100]. The searches considered oxidation of methionine and modification of cysteine by carbamidomethylation as well as partial cleavage leaving one internal cleavage site. Of the results given by the MASCOT software, those having a probability score value higher than 53 were considered for successful protein identification. To find out putative functions, protein accession numbers of the identified spots were searched in the JGI website for *P. chrysosporium*. For the identified proteins, the functional classification was made by consulting to the functional categories list contained in the same website.

Protein subcellular localization predictions were obtained from WoLF PSORT web server [101].

Author details

¹Department of Biological Sciences, Middle East Technical University, Ankara, Turkey. ²Department of Biology, Erciyes University, Kayseri, Turkey. ³Institut für Mikrobiologie, Ernst-Moritz Arndt-Universität Greifswald, Greifswald, Germany.

Authors' contributions

VY carried out the 2DE experiments for time dependent Pb exposure studies. SÖ carried out the 2DE experiments for dose dependent Pb exposure studies. DB carried out the mass spectrometry analyses and protein identifications. KB participated in the optimization of 2DE protocol for *P. chrysosporium*. MH participated in the design of the study. GÖ conceived of the study, and participated in its design and coordination. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 13 August 2010 Accepted: 9 March 2011

Published: 9 March 2011

References

1. Piechalak A, Tomaszewska B, Baralkiewicz D, Malecka A: Accumulation and detoxification of lead ions in legumes. *Phytochemistry* 2002, **60**:153-162.

2. Sharma P, Dubey RS: **Lead toxicity in plants.** *Braz J Plant Physiol* 2005, **17**:35-52.
3. Ercal N, Gurer-Orhan H, Aykin-Burns N: **Toxic metals and oxidative stress part I: mechanisms involved in metal-induced oxidative damage.** *Curr Top Med Chem* 2001, **1**:529-539.
4. Kirk TK, Farrell RL: **Enzymatic "combustion": The microbial degradation of lignin.** *Ann Rev Microbiol* 1987, **41**:465-501.
5. Martinez AT: **Molecular biology and structure-function of lignin-degrading heme peroxidases.** *Enzyme Microb Technol* 2002, **30**:425-444.
6. Yetiş Ü, Dölek A, Dilek FB, Özcengiz G: **The removal of Pb(II) by *Phanerochaete chrysosporium*.** *Water Res* 2000, **34**:4090-4100.
7. Iqbal M, Edyvean RGJ: **Biosorption of lead, copper and zinc ions on loofa immobilized biomass of *Phanerochaete chrysosporium*.** *Miner Eng* 2004, **17**:217-223.
8. Li Q, Wu S, Liu G, Liao X, Deng X: **Simultaneous biosorption of cadmium (II) and lead (II) ions by pretreated biomass of *Phanerochaete chrysosporium*.** *Sep Purif Technol* 2004, **34**:135-142.
9. Yetiş Ü, Özcengiz G, Dilek FB, Ergen N, Erbay A, Dolek A: **Heavy metal biosorption by white-rot fungi.** *Water Sci Technol* 1998, **38**:323-330.
10. Huang DL, Zeng GM, Jiang XY, Feng CL, Yu HY, Huang GH, Liu HL: **Bioremediation of Pb-contaminated soil by incubating with *Phanerochaete chrysosporium* and straw.** *J Hazard Mater* 2006, **134**:268-276.
11. Zeng G, Huang D, Huang G, Hu T, Jiang X, Feng CL, Chen YN, Tang L, Liu HL: **Composting of lead-contaminated solid waste with inocula of white-rot fungus.** *Bioresour Technol* 2007, **98**:320-326.
12. Chen C, Wang J: **Response of *Saccharomyces cerevisiae* to lead ion stress.** *Appl Microbiol Biot* 2007, **74**:683-687.
13. Ustun GE: **Occurrence and removal of metals in urban wastewater treatment plants.** *J Hazard Mater* 2009, **172**:833-838.
14. Karvelas M, Katsoyiannis A, Samara C: **Occurrence and fate of heavy metals in the wastewater treatment process.** *Chemosphere* 2003, **53**:1201-1210.
15. Abdel-Halim SH, Shehata AM, El-Shahat MF: **Removal of lead ions from industrial waste water by different types of natural materials.** *Water Res* 2003, **37**:1678-1683.
16. Sthianopkao S, Sreesai S: **Utilization of pulp and paper industrial wastes to remove heavy metals from metal finishing wastewater.** *J Environ Manage* 2009, **90**:3283-3289.
17. Namasivayam C, Yamuna RT: **Waste biogas residual slurry as an adsorbent for the removal of Pb(II) from aqueous solution and radiator manufacturing industry wastewater.** *Bioresour Technol* 1995, **52**:125-131.
18. Nesatyy VJ, Suter MJ: **Proteomics for the analysis of environmental stress responses in organisms.** *Environ Sci Technol* 2007, **41**:6891-6900.
19. Washburn MP, Yates JR: **Analysis of the microbial proteome.** *Curr Opin Microbiol* 2000, **3**:292-297.
20. Rabilloud T, Chevallet M, Luche S, Leize-Wagner E: **Oxidative stress response: a proteomic view.** *Expert Rev Proteomics* 2005, **2**:949-956.
21. Vido K, Spector D, Lagniel G, Lopez S, Toledano MB, Labarre J: **A proteome analysis of the cadmium response in *Saccharomyces cerevisiae*.** *J Biol Chem* 2001, **276**:8469-8474.
22. Hu Y, Wang G, Chen GY, Fu X, Yao SQ: **Proteome analysis of *Saccharomyces cerevisiae* under metal stress by two-dimensional differential gel electrophoresis.** *Electrophoresis* 2003, **24**:1458-70.
23. Bae W, Chen X: **Proteomic study for the cellular responses to Cd²⁺ in *Schizosaccharomyces pombe* through amino acid-coded mass tagging and liquid chromatography tandem mass spectrometry.** *Mol Cell Proteomics* 2004, **3**:596-607.
24. Gillet S, Decottignies P, Chardonnet S, Le Maréchal P: **Cadmium response and redoxin targets in *Chlamydomonas reinhardtii*: a proteomic approach.** *Photosynth Res* 2006, **89**:201-211.
25. Özcan S, Yıldırım V, Kaya L, Albrecht D, Becher D, Hecker M, Özcengiz G: ***Phanerochaete chrysosporium* soluble proteome as a prelude for the analysis of heavy metal stress response.** *Proteomics* 2007, **7**:1249-1260.
26. Gourlay CW, Ayscough KR: **Actin-induced hyperactivation of the Ras signaling pathway leads to apoptosis in *Saccharomyces cerevisiae*.** *Mol Cell Biol* 2006, **26**:6487-501.
27. Hall PA, Russell SHE: **The pathobiology of the septin gene family.** *J Pathol* 2004, **204**:489-505.
28. Phillips AJ, Crowe JD, Ramsdale M: **Ras pathway signaling accelerates programmed cell death in the pathogenic fungus *Candida albicans*.** *Proc Natl Acad Sci USA* 2006, **103**:726-731.
29. Shyu AB, Wilkinson MF: **The double lives of shuttling mRNA binding proteins.** *Cell* 2000, **102**:135-138.
30. Chinnusamy V, Gong Z, Zhu JK: **Nuclear RNA export and its importance in abiotic stress responses of plants.** *Curr Top Microbiol Immunol* 2008, **326**:235-255.
31. Yang C, Carrier F: **The UV-inducible RNA-binding protein A18 (A18 hnRNP) plays a protective role in the genotoxic stress response.** *J Biol Chem* 2001, **276**:47277-47284.
32. Sakashita E, Tatsumi S, Werner D, Endo H, Mayeda A: **Human RNPS1 and its associated factors: a versatile alternative pre-mRNA splicing regulator in vivo.** *Mol Cell Biology* 2004, **24**:1174-1187.
33. Linder P: **Dead-box proteins: a family affair—active and passive players in RNP-remodeling.** *Nucleic Acids Res* 2006, **34**:4168-4180.
34. Briolat V, Reyssat G: **Identification of the *Clostridium perfringens* genes involved in the adaptive response to oxidative stress.** *J Bacteriol* 2002, **184**:2333-2343.
35. Albrecht D, Guthke R, Brakhage AA, Kniemeyer O: **Integrative analysis of the heat shock response in *Aspergillus fumigatus*.** *BMC Genomics* 2010, **11**:32.
36. Nakamura T, Muramoto Y, Yokota S, Ueda A, Takabe T: **Structural and transcriptional characterization of a salt-responsive gene encoding putative ATP-dependent RNA helicase in barley.** *Plant Sci* 2004, **167**:63-70.
37. Li D, Liu H, Zhang H, Wang X, Song F: **OsBIRH1, a DEAD-box RNA helicase with functions in modulating defence responses against pathogen infection and oxidative stress.** *J Exp Bot* 2008, **59**:2133-2146.
38. Liu HH, Liu J, Fan SL, Song MZ, Han XL, Liu F, Shen FF: **Molecular cloning and characterization of a salinity stress-induced gene encoding DEAD-box helicase from the halophyte *Apocynum venetum*.** *J Exp Bot* 2008, **59**:633-644.
39. Dastoor Z, Dreyer JL: **Potential role of nuclear translocation of glyceraldehyde-3-phosphate dehydrogenase in apoptosis and oxidative stress.** *J Cell Sci* 2001, **114**:1643-1453.
40. Weeks ME, Sinclair J, Butt A, Chung YL, Worthington JL, Wilkinson CRM, Griffiths J, Jones N, Waterfield MD, Timms JF: **A parallel proteomic and metabolomic analysis of the hydrogen peroxide- and Sty1p-dependent stress response in *Schizosaccharomyces pombe*.** *Proteomics* 2006, **6**:2772-2796.
41. Magherini F, Tani C, Gamberi T, Caselli A, Bianchi L, Bini L, Modesti A: **Protein expression profiles in *Saccharomyces cerevisiae* during apoptosis induced by H₂O₂.** *Proteomics* 2007, **7**:1434-1445.
42. Kim IS, Yun RS, Kwak SH, Jin IN: **The physiological role of CPR1 in *Saccharomyces cerevisiae* KNU5377 against menadione stress by proteomics.** *J Microbiol* 2007, **45**:326-332.
43. Morigasaki S, Shimada K, Ikner A, Yanagida M, Shiozaki K: **Glycolytic enzyme GAPDH promotes peroxide stress signaling through multistep phosphorelay to a MAPK cascade.** *Mol Cell* 2008, **11**:108-113.
44. Yang Y, Kwon HB, Peng HP, Shih MC: **Stress responses and metabolic regulation of glyceraldehyde-3-phosphate dehydrogenase genes in Arabidopsis.** *Plant Physiol* 1993, **101**:209-216.
45. Baek D, Jin Y, Jeong JC, Lee HJ, Moon H, Lee J, Shin D, Kang CH, Kim DH, Nam J, Lee SY, Yun DJ: **Suppression of reactive oxygen species by glyceraldehyde-3-phosphate dehydrogenase.** *Phytochemistry* 2008, **69**:333-338.
46. Carpentier SC, Witters E, Laukens K, Van Onckelen H, Swennen R, Panis B: **Banana (*Musa spp.*) as a model to study the meristem proteome: acclimation to osmotic stress.** *Proteomics* 2007, **7**:92-105.
47. Varón D, Boylan SA, Okamoto K, Price CW: ***Bacillus subtilis* gtaB encodes UDP-glucose pyrophosphorylase and is controlled by stationary-phase transcription factor sigma B.** *J Bacteriol* 1993, **175**:3964-3971.
48. Yan S, Tang Z, Su W, Sun W: **Proteomic analysis of salt stress-responsive proteins in rice root.** *Proteomics* 2005, **5**:235-244.
49. Li L, Wright SJ, Krystofova S, Park G, Borkovich KA: **Heterotrimeric G protein signaling in filamentous fungi.** *Annu Rev Microbiol* 2007, **61**:423-452.
50. Chen D, Toone WM, Mata J, Lyne R, Burns G, Kivinen K, Brazma A, Jones N, Bahler J: **Global transcriptional responses of fission yeast to environmental stress.** *J Mol Biol Cell* 2003, **14**:214-229.

51. Perfus-Barbeoch L, Jones AM, Assmann SM: **Plant heterotrimeric G protein function: insights from *Arabidopsis* and rice mutants.** *Curr Opin Plant Biol* 2004, **7**:719-731.
52. Joo JH, Wang S, Chen JG, Jones AM, Fedoroff NV: **Different signaling and cell death roles of heterotrimeric g protein a and b subunits in the arabidopsis oxidative stress response to ozone.** *The Plant Cell* 2005, **17**:957-970.
53. Suvarna K, Bartiss A, Wong B: **Mannitol-1-phosphate dehydrogenase from *Cryptococcus neoformans* is a zinc-containing long-chain alcohol/polyol dehydrogenase.** *Microbiology* 2000, **146**:2705-2713.
54. Hondorp ER, Matthews RG: **Oxidative stress inactivates cobalamin-independent methionine synthase (MetE) in *Escherichia coli*.** *PLoS Biol* 2004, **2**:e336.
55. Hohmann S: **Osmotic stress signaling and osmoadaptation in yeasts.** *Microbiol Mol Biol Rev* 2002, **66**:300-372.
56. García-Leiro A, Cerdán ME, González-Siso MI: **Proteomic analysis of the oxidative stress response in *Kluyveromyces lactis* and effect of glutathione reductase depletion.** *J Proteome Res* 2010, **9**:2358-2376.
57. Reddy AM, Kumar SG, Jyonthsnakumari G, Thimmanai S, Sudhakar C: **Lead induced changes in antioxidant metabolism of horsegram (*Macrotyloma uniflorum* (Lam.) Verdc.) and bengalgram (*Cicer arietinum* L.).** *Chemosphere* 2005, **60**:97-104.
58. Simpson JP, Di Leo R, Dhanoa PK, Allan WL, Makhmoudova A, Clark SM, Hoover GJ, Mullen RT, Shelp BJ: **Identification and characterization of a plastid-localized Arabidopsis glyoxylate reductase isoform: comparison with a cytosolic isoform and implications for cellular redox homeostasis and aldehyde detoxification.** *J Exp Bot* 2008, **59**:2545-2554.
59. Easton JA, Thompson P, Crowder MW: **Time-dependent translational response of *E. coli* to excess Zn (II).** *J Biomed Tech* 2006, **17**:303-307.
60. Hwang EW, Kim KA, Park SC, Jeong MJ, Byun MO, Kwon HB: **Expression profiles of hot pepper (*Capsicum annuum*) genes under cold stress conditions.** *J Biosci* 2005, **30**:657-667.
61. Kim YK, Yoo WI, Lee SH, Lee MY: **Proteomic analysis of cadmium-induced protein profile alterations from marine alga *Nannochloropsis oculata*.** *Ecotoxicology* 2005, **14**:589-596.
62. Inui M, Suda M, Okino S, Nonaka H, Puskás LG, Vertes AA, Yukawa H: **Transcriptional profiling of *Corynebacterium glutamicum* metabolism during organic acid production under oxygen deprivation conditions.** *Microbiology* 2007, **153**:2491-2504.
63. Singh R, Lemire J, Mailloux RJ, Appanna VD: **A novel strategy involved anti-oxidative defense: The conversion of NADH into NADPH by a metabolic network.** *PLoS ONE* 2008, **3**:1-7.
64. Kumari M, Taylor GJ, Deyholos MK: **Transcriptomic responses to aluminum stress in roots of *Arabidopsis thaliana*.** *Mol Genet Genomics* 2008, **279**:339-357.
65. Marchitti SA, Brocker C, Stagos D, Vasiliou V: **Non-P450 aldehyde oxidizing enzymes: the aldehyde dehydrogenase superfamily.** *Expert Opin Drug Metabol Toxicol* 2008, **4**:697-720.
66. Parvatiyar K, Alsabbagh EM, Ochsner UA, Stegemeyer MA, Smulian AG, Hwang SH, Jackson CR, McDermott TR, Hassettl DJ: **Global analysis of cellular factors and responses involved in *Pseudomonas aeruginosa* resistance to arsenite.** *J Bacteriol* 2005, **187**:4853-4864.
67. Bro C, Regenber B, Lagniel G, Labarre J, Montero-Lomeli M, Nielsen J: **Transcriptional, proteomic, and metabolic responses to lithium in galactose-grown yeast cells.** *J Biol Chem* 2003, **278**:32141-32149.
68. Sunkar R, Bartels D, Kirch H: **Overexpression of a stress-inducible aldehyde dehydrogenase gene from *Arabidopsis thaliana* in transgenic plants improves stress tolerance.** *Plant J* 2003, **35**:452-464.
69. Kotchoni SO, Kuhns C, Ditzler A, Kirch HH, Bartels D: **Over-expression of different aldehyde dehydrogenase genes in *Arabidopsis thaliana* confers tolerance to abiotic stress and protects plants against lipid peroxidation and oxidative stress.** *Plant Cell Environ* 2006, **29**:1033-1048.
70. Norbeck J, Blomberg A: **Metabolic and regulatory changes associated with growth of *Saccharomyces cerevisiae* in 1.4 M NaCl. Evidence for osmotic induction of glycerol dissimilation via the dihydroxyacetone pathway.** *J Biol Chem* 1997, **272**:5544-5554.
71. Boy-Marcotte E, Lagniel G, Perrot M, Bussereau F, Boudsocq A, Jacquet M, Labarre J: **The heat shock response in yeast: differential regulations and contributions of the Msn2p/Msn4p and Hsf1p regulons.** *Mol Microbiol* 1999, **33**:274-283.
72. Rep M, Krantz M, Thevelein JM, Hohmann S: **The transcriptional response of *Saccharomyces cerevisiae* to osmotic shock: Hot1p and Msn2p/Msn4p are required for the induction of subsets of high osmolarity glycerol pathway-dependent genes.** *J Biol Chem* 2000, **275**:8290-8300.
73. Godon C, Lagniel G, Lee J, Buhler JM, Kieffer S, Perrot M, Boucherie H, Toledano MB, Labarre J: **The H₂O₂ stimulon in *Saccharomyces cerevisiae*.** *J Biol Chem* 1998, **273**:22480-22489.
74. Causton HC, Ren B, Koh SS, Harbison CT, Kanin E, Jennings EG, Lee TI, True HL, Lander ES, Young RA: **Remodeling of yeast genome expression in response to environmental changes.** *Mol Biol Cell* 2001, **12**:323-337.
75. Inoue Y, Tsujimoto Y, Kimura A: **Expression of the glyoxalase I gene of *Saccharomyces cerevisiae* is regulated by high osmolarity glycerol mitogen-activated protein kinase pathway in osmotic stress response.** *J Biol Chem* 1998, **273**:2977-2983.
76. Wang Y, Cao YY, Jia XM, Cao YB, Gao PH, Fu XP, Ying K, Chen WS, Jiang YY: **Cap1p is involved in multiple pathways of oxidative stress response in *Candida albicans*.** *Free Radic Biol Med* 2006, **40**:1201-1209.
77. Espartero J, Sanchez-Aguayo I, Pardo JM: **Molecular characterization of glyoxalase-1 from a higher plant: upregulation by stress.** *Plant Mol Biol* 1995, **29**:223-1233.
78. Shiozaki N, Yamada M, Yoshida Y: **Analysis of salt-stress-inducible ESTs isolated by PCR-subtraction in salt-tolerant rice.** *Theor Appl Genet* 2005, **110**:1177-1186.
79. Lee DG, Ahsan N, Lee SH, Lee JJ, Bahk JD: **Chilling stress-induced proteomic changes in rice roots.** *J Plant Physiol* 2009, **166**:1-11.
80. Sommer A, Fischer P, Krause K, Boettcher K, Brophy PM, Walter RD, Liebau E: **A stress-responsive glyoxalase I from the parasitic nematode *Onchocerca volvulus*.** *Biochem J* 2001, **353**:445-452.
81. Gdula DA, Sandaltzopoulos R, Tsukiyama T, Wu C: **Inorganic pyrophosphatase is a component of the Drosophila nucleosome remodeling factor complex.** *Gene Dev* 1998, **12**:3206-3216.
82. Schwanbeck R, Xiao H, Wu C: **Spatial contacts and nucleosome step movements induced by the NURF chromatin remodeling complex.** *J Biol Chem* 2004, **279**:39933-39941.
83. Rodríguez-Gabriel MA, Remacha M, Ballesta JP: **The RNA interacting domain but not the protein interacting domain is highly conserved in ribosomal protein P0.** *J Biol Chem* 2000, **275**:2130-2136.
84. Yacoub A, Kelley MR, Deutsch WA: **Drosophila ribosomal protein P0 contains apurinic/aprimidinic endonuclease activity.** *Nucl Acids Res* 1996, **24**:4298-4303.
85. Ji H, Moritz RL, Kim YS, Zhu HJ, Simpson RJ: **Analysis of Ras-induced oncogenic transformation of NIH-3T3 cells using differential-display 2-DE proteomics.** *Electrophoresis* 2007, **28**:1997-2008.
86. Yuan X, Kuramitsu Y, Furumoto H, Zhang X, Hayashi E, Fujimoto M, Nakamura K: **Nuclear protein profiling of Jurkat cells during heat stress-induced apoptosis by 2-DE and MS/MS.** *Electrophoresis* 2007, **28**:2018-2026.
87. Singh OV: **Proteomics and metabolomics: the molecular make-up of toxic aromatic pollutant bioremediation.** *Proteomics* 2006, **6**:5481-5492.
88. Zhao B, Poh CL: **Insights into environmental bioremediation by microorganisms through functional genomics and proteomics.** *Proteomics* 2008, **8**:874-881.
89. Sharon A, Finkelstein A, Shlezinger N, Hatam I: **Fungal apoptosis: function, genes and gene function.** *FEMS Microbiol Rev* 2009, **33**:833-854.
90. Lorin S, Dufour E, Sainsard-Chanet A: **Mitochondrial metabolism and aging in the filamentous fungus *Podospora anserina*.** *Biochim Biophys Acta* 2006, **1757**:604-610.
91. Zamaraeva MV, Sabirov RZ, Maeno E, Ando-Akatsuka Y, Bessonova S, Okada Y: **Cells die with increased cytosolic ATP during apoptosis: a bioluminescence study with intracellular luciferase.** *Cell Death Differ* 2005, **12**:1390-1397.
92. Rodrigues-Pousada CA, Nevitt T, Menezes R, Azevedo D, Pereira J, Amaral C: **Yeast activator proteins and stress response: an overview.** *FEBS Letters* 2004, **567**:80-85.
93. Vivancos AP, Jara M, Zuin A, Sansó M, Hidalgo E: **Oxidative stress in *Schizosaccharomyces pombe*: different H₂O₂ levels, different response pathways.** *Mol Genet Genomics* 2006, **276**:495-502.
94. Prouty AL: **Bench-scale development and evaluation of a fungal bioreactor for color removal from bleach effluents.** *Appl Microbiol Biotechnol* 1990, **32**:490-493.

95. Damerval C, de Vienne D, Zivy M, Thiellement H: **The technical improvements in two-dimensional electrophoresis increase the level of genetic variation detected in wheat-seedling proteins.** *Electrophoresis* 1986, **7**:52-54.
96. Ramagli LS, Rodriguez LV: **Quantitation of microgram amounts of protein in two-dimensional polyacrylamide gel electrophoresis sample buffer.** *Electrophoresis* 1985, **6**:559-563.
97. Görg A, Postel W, Günther S, Weser J, Strahler JR, Hanash SM, Somerlot L, Kuick R: **Approach to stationary two-dimensional pattern: influence of focusing time and immobiline/carrier ampholytes concentrations.** *Electrophoresis* 1988, **9**:37-46.
98. Neuhof VN, Arold N, Taube D, Erhardt W: **Improved staining of proteins in polyacrylamide gels including isoelectric focusing gels with clear background at nanogram sensitivity using Coomassie Brilliant Blue G-250 and R-250.** *Electrophoresis* 1988, **9**:255-262.
99. Joint Genome Institutes (JGI) *P. chrysosporium* protein sequences database. [ftp://ftp.jgi-psf.org/pub/JGI_data/Phanerochaete_chrysosporium/v2.0/BestModels2.1.prot.gz].
100. MASCOT software. [<http://www.matrixscience.com>].
101. WoLF PSORT web server. [<http://wolfsort.seq.cbrc.jp>].

doi:10.1186/1477-5956-9-12

Cite this article as: Yıldırım *et al.*: Characterization of proteome alterations in *Phanerochaete chrysosporium* in response to lead exposure. *Proteome Science* 2011 **9**:12.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

