

SCIENTIFIC REPORTS



OPEN

Optomechanically induced transparency in multi-cavity optomechanical system with and without one two-level atom

Received: 05 January 2016

Accepted: 09 June 2016

Published: 28 June 2016

Amjad Sohail, Yang Zhang, Jun Zhang & Chang-shui Yu

We analytically study the optomechanically induced transparency (OMIT) in the N -cavity system with the N th cavity driven by pump, probing laser fields and the 1st cavity coupled to mechanical oscillator. We also consider that one atom could be trapped in the i th cavity. Instead of only illustrating the OMIT in such a system, we are interested in how the number of OMIT windows is influenced by the cavities and the atom and what roles the atom could play in different cavities. In the resolved sideband regime, we find that, the number of cavities precisely determines the maximal number of OMIT windows. It is interesting that, when the two-level atom is trapped in the even-labeled cavity, the central absorptive peak (odd N) or dip (even N) is split and forms an extra OMIT window, but if the atom is trapped in the odd-labeled cavity, the central absorptive peak (odd N) or dip (even N) is only broadened and thus changes the width of the OMIT windows rather than induces an extra window.

Cavity optomechanical system (OMS) has recently attracted increasing interest in both theory and experiment (ref. 1 and references therein). It usually composed of two mirrors with one fixed and the other movable or a micro-mechanical membrane oscillating inside two fixed mirrors. Such a system demonstrates the interaction between the movable oscillator and the optical field in the cavity via the radiation pressure and becomes a platform for the study^{2–16} of quantum ground-state cooling^{17–20}, strong coupling dynamics^{1,7,21,22} and other coherent dynamics in microscopic and macroscopic domains^{23–25}. When a strong laser field drives the optomechanical cavity, an analogue of electromagnetically induced transparency (EIT) for the output at the frequency of the weak detecting field could appear^{26,27}. Such an EIT-like phenomenon is usually called as the optomechanically induced transparency (OMIT) which is equivalent to the case of two coupled harmonic oscillators²¹ and has been demonstrated in experiments^{28–30}. OMIT has also been widely investigated in diverse aspects including the cases with higher-order sidebands³¹ or in the nonlinear regime^{32–34}, OMIT in the cavity with membranes^{35,36} and so on. In particular, OMIT has shown many potential applications in control of light speed²⁹, charge measurement³⁷, single photon router³⁸ and so on, which forms the further motivations to study OMIT.

Introducing the atomic freedom into OMS can not only strengthen the coupling but also allow rich physics via enhanced nonlinearities^{39–41}. It has been applied to improve optomechanical cooling^{42–44} and even the ground-state cooling outside the resolved sideband regime⁴⁵. In particular, it is shown⁴⁶ that a two-level atomic ensemble coupled to OMS can both enhance the photon-phonon coupling through radiation pressure and broaden the transparency windows. In addition, coupled-cavity array related to the 1D waveguide or atoms has been widely studied in the control of photon transport such as quantum router^{47–50}. Does the multiple-cavity quantum optomechanics bring new insight into OMIT? How can the OMIT be controlled if introducing the atomic freedom into the multiple-cavity system?

In this paper, we address the above questions by investigating the OMIT phenomenon in multiple-cavity optomechanical system coupled to one two-level atom. *Here instead of only illustrating the OMIT in such a system, we are especially interested in how the number of OMIT windows is related to the number of the cavities as well as the potential trapped atom and what roles the atom could play in different cavities.* Through our analytic calculations, it is shown that the maximal number of OMIT windows is precisely determined by the cavity number, if there

School of Physics and Optoelectronic Technology, Dalian University of Technology, Dalian 116024, P.R. China. Correspondence and requests for materials should be addressed to C.-s.Y. (email: quaninformation@sina.com or ycs@dlut.edu.cn)




Figure 1. Schematic diagram of N cavities connected through tunneling parameters g_n . A strong driving field and a weak probing field are injected into cavity N while the first cavity is coupled with mechanical resonator.

does not exist any atom in the multi-cavity system. In particular, we find that the atom trapped in different cavities will play different roles in OMIT. When one atom is trapped in even-labeled cavity, the central absorptive peak (odd N) or dip (even N) is split and forms an extra OMIT window, but when the atom is trapped in odd-labeled cavity, the central absorptive peak (odd N) or dip (even N) is only broadened and thus changes the width of the OMIT windows instead of inducing the extra window. In addition, we also find that the multiple OMIT windows are the result of the coupling of multiple cavities irrespective of the participation of the mechanical oscillator. A numerical simulation is also given to support our results.

Results

The model. The optomechanical system under consideration is shown schematically in Fig. 1. The system includes N cavities labelled by $1, 2, \dots, N$ with the frequency of j th cavity denoted by ω_j . The n th and $(n+1)$ th cavities with $n \neq N$ are connected through tunneling parameters (hopping rates) g_n . Such a coupled cavity array (2D) has been systematically studied in various cases in ref. 51 and later considered in the single-photon router^{49,50}. Here we only consider 1D cavity chain, in particular, we let one end mirror of Cavity 1 be movable as shown in Fig. 1. Thus it forms an optomechanical system. Cavity N is separately driven by one coupling field ε_c and one probing field ε_p . In addition, we assume that one two-level atom could be trapped in the i th cavity $1 \leq i \leq N$ with g_a denoting the atom-cavity coupling strength. In this model, the optical modes are described by annihilation (creation) operators $c_n (c_n^\dagger)$ and the mechanical mode is represented by $b (b^\dagger)$ which is equivalent to the description by x_m and p_m . This similar description can be found in ref. 41. Let the frequency of the coupling field be ω_c , so in the rotating frame at ω_c , the Hamiltonian of our system reads

$$H = \sum_j \Delta_j c_j^\dagger c_j + \omega_m b^\dagger b + \frac{1}{2} \Delta_a \sigma_z + i\varepsilon_c (c_N^\dagger - c_N) + i\varepsilon_p (c_N^\dagger e^{-i\Delta t} - c_N e^{i\Delta t}) + g_a (c_i \sigma_+ + c_i^\dagger \sigma_-) - g c_1^\dagger c_1 (b^\dagger + b) + \sum_{n=1}^{N-1} g_n (c_{n+1}^\dagger c_n + c_n^\dagger c_{n+1}) \quad (1)$$

with ω_p, ω_a representing the frequency of the probing field and the atomic transition frequency. In Eq. (1) the first three terms, respectively, denote the free Hamiltonian for the cavities, the movable mirror and the trapped atom with $\Delta_j = \omega_j - \omega_c$, $\Delta = \omega_p - \omega_c$ and $\Delta_a = \omega_a - \omega_c$, the last two terms in first line corresponds to the interaction of the N th cavity driven by the coupling field ε_c and the probing field ε_p . The first term in the second line of Eq. (1) describes the interaction between the atom and the i th cavity, the second term corresponds to the interaction between the 1st cavity and the movable mirror via the radiation pressure, and the last term describes the hopping between the two adjacent cavities. In addition, g in Eq. (1) denotes the coupling strength between the 1st cavity and the mechanical oscillator. It is obvious that $\Delta_a = g_a = 0$ means no atom in the cavities.

The dynamics. Based on the above Hamiltonian, one can easily obtain the Langevin Equations for the operators. So the corresponding equations for the mean value of operators in the mean-field approximation, viz, $\langle \dot{s}t \rangle = \langle s \rangle \langle t \rangle$, can be given by

$$\langle \dot{c}_N \rangle = -(\kappa_N + i\Delta_N) \langle c_N \rangle - ig_{N-1} \langle c_{N-1} \rangle + \varepsilon_c + \varepsilon_p e^{-i\Delta t}, \quad (2)$$

$$\langle \dot{c}_n \rangle = -(\kappa_n + i\Delta_n) \langle c_n \rangle - i(g_{n-1} \langle c_{n-1} \rangle + g_n \langle c_{n+1} \rangle), \quad n \neq 1, i, N, \quad (3)$$

$$\langle \dot{c}_i \rangle = -(\kappa_i + i\Delta_i) \langle c_i \rangle - ig_a \langle \sigma_- \rangle - i(g_{i-1} \langle c_{i-1} \rangle + g_i \langle c_{i+1} \rangle), \quad i \neq 1, N, \quad (4)$$

$$\langle \dot{c}_1 \rangle = -(\kappa_1 + i\tilde{\Delta}_1) \langle c_1 \rangle - ig_1 \langle c_2 \rangle + ig \langle c_1 \rangle (\langle b^\dagger \rangle + \langle b \rangle), \quad (5)$$

$$\langle \dot{b} \rangle = -(\gamma_m + i\omega_m) \langle b \rangle + ig | \langle c_1 \rangle |^2, \quad (6)$$

$$\langle \dot{\sigma}_- \rangle = -(\gamma_a + i\Delta_a) \langle \sigma_- \rangle + ig_a \langle c_i \rangle \langle \sigma_z \rangle, \quad (7)$$

$$\langle \dot{\sigma}_z \rangle = -2(1 + \langle \sigma_z \rangle) \gamma_a + 2ig_a (\langle c_i^\dagger \rangle \langle \sigma_- \rangle + \langle c_i \rangle \langle \sigma_+ \rangle). \quad (8)$$

Here κ_n denotes the leakage of n th cavity and γ_m and γ_a , respectively, represent the decay rates of the mechanical oscillator and the atom. If the atom is trapped in the first cavity, Eq. (5) should be replaced by

$$\langle \dot{c}_1 \rangle = -(\kappa_1 + i\tilde{\Delta}_1)\langle c_1 \rangle - ig_1\langle c_2 \rangle - ig_a\langle \sigma_- \rangle + ig\langle c_1 \rangle(\langle b^\dagger \rangle + \langle b \rangle). \tag{9}$$

If the atom is trapped in N th cavity, Eq. (2) should be replaced by

$$\langle \dot{c}_N \rangle = -(\kappa_N + i\Delta_N)\langle c_N \rangle - ig_{N-1}\langle c_{N-1} \rangle - ig_a\langle \sigma_- \rangle + \varepsilon_c + \varepsilon_p e^{-i\Delta t}. \tag{10}$$

In order to solve the dynamics, we suppose

$$\langle \hat{O}(t) \rangle = \bar{O} + \delta O(t) = \bar{O} + O_- e^{-i\Delta t} + O_+ e^{i\Delta t}, \tag{11}$$

for any operator $\hat{O}(t)$ with \bar{O} denoting the steady-state value without ε_p and $\delta O = O_- e^{-i\Delta t} + O_+ e^{i\Delta t}$ induced by the weak probing field. Substituting Eq. (11) into Eqs (2–10), one can obtain an equation array for \bar{O} which has the same form as Eqs (2–10) except $\varepsilon_p = 0$ and $\bar{O} = 0$. This equation arrays are omitted here. In addition, one can also obtain an equation array for $\delta O(t)$ which is given in the Methods (Eqs (17–22)). By solving the equations for \bar{O} , one can find that

$$\bar{\lambda} = \bar{b} + \bar{b}^* = \frac{2\omega_m g}{\omega_m^2 + \gamma_m^2} |\bar{c}_1|^2, \tag{12}$$

and

$$\bar{\sigma}_z = \frac{\gamma_a(\gamma_a^2 + \Delta_a^2)}{2i\Delta_a g_a^2 |c_i|^2 - \gamma_a(\gamma_a^2 + \Delta_a^2)}. \tag{13}$$

In addition, considering Eq. (11), one can easily find the equations for O_\pm . However, for the purpose of this paper, we only provide the equations for O_- in the Methods Eqs (23–28) within the resolved sideband regime, i.e., $\omega_m \gg \kappa$ and $\Delta_n = \tilde{\Delta}_1 = \Delta_a = \omega_m$ where $\tilde{\Delta}_1 = \Delta_1 - g\bar{\lambda}$. These equations provide the fundamental description of the dynamics of the model considered here.

Output field. In order to reveal the OMIT, we will have to find out the response of the system to the probing frequency, which can be detected by the output field. Based on the input–output theory⁵², we can obtain

$$\varepsilon_{out,p} e^{-i\Delta t} + \varepsilon_p e^{-i\Delta t} + \varepsilon_c = 2\kappa_N \langle c_N \rangle. \tag{14}$$

Substituting Eq. (11) into Eq. (14), one can find that the total output field at the probing frequency ω_p can be given by

$$\varepsilon_T = \frac{\varepsilon_{out,p}}{\varepsilon_p} + 1 = \frac{2\kappa_N c_{N,-}}{\varepsilon_p} = \chi_p + i\tilde{\chi}_p. \tag{15}$$

It is clear that $\chi_p = \text{Re}(\varepsilon_T)$ and $\tilde{\chi}_p = \text{Im}(\varepsilon_T)$ are the in-phase and out-of-phase quadratures of the output probing field, representing the absorptive and dispersive behavior of the output probing field, respectively. The quadrature can be measured via the homodyne technique⁵². So the next task is to find $c_{N,-}$. In order to gain more physical insight, we only consider the system in the sideband resolved regime. Thus $c_{N,-}$ can be easily obtained by solving Eqs (23–28). So the output field ε_T can be directly given by

$$\varepsilon_T = 2\kappa_N c_{N,-} = \frac{2\kappa_N}{(\kappa_N - ix) + \frac{g_{N-1}^2}{\kappa_{N-1} - ix + \frac{g_{N-2}^2}{\kappa_{N-2} - ix + \frac{g_{N-3}^2}{\dots + \frac{g_1^2}{\kappa_1 - ix + \frac{g_a^2 |\bar{\sigma}_z|^2}{\gamma_a - ix}}}}}, \tag{16}$$

where $x = \Delta - \omega_m$ and $G = g\bar{c}_1$ is the effective optomechanical rate. In above equation, the first line of the denominator represents two cavities with radiative decays κ_N and κ_{N-1} are connected through their coupling strength g_{N-1} . Second line represents two cavities with radiative decays κ_{N-1} and κ_{N-2} are connected through their coupling strength g_{N-2} and so on. The 1st cavity in the last line is coupled to the mechanical oscillator by an effective coupling G . In addition, an extra term $\frac{g_a^2 |\bar{\sigma}_z|^2}{\gamma_a - ix}$ corresponding to the atomic contribution appears in the κ_i line which denotes the atom is coupled to the i th cavity with an effective coupling $g_a |\bar{\sigma}_z|$. Certainly, if the atom is trapped in the first cavity, this term will appear in the last line. If the atom is placed in the N th cavity, it will appear in the first line of the denominator. It is obvious that the output field depends on both the parameters of the system and the steady-state values of c_1 and c_i . These two values can be determined by solving the equations for all \bar{O}




Figure 2. The Real part $\text{Re}(\varepsilon_T)$ in the absence of atom-field coupling, as a function of x/κ_N for (a) two cavities, (b) three cavities and (c) four cavities with the parameters $\omega_m = 2\pi \times 51.8 \text{ MHz}$, $\gamma_m = 2\pi \times 41 \text{ KHz}$, $G = 2\pi \times 10 \text{ MHz}$, $\kappa_N = 2\pi \times 15 \text{ MHz}$, $\kappa_1 = \kappa_2 = \kappa_3 = \dots = 2\pi \times 0.027 \text{ MHz}$, and the coupling rates $g_1 = g_2 = g_3 = \kappa_N$. The solid lines show the analytic expressions given by Eq. (16), but the dotted lines represent the solution numerically solved from Eqs (23–28), which guarantees the validity of our analytical result Eq. (16).

which have been omitted here. But the concrete expressions of c_1 and c_i are quite complicated, so it is impossible to present the concrete forms. It is fortunate that this does not influence our understanding on the OMIT window numbers. One can find from the latter part that the values of c_1 and c_i only affect the width of the OMIT windows. In this sense, it doesn't matter whether they can be simultaneously assigned by some values. Therefore, for simplicity, one can select that $\bar{\sigma}_z = -1$ and G can be given by any reasonable and convenient assignment.

OMIT windows. The OMIT is signaled by the simultaneously vanishing absorption and dispersion, which is further related to the simultaneously vanishing χ_p and $\tilde{\chi}_p$, that is ε_T . In order to show the OMIT windows as many as possible, we restrict ourselves to the weak dissipative regime, i.e., $g_i \gtrsim \kappa_N \gg \kappa_i, \gamma_{m/a}$, to discuss the points where ε_T vanishes. This is also supported by our latter numerical procedures.

Without atom. If there does not exist any atom in the optomechanical system, the term with g_a vanishes due to $g_a = 0$. In this case, the vanishing ε_T means that the denominator approaches infinity which can be further determined by the vanishing denominator corresponding to the numerator $|g_{N-1}|^2$. It is obvious that the condition with such a vanishing denominator corresponds to an equation with N degrees. Therefore it has at most N different roots. This means that such an optomechanical system has at most N OMIT windows. To give an intuitive illustration of the OMIT, we numerically evaluate OMIT and demonstrate the multiple transparency windows due to the interaction between cavity fields and the mechanical oscillator. We take the parameters from^{53–55} where, the damping rate of mechanical oscillator $\gamma_m = 2\pi \times 41 \text{ kHz}$, decay rate of the driven cavity field $\kappa_N = 2\pi \times 15 \text{ MHz}$ and the frequency of oscillator $\omega_m = 2\pi \times 51 \text{ MHz}$. For the case of the resolved sideband regime, i.e. the mechanical frequency is much greater than the decays and $\Delta_n = \Delta_1 = \Delta_a = \omega_m$, we plot the phase quadratures of the output probing fields for a system with two, three and four cavities in Fig. 2 which exhibits two windows, three windows and four windows respectively. We assume that the 1st cavity coupled to mechanical oscillator with $G = 2\pi \times 12 \text{ MHz}$. The multiple transparency windows display that the optomechanical system becomes simultaneously transparent to the probing field at multiple different frequencies, which is the result of the destructive interferences between the input probing field and the anti-Stokes fields generated by the interactions of the coupling field with the multiple cavities. In addition, in order to show the effects of G , we plot Fig. 3 with different choices of G . One can find that the larger G corresponds to the wider central absorptive peak (or dip for odd number of cavities) in the valid range of G . Numerical results show that the interval that the OMITs occur (from about -2 to 2 in all the figures) is almost independent of the numbers of cavities. In fact, the width is determined by all the hopping rate g_n . Here in order to find out many enough OMIT windows, we let all $g_n = \kappa_N$, so the interval (if defined by the half width) is slightly changed. Under this condition, by numerical demonstrations, we find that the half width is increased with N . In particular, one can easily prove that when N tends to infinity, the half width is just 4. So when the central absorptive peak or dip gets wider, and the others get narrower due to the fixed interval. In one word, the value of G only affects the width of the transparency window instead of the maximal number of the OMIT window.

One atom in one cavity. Since we have set $g_i \gtrsim \kappa_N \gg \kappa_i, \gamma_i$, for an intuitive understanding of the number of OMIT windows, one can safely neglect the dissipative constants which contributes to the level width of the cavity as well as the atom. Under such a condition, one can find that there exist two cases in our optomechanical system.

1) **The atom is trapped in the odd-labeled cavity.** In this case, one can see that the extra term $\frac{g_a^2 |\bar{\sigma}_z|^2}{\gamma_a - ix}$ can only exist in the lines corresponding to $\kappa_1, \kappa_3, \dots$. The contribution of such an extra atomic term is mathematically to increase the numerator of the same line and physically to directly broaden the central absorptive peak for even N (or absorptive dip for odd N) and then to change the width of the OMIT windows, which is analogous to increasing G in the case without atom. The most obvious example is when the atom is trapped in the first cavity. One can easily find that for weak γ_a and γ_m , the atomic term can be approximately absorbed in the term corresponding to the mechanical oscillator and the net result is equivalent to increasing $|g_1|^2$. 2) **The atom is trapped in the**




Figure 3. Real part $\text{Re}(\varepsilon_T)$ (solid red) and the imaginary part $\text{Im}(\varepsilon_T)$ (dashed black) as a function of x/κ_4 for four cavities. The three subplots from above to bottom, respectively, correspond to $G = 8 \text{ MHz}$, $G = 10 \text{ MHz}$ and $G = 12 \text{ MHz}$. The other parameters are the same as in Fig. 2. One can see that the width of the central absorptive peak becomes wide with the increasing of G .




Figure 4. Real part $\text{Re}(\varepsilon_T)$ of the output field as a function of $x/\kappa_{3/4}$ with $\gamma_a = 2\pi \times 0.01 \text{ MHz}$ and $g_a = 2\pi \times 10 \text{ MHz}$. The other parameters are the same in Fig. 2. (a,c) illustrate cases with four cavities, where (a) corresponds to the atom trapped in cavity 1 (solid red) and cavity 3 (dashed black) and (c) corresponds to the atom in cavity 2 (solid red) and cavity 4 (dashed black). (b,d) correspond to the cases with three cavities, where (b) illustrates the atom trapped in cavity 1 (solid red) and cavity 3 (dashed black) and (d) shows the atom trapped in cavity 2 (solid red). The dashed blue lines in all the figures mean no trapped atom.

even-labeled cavity. In this case, the extra atomic term can lead to that the degree of the equation of the vanishing denominator corresponding to the numerator $|g_{N-1}|^2$ is added by 1. So when the atom is trapped in the even-labeled cavity, one can find one more extra OMIT window compared with the case without any atom. Similarly, in order to give an illustration of these different cases, we numerically evaluate the OMIT. We plot the figure in Fig. 4 with $g_a = 2\pi \times 10 \text{ MHz}$ and $\gamma_a = 2\pi \times 0.01 \text{ MHz}$. However, we don't plot the imaginary part $\text{Im}(\varepsilon_T)$ for the sharp illustration. We observe that, in four-cavity system, the width of the central absorptive peak tends to become wide through embedding the atom into the cavity 1 or cavity 3 as shown in Fig. 4(a). But, when the atom is placed in cavity 2 or 4, we have found the resonant character of the weak probing field changes and the central absorptive peak splits. Hence four OMIT windows transfigure to a penta OMIT window, as shown in Fig. 4(c). Similarly, in Fig. 4(b,d) that correspond to the cases of three cavities, one can find that the atom will directly lead to the broadening or splitting of the central absorptive dips instead of absorptive peaks.

The role of the mechanical oscillator. Actually the physical mechanism of the mechanical oscillator about the production of OMIT has been well known^{26,27}. In this part, we are only interested in how the existence of the mechanical oscillator affects the number of OMIT windows. If there does not exist any mechanical oscillator, that means $G = 0$. If the atom is trapped in the first cavity under this condition, the number of the OMIT windows will keep invariant, but the width of the OMIT window will become narrow. This could be equivalently understood as the case without atom in the optomechanical system. That is, the role of the mechanical oscillator is to broaden the OMIT window in this case. In other cases, that is, no atom exists or the atom is only trapped in the even-labeled cavity and so on, one can easily find that the OMIT windows will be decreased by 1. In this case, one can draw the conclusion that the mechanical oscillator contributes an OMIT window. In this sense, we can say that the multiple OMIT windows should come from the coupling of the multiple cavities instead of the direct participation of the mechanical oscillator.

Discussions and Conclusion

Before the end, we would like to emphasize that similar to multiple EIT windows, the multiple OMIT windows permit the probing light with different frequencies to transmit simultaneously. So the OMIT with multiple windows could also be used in multi-channel optical communication and multichannel quantum information processing⁵⁶. OMIT is also closely related to the superluminal and ultraslow light propagation^{18,57}, the quantum router³⁸, charge measurement³⁷ and so on. Hence, OMIT with multiple transparency windows could mean wider applications. In addition, the experimental realization of coupled cavity array is systematically reviewed in ref. 51. The parameters we used are mainly taken from refs 18, 28 and 29 which report the current experiments about the optomechanical system and OMIT. These can be used to well evaluate the feasibility.

We would also like to mention that one can also consider an atomic ensemble instead of a single atom in the system. We think that the net effect is equivalent to increasing the coupling between the single atom and cavity if the atomic ensemble is considered in the limit of large atomic number. In addition, if two or more atoms are trapped in different even-labeled cavities, respectively, we think multiple extra windows will occur. If they are trapped in different odd-labeled cavities, the OMIT window will change much greater. In addition, the entanglement in this optomechanical system is an interesting topic. Our preliminary results have shown the entanglement can be produced between the different components of this optomechanical system (such as between two cavities, or between one cavity and the movable mirror). It is interesting that the entanglement between the mirror and the N th cavity could be enhanced by the multiple cavities, but the entanglement of other components could also be reduced. In particular, the existence of the atom could play different roles in the control of the generation of various entanglement. All the detailed results will be reported in the latter papers.

In summary, we have theoretically discussed the response of an optomechanical system which includes N cavities. We have given a general analytical expression of the generation of multiple OMIT windows. The mechanism of OMIT could have been well understood and even one could have known that an atom or atomic ensemble could broaden the width of OMIT window. However, it is shown here that the number of the OMIT windows directly depend upon the number of cavities. In particular, we find that, when the atom is trapped in even-labeled cavity, the number of the OMIT windows will be increased by one; if the atom is trapped in the odd-labeled cavity, the only the width of the OMIT windows could be changed. In addition, we also find that the multiple OMIT windows are only attributed to the coupling of the multiple cavities and irrespective of the coupling to the mechanical oscillator, because the mechanical oscillator could produce only one additional OMIT window or change the width of the OMIT windows which depends on the even- or odd- labelled cavity that the atom is trapped in.

Methods

In this section, we will give a brief introduction of the derivation of the equations used in the main text. Substituting $\langle \hat{O}(t) \rangle = \bar{O} + \delta O(t)$ given in Eq. (11) into Eqs (2–10), Eqs (2–10) can be rewritten by \bar{O} and $\delta O(t)$. Since $\delta O(t)$ is small and depends on time and \bar{O} is independent of time. One can separate equations into one related to time and the other irrelevant of time. The equation array irrelevant of time corresponding to \bar{O} has the same form as Eqs (2–10) except setting $\varepsilon_p = 0$ and $\bar{O} = 0$. In other words, if we replace $\langle O \rangle$ in Eqs (2–10) by \bar{O} and let $\varepsilon_p = 0$ and $\bar{O} = 0$, we will obtain the equations for \bar{O} . Our Eqs (12) and (13) are solved from these equations, but for avoiding repetition, these equations are omitted here. The equations with time corresponding to $\delta O(t)$ should obviously include the term $\varepsilon_p e^{-i\Delta t}$. They can be directly given as follows.

$$\delta \dot{c}_N = -(\kappa_N + i\Delta_N)\delta c_N - ig_{N-1}\delta c_{N-1} + \varepsilon_p e^{-i\Delta t}, \quad (17)$$

$$\delta \dot{c}_n = -(\kappa_n + i\Delta_n)\delta c_n - i(g_{n-1}\delta c_{n-1} + g_n\delta c_{n+1}), \quad n \neq 1, i, N, \quad (18)$$

$$\delta \dot{c}_i = -(\kappa_i + i\Delta_i)\delta c_i - ig_a\delta\sigma_- - i(g_{i-1}\delta c_{i-1} + g_i\delta c_{i+1}), \quad i \neq 1, N, \quad (19)$$

$$\delta \dot{c}_1 = -(\kappa_1 + i\tilde{\Delta}_1)\delta c_1 - ig_1\delta c_2 + iG(\delta b^* + \delta b), \quad (20)$$

$$\delta \dot{b} = -(\gamma_m + i\omega_m)\delta b + i(G\delta c_1^* + G^*\delta c_1), \quad (21)$$

$$\delta \dot{\sigma}_- = -(\gamma_a + i\Delta_a)\delta\sigma_- + ig_a\delta c_i\bar{\sigma}_z, \quad (22)$$

where $G = g\bar{c}_1$ is the effective optomechanical rate. As mentioned in the text, we consider the system in the resolved sideband regime in order to gain more physical insight. That is, we let $\omega_m \gg \kappa$ and $\Delta_n = \tilde{\Delta}_1 = \Delta_a = \omega_m$. In such a resolved sideband regime, the lower sideband, far off-resonance can be safely neglected. This means that in Eq. (11), $O_+ \approx 0$ which is the same as²⁸. Thus, Eqs (17–22) can be rewritten for O_- as

$$0 = -(\kappa_N + ix)c_{N,-} - ig_{N-1}c_{N-1,-} + \varepsilon_p, \quad (23)$$

$$0 = -(\kappa_n + ix)c_{n,-} - i(g_{n-1}c_{n-1,-} + g_n c_{n+1,-}), \quad n \neq 1, i, N, \quad (24)$$

$$0 = -(\kappa_i + ix)c_{i,-} - ig_a \sigma_{-,i} - i(g_{i-1}c_{i-1,-} + g_i c_{i+1,-}), \quad i \neq 1, N, \quad (25)$$

$$0 = -(\kappa_1 + ix)c_{1,-} - ig_1 c_{2,-} + iGb_-, \quad (26)$$

$$0 = -(\gamma_m + ix)b_- + iG^*c_{1,-}, \quad (27)$$

$$0 = -(\gamma_a + ix)\sigma_{-,a} + ig_a c_{i,-} \bar{\sigma}_z, \quad (28)$$

where $x = \Delta - \omega_m$ is again the detuning from the center line of the sideband.

References

1. Aspelmeyer, M., Kippenberg, T. J. & Marquardt, F. Cavity optomechanics. *Rev. Mod. Phys.* **86**, 1391 (2014).
2. Mancini, S. *et al.* Entangling macroscopic oscillators exploiting radiation pressure. *Phys. Rev. Lett.* **88**, 120401 (2002).
3. Vitali, D. *et al.* Optomechanical entanglement between a movable mirror and a cavity field. *Phys. Rev. Lett.* **98**, 030405 (2007).
4. Hartmann, M. J. & Plenio, M. B. Steady state entanglement in the mechanical vibrations of two dielectric membranes. *Phys. Rev. Lett.* **101**, 200503 (2008).
5. Liao, J. Q., Wu, Q. Q. & Nori, F. Entangling two macroscopic mechanical mirrors in a two-cavity optomechanical system. *Phys. Rev. A* **89**, 014302 (2014).
6. Nunnenkamp, A., Børkje, K. & Girvin, S. M. Single-photon optomechanics. *Phys. Rev. Lett.* **107**, 063602 (2011).
7. Rabl, P. Photon Blockade Effect in Optomechanical Systems. *Phys. Rev. Lett.* **107**, 063601 (2011).
8. Purdy, T. P. *et al.* Strong optomechanical squeezing of light. *Phys. Rev. X* **3**, 031012 (2013).
9. Tian, L. Optoelectromechanical transducer: Reversible conversion between microwave and optical photons. *Ann. Phys. (Berlin)* **527**, 1 (2015).
10. Chang, D. E., Ni, K. K., Painter, O. & Kimble, H. J. Ultrahigh-Q mechanical oscillators through optical trapping. *New J. Phys.* **14**, 045002 (2012).
11. Akram, M. J. & Saif, F. Adiabatic population transfer based on a double stimulated raman adiabatic passage. *J. Russ. Laser Res.* **35**(6), 547 (2014).
12. Wang, Y. D. & Clerk, A. A. Using Interference for high fidelity quantum state transfer in optomechanics. *Phys. Rev. Lett.* **108**, 153603 (2012).
13. Karuza, M. *et al.* Tunable linear and quadratic optomechanical coupling for a tilted membrane within an optical cavity: theory and experiment. *J. Opt.* **15**, 025704 (2013).
14. Hammerer, K., Sørensen, A. S. & Polzik, E. S. Quantum interface between light and atomic ensembles. *Rev. Mod. Phys.* **82**, 1041 (2010).
15. Saif, F., LeKien, F. & Zubairy, M. S. Quantum theory of a micromaser operating on the atomic scattering from a resonant standing wave. *Phys. Rev. A* **64**, 043812 (2001).
16. Islam, R. U., Ikram, M. & Saif, F. Engineering maximally entangled N-photon NOON field states using an atom interferometer based on Bragg regime cavity QED. *J. Phys. B: At. Mol. Opt. Phys.* **40**, 1359 (2007).
17. Bhattacharya, M. & Meystre, P. Trapping and cooling a mirror to its quantum mechanical ground state. *Phys. Rev. Lett.* **99**, 073601 (2007).
18. Teufel, J. D. *et al.* Sideband cooling of micromechanical motion to the quantum ground state. *Nature* **475**, 359 (2011).
19. Chan, J. *et al.* Laser cooling of a nanomechanical oscillator into its quantum ground state. *Nature (London)* **478**, 89 (2011).
20. Meystre, P. A short walk through quantum optomechanics. *Ann. Phys. (Berlin)* **525**, 215 (2013).
21. O'Connell, A. D. *et al.* Quantum ground state and single-phonon control of a mechanical resonator. *Nature* **464**, 697 (2010).
22. Akram, U., Kiesel, N., Aspelmeyer, M. & Milburn, G. J. Single-photon opto-mechanics in the strong coupling regime. *New J. Phys.* **12**, 083030 (2010).
23. Rogers, B. *et al.* Hybrid optomechanics for quantum technologies. *Quantum Measurements and Quantum Metrology* **2**, 11 (2014).
24. Kippenberg, T. J. & Vahala, K. J. Cavity optomechanics: Back-Action at the Mesoscale. *Science* **321**, 1172 (2008).
25. Restrepo, J., Ciuti, C. & Favero, I. Single-polariton optomechanics. *Phys. Rev. Lett.* **112**, 013601 (2014).
26. Agarwal, G. S. & Huang, S. Electromagnetically induced transparency in mechanical effects of light. *Phys. Rev. A* **81**, 041803 (2010).
27. Huang, S. & Agarwal, G. S. Electromagnetically induced transparency with quantized fields in optocavity mechanics. *Phys. Rev. A* **83**, 043826 (2010).
28. Weis, S. *et al.* Optomechanically induced transparency. *Science* **330**, 1520 (2010).
29. Safavi-Naeini, A. H. *et al.* Electromagnetically induced transparency and slow light with optomechanics. *Nature (London)* **472**, 69 (2011).
30. Dong, C. *et al.* Transient optomechanically induced transparency in a silica microsphere. *Phys. Rev. A* **87**, 055802 (2013).
31. Xiong, H. *et al.* Higher-order sidebands in optomechanically induced transparency. *Phys. Rev. A* **86**, 013815 (2012).
32. Lemonde, M. A., Didier, N. & Clerk, A. A. Nonlinear interaction effects in a strongly driven optomechanical cavity. *Phys. Rev. Lett.* **111**, 053602 (2013).
33. Børkje, K. *et al.* Signatures of Nonlinear Cavity Optomechanics in the weak Coupling regime. *Phys. Rev. Lett.* **111**, 053603 (2013).
34. Kronwald, K. & Marquardt, F. Optomechanically induced transparency in the nonlinear quantum regime. *Phys. Rev. Lett.* **111**, 133601 (2013).
35. Karuza, M. *et al.* Optomechanically induced transparency in a membrane-in-the-middle setup at room temperature. *Phys. Rev. A* **88**, 013804 (2013).
36. Hou, B. P., Wei, L. F. & Wang, S. J. Optomechanically induced transparency and absorption in hybridized optomechanical systems. *Phys. Rev. A* **92**, 033829 (2015).

37. Zhang, J. Q. *et al.* Precision measurement of electrical charge with optomechanically induced transparency. *Phys. Rev. A* **86**, 053806 (2012).
38. Agarwal, G. S. & Huang, S. Optomechanical systems as single-photon routers. *Phys. Rev. A* **85**, 021801 (2012).
39. Pirkkalainen, J. M. *et al.* Cavity optomechanics mediated by a quantum two-level system. *Nature Communications* **6**, 6981 (2014).
40. Qu, K. & Agarwal, G. S. Phonon-mediated electromagnetically induced absorption in hybrid opto-electromechanical systems. *Phys. Rev. A* **87**, 031802(R) (2013).
41. Wang, H. *et al.* Optomechanical analog of two-color electromagnetically induced transparency: Photon transmission through an optomechanical device with a two-level system. *Phys. Rev. A* **90**, 023817 (2014).
42. Genes, C., Ritsch, H. & Vitali, D. Micromechanical oscillator ground-state cooling via resonant intracavity optical gain or absorption. *Phys. Rev. A* **80**, 061803(R) (2009).
43. Hammerer, K. *et al.* Optical lattices with micromechanical mirrors. *Phys. Rev. A* **82**, 021803(R) (2010).
44. Camerer, S. *et al.* Realization of an optomechanical interface between ultracold atoms and a membrane. *Phys. Rev. Lett.* **107**, 223001 (2011).
45. Bariani, F. *et al.* Hybrid optomechanical cooling by atomic Λ systems. *Phys. Rev. A* **90**, 033838 (2014).
46. Ian, H. *et al.* Cavity optomechanical coupling assisted by an atomic gas. *Phys. Rev. A* **78**, 013824 (2008).
47. Neumeier, L., Leib, M. & Hartmann, M. J. Single-photon transistor in circuit quantum electrodynamics. *Phys. Rev. Lett.* **111**, 063601 (2013).
48. Yan, C. H., Jia, W. Z. & Wei, L. F. Controlling single-photon transport with three-level quantum dots in photonic crystals. *Phys. Rev. A* **89**, 033819 (2014).
49. Zhou, L., Yang, L. P., Li, Y. & Sun, C. P. Quantum routing of single photons with a cyclic three-level system. *Phys. Rev. Lett.* **111**, 103604 (2013).
50. Lu, J., Zhou, L., Kuang, L. M. & Nori, F. Single-photon router: coherent control of multichannel scattering for single photons with quantum interferences. *Phys. Rev. A* **89**, 013805 (2014).
51. Hartmann, M. J., Brandal, F. G. S. L. & Plenio, M. B. Quantum many-body phenomena in coupled cavity arrays *Laser & photon. Rev.* **2**, 527 (2008).
52. Walls, D. F. & Milburn, G. J. *Quantum Optics* CH 7, 128–131 (Springer, Berlin, 1994).
53. Lin, Q. *et al.* Coherent mixing of mechanical excitations in nano-optomechanical structures. *Nat. Photonics* **4**, 236 (2010).
54. Zheng, C. *et al.* Controllable optical analog to electromagnetically induced transparency in coupled high-Q microtoroid cavities. *Opt. Express.* **20**, 18319 (2012).
55. Astafiev, O. *et al.* Single artificial-atom lasing. *Nature* **449**, 588 (2007).
56. Huang, S. & Tsang, M. Electromagnetically induced transparency and optical memories in an optomechanical system with N membranes. *arXiv:1403.1340v1*.
57. Tarhan, D., Huang, S. & Müstecaplıoğlu, Ö. E. Superluminal and ultraslow light propagation in optomechanical systems. *Phys. Rev. A* **87**, 013824 (2013).

Acknowledgements

This work was supported by the National Natural Science Foundation of China, under Grant No. 11375036, the Xinghai Scholar Cultivation Plan and the Fundamental Research Funds for the Central Universities under Grant No. DUT15LK35 and No. DUT15TD47. A.S. is supported by China Scholarship Council (CSC) for the Research Fellowship.

Author Contributions

A.S. conceived the idea, A.S. performed the calculations, C.-s.Y. analyzed the results and wrote the main manuscript text, Y.Z. and J.Z. participated in the discussions. All authors reviewed the manuscript.

Additional Information

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Sohail, A. *et al.* Optomechanically induced transparency in multi-cavity optomechanical system with and without one two-level atom. *Sci. Rep.* **6**, 28830; doi: 10.1038/srep28830 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>