


Antepartum Depression and Associated Factors Among Pregnant Women Attending ANC Clinics in Gurage Zone Public Health Institutions, SNNPR, Ethiopia, 2019

This article was published in the following Dove Press journal:
Psychology Research and Behavior Management

Solomon Shitu Ayen
Selamawit Alemayehu
Fentahun Tamene 

Department of Midwifery, Wolkite
University College of Health and Medical
Sciences, Wolkite University, Wolkite,
Ethiopia

Background: Antepartum depression is a form of clinical depression that can be caused by the stress and worry that pregnancy can bring a more severe level. It can be triggered by an unplanned pregnancy, a history of substance abuse, and low economic status. Antepartum depression is critical and has an intergenerational impact on children in the developing world and it can have negative effects on fetal development. Therefore, this study aimed to assess its magnitude and associated factors among pregnant women attending antenatal clinics in Gurage zone public health institutions, SNNPR, Ethiopia, 2019.

Methods: Intuition-based cross-sectional study was employed in Gurage zone public health institutions from February 14 to April 14/2019. A systematic random sampling technique was used. The data were collected and then entered into EpiData and processed by SPSS version 24.0 for analysis. P values <0.05 with 95% confidence level were used to declare statistical significance.

Results: In this study, a total of 343 pregnant mothers were participated by making a response rate of 96%. The magnitude of antepartum depression among pregnant women was 27.6% (95% CI: 22.4–33.2). The multivariable analysis showed that respondents who had unplanned pregnancy [AOR=2.11 (95% CI: 1.05–4.44)], having complications during their previous labor and delivery [AOR=4.42 (95% CI: 2.06–9.48)], previous history of child hospitalization [AOR=3.34 (95% CI: 1.48–7.51)], and satisfaction in their marriage [AOR=3.9 (95% CI: 1.15–13.21)] were associated with antepartum depression.

Conclusion: In this study, about one in four women during pregnancy develop antepartum depression. Unplanned pregnancy, complications during labor and delivery of the last baby, women having a history of child hospitalization, and maternal satisfaction with marriage were statistically associated factors with APD. To prevent further consequences, all concerned bodies need to take action by making targeted intervention and early screening of all pregnant mothers for depression.

Keywords: antepartum depression, pregnant women, Gurage, Ethiopia

Background

World health organization (WHO) defined depression as a common mental disorder that presents with depressed mood, loss of interest or happiness, decreased energy, feelings of guilt, or low self-worth, disturbed sleep or appetite, and poor concentration.^{1–3} Antepartum depression (APD) is a form of clinical depression that can be caused by the stress and worry that pregnancy can bring more severe

Correspondence: Solomon Shitu Ayen
Department of Midwifery, Wolkite
University College of Health and Medical
Sciences, Wolkite University, Wolkite 07,
Ethiopia
Tel +251924704053
Email solomonsht7@gmail.com

level. Any form of prenatal stress felt by the mother can have negative effects on various aspects of fetal development, which can cause harm to the mother and child.^{1,4,5}

Pregnancy causes significant strain on a women's body like mood changes, memory loss, eating habits, and sleeping changes.^{1,6} Symptoms of APD are; inability to concentrate, difficulty remembering, feeling emotional, extreme irritability, sleeping too much or not enough or restlessness, extreme or unending fatigue, desire to overeat or, weight loss/gain unrelated to pregnancy, loss of interest in sex, a sense of dread about everything including the pregnancy, feelings of guilt, persistent sadness, thoughts of suicide, or death.^{2,7,8}

Antepartum depression can be caused by various reasons including a combination of hormonal changes and psychological disturbances associated with pregnancy. Relationship problems, family or personal history of depression, maternal anxiety, life stress, and lack of social support, unintended pregnancy, domestic violence, low income, single status, previous pregnancy loss are some of the causes.^{2,9,10} Untreated depression in pregnancy has been also associated with poor pregnancy and birth outcomes such as pre-eclampsia, low birth weight, increased risk of premature delivery, postpartum depression.^{5,11,12}

Antepartum depression contributes to 1/3 of work absence due to diseases and 1/5 of all primary care visits. It is estimated that 25–35% of pregnant women have depressive symptoms and that 20% of them may meet the diagnostic criteria for major depression. According to the WHO, depression is projected to become the second leading contributor to the global burden of disease by 2020.^{1,13–16}

In terms of regional distributions, high rates of mental health problems in pregnant women and mothers have been reported from Africa. One in three to one in five women in developing countries, and about one in ten in developed countries, and 13% of new mothers experience depression.^{17–20}

Even though APD is the health problem of the mothers, as knowledge of the investigators, there is limited information in southern Ethiopia and our locality. So, this study is aimed to show the magnitude and factors associated with APD among pregnant mothers who attend ANC in Gurage zone public health institutions, Ethiopia.

Methods

Study Design, Area, and Period

An institution-based cross-sectional study was conducted from February 14 to April 14/2019 in Gurage zone public

health institutions. The Gurage zone is one of the administrative zones in South Ethiopia. This is located southwest of Addis Ababa. The capital city of the Gurage zone is Wolkite which is located 153km from Addis Ababa capital city of Ethiopia. It has 16 districts and 5 town administrations. There are seven hospitals (five public and two non-governmental) and 72 health centers serving the total population in the zone.

Source Population

All pregnant women who visit ANC clinic in Gurage zone public health institutions.

Study Population

All pregnant women who attend ANC clinic in Gurage zone public health institutions during the study period.

Inclusion Criteria

All pregnant women attending ANC clinic in Gurage zone public health institutions during the data collection period were included.

Exclusion Criteria

Pregnant women who were seriously ill and unable to respond during the data collection period.

Sample Size Determination

The sample size was determined by using a single population proportion formula by considering assumptions of the proportion of depression 31.1% in a study done in Maichew.³⁵ A design effect of 5% at a 95% CI.

Considering a 10% non-response rate, the total study subjects were $328+33=361$ pregnant women.

A systematic random sampling technique was used to select pregnant women coming to the Gurage zone public health institutions during the data collection period.

Data Collection Method

The interviewer-administered questionnaire was prepared in such a way that it includes socio-demographic and economic characteristics, obstetric variables, sex of the last baby, desired sex of the last baby, number of living children by reviewing relevant literature. Before interviewing data collectors gave information about the aim of the study, purposes, possible risks, and benefits, the right and refusal of mothers, and the confidentiality issues. The data were collected by 8 data collectors and supervised by 2 health professionals.

The data collectors were trained for summation of the result and cut off point whether the women have depressive symptoms or not then if she has depressive symptoms data collectors were informed for health professionals working at ANC unit confidentially and they council her and appoint after a week to see changes.

Dependent Variables

Antepartum depression.

Independent Variables

Social demographic variable (age, income, educational status of mothers, husbands' educational status).

Obstetric variable (unplanned pregnancy, mode of delivery, complications during labor and delivery).

Substance abuse (smoke during pregnancy and before pregnancy, khat chewing).

Previous psychiatric history, violence, Social support.

Operational Definitions

Depressed: Those pregnant mothers who score ≥ 13 cuts off points of EPDS. From ten EPDS questions each of which has four choices resulting maximum score of 30 and minimum 0.³⁶

Not Depressed: Those pregnant mothers who score <13 cuts off point of EPDS. From ten EPDS question's each of which has four choices resulting maximum score of 30 and a minimum of 0.³⁶

Data Quality Control

The questionnaire was designed first in the English language and it was translated to local language Amharic language by a translator and again it was translated back to English. The questionnaire was adopted from different studies in the locality but not validated. A pretest was conducted on 5% of the total sample size at one health center which is not selected as a study area by data collectors and then the questionnaire was assessed for its clarity and a necessary correction was done accordingly. During data, collection data collectors were supervised by supervisors, and overall activities were controlled by principal investigators, and finally after data collection before entry all collected data were checked for completeness.

Data Processing and Analysis

Data were coded, cleaned, edited, entered into EpiData version 3.1 to minimize logical errors, and then the data was exported to SPSS version 24 for analysis. Descriptive

statistical analysis was carried out to compute frequency and percentage and presented by text and tables. Binary logistic regression analysis was used to ascertain the association between explanatory and outcome variables. Variables with significant ($P < 0.25$) association in the bivariate analysis were entered into multivariate analysis and those variables with $P < 0.05$ in multivariate analysis were considered to be statistically significant. Finally, the result of the study was presented in texts, tables and it was discussed using the odds ratio and 95% confidence level.

Results

Socio-Demographic Characteristics

In this study, a total of 343 pregnant mothers were participated by making a response rate of 96%. The majority 141 (41%) of respondents were in the age range between 25–29 years old with a mean of 27.3 (SD \pm 5.6) years. Out of the total respondents, 142 (41.4%) were Orthodox followed by Muslim 84 (24.5), protestant 65 (19%), and catholic 52 (15.2%). About the educational status of the women 209 (60%) was educated above secondary school. The marital status of the participants 282 (82.2%) was married. More than half (60.0%) of the respondents were a housewife and 187 (54.5%) were urban residents. One hundred thirty-one (38%) of the husband's occupation were a merchant. About three fourth 225 participants monthly income was ≤ 1000 (Table 1).

Obstetric Characteristics of Pregnant Mothers

Of respondents, 116 (33.8%) were get pregnant two times. Sixty-three percent of the participant's current pregnancy was planned. More than three fourth of participants had no history of abortion. Nearly two-thirds of the participant's previous baby's sex was male. Sixty-five percent of the respondent's place of delivery for the previous pregnancy was spontaneous delivery (Table 2).

Substance Abuse of Pregnant Women

In this study, 81 (23.6%) of respondents had drunk alcohol before pregnancy, whereas 73 (21.3%) respondents had a history of drinking alcohol during pregnancy. Twenty-nine (8.45%) respondents had a history of smoking before or during pregnancy.

Psychosocial History

More than half (53.2%) of respondents had good support from their husbands, most 136 (54.4%) of the respondents

Table 1 Sociodemographic Characteristics of Pregnant Women Who Attained Antenatal Care Services in Gurage Zone Public Health Institutions, 2019 (n=343)

Variables	Category	Frequency	Percent
Age	≤19	18	5.2
	20–24	91	26.6
	25–29	141	41.1
	30–34	53	15.5
	35–39	23	6.7
	≥40	17	5
Religion	Orthodox	142	41.4
	Catholic	52	15.2
	Protestant	65	19
	Muslim	84	24.5
Marital status	Married	282	82.2
	Single	19	5.5
	Divorce	14	4.1
	Widowed	28	8.2
Educational status of the mother	No formal education	47	13.7
	Primary school (1–8)	87	25.4
	Secondary school(9–12)	137	39.9
	Diploma and above	72	21
Occupation of the mother	Housewife	208	60.6
	Merchant	56	16.3
	Farmer	33	9.6
	Daily labor	23	6.7
	Student	23	6.7
Husbands educational level	No formal education	40	11.7
	Primary school (1–8)	72	21
	Secondary school(9–12)	99	28.9
	Diploma and above	71	20.7
Occupation of the mother	Gov.t worker	131	38
	Merchant	63	18.4
	Farmer	49	14.3
Daily labor Household income	39	11.4	
	≤1000	225	74.3
	≥2000	88	25.8

had satisfied with their relationship with the husband's family. More than half (63.2%) of the respondents had no previous history of depression whereas, 84 (24.5%) of

Table 2 Obstetric Characteristics of Pregnant Women Who Attained Antenatal Care Services in Gurage Zone Public Health Institutions, 2019 (n=343)

Variables	Category	Number	Percent
Number of gravidities	One	67	19.5
	Two	116	33.8
	Three	61	17.8
	Four and above	99	28.9
Number ANC follow-ups on current pregnancy	First visit	76	22.2
	Second visit	145	42.3
	Third visit	100	29.2
	Fourth visit	22	6.4
Number of alive children (n=277)	One	65	23.5
	Two	100	36.1
	Three and above	112	40.4
History of abortion	Yes	62	18.1
	NO	281	81.9
Sex of last baby (n=277)	Male	173	62.5
	Female	104	37.5
Desired sex of the last baby (n=277)	Male	165	59.6
	Female	112	40.4
Mode of delivery for your last pregnancy (n=277)	Vaginal	207	74.7
	C/section (C/S)	27	9.7
	Instrumental delivery	43	15.5
Any complication during delivery (n=277)	Yes	81	29.2
	No	196	70.8
Any of your children hospitalized (n=277)	YES	50	18
	NO	227	82
Was your pregnancy planned	Yes	237	69.1
	NO	106	30.9

participants had practiced violence by their husband from those 54.8%, 20.2% and 25% of respondents had experienced verbal, physical, and both (verbal and physical) abuse in their home respectively (Table 3).

Magnitude of Depression

In this study, the proportion of women who had APD was 91 (26.5%) with 95% CI: 22.0–31.3.

Factors Associated with Antepartum Depression

The result revealed that age, educational level of the women, unplanned pregnancy, mode of delivery,

Table 3 Psychosocial History of Pregnant Women Who Attained Antenatal Care Services in Gurage Zone Public Health Institutions, 2019 (n=343)

Variables	Category	Number	Percent
Any of your relatives suffered from mental illness	Yes	91	16.6
	No	252	73.5
Have you ever experienced any abuse in your home	Yes	84	24.5
	No	259	75.5
What kind of abuse do ever experience (n=84)	Verbal	46	54.8
	Physical	17	20.2
	Verbal and physical	21	25
Are you satisfied with your marriage	Yes	186	54.2
	No	58	16.9
	More or less	99	28.9
Does your husband support you and your pregnancy	Yes	184	53.6
	No	78	22.7
	More or less	81	23.6
Are you satisfied with your relationship with your husband's family	Yes	186	54.1
	No	58	16.9
	More or less	99	29

complication during labor and delivery, child hospitalization, previous history of depression, maternal satisfaction with marriage, violence, and smoking during pregnancy were considered for the multivariate logistic regression model. In multivariable logistic regression unplanned pregnancy, complications during labor and delivery, child hospitalization, and maternal satisfaction with mirage were significantly associated with APD.

The odds of depression are increased by 2.11 times for women who have an unplanned pregnancy than their counterparts. Those pregnant women who have a previous history of any complication during labor and delivery were about four times more at risk of depression than women who had no previous history of complications. Those women who have a history of their child hospitalized were 3.34 times the risk of developing depression than their counterparts and those pregnant women who were not satisfied with their mirage were also 3.9 times more likely to have increased the risk of depression than those women who were satisfied in their marriage (Table 4).

Discussion

This study showed that 26.5% of respondents had antepartum depression with 95% CI: 22.0–31.3. Unplanned

pregnancy, complications during delivery, child hospitalization, and maternal satisfaction with mirage were significantly associated with APD.

In this study 26% of respondents suffer from antepartum depression, the finding was in line with the study done in Gondar 25.8%.^{26,33} Whereas, the finding of the study is lower than the study done in Maichew (31.1%) North Ethiopia.^{21,35} The difference might be due to the usage of different tools and sample size. Both studies used a self-reporting questioner of 20 yes-no questions. On the other hand, this figure was higher when compared to other similar studies done in Brazil (14.2%) Turkey (10.9%), Debra tabor (11.8%), and Addis Ababa (11.7%).^{23,24,28,31,32,36} This discrepancy might be due to the different tools, assessment period, methods, and the difference in the study area.

This study found that antepartum depression was four times higher among women who are not satisfied with their marriage than those who were satisfied with marriage AOR 3.9, (95% CI: 1.15–13.21). This result is consistent with the study conducted in Gondar; the Amhara region in Ethiopia.^{19,22,30,33} This might be since during the period of pregnancy and delivery women were handling the situation alone and did not have the needed support from their partner.

Unplanned pregnancy was also associated with higher odds of having postpartum depression AOR 2.11 (95% CI: 1.0, 4.44). This result was also observed in most of the previous studies carried out in Turkey and northwestern Brazil and Addis Ababa Ethiopia.^{25,28,31,36} This may be because the woman with an unplanned pregnancy may be related to the economic problem and she became stressed due to how she gives care for the newborn and health care.

The women with a previous history of complications during delivery were 4.4 times developed APD than those counterparts AOR 4.42 (95% CI: 2.06–9.48). This result is consistent with the study conducted in North Ethiopia Maichew.^{29,34,35} This might be because negative life events in the previous history are most influential on an individual's mental status it also might be because they are frightened they might face the same problem on their current labor and delivery as well.

Participants whose baby hospitalized were three times develop APD than those who were not hospitalized AOR=3.34 (95% CI: 4.8, 7.51). The finding was similar to studies conducted in Brazil, and Debar tabor in Ethiopia.^{27,28,32} This might be due to women who had experienced child hospitalization problems before was

Table 4 Factors Affecting Pregnant Women Who Attended the ANC Clinic in Gurage Zone Public Health Institutions, 2019 (n=343)

Types of Variables		Depression		OR(95% CI)		
		Depressed%	Not Depressed%	COR	AOR	P-Value
Age	≤19	6(6.6)	12(4.8)	1		
	20–24	20(22)	71(27.2)	0.16(-1.89–0.91)	0.06(-0.99–5.22)	0.6
	25–29	33(36.3)	108(42.9)	0.82(-1.19–1.41)	0.08(-2.12–6.45)	2.2
	30–34	17(18.70)	36(14.3)	0.4(-2.61–0.98)	0.32 (-5.63–8.56)	0.39
	35–39	6(6.6)	17(6.7)	0.26(-0.59–2.81)	0.15(-1.24–2.89)	5.87
	≥40	9(9.9)	8(3.2)	0.15(-2.36–0.12)	0.45(0.02–15.36)	0.91
The educational level of women	No formal education	16(17.6)	31(12.3)	0.88(0.38–2.05)	1.29(0.40–4.14)	0.670
	Primary school(1–8)	21(23.1)	66(26.2)	0.376(0.17-0.84)	0.44(0.14–1.38)	0.158
	Secondary school	26(28.6)	111(44)	0.258(0.12-0.55)	0.53(0.19–1.53)	0.243
	Diploma and above	28(30.8)	44(17.5)	1	1	
Was the pregnancy Planned?	Yes	57(62.6)	180(71.4)	1	1	
	No	34(37.4)	72(28.6)	2.46(1.29–4.66)	2.11(1.0–4.44)*	0.049
Mode of delivery of last baby (n=277)	Vaginal	45(61.6)	162(79.4)	1	1	
	C/section(C/S)	11(15.1)	16(7.8)	2.046(0.79–5.32)	1.41(0.44–4.52)	0.569
	Instrumental	17(23.3)	26(12.7)	2.28(1.05–4.97)	1.13(0.43–2.99)	0.805
Any complication during delivery (n=277)	Yes	35(47.9)	46(22.5)	3.47(1.8–6.67)	4.42(2.06–9.48)*	0.000
	No	38(52.1)	158(77.5)	1	1	
Previous history of depression	Yes	53(58.2)	72(28.2)	1.61(0.91–2.83)	0.67(0.32–1.43)	0.302
	No	38(41.8)	180(71.4)	1	1	
Any children hospitalized (n=277)	Yes	21(28.8)	86(42.2)	1.83(1.94–3.59)	3.34(1.48–7.51)*	0.004
	No	52(71.2)	118(57.8)	1	1	
Any abuse in her home	Yes	27(29.7)	57(22.6)	1.55(0.84–2.86)	1.02(0.42–2.48)	0.970
	No	64(70.3)	195(77.4)	1	1	
Satisfaction on marriage	Yes	48(52.7)	138(54.8)	1	1	
	No	22(24.2)	36(14.3)	0.33(0.16-0.67)	0.49(0.18–1.35)	0.171
	More or less	21(23.1)	78(31)	1.38(1.52–3.65)	3.9(1.15–13.21)*	0.029
Smoking during pregnancy	Yes	3(3.3)	26(10.3)	0.15(0.02–1.17)	0.38(0.05–3.19)	0.371
	No	88(96.7)	226(89.7)	1	1	

Note: *statistically significant at p-value ≤0.05.

that, they might stress themselves thinking about the current baby might face the same problem. And worrying themselves that, there could be a chance they lose their baby.

Using a cross-sectional study design that hinders the researcher from establishing cause and effect relationship between the possible determinant of antepartum depression and the outcome variable and using EPDS may expose participants to bias.

Conclusion

More than one in four pregnant women develops APD from those who receive ANC in Gurage zone public health institutions. Unplanned pregnancy, complications during labor and delivery of the previous baby, women having

a history of child hospitalization, and maternal satisfaction with marriage were statistically associated factors that determine antepartum depression. This implies APD is a concern of government and other stakeholders to work on this to perform the regular preventive screening during antenatal follow-ups for those pregnant women.

Limitation of the Study

A cross-sectional study design hinders the researcher from establishing cause and effect relationship between the possible determinants and the outcome of interest.

Institutional study design may not show the real prevalence because there are mothers who did not visit health institution for ANC so minimize the real incidence.

Abbreviations

ANC, Antenatal Care; AOR, Adjusted Odds Ratio; APD, Antepartum depression; CI, Confidence Interval; EPDS, Edinburg postnatal depression; PI, Principal investigator; SPSS, Statistical package for social sciences; WHO, World health organization.

Data Sharing Statement

The datasets used and/or analyzed during the current study available from the corresponding author on reasonable request.

Ethical Consideration and Consent to Participate

Ethical clearance was obtained from Wolkite University, College of Health and Medical Sciences, Institutional Health Research Ethics Review Committee (IHRERC). A formal letter for permission and support was written to the zonal health department of Gurage from Wolkite University and official permission to undertake the study was obtained and permission to conduct the study was asked. The respondents were informed about the objective, purpose, risks, and benefits of the study and the right to refuse to participate, and then informed written and signed consent was taken. The study posed a low or no more than minimal risk to the study participants. Also, the study did not involve any invasive procedures. Moreover, the confidentiality of information was guaranteed by using code numbers rather than personal identifiers and by keeping the data locked. The study was conducted in accordance with the Declaration of Helsinki.

Consent for Publication

Not applicable.

Declaration

The content of the study is solely the responsibility of the authors.

Acknowledgment

We would like to thank Wolkite University for giving technical support to prepare this research.

We would like to forward our deepest appreciation to in Gurage zone public health office and staff for their special support. Also, special gratitude goes to data collectors and our study participants.

Author Contributions

All authors made a significant contribution to the work reported, whether that is in the conception, study design, execution, acquisition of data, analysis, and interpretation, or in all these areas; took part in drafting, revising, or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted; and agree to be accountable for all aspects of the work.

Funding

There is no source of funding for this research. All costs were covered by researchers.

Disclosure

The authors would like to declare that we have no conflicts of interest for this work.

References

1. World health organization and World organization of family doctors- (Wonca). *Integrating Mental Health into a Primary Care-A Global Perspective*. Geneva, Switzerland: WHO; 2008.
2. Gelaye B, Lemma S, Deyassa N, et al. Prevalence and correlates of mental distress among working adults in Ethiopia. *Clin Pract Epidemiol Mental Health*. 2012;8(1):126–133.
3. Kastrup CM. Global mental health secondary publication. *Dan Med Bull*. 2007;54(1):42–43.
4. Antenatal depression. 2013. Available from: www.nct.org.uk. Accessed December 29, 2020.
5. Gurage zone socio-Economic profile Archived 2007-10-07at the Wayback Machine Gurage zone government webpage
6. Prevalence of Anxiety, Depression and Associated Factors among Pregnant Women of Hyderabad, Pakistan NIHPA Author Manuscripts 2009; 55(5).
7. Alem A, Araya M, Melaku Z, Wendimagegn D, Abdulahi A. Mental distress in medical students of Addis Ababa University. *Ethiop Med J*. 2005;43(3):159–166.
8. Dessa N, Berhane Y, et al. Intimate partner violence and depression among women in rural Ethiopia: a cross-sectional study. *Clin Pract Epidemiol Mental Health*. 2009;5:8. doi:10.1186/1745-0179-1185-1188
9. Vesga-Lopez, et al. Antepartum depression in Brazile. *Tealilo Aende Psychiatry*. 2013;65(7):805.
10. Murray J, Lopez AD. *The Global Burden of Disease: A Comprehensive Assessment of Mortality Disability from Disease, Injuries, and Risk Factors in 1990 and Projected to 2020*. Boston: Harvard School of Public Health, World Health Organization; 1996.
11. Tesfaye A. Prevalence and correlates of mental distress among regular undergraduate students ofHawassa University: a cross-sectional survey. *East Afr J Public Health*. 2009;6(1):85–94. doi:10.4314/eajph.v6i1.45755
12. Bennett HA, Einarson A, Taddio A, Koren G, Einarson TR. Prevalence of depression during pregnancy: a systematic review. *Obstet Gynecol*. 2004;103:698–708. doi:10.1097/01.AOG.0000116689.75396.5f
13. Kebede. D, Alem. A, Rashid E. The prevalence and socio-demographic correlates of mental distress in Addis Ababa, Ethiopia. *ActapsychiatricaScandinevica*. 1999;100(Suppl.397):5–10.

14. Audu MD, Tesfaye M, Gurmu S, Mela M. Lifetime prevalence of substance abuse and mental distress among homicide offenders in Jimma prison, Ethiopia. *Afr J Appl Psychol*. 2008;2(3):15–26.
15. Alem A, Kebede D, Woldeesmiat G, Jacobsson L, Kullgren G. The prevalence and socio-demographic correlates of mental distress in Butajira, Ethiopia. *Acta Psychiatr Scand Suppl*. 1999;(397):48–55.
16. Rich-Edwards JW, Kleinman K, Abrams A, et al. Sociodemographic predictors of antenatal and postpartum depressive symptoms among women in a medical group practice. *J Epidemiol Community Health*. 2006;60(3):221–227. doi:10.1136/jech.2005.039370
17. BizuGelaye S, Lemma N, YonasBahretibeb M, et al. Prevalence and correlates of mental distress among working adults in Ethiopia. *Clin Pract Epidemiol Mental Health*. 2012;8:126–133.
18. Demena T, Mossie A, Tesfaye M. Khat chewing and mental distress: a community based study, in Jimma City, Southwestern Ethiopia. *Ethiop J Health Sci*. 2011;21(1):37–45. doi:10.4314/ejhs.v21i1.69042
19. Fekadu A, Alem A, et al. Utility of the concept of minor depressive disorder: evidence from a largely rural community sample in a developing country setting. *J Affect Disord*. 2007;104(1–3):111–118. doi:10.1016/j.jad.2007.03.008
20. Dessa N, Berhane Y, et al. Depression among women in rural Ethiopia as related to socioeconomic factors: a community-based study on women in reproductive age groups. *Scand J Public Health*. 2008;36(6):589–597. doi:10.1177/1403494808086976
21. Marcus M, TaghiYasamy M, van Ommeren M, Chisholm D, Saxena S. *Depression: A Global Public Health Concern*. WHO Department of Mental Health and Substance Abuse; 2012.
22. Kalifa, D S., Glavin, K., Bjertness, E., et al. Antepartum depression among Turkey woman. *Int J Women's Health*. 2015;7:677–684. doi:10.2147/IJWH.S81401
23. Hailemariam S, Tessema, F., Asefa, M. et al. The prevalence of depression and associated factors in Ethiopia: findings from the National Health Survey. *Int J Ment Health Syst*. 2012;6:23. doi:10.1186/1752-4458-6-23
24. World Health Organization, World suicide prevention day 2012. Available from: http://www.who.int/mediacentre/events/annual/world_suicide_prevention_day/en/. Accessed December 29, 2020.
25. Pantelis C, Hindler CG, Taylor JC. Use and abuse of khat (*Catha edulis*): a review of the distribution, pharmacology, side effects and a description of psychosis attributed to khat chewing. *Pharmacol Med*. 1989;19:657–668.
26. Wilson Pamela”Antenataldepression” health.ninemsn.com 4 April 2013
27. Wilder P, Mathys K, Brenneisen R, Kalix P, Fisch HU. Pharmacodynamics and pharmacokinetics of khat: a controlled study. *Clin Pharmacol Ther*. 1994;5595:556–562.
28. Weyerer S, Gorfer ES, Köhler L, et al. Prevalence and risk factors for depression in non-demented primary care attenders aged 75 years and older. *J Affect Disord*. 2008;111:153–163. doi:10.1016/j.jad.2008.02.008
29. Vink D, Aartsen JM, Schoevers AR. Risk factors for anxiety and depression in the elderly: A review. *J Affect Disord*. 2008;106:29–44. doi:10.1016/j.jad.2007.06.005
30. Teixeira C, Figueiredo B, Conde A, Pacheco A, Costa R. Anxiety and depression during pregnancy in women and men. *J Affect Disord*. 2009;119:142–148. doi:10.1016/j.jad.2009.03.005
31. Pereira PK, Lovisi GM, Pilowsky DL, Lima LA, Legacy LF. Depression during pregnancy: prevalence and risk factors among women attending a public health clinic in Rio de Janeiro, Brazil. *Cad Saude Publica*. 2009;25:2725–2736. doi:10.1590/S0102-311X2009001200019
32. Karacam Z, Ance G. Depression, anxiety, and influencing factors in pregnancy: a study in a Turkish population. *Midwifery*. 2009;25:344–356. doi:10.1016/j.midw.2007.03.006
33. Bisetegn TA, Mihretie G, Muche T. Prevalence and predictors of depression among pregnant women in Debretabor town, northwest Ethiopia. *PLoS One*. 2016;11(9). doi:10.1371/journal.pone.0161108
34. Ayele TA, Azalea T, Alemu K, Abdissa Z, Mulat H, Fekadu A. Prevalence and associated factors of antenatal depression among women attending antenatal care service at Gondar University Hospital, Northwest Ethiopia. *PLoS One*. 2016;11:5. doi:10.1371/journal.pone.0155125
35. Tilahun B. Prevalence of antenatal depression and associated factors among pregnant women in Maichew, North Ethiopia. 2016;11:2.
36. Demewoze H Prevalence and associated factors of antenatal depression among women attending antenatal care service at Addis Ababa Ethiopia.

Psychology Research and Behavior Management

Dovepress

Publish your work in this journal

Psychology Research and Behavior Management is an international, peer-reviewed, open access journal focusing on the science of psychology and its application in behavior management to develop improved outcomes in the clinical, educational, sports and business arenas. Specific topics covered in the journal include: Neuroscience, memory and decision making; Behavior modification and management; Clinical

applications; Business and sports performance management; Social and developmental studies; Animal studies. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/psychology-research-and-behavior-management-journal>