

Research Article

Reactive Oxygen Species Mediated Prostaglandin E₂ Contributes to Acute Response of Epithelial Injury

Yi-Ping Hu,^{1,2} Yin-Bo Peng,¹ Yi-Fan Zhang,¹ Ying Wang,¹
Wei-Rong Yu,¹ Min Yao,¹ and Xiu-Jun Fu¹

¹Department of Burns and Plastic Surgery, No. 9 People's Hospital, and Institute of Traumatic Medicine, Shanghai Jiao Tong University School of Medicine, Shanghai 201900, China

²Department of Burns and Plastic Surgery, Guangzhou Red Cross Hospital, Jinan University, Guangzhou 510220, China

Correspondence should be addressed to Xiu-Jun Fu; fxver159@yahoo.com

Received 22 October 2016; Revised 20 December 2016; Accepted 28 December 2016; Published 9 February 2017

Academic Editor: Liang-Jun Yan

Copyright © 2017 Yi-Ping Hu et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Reactive oxygen species (ROS) generated after tissue injury play a crucial role during wound healing through initiating acute inflammation, clarifying infection and dead tissue, and mediating various intracellular signal transduction. Prostaglandin E₂ (PGE₂) has been identified as one of the major factors responsible for inflammation and tissue repair. In this study, we tested our hypothesis that ROS produced by damaged human keratinocytes induces the synthesis of PGE₂. In vitro epithelial wounding model was used to observe the production of ROS and secretion of PGE₂ as well as the involved signal pathway. The mechanical injury caused the rapid production of ROS in in vitro cultured keratinocytes, which was significantly blocked by an inhibitor of nicotinamide adenine dinucleotide phosphate oxidase. The increased intracellular ROS caused by mechanical injury stimulates PGE₂ production in a time-dependent manner via the activation of cyclooxygenase-2 (COX-2), which was stimulated by phosphorylation of extracellular signal-regulated protein kinase (ERK). These results indicate ROS-induced ERK activation leading to the activation of COX-2 and the synthesis of PGE₂ in human keratinocytes responding to mechanical injury in the acute phase.

1. Introduction

Cutaneous wound healing is a complex yet well-organized process, which includes three interactive phases of inflammation, proliferation, and tissue reconstruction. Injury, as the signal for initiating wound healing, triggers the process of tissue repair by activating the directly or indirectly wound-involved cells, for example, epithelial cells, endothelial cells, fibroblasts, and inflammatory cells. The traditional concept widely accepts that the chemotaxis of leukocytes to the wound site successively follows the activation of hemostasis after injury, and reactive oxygen species (ROS) are mainly produced by leukocytes through “respiratory burst” for disinfection and debridement. However, a recent study revealed that a fast and dramatic increase of ROS at the wound margin was essential for rapid recruitment of leukocytes to the wound site [1]. ROS produced by injured epithelial cells through the activation of nicotinamide adenine dinucleotide phosphate (NADPH) oxidase (NOX) play a critical

role in initiating inflammatory response of wound healing. Additional studies also show that, in contrast to oxidative damage at high concentration, ROS serve as intracellular and extracellular signaling messengers and regulate numerous downstream signal transduction and gene expression at low concentration [2, 3]. ROS have been determined to be able to activate mitogen-activated protein kinases (MAPKs) cascades [4–7]. Comprised of extracellular signal-regulated protein kinase (ERK), c-Jun N-terminal kinase (JNK), and p38 kinase, MAPKs are the core molecules of cell stress-response signaling network. MAPKs are protein kinases specific to the amino acids serine, threonine, and tyrosine, which are involved in directing the cellular response to a diverse array of stimuli including mechanical damage. Activated by the injury signals, MAPKs coordinate cell functions including proliferation, differentiation, migration, survival or apoptosis, and gene expression during wound healing [6, 8, 9].

Cyclooxygenase (COX) enzyme and its enzymatic product prostaglandin E₂ (PGE₂) are known to be critical inflammatory factors in the early phase of wound healing [10]. Among the two isoforms of COX, COX-1 and COX-2, COX-1 is expressed constitutively in most tissues and may be responsible for housekeeping functions. In contrast, COX-2 is not detectable in most normal tissues, but its expression can be induced by endotoxin, cytokines, growth factors, and carcinogens [11, 12]. PGE₂ is believed to be strongly associated with the signs of redness, swelling, heat, and pain in the wound area, which indicates its role as a pivotal proinflammatory factor. It has been confirmed that PGE₂ has a profound influence on wound repair by affecting proliferation and migration of epithelial cells, vascular tone, regional blood flow, vascular permeability and remodeling, and angiogenesis [13, 14].

As ROS have been demonstrated to activate many intracellular molecules and upregulate oxidative stress-related genes during wound healing, we hypothesized that ROS might modulate the production of PGE₂ through MAPKs pathway on mechanical damaged human keratinocytes, and the increased PGE₂ might then be responsible for cell injury and repair and inflammatory cell recruitment. In the present study, we sought to investigate the effects of ROS on PGE₂ production in human keratinocytes after mechanical injury, as well as the possible mechanisms.

2. Materials and Methods

2.1. Cell Culture and Treatment. The immortalized human skin keratinocyte cell line, HaCat, from the Institute of Basic Medical Sciences, Chinese Academy of Medical Sciences, was grown in RPMI-1640 medium (Gibco, BRC) supplemented with 10% FBS at 37°C with 5% CO₂. Cells were seeded at 6-well plates or 60 mm dishes for further experiments. After 48 hr of incubation, cells grew to 70%–80% confluence, and the medium was replaced with RPMI-1640 containing 1% FBS for overnight before the indicated treatment. The undamaged cells worked as control. When assessing the effects of ROS inhibitor, ERK inhibitor, or COX-2 inhibitor, cells were pretreated with diphenyleiiodonium (DPI) (Sigma, MO, USA) for 30 min, PD98059 (Beyotime, Shanghai, China) for 1 hr, or NS398 (Beyotime, Shanghai, China) for 1 hr, respectively, before injury.

2.2. In Vitro Wounding Model. Freshly isolated HaCat cells growing to 70%–80% confluence in 6-well plate or 60 mm dish were used for wounding model. Briefly, cells were mechanically removed from the plate or dish by dragging a 20 μ L pipette tip linearly on the confluent monolayer as the plate rested over a template. The spacing between the scratching lines was 3 mm.

2.3. ROS Detection. Cells grown in 60 mm culture dishes were incubated for 15 min at 37°C in dark with 10 μ M of carboxy-H₂DCFDA (Invitrogen, CA, USA). The cells were then washed three times with Phosphate-Buffer Saline (PBS) and exposed with scratching injury for indicated time at 37°C. When assessing the effect of ROS inhibitor, cells

were pretreated with 10 μ M DPI for 30 min. The cells were harvested and centrifuged for 5 min at 10,000 rpm at 4°C to remove the supernatants. After the pellets were resuspended in PBS, levels of ROS were detected using a fluorospectrophotometer (Molecular Devices, USA) with 480 nm excitation and 530 nm emission.

2.4. Western Blot Analysis. Cells were washed with PBS, harvested using a scraper, and solubilized in cold cell lysis buffer (Beyotime, Shanghai, China). Aliquots of lysate were heated for 5 min at 95°C. Equal amounts of lysate were subjected to SDS-polyacrylamide gel electrophoresis on 10% gels and were transferred onto a PVDF membrane (Millipore, USA). The membranes were blocked with 5% nonfat dry milk in 0.01 M Tris-buffered saline (PH 7.4) containing 0.05% Tween-20 (TBST) at room temperature for 1 hr. The membranes were then incubated with primary antibodies of ERK, phosphorylated ERK (p-ERK) rabbit monoclonal antibody (mAb), or COX-2 rabbit mAb (1:1000) (Cell Signaling Technology, MA, USA) overnight at 4°C after they were incubated with appropriate HRP-conjugated secondary antibodies. The protein bands on the blots were detected with enhanced chemiluminescence detection kit (Thermo Scientific, IL, USA) according to the manufacturer's instructions.

2.5. COX Activity Assessment. More than 1×10^9 cells in culture were harvested using a scraper and resuspended in 100 μ L of a cell lysis buffer containing 1 mM EDTA. The lysed cells were centrifuged at 2000 rpm at 4°C for 10 min, and the supernatant was discarded. Cell pellets were resuspended in 400 μ L of a lysis buffer containing 0.1 M Tris-HCl, 1 mM EDTA, homogenized with ice ultrasonic for 30 s, and then centrifuged at 10000 rpm at 4°C for 15 min. The supernatant was assayed with the COX fluorescence activity assay kit (Cayman Chemical CO, MI, USA), following the manufacturer's instructions.

2.6. Measurement of PGE₂ Release. Supernatants of HaCat culture were collected and concentrated by centrifugation. The concentration of PGE₂ was determined by PGE₂ ELISA Kit (Cayman Chemical Co., MI, USA), according to the manufacturer's instructions.

2.7. Statistical Analysis. Results were expressed as means \pm SEMs. Statistical analysis was performed using Student's *t*-test and ANOVA. Those *p* values that were less than 0.05 were considered statistically significant.

3. Results

3.1. Scratching Injury Induces Generation of ROS. In order to understand the effects of ROS on epithelial wound healing, the levels of intracellular ROS in HaCat cells were measured after injury. HaCat cells were injured by manual scratches, and the assessments of ROS levels were made using fluorescence probe carboxy-H₂DCFDA, which can be oxidized to fluorescent fluorescein by ROS in cells (Figure 1(a), center column). As shown in Figure 1, scratching injury caused a




FIGURE 1: Rapid formation of ROS induced by scratching injury in HaCat cells. (a) The intracellular ROS were visualized by the reaction with probe carboxy-H₂DCFDA under fluorescent microscopy. Cells without damage worked as negative control, while H₂O₂ treated cells were used as positive control. (b) The quantities of ROS were shown after scratching for undamaged control cells, damaged cells, and damaged cells plus DPI treatment. Data are representative of three independent experiments with triplicate samples. * and # indicate $p < 0.05$ compared with undamaged control cells and with scratched cells, respectively.

rapid increase in intracellular ROS (Figure 1(a), right column) in a time-dependent manner with a maximal response within 30 min (3.5-fold) and sustained to 60 min (3-fold) (Figure 1(b)). In addition, pretreatment with a NOX inhibitor, DPI, for 30 min dramatically blocked the scratching-induced production of ROS (1.8-fold after 30 min and 1.5-fold after 60 min) in HaCat cells (Figure 1(b)). These data suggest that mechanical injury can induce rapid production of ROS during wound healing.

3.2. Scratching Injury Increases Synthesis of PGE₂ via ROS. Recent evidence has shown that ROS have multiple downstream targets and play a key role in triggering production of proinflammatory factors, such as PGE₂. Thus, to determine whether there is change of the level of PGE₂ after epithelial damage, we investigated the release of PGE₂ after scratching injury on HaCat cells in vitro. The level of PGE₂ was significantly increased at 6 hr after injury and remained greater than the undamaged control for at least 24 hr in HaCat after injury (Figure 2(a)).

In order to know the relationship between the generation of ROS and the secretion of PGE₂ after scratching injury, the inhibitor of NOX, DPI, was used to pretreat the cells, and then the levels of PGE₂ were measured. The level of PGE₂ in the DPI treatment plus scratching injury cells was significantly lower than that in the injury only cells at 4 hr or 6 hr after injury, respectively (Figure 2(b)). Our data indicate that scratching injury-induced PGE₂ synthesis is mediated by ROS.

3.3. ERK Activation Is Involving in Scratching-Injury-Induced Enhancements of ROS and PGE₂. ROS have also been reported to play a major role as second messengers and to

contribute to the activation of signaling pathway. In an effort to understand the mechanism underlying the injury-induced generation of ROS and synthesis of PGE₂, ERK activation after scratching injury was investigated. As illustrated in Figure 3(a), scratching injury increased the extent of ERK phosphorylation, which peaked at 15 min and returned to basal levels after 60 min. No significant change in total ERKs was found after injury. These data suggest that scratching injury can activate ERK signaling in HaCat cells.

To examine whether ROS generation is involved in injury-stimulated activation of ERK, we tested the effect of DPI (NOX/ROS inhibitor) on ERK phosphorylation after injury. HaCat cells were or were not pretreated with DPI for 30 min before scratches, and then phosphorylated ERK was determined using Western blotting. Figure 3(b) showed that 1 μM or 5 μM DPI pretreatment did not significantly attenuate injury-induced ERK activation 15 min after injury. However, 10 μM DPI pretreatment almost completely blocked injury-induced phosphorylation of ERK. Taken together, the data indicated that ROS are critical mediators of the injury-induced activation of ERK in HaCat cells.

Further, the effect of ERK activation on the injury-induced release of PGE₂ was investigated. Incubation of HaCat cells with ERK inhibitor, PD98059, reduced injury-induced PGE₂ production (Figure 3(c)). This suggested that ERK activation is located upstream of injury-induced PGE₂ release in HaCat cells. However, it should be noted that the ERK is only partially responsible for PGE₂ release induced by injury, since ERK inhibitor did not completely block PGE₂ release.

3.4. Injury-Induced Synthesis of PGE₂ Is Dependent on COX-2, Not COX-1. Several reports have illustrated that COX-




FIGURE 2: Scratching-injury-induced PGE₂ production and the dependence on ROS. (a) PGE₂ secretion by HaCat cells after scratches. PGE₂ was measured by ELISA in different periods of time as indicated. (b) Suppressed secretion PGE₂ by NOX/ROS inhibitor. Values are means \pm SEMs of three replicates. * and # indicate $p < 0.05$ compared with undamaged control cells and with scratched cells, respectively.

the major rate-limiting enzyme, involved in the synthesis of PGE₂. For understanding the activity COX after injury, we tested its activity by COX activity assay. As illustrated in Figure 4(a), injury significantly upgraded COX-2 activity at 2 hr after scratching. However, low level of COX-1 was detected after injury and had no difference compared to the level of undamaged control cells. After that, COX-2 protein expression levels were evaluated by Western blotting. Enhanced expression of COX-2 was examined just 1 hr after scratching in HaCat cells, with a continuous enhancement 2–8 hr after injury (Figure 4(b)). These data suggested that COX-2, not COX-1, is dramatically induced by injury. To corroborate the link between injury-induced ROS production, ERK phosphorylation, and COX-2 generation, we used DPI or PD98059 to treat HaCat cells before injury. Figures 4(c) and 4(d) showed that incubation of HaCat cells with the different dose of DPI or PD98059 decreased injury-induced COX-2 generation. Our data illuminated that ROS production and ERK activation are responsible for COX-2 induction. To further confirm a role of COX-2 for injury-induced PGE₂ synthesis, NS398, a specific COX-2 inhibitor, was used to determine PGE₂ production after injury in HaCat cells. A decrease in injury-induced PGE₂ induction by the addition of NS398 was detected via ELISA (Figure 4(e)). In summary, the stimulation of PGE₂ was mediated by the activation of COX-2 but not COX-1 after injury. It should be noted, however, that COX-2 is not the only molecule responsible for PGE₂ secretion induced by injury as COX-2 inhibitor did not fully block PGE₂ release.

4. Discussion

In this study, we investigated the molecular mechanisms by which ROS stimulated the production of proinflammatory mediator PGE₂ in human keratinocytes with an in vitro injury model. We found that the increased intracellular ROS caused by mechanical injury stimulates PGE₂ production in a time-dependent manner via the activation of COX-2 in human keratinocytes. Additionally, this stimulation requires activation of the signal pathway through phosphorylation of ERK, which contributes to COX-2 induction and PGE₂ synthesis. These results suggest that ROS are not only potent oxidant involved in cellular injury response, but also second messengers activating signal pathway and modulating inflammatory mediators. To our knowledge, this is the first demonstration of ROS-induced ERK phosphorylation leading to the activation of COX-2 and the synthesis of PGE₂ in mechanically injured human keratinocytes.

Cutaneous wound healing commences with blood coagulation followed by infiltration of neutrophils and macrophages to the wound site, which then release a large amount of ROS for anti-infection and debridement. Deficiency of ROS production in phagocytes is responsible for the chronic granulomatous disease, which can cause impaired wound healing in human [15]. Nevertheless, the excess of ROS including superoxide anion (O₂⁻), hydrogen peroxide (H₂O₂), and hydroxyl radical ([•]OH) is primarily regarded as damage to cells during tissue regeneration. In contrast to a large amount of ROS produced in phagocytes,




FIGURE 3: Participation of ERK activation in scratching injury-induced release of ROS and PGE₂. (a) The change of p-ERK and ERK in HaCat cells treated with scratching injury for different periods of time as indicated. The protein levels of p-ERK and ERK were determined by immunoblot analysis and normalized to β-tubulin. (b) The level of p-ERK was decreased in HaCat cells preincubated with different concentrations of DPI for 30 min followed by scratches for 0.5 hr. (c) The dependence of PGE₂ production caused by scratching injury on ERK activation. Values are means ± SEMs of three replicates. * and # indicate *p* < 0.05 compared with undamaged control cells and with scratched cells, respectively.

low level of ROS secreted in a sustained manner has been reported in multiple wound healing cells including keratinocytes [16], fibroblasts [17], and endothelial cells [18]. These low concentrations of ROS participate in a variety of biological processes, including epithelialization, angiogenesis, and granulomatous tissue formation [2, 4]. This study indicates that mechanical injury induces rapid production of ROS in human keratinocyte, which is one of the major cells responsible for wound healing. And the rapidly increased ROS in injured keratinocyte are mainly

produced by NOX, as the specific inhibitor of this enzyme significantly blocked the scratching-induced production of ROS. This result is consistent with previous in vivo observation of enriched H₂O₂ production in the wound site of mouse dermal wound model [2].

Using a full-thickness incisional model of normal wound in mice, the previous study has shown that, during the acute phase of wound healing, PGE₂ is the predominant proinflammatory mediator derived from metabolite [10]. PGE₂ may play a key role in initiating the early inflammation




FIGURE 4: Injury-induced PGE₂ release mediated by COX-2, not COX-1. (a) COX-1 and COX-2 activity of uninjured or injured HaCat cells were measured by COX activity assay. (b) COX-2 expression was increased in HaCat cells treated with scratching injury for different periods of time as indicated. The protein levels of COX-2 were determined by immunoblot analysis and normalized to β-tubulin. (c) The induced expression of COX-2 in HaCat cells was suppressed by NOX/ROS inhibitor DPI. (d) The induced expression of COX-2 in HaCat cells was suppressed by ERK inhibitor. (e) The induced expression of COX-2 in HaCat cells was suppressed by a COX-2 inhibitor. Values are means ± SEMs of three replicates. * and # indicate *p* < 0.05 compared with undamaged control cells and with scratched cells, respectively.

and recruiting cells of the immune system to the site of injury. The increased level of PGE₂ after wounding was confirmed in the in vitro cultured human keratinocytes induced by scratching injury. In addition, the production of PGE₂ after injury in keratinocytes was significantly attenuated by NOX/ROS inhibitor DPI. Similar findings have been demonstrated in UVA radiated keratinocytes [19] and H₂O₂ treated endothelial cells [13]. COX-2 was established as the key enzyme involved in the production of PGE₂ under abnormal conditions. COX-2 rather than COX-1 was found, in this study, to be responsible for the increased PGE₂ in injured keratinocytes based on the evidence of dramatically enhanced expression of COX-2 protein. These indicate that ROS-COX2/PGE₂ is one of the critical intracellular signal pathways accounting for acute response to mechanical injury in keratinocytes. Multiple prostanoid and leukotriene receptors have been identified on keratinocytes [14], indicating that PGE₂ produced by keratinocytes can influence the proliferation, migration, and reepithelialization during wound healing in a manner of autocrine. In addition, PGE₂ secreted by keratinocytes could influence the process of wound healing by modulating the proliferation of fibroblast [20] and affecting the vascular tone, regional blood flow, and angiogenesis [13] through paracrine.

A very common targeted molecule by ROS inside cell is protein tyrosine phosphorylation (PTP), which controls the phosphorylation of tremendous proteins involving cellular signal transduction. MAPKs, as well as phosphoinositide 3-kinase (PI3K), have been identified to be involved ROS-mediated intracellular activities [4]. The activation of ERK (one of the MAPKs) by ROS is related to cell propagation and migration, which contributes to wound healing [8, 9] and the invasion and metastasis of malignant tumors [11]. Increased extent of ERK phosphorylation was identified in scratching injured HaCat, while the total ERKs did not increase significantly. ERK phosphorylation was further demonstrated to be suppressed by NOX/ROS inhibitor DPI dramatically, which confirms that NOX/ROS is indispensable for damage-induced activation of ERK in HaCat. ERK phosphorylation was reported in macrophage treated with oxidized low density lipoprotein, and p-ERK further upregulated COX-2 mRNA and protein expression [21, 22]. In the current study, COX-2 rather than COX-1 was confirmed to be activated p-ERK in HaCat. The important inflammatory mediator PGE₂ production in HaCat was declined by ERK inhibitor, PD98059, which suggests the upstream mediation of ERK on the PGE₂ release after mechanical injury.

In summary, the NOX/ROS-p-ERK-COX-2/PGE₂ pathway is proved by the present study to be involved in the reaction of human keratinocytes to acute mechanical injury. Our results shed a light on the mechanisms accounting for cellular signal transduction in the acute phase of epithelial wound injury, which may be beneficial for developing future therapeutic approaches for tissue repair.

Competing Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Authors' Contributions

Yi-Ping Hu and Yin-Bo Peng contributed equally to this manuscript.

Acknowledgments

The authors are grateful to Dr. Manghua Xu for technical support. The authors appreciate Dr. Zhiyin Yang for discussion and technical support. This study was supported by the National Natural Science Foundation of China (Grants nos. 81272113 for Min Yao, 81201469 for Ying Wang, and 81501656 for Xiu-Jun Fu).

References

- [1] P. Niethammer, C. Grabher, A. T. Look, and T. J. Mitchison, "A tissue-scale gradient of hydrogen peroxide mediates rapid wound detection in zebrafish," *Nature*, vol. 459, no. 7249, pp. 996–999, 2009.
- [2] S. Roy, S. Khanna, K. Nallu, T. K. Hunt, and C. K. Sen, "Dermal wound healing is subject to redox control," *Molecular Therapy*, vol. 13, no. 1, pp. 211–220, 2006.
- [3] O. N. Oktyabrsky and G. V. Smirnova, "Redox regulation of cellular functions," *Biochemistry (Moscow)*, vol. 72, no. 2, pp. 132–145, 2007.
- [4] X.-J. Fu, Y.-B. Peng, Y.-P. Hu, Y.-Z. Shi, M. Yao, and X. Zhang, "NADPH oxidase 1 and its derived reactive oxygen species mediated tissue injury and repair," *Oxidative Medicine and Cellular Longevity*, vol. 2014, Article ID 282854, 10 pages, 2014.
- [5] F. Jiang, Y. Zhang, and G. J. Dusting, "NADPH oxidase-mediated redox signaling: roles in cellular stress response, stress tolerance, and tissue repair," *Pharmacological Reviews*, vol. 63, no. 1, pp. 218–242, 2011.
- [6] A. Matsuzawa and H. Ichijo, "Redox control of cell fate by MAP kinase: physiological roles of ASK1-MAP kinase pathway in stress signaling," *Biochimica et Biophysica Acta*, vol. 1780, no. 11, pp. 1325–1336, 2008.
- [7] J. Zhang, X. Wang, V. Vikash et al., "ROS and ROS-mediated cellular signaling," *Oxidative Medicine and Cellular Longevity*, vol. 2016, Article ID 4350965, 18 pages, 2016.
- [8] G.-D. Sharma, J. He, and H. E. P. Bazan, "p38 and ERK1/2 coordinate cellular migration and proliferation in epithelial wound healing. Evidence of cross-talk activation between map kinase cascades," *Journal of Biological Chemistry*, vol. 278, no. 24, pp. 21989–21997, 2003.
- [9] Y. Matsubayashi, M. Ebisuya, S. Honjoh, and E. Nishida, "ERK activation propagates in epithelial cell sheets and regulates their migration during wound healing," *Current Biology*, vol. 14, no. 8, pp. 731–735, 2004.
- [10] M. Kapoor, F. Kojima, L. Yang, and L. J. Crofford, "Sequential induction of pro- and anti-inflammatory prostaglandins and peroxisome proliferators-activated receptor-gamma during normal wound healing: a time course study," *Prostaglandins Leukotrienes and Essential Fatty Acids*, vol. 76, no. 2, pp. 103–112, 2007.
- [11] W.-T. Chiu, S.-C. Shen, J.-M. Chow, C.-W. Lin, L.-T. Shia, and Y.-C. Chen, "Contribution of reactive oxygen species to migration/invasion of human glioblastoma cells U87 via ERK-dependent COX-2/PGE₂ activation," *Neurobiology of Disease*, vol. 37, no. 1, pp. 118–129, 2010.

- [12] I. Goren, S.-Y. Lee, D. Maucher et al., "Inhibition of cyclooxygenase-1 and -2 activity in keratinocytes inhibits PGE₂ formation and impairs vascular endothelial growth factor release and neovascularisation in skin wounds," *International Wound Journal*, vol. 14, no. 1, pp. 53–63, 2017.
- [13] S. Eligini, I. Arenaz, S. S. Barbieri et al., "Cyclooxygenase-2 mediates hydrogen peroxide-induced wound repair in human endothelial cells," *Free Radical Biology and Medicine*, vol. 46, no. 10, pp. 1428–1436, 2009.
- [14] R. K. Sivamani, "Eicosanoids and keratinocytes in wound healing," *Advances in Wound Care*, vol. 3, no. 7, pp. 476–481, 2014.
- [15] J. Bylund, K. L. Brown, C. Movitz, C. Dahlgren, and A. Karlsson, "Intracellular generation of superoxide by the phagocyte NADPH oxidase: how, where, and what for?" *Free Radical Biology and Medicine*, vol. 49, no. 12, pp. 1834–1845, 2010.
- [16] C. K. Sen, S. Khanna, B. M. Babior, T. K. Hunt, E. Christopher Ellison, and S. Roy, "Oxidant-induced vascular endothelial growth factor expression in human keratinocytes and cutaneous wound healing," *Journal of Biological Chemistry*, vol. 277, no. 36, pp. 33284–33290, 2002.
- [17] W. J. O'Brien, T. Heimann, and F. Rizvi, "NADPH oxidase expression and production of superoxide by human corneal stromal cells," *Molecular Vision*, vol. 15, pp. 2535–2543, 2009.
- [18] T. Tojo, M. Ushio-Fukai, M. Yamaoka-Tojo, S. Ikeda, N. Patrushev, and R. W. Alexander, "Role of gp91phox (Nox2)-containing NAD(P)H oxidase in angiogenesis in response to hindlimb ischemia," *Circulation*, vol. 111, no. 18, pp. 2347–2355, 2005.
- [19] A. Valencia and I. E. Kochevar, "Nox1-based NADPH oxidase is the major source of UVA-induced reactive oxygen species in human keratinocytes," *Journal of Investigative Dermatology*, vol. 128, no. 1, pp. 214–222, 2008.
- [20] T. A. Wilgus, V. K. Bergdall, K. L. Tober et al., "The impact of cyclooxygenase-2 mediated inflammation on scarless fetal wound healing," *American Journal of Pathology*, vol. 165, no. 3, pp. 753–761, 2004.
- [21] K. Taketa, T. Matsumura, M. Yano et al., "Oxidized low density lipoprotein activates peroxisome proliferator-activated receptor- α (PPAR α) and PPAR γ through MAPK-dependent COX-2 expression in macrophages," *Journal of Biological Chemistry*, vol. 283, no. 15, pp. 9852–9862, 2008.
- [22] X. Yu, X. Li, G. Zhao et al., "OxLDL up-regulates Niemann-Pick type C1 expression through ERK1/2/COX-2/PPAR α -signaling pathway in macrophages," *Acta Biochimica et Biophysica Sinica*, vol. 44, no. 2, pp. 119–128, 2012.