



Review article

The use of plant extracts and their phytochemicals for control of toxigenic fungi and mycotoxins

Rhulani Makhuvele^{a,*}, Kayleen Naidu^a, Sefater Gbashi^a, Velaphi C. Thipe^{a,b}, Oluwafemi A. Adebo^a, Patrick B. Njobeh^{a,**}^a Department of Biotechnology and Food Technology, Faculty of Science, University of Johannesburg, Doornfontein Campus, P.O. Box 17011, Doornfontein Campus, Gauteng, 2028, South Africa^b Laboratório de Ecotoxicologia - Centro de Química e Meio Ambiente - Instituto de Pesquisas Energéticas e Nucleares (IPEN) - Comissão Nacional de Energia Nuclear-IPEN/CNEN-SP, Av. Lineu Prestes, 2242 - Butantã, 05508-000, São Paulo, Brazil

ARTICLE INFO

Keywords:

Food science
Food safety
Mycotoxin
Fungal invasion
Detoxification
Plant extracts
Nanosponge
Food and feed

ABSTRACT

Mycotoxins present a great concern to food safety and security due to their adverse health and socio-economic impacts. The necessity to formulate novel strategies that can mitigate the economic and health effects associated with mycotoxin contamination of food and feed commodities without any impact on public health, quality and nutritional value of food and feed, economy and trade industry become imperative. Various strategies have been adopted to mitigate mycotoxin contamination but often fall short of the required efficacy. One of the promising approaches is the use of bioactive plant components/metabolites synergistically with mycotoxin-absorbing components in order to limit exposure to these toxins and associated negative health effects. In particular, is the fabrication of β -cyclodextrin-based nanosponges encapsulated with bioactive compounds of plant origin to inhibit toxigenic fungi and decontaminate mycotoxins in food and feed without leaving any health and environmental hazard to the consumers. The present paper reviews the use of botanicals extracts and their phytochemicals coupled with β -cyclodextrin-based nanosponge technology to inhibit toxigenic fungal invasion and detoxify mycotoxins.

1. Introduction

Food and feed contamination by toxigenic fungi is of serious health concern and has been recorded since mankind initiated the cultivation and storage of agricultural commodities. These toxigenic fungi belong mainly to the genera *Aspergillus*, *Fusarium*, *Claviceps*, *Penicillium*, *Stachybotrys*, and *Alternaria*, etc. They produce toxic substances, some of which are referred to as mycotoxins. Cumulative evidence from the literature suggest that these mycotoxins do not present any apparent biochemical importance in fungal development or growth. However some secondary metabolites play a role in virulence, development and pathogenicity (Perincherry et al., 2019; Venkatesh and Keller, 2019). Mycotoxins have adverse effects on humans and animals even at low concentrations (Vila-Donat et al., 2018; Adekoya et al., 2019), and the proliferation of toxigenic fungi on food and feed is always favoured by certain environmental factors such as humidity and temperature, as well as certain biotic conditions (i.e. substrate composition). High incidences of mycotoxin

occurrence are usually reported in tropical regions such as Asia and sub-Saharan Africa, where conditions for their proliferation are optimal. Mycotoxins contaminate mainly commodities such as cereals, grains, nuts, and their by-products during production, pre-harvest and post-harvest (Milicevic et al., 2010; Zain, 2011; Nleya et al., 2018; Balendres et al., 2019; Gbashi et al., 2019). These toxins often enter the body mainly via ingestion, as well as by inhalation, parental and dermal exposure routes. The toxins can also enter the food chain through infected crops which are consumed either directly or indirectly by humans or animals as feed sources. As such, they appear in meat, milk and eggs (Hojnik et al., 2017).

Mycotoxin contamination is of serious concern for food safety and security worldwide. These toxins account for enormous economic losses in agricultural productivity and trade, with more severity in poor and developing countries. It is estimated that about 60–80% of crops worldwide can be contaminated by mycotoxins, thus resulting in significant economic losses (Eskola et al., 2019). The toxic effects of

* Corresponding author.

** Corresponding author.

E-mail addresses: rhulanim@uj.ac.za (R. Makhuvele), pnjobeh@uj.ac.za (P.B. Njobeh).<https://doi.org/10.1016/j.heliyon.2020.e05291>

Received 1 July 2020; Received in revised form 16 September 2020; Accepted 14 October 2020

2405-8440/© 2020 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

mycotoxins are persistent, and they are difficult to completely eradicate once they enter the food/feed chain. In the agricultural industry, mycotoxins cause loss in livestock production due to reduced growth rates, decreased immunity and fertility, reduced eggs, meat and milk production, and increased mortality (Thipe et al., 2020). Furthermore, the effects associated with mycotoxins cause veterinary and healthcare costs (Marroquin-Cardona et al., 2014; Kagot et al., 2019). This contamination of food and feedstuffs by mycotoxins equally reduces the nutritional value, quality and safety of food and feed (Luo et al., 2018). Many countries have established regulatory limits on mycotoxins in agricultural commodities to limit human and animal health risks associated with them (Marroquin-Cardona et al., 2014; Haque et al., 2020). Mycotoxins cause a disease known as mycotoxicosis (Bennett and Klich, 2003; Liew and Mohd-Redzwan, 2018). The health effects of mycotoxins are extensive as they potentiate some hepatotoxic, nephrotoxic, mutagenic, genotoxic, carcinogenic, immunosuppressive and teratogenic properties. The most prevalent mycotoxins of agricultural importance include the aflatoxins (AFs), ochratoxins (OTA), fumonisins (FBs), trichothecenes and zearalenone (ZEN). These have been given serious attention due to the health risks that they pose in humans and animals (Dikhoba et al., 2019; Celik, 2020).

Various approaches to control and prevent mycotoxins in food and feed have been developed. These approaches are categorised as chemical and micro-biological methods (Adebo et al., 2017; Adebiyi et al., 2019). These methods were shown to be effective in the prevention of growth of toxigenic fungi and production of associated mycotoxins during pre-harvest, after harvest food and during storage of food commodities. Chemical methods involve the use of chemicals such as ammonia, sodium hydroxide, hydrochloric acid, butylated hydroxytoluene, butylated hydroxyanisole and oltipraz to decontaminate mycotoxins (Galvano et al., 2001; Karlovsky et al., 2016; Ćolović et al., 2019). In addition to their inefficacy in decontamination of mycotoxins, their long-term extreme use is still limited due to their residual toxic product/s, public health and environmental concerns (Abdel-Fattah et al., 2018; Alberts et al., 2019; Sadhasivam et al., 2019; Meng et al., 2020) coupled with the interference with nutrients and organoleptic properties of food and feed (Celik, 2020).

Physical methods include cleaning, dehulling, sorting, milling and ultra-violet light, pulsed light, cold plasma as well as irradiation. Other physical methods involve the use of adsorbents or binders such as activated charcoal, bentonite, zeolites and sepiolite clay. The methods have been effectively applied in decontaminating mycotoxins. However, technology implementation costs, potential residual toxic effects, poor adsorption and low specificity (i.e. selective action) against some mycotoxins still hamper their routine application (Mahato et al., 2019; Celik, 2020). Microbiological methods entail the use of probiotic bacteria, yeasts and their enzymes, which are effective in reducing mycotoxins in food and feed (Tian et al., 2016; Adebo et al., 2017; Udomkun et al., 2017). However, these microorganisms and their enzymes often interfere with nutrient supply resulting in undesirable products. Apart from this, enzymatic degradation by-products still limit their uses (Juodeikiene et al., 2012; Adebiyi et al., 2019; Lyagin and Efremenko, 2019). Therefore, there is a need to discover alternative methods that can prevent fungal colonization of agricultural commodities, detoxify or bio-transform mycotoxin residues to less- or non-toxic forms without any limitations (Iram et al., 2015; Powers et al., 2019; Haque et al., 2020).

Botanicals such as essential oils, spices, herbs and crude extracts present outstanding alternatives for the discovery of biofungicides and nutraceuticals for mitigating mycotoxicosis and related infections. Botanicals are generally regarded as environmentally friendly and safer alternative sources of bioagents for the control of fungi and mycotoxins in food and feed (Iram et al., 2016; Adebo et al., 2020; Prakash et al., 2020). They are more affordable as opposed to other materials used for the same purpose, they provide a synergistic approach as protectants of fungal/mycotoxin contamination and further stimulate pathways that elicit the natural defence systems in plant tissues (da Cruz Cabral et al.,

2013; Alberts et al., 2019; Gacem et al., 2020; Meng et al., 2020). They contain various phytochemicals with pharmacological properties against various diseases. Recent studies have been conducted on the possible application of botanicals as bio-fungicides and nutraceuticals to ameliorate the proliferation of toxigenic fungi and mycotoxin contamination in food and feed (Dikhoba et al., 2019; Ponzilacqua et al., 2019; Kavitha et al., 2020). This review aimed to elucidate the use of botanical extracts and their phytochemical compounds to prevent and detoxify mycotoxins without any adverse effect on the nutritional value of food and feed. We also review nano-encapsulation technology which can be adopted to improve bioavailability and solubility of phytochemicals used as biofungicides and nutraceuticals to prevent growth of mycotoxigenic fungi and subsequent mycotoxin contamination.

2. Mitigation of toxigenic fungi and mycotoxins using botanical methods

2.1. Importance of botanicals

Plants have been widely used in folk medicine since time immemorial in order to treat and prevent various ailments from one generation to another (Horn and Vargas, 2008; Street and Prinsloo, 2013). Despite the advancement made in modern medicine, many populated groups in developing countries still depend on traditional medicine for preventing and treating various ailments. This is due to cultural beliefs, low cost and effectiveness (Moura-Costa et al., 2012; Ayele, 2018). According to the World Health Organization (WHO, 2001), approximately 80% of the population of the world still depend on traditional medicine for primary healthcare (Prakash et al., 2020). Recent studies have also created renewed interest in the use of botanicals and their compounds as nutraceuticals in this regard in both developed and developing countries (Galvano et al., 2001; Reddy et al., 2010; Anjorin et al., 2013; Dikhoba et al., 2019). The advantage of using plants for drug discovery is due to their abundance in nature and wide distribution geographically. A considerable number of drugs have been synthesized from plants based on their use in traditional medicine (Van Wyk et al., 1997; Dias et al., 2012). Africa is richly endowed with a wealth of medicinal plants. However, very few studies have looked into exploiting their constituents for mycotoxin detoxification (Stoev et al., 2019; Makhuvele et al. unpublished data), and thus necessitate studies in this regard. The following sections provide an overview of various botanicals that exhibit antifungal activities. In addition, we also follow a nanoencapsulation approach through the fabrication and use of nanosponges to improve the stability from oxidation/degradation, availability, and demonstrate the most pronounced efficacy of botanicals.

2.2. Phytochemicals as a source of therapeutics and nutraceuticals

Plants produce secondary metabolites as a defence mechanism against pathogenic microorganisms, insects and adverse environmental conditions. These metabolites are known as phytochemicals, which are non-nutritive (Prakash et al., 2020) and to some extent essential oils. However, they can protect humans and animals against certain diseases caused by microorganisms or toxins associated with them due to the antimicrobial properties they possess (Palombo, 2011; Shin and Park, 2018; Redondo-Blanco et al., 2019). The metabolites are the most promising chemo-preventative agents for future drug discovery and development (Alabi et al., 2011). There are various major groups of phytochemical compounds discovered to date and differ according to their chemical structures (Das et al., 2020). These major groups include phenolic compounds, flavonoids, phytosterols, carotenoids, tocopherols, terpenoids, alkaloids, saponins, tannins, aromatic acids, glucosinolates, carotenoids, essential oils, chlorophyll and organic acids as well as proteases inhibitors (Bhattachar, 2011; Adebo and Gabriela Medina-Meza, 2020; Loi et al., 2020). Such compounds may act directly or indirectly to protect against ailments or pathogens, because they contain

Table 1. Studies on the effects of plant extracts and their compounds on mycotoxin-induced toxicity.

| Plant source | Protective agent | Model | Mechanism of action | References |
|---|-----------------------------------|--|--|---|
| Native phytochemicals | | | | |
| <i>Curcuma longa L. (Turmeric)</i> | Curcumin | <i>In vitro</i> Ames assay, rodent & chicken | Hepatoprotective effects against AFB ₁ toxicity <i>in vitro</i> and <i>in vivo</i> | Soni et al. (1997); Smerak et al. (2006a); Nayak and Sashidhar (2010); Limaye et al. (2018) |
| | Turmeric | Chicken broiler | Protected against AFB ₁ oxidative damage | Gowda et al. (2008); Nayak and Sashidhar (2010) |
| <i>Curcuma amada</i> (Ginger) | Ginger | HepG2 cells & rats | Inhibited AFB ₁ , STE and PAT toxicity | Yang et al. (2011); Vipin et al. (2017) |
| | 6-Gingerol | HepG2 cells <i>In vitro</i> | Reduced PAT induced DNA damage in HepG2 | El-Shayeb (1980); Yang et al. (2011) |
| <i>Thymus vulgaris</i> (Thyme) | Thyme oil | Rats | Ameliorated oxidative stress and genotoxicity effects of AFB ₁ | Abdel-Fattah et al. (2010) |
| <i>Syzygium aromaticum</i> (Clove) | Clove | <i>In vitro</i> | Inhibited AF, STE, & CIT toxicity | Hitokoto et al. (1980); Azzouz (1981); Hussain et al. (2012) |
| | Quercetin | <i>In vitro</i> | Prevented AFB ₁ carcinogenesis | Bhattacharya & Firozi (1988) |
| <i>Camellia sinensis L.</i> (green tea) and a variety of plants | Epigallocatechin-3-gallate (EGCG) | Mice | Protected against DON & HT-2 toxin toxicity | Smerak et al. (2006b); Yang et al. (2011); Do et al. (2015) |
| <i>Solanum lycopersicum</i> (tomatoes) | Lycopene | Chicken broiler, Mice & rats | Protected against oxidative, inflammatory, hormonal & reproductive damage induced by AFB ₁ , OTA & ZEN in mice, also inhibited T-2 toxin induced oxidative stress | Leal et al. (1999); Aydin et al. (2013); Palabiyik et al. (2013); Wu et al. (2017a); Hedayati et al. (2018) |
| | Cyanidin | <i>In vitro</i> HepG2 & Caco-2 cells | Protected against AFB ₁ & OTA-induced oxidative stress | Sorrenti et al. (2012); Do et al. (2015) |
| | Rutin | Rats | Reversed T-2 toxin-induced lipid peroxidation in liver homogenate | El-sawi and Al-Seeni. (2009); Wu et al. (2017a) |
| Crude extracts | | | | |
| <i>Premna integrifolia</i> | | Mice | Inhibited AFB ₁ toxicity | Singh et al. (2019) |
| <i>Silybum marianu</i> | | Broiler chicken | Inhibited OTA toxicity | Stoev et al. (2019) |
| <i>Withania somnifera</i> | | Broiler chicken | Inhibited OTA toxicity <i>in vivo</i> | Stoev et al. (2019) |
| <i>Ammona senegalensis</i> | | <i>In vitro</i> | Inhibited AFB ₁ genotoxicity in Ames, Vitotox and comet assays | Makhuvele et al. (2018a) |
| <i>Monanthonotaxis caffra</i> | | <i>In vitro</i> | Inhibited AFB ₁ genotoxicity <i>in vitro</i> | Makhuvele et al. (2018b) |

antimutagenic, antigenotoxic, antimicrobial, anthelmintic, anticarcinogenic, antiproliferative and anti-inflammatory as well as antioxidant properties (Galvano et al., 2001; Makhafola et al., 2017; Velu et al., 2018; Lahlou et al., 2019).

2.3. Antifungal effects and detoxification abilities of plant extracts and their phytochemicals compounds on toxigenic fungi and mycotoxin toxicity

Plants possess antimutagens, antimicrobial, antioxidants or anticarcinogens capable of compromising the toxic and genotoxic effects of mycotoxins (Madrigal-Santillan et al., 2010; Anjorin et al., 2013; Powers et al., 2019). Antioxidants protect the cell membranes and macromolecules by scavenging free radicals (Wu et al., 2017a). Furthermore, phytochemicals induce cytotoxicity in fungi by disrupting cell membrane permeability and functions; inhibiting cytoplasmic and mitochondrial enzymes; inhibiting enzymes involved in cell wall components synthesis; and altering the cell compartment, osmotic and the redox balance (Loi et al., 2020). However, plant extracts and their compounds also act by inducing xenobiotic detoxification and biotransformation pathways (Gross-Steinmeyer and Eaton, 2012; Wu et al., 2017b). Phytochemicals are capable of inhibiting enzymes that activate Phase I carcinogens as well as induce enzymes for Phase II detoxification (Galvano et al., 2001; Wu et al., 2017b). The bioactive compounds in plants have been used as additives to prevent fungal growth and aflatoxin (AF) contamination in food and feed (Table 1), thus reducing the risks of mutagenicity and carcinogenicity of such mycotoxins as AFs (Mathuria and Verma, 2007).

Antifungal and anti-mycotoxigenic activities of herbal plants with potential antioxidant properties were investigated against fungal strains that are phytopathogenic, i.e. *Fusarium verticillioides*, *A. flavus* and *A. ochraceous*. The results reported potentials of the selected medicinal plants to be used for the discovery of biofungicides that may prevent oxidative related food spoilage (Dikhoba et al., 2019). A study by Abdel-Fattah et al. (2018) reported the antioxidant, antifungal and anti-mycotoxigenic potentials of wild stevia extracts against *A. flavus*, *A. ochraceus*, *A. niger*, and *F. moniliforme*. Furthermore, essential oils have been found to effectively modulate the growth of mycotoxigenic fungi such as *A. flavus*, *A. oryzae*, *A. niger*, *Alternaria alternata*, *F. moniliforme*, *F. graminearum*, *Penicillium citrinum* and *P. viridicatum*, etc., and their associated mycotoxins (da Cruz Cabral et al., 2013; Prakash et al., 2015; Powers et al., 2019). A study by Kocic-Tanackov et al. (2019) reported how essential oil of *Carum carvi* L applied at a concentration from 1.5 µl/g demonstrated complete inhibition of the growth of *A. parasiticus*, and likewise 4.5 µl/g exhibited complete inhibition of the growth of *A. flavus* as well as the secretion of aflatoxins by the same strain in polenta. Also noted was that a concentration of 35.0 µl/g of *Juniperus communis* L. essential oil showed strong potency against *A. parasiticus* and *A. flavus* IKB (i.e. by significantly inhibited their growth) with percentage inhibition between 42.4 and 79.8%, while 50.0 µl/g of *J. communis* impeded aflatoxin production by *A. flavus* IKB in polenta completely.

Curcumin and ellagic acid are examples of compounds isolated from plants which are used as food and feed supplements. These compounds prevent metabolism of aflatoxin B₁ (AFB₁) and increase the activity of glutathione-S-transferase involved in the detoxification of xenobiotics. They were equally found via Ames assay and in rat and chicken models to protect against mutagenicity induced by AFB₁ with *Salmonella typhimurium* strain TA98 and TA100 (Soni et al., 1997; Gowda et al., 2008). Smerak et al. (2006a) studied the effect of curcumin against AFB₁ mutagenicity by Ames, *in vivo* micronucleus, comet and chemiluminescence and blastic transformation methods. The results showed that curcumin was able to protect cells against mutagenicity. They also demonstrated a significant reduction in DNA damage through stimulation of DNA repairs. Curcumin has earlier been demonstrated by many authors to exhibit anticarcinogenic, antiproliferative and antimutagenic effects against various mutagens *in vitro* and *in vivo* (El-Hamss et al., 1999; Inano and Onoda, 2002; Polasa et al., 2004; Hosseini and Hosseinzadeh, 2018; Khan et al., 2019). Resveratrol is another natural

product isolated from grape skin which suppresses the proliferation of many tumour cells such as breast, pancreatic and prostate cancers by inhibiting xenobiotic metabolism and inducing detoxification pathways (Farombi, 2004; Thiye et al., 2019). Studies have showed that this product can protect mycotoxin-induced toxicity *in vitro* and *in vivo* (Do et al., 2015; Sridhar et al., 2015; Tabeshpour et al., 2018). Furthermore, 6-gingerol, a natural bioactive compound extracted from ginger was found to possess strong protective effects against PAT-induced genotoxicity in HepG2 cells *in vitro* (Yang et al., 2011).

Lycopene is a natural product found in tomatoes, papaya and other red fruits and vegetables that showed to have protective effects against ZEN oxidative, reproductive and hormonal damage in mice (Aydin et al., 2013; Palabiyik et al., 2013). Lycopene also prevented T-2 toxin-induced oxidative stress and maintained GSH cellular levels *in vivo* (Leal et al., 1999). Furthermore, lycopene reduced AFB₁ and OTA-induced oxidative stress and apoptosis in rats (Hedayati et al., 2018). Cyanidin, a phytochemical found in various medicinal herbs, fruits and vegetables including grapes, blackberry, cherry, cranberry, raspberry, red cabbage, red onion, etc., showed a protective effect against AFB₁ and OTA-induced toxicity in hepatocytes and enterocytes (Sorrenti et al., 2012). El-Sawi and Al-Seeni (2009) reported that rutin displayed strong antioxidant activity against T-2 toxin in rat liver and also decreased lipid peroxidation induced by T-2 toxin. Shehata et al. (2017) demonstrated that oil-bioactive films from three extracts of immature fig fruit, leaves, and pomegranate husks to be a novel method for post-harvest grain management against mycotoxins.

Furthermore, Negera & Washe (2019) evaluated AF degradation abilities of some selected natural food spices including garlic, ginger, black cumin, clove, sacred basil, lemon grass, thyme, fenugreek and lemon, traditionally used by the Ethiopian community for food flavoring and preservation. Electrochemical and LC-MS/MS methods were used to investigate aflatoxin degradation efficacy of the spice extracts by determining the toxin in extract-treated and non-treated samples. The results revealed that garlic had maximum AFB₁ degradation activity followed by lemon and the other dietary spices during 1-hour exposure to AFB₁ standard at 25 °C. The results also showed that the possible mechanism of AFB₁ degradation is through chemical transformation of AFB₁ parent compound to another compound by the plant extracts. Iram et al. (2015) investigated the ability of *Corymbia citriodora* plant extract to detoxify AFB₁ and AFB₂ both *in vitro* and *in vivo*. They observed that the leaf extracts had maximum detoxification at pH 8 and 30 °C temperature after 72 h of incubation. Iram et al. (2016) demonstrated that *Trachyspermum ammi* seed extracts can be used for the development of biologically safe herbal additives in food and feed. Ponzilacqua et al. (2019) evaluated degradation capabilities of aqueous plant extracts of *Rosmarinus officinalis*, *Origanum vulgare*, *Psidium cattleianum* and *Passiflora alata* against AFB₁, with *R. officinalis* extract exhibiting the highest degradation percentage of AFB₁ (range: 49.0–60.3%) at 24–48 h, followed by *O. vulgare* with (range of 30.7%–38.3%) after 48 h of incubation.

Makhuvele et al. (2018a) investigated the antigenotoxicity of plant extracts against AFB₁-induced genotoxicity. The results showed that most plant extracts from *Artabotrys brachypetalus*, *Helichrysum petiolare*, *Hexalobus monopterus*, *Friesodielsia obovate*, *Monanthonotaxis caffra*, *Protea hybrid*, *Protea roupelliae*, *Monodora junodis*, *Uvaria caffra*, *Xylopia parviflora*, *Rhoicissus sekhukhuniensis*, *Podocarpus henkelii*, *Podocarpus elongatus* and *Agapanthus praecox* had moderate to strong antimutagenic potency in both Vitotox and Ames assays. Ethanolic extract of *Premna integrifolia* leaves demonstrated a protective effect in mice liver against AFB₁-induced toxicity by inhibiting apoptosis and oxidative stress (Singh et al., 2019). Furthermore, extracts of *Schisandra chinensis* and *Thonninga sanguinea* were shown to possess hepatoprotective effects against AFB₁ by enhancing antioxidant and detoxification in rats (Gyamfi and Aniya, 1998; Ip et al., 1999). Feed additives from South African medicinal plants, i.e., *Silybum marianum*, *Withania somnifera* and *Centella asiatica* were tested and except for *Centella asiatica*, the extracts were found to give partial protection against the growth inhibitory activity of OTA and

associated immunosuppression in broiler chicks (Stoev et al., 2019). The same extracts also showed some hepatoprotective effects on broiler chicks exposed to OTA with extract from *S. marianum* exhibiting nephroprotective effect against OTA toxicity. Very recently, Nerilo et al. (2020) evaluated the use of ginger essential oils (GEO) as a fumigant agent for stored maize grains. They reported that GEO was mainly composed of α -zingiberene (23.85%) and geranial (14.16%). Furthermore, their results revealed that 25 and 50 $\mu\text{g/g}$, respectively, exhibited antifungal activity against *A. flavus* and inhibited AFB₁ and AFB₂ production. Despite the efficacy of plant extracts and their phytochemicals in the management of toxigenic fungi and their toxins, there are various limitations to the use of phytochemicals and their crude extracts as biofungicides and nutraceuticals. The subsequent section elaborates on these limitations.

2.4. Current limitations on the use of phytochemicals as biofungicide

There is an increased demand for the use of plant extracts and their compounds and this raises concerns about the safety, toxicity and quality of these products. Studies have reported the contamination of plant materials with medicinal properties by mycotoxins (Ashiq et al., 2014; Mulaudzi, 2019; Altyn and Twaruzek, 2020; Thiye et al., 2020). Furthermore, research has shown that many plant extracts used in traditional medicine and as food ingredients are toxic and mutagenic. As such, a thorough screening of their toxicological properties is necessary. Those that show no signs of toxicity are given high priority (Verschaeve and Van Staden, 2008; Sajid et al., 2019). The use of plant extracts, especially in exploiting their bioactive compounds, are exploited in the discovery and development of new antifungal and nutraceutical agents.

The high frequency of toxigenic fungal manifestation warrants the consistent evaluation of bioactive phytochemicals and their derivatives that exhibit high antifungal and anti-mycotoxigenic activities as alternatives to conventional fungicides. Table 2 briefly presents phytochemicals that have shown antifungal activities and their mode of action, thus inhibiting their respective mycotoxin biosynthesis. Moreover, nano-encapsulation formulations are also presented, exhibiting a synergistic effect between encapsulate and phytochemical with low side effects. The use of natural formulations antifungal agents for safe, effective, and eco-friendly features against fungi and mycotoxins is ideal for agricultural practices compared to standard antifungal agents (Rai et al., 2020). Literature precedence has proposed the notion of fully understanding the exact principles underlying the antifungal and anti-mycotoxigenic mode of action of phytochemicals against mycotoxigenic fungi (Chaudhari et al., 2020; Xu et al., 2020), facilitated through the following:

- (i) inhibition of ergosterol biosynthesis, a major sterol that regulates plasma membrane biogenesis;
- (ii) disruption of fungal cell membrane; and
- (iii) production of reactive oxygen species (ROS) which results in oxidative stress.

The nanoencapsulation (e.g. nanoemulsion, nanofibre, nanogel, nanoliposome, nanoparticle, nanotube, and nanosponge) coupled with phytochemicals provides promising avenues to explore possible strategies for enhancing efficacy and combating fungal resistance where conventional antifungal drugs prove to be ineffective (Dhakar et al., 2019; Redondo-Blanco et al., 2019). Powers et al. (2019) emphasized on the priority and importance of big data approaches for expediting the identification of active components. They utilized a huge data to evaluate antifungal activity of 82 essential oils against *A. niger*, *C. albicans*, and *C. neoformans*. The results obtained demonstrated that antifungal susceptibility to the essential oils was as follows: *C. neoformans* > *C. albicans* > *A. niger*. However, the use of phytochemical compounds alone limits their full applications. This limitation is due to their instability, insolubility, high volatility, low bioavailability, ability to change organoleptic characteristics of food/feed, and limited of facilities and resources for

their extraction and purification (Sajid et al., 2019; Adebo et al., 2020; Loi et al., 2020). To overcome these factors, cohesive research collaborations between academics, research institutions, government agencies, food and pharmacological industries, as well as international stakeholders are required. In addition, different emerging technologies such as nanotechnology can, in part, provide solutions to encounter some of these limitations (Prakash et al., 2020; Thiye et al., 2020). Phytochemical compounds can be encapsulated in edible coatings or with nanoparticles such as nanosponges to elicit the factors listed herein (Alotaibi et al., 2019; Loi et al., 2020; Thiye et al., 2020). The subsequent section gives brief information on nanocarriers and their encapsulated bioactive compounds for use as antifungal and mycotoxin detoxification agents.

3. Innovative technology for mycotoxin detoxification

Frontiers in the field of nanotechnology have led to their vast applications in the field of nanomedicine, extending to the agricultural sector. Great strides have been made in the implementation of nanotechnology in agriculture for controlling mycotoxin contamination in food and feed supply chain (Thiye et al., 2018). The era of green nanotechnology, with the use of plant phytochemicals for the production of nanomaterials, has greatly improved their safety for use in agriculture as mycotoxin detoxifying agents (El-Desouky & Ammar, 2016; Thiye et al., 2020). A combination of nanotechnology and botanical extracts, as well as their phytochemicals, has shown significant results in the pharmacological, agricultural and cosmetic industries. Nano-encapsulated phytochemicals demonstrated strong efficacy over the free form because of the increased surface area, protection of encapsulated compounds from internal and external environmental conditions (Prakash et al., 2020). Nanocarriers can protect bioactive phytochemical compounds against thermal and photodegradation and further provide controlled release of antifungal compounds for the development of active packaging for maintaining the integrity of food/feed during storage and protection from fungal growth and mycotoxin contamination (Bahrami et al., 2020). They also caused reduced toxic effects of these plant-based drugs (Pushpalatha et al., 2018; Sajid et al., 2019). To date, there are different types of nanocarriers used for drug delivery, including liposomes, metal nanoparticles, polymeric nanoparticles, polymeric micelles and nanosponges. All these nanocarriers have been reported to be effective as drug delivery systems for plant-based products in cosmetics, agriculture and medicine (Pushpalatha et al., 2018; Thiye et al., 2020; Udomkun and Njukwe, 2020). However, there is limited information in the literature on the use of nanosponge encapsulated biofungicides, and for that, this paper highlights the use of cyclodextrin nanosponges as carrier vehicles to encapsulate phytochemicals.

3.1. Nanosponges as encapsulation system for antifungal and detoxifying phytochemicals

Nanosponges are emerging innovative drug delivery or encapsulation systems that result from the advancement of nanotechnology. They are microscopic, sponge-like particles with nanometer cavities, which can encapsulate both hydrophilic and lipophilic substances including toxins, but when doing so, enhance their stability, bioavailability and solubility (Ananya et al., 2020). Nanosponges are non-irritant, non-toxic, non-mutagenic and non-allergenic agents made-up of a polyester mixed with crosslinkers in a solution. The polyester is biodegradable and effective in delivering the drug to the targeted site (Pawar et al., 2016; Bhowmik et al., 2018). Nanosponges come in different forms including cyclodextrin which is most widely used and promising in encapsulation of phytochemicals. Cyclodextrin nanosponges are natural polymers formed from enzymatic degradation of starch and consist of a ring of oligosaccharides molecules. They are characterized by a highly porous spherical, amorphous or crystalline structure (Sherje et al., 2017). An interesting property of cyclodextrin-based nanosponges is that some important physicochemical properties of the nanomaterials such as

Table 2. Antifungal effects of plant extracts, their compounds, and nanoformulations against mycotoxigenic fungi.

| Plant source | Protective agent | Model | Mechanism of action | References |
|---|--|--|--|--|
| Native phytochemicals | | | | |
| <i>Curcuma longa</i> L. (Turmeric) | Curcumin | <i>F. solani</i> , <i>C. albicans</i> , <i>P. expansum</i> , <i>A. flavus</i> , and <i>A. parasiticus</i> | Downregulation of $\Delta^{5,6}$ desaturase gene (<i>ERG3</i>) resulting in reduced ergosterol biosynthesis leads to cell death attributed by elevated levels of reactive oxygen species (ROS) production, Reduced proteinase secretion and inhibition of H^+ -ATPase activity induces acidification of extracellular and intracellular matrix inhibition of hyphae development through the suppression of thymidine uptake 1 (TUP1), Curcumin is a photosensitizer (OPS): Where $OPS + \gamma \rightarrow$ singlet excited state (1PS*) \rightarrow triplet excited state (3PS*) $\rightarrow H^+$ to a biomolecule = radicals (anion superoxide (O_2^-), hydrogen peroxide (H_2O_2), hydroxyl radical ($\bullet OH$), and singlet oxygen (1O_2) resulting in cell death by apoptosis, necrosis, or autophagy. | Sharma et al. (2010); Neelofar et al. (2011); Moghadamtousi et al. (2014); Chen et al. (2018); Song et al. (2020); Narayanan et al. (2020) |
| | Turmeric essential oil (e.g. β -pinene, camphor, and eucalyptol) | <i>A. flavus</i> | Fungicidal activity to destroy cellular integrity and induced alteration in mitochondrial membrane potential. | Hu et al. (2015) |
| <i>Curcuma amada</i> (Ginger) | α -zingiberene, geraniol, 6-gingerol, and 6-shogaol | <i>A. flavus</i> , <i>A. parasiticus</i> and <i>F. verticillioides</i> | Anti-mycotoxinogenic activities through induced alteration in mitochondrial membrane potential which causes loss of membrane integrity, leakage of cellular material, and inhibition of ergosterol biosynthesis. | Kavitha et al. (2020); Nerilo et al. (2020); Gacem et al. (2020) |
| | Chitosan | <i>A. ochraceus</i> | Inhibition of gene expressions that are critical for cell wall and plasma membrane homeostasis, ribosome biogenesis and other biological processes through the distraction of the integrity of cell surface architecture and protein biosynthesis. | Meng et al. (2020) |
| <i>Rosmarinus officinalis</i> L. essential oil | 1,8-cineole (eucalyptol), camphor and α -pinene | <i>A. flavus</i> | Reduces ergosterol production and biomass of mycelium. | da Silva Bomfim et al. (2019) |
| <i>Syzygium aromaticum</i> (Clove) | Eugenyl acetate, eugenol, and β -caryophyllene | <i>A. flavus</i> and <i>A. niger</i> | Induces cell death through early apoptosis (nuclear condensation) and late apoptosis (damage of plasma membrane) in hyphae, Downregulation of metabolic genes [secondary metabolism global regulator (<i>laeA</i>), lipase (<i>lipA</i>), and metalloprotease (<i>metP</i>)] responsible for fungal lipid and protein metabolism. | Oliveira et al. (2020); Castellanos et al. (2020) |
| <i>Punica granatum</i> (pomegranate) | Tannins | <i>Alternaria alternata</i> , <i>A. niger</i> <i>F. oxysporum</i> , <i>F. culmorum</i> , <i>F. graminearum</i> , and <i>P. digitatum</i> | Generation of ROS resulting in the destruction of the plasma membrane and mitochondrial dysfunction. | Zhu et al. (2019) |
| <i>Camellia sinensis</i> L. (green tea) and a variety of plants | Epigallocatechin 3-O-gallate (EGCG) | <i>Candida spp.</i> | Formation of lesions on the cell membrane caused by loss of cell membrane integrity, cellular and plasma membrane damage, Increased membrane permeability that causes osmotic imbalance which ultimately results in cell death. | Behbehani et al. (2019) |
| <i>Solanum lycopersicum</i> (tomatoes) | Lycopene | <i>C. albicans</i> | Plasma membrane depolarization and cell cycle arrest (G2/M) through increased intracellular ROS, Elevated levels of cytosolic and mitochondrial Ca^{2+} homeostasis causes mitochondrial dysfunction, Facilitates cytochrome c release that results in caspase activation. | Choi and Lee (2015) |
| <i>Piper nigrum</i> L. (Pepper) | limonene, sabinene, and β -caryophyllene | <i>F. oxysporum</i> and <i>A. niger</i> . | Disruption of cell wall and plasma membrane, coagulation of the cytoplasm, damage cellular organelles and ergosterol biosynthesis. | Castellanos et al. (2020) |

(continued on next page)

Table 2 (continued)

| Plant source | Protective agent | Model | Mechanism of action | References |
|---|--|---|---|--|
| Nanoformulations | | | | |
| <i>Curcuma longa</i> L. and various fruits and vegetables | Nanovesicles (curcumin and quercetin co-encapsulation) | <i>A. niger</i> , <i>A. fumigates</i> , and <i>Candida spp.</i> , | Causes the excess formation of lomasomes and plasmalemmasomes which results in cell wall and plasma membrane expansion imbalance. | Sadeghi-Ghadi et al. (2020); Rai et al. (2020) |
| <i>Streptomyces natalensis</i> | Nanoemulsions (curcumin, piperine, and tualang honey) combinations | <i>Candida spp.</i> | Nanoemulsions (curcumin + piperine + honey) possessed favorable antifungal activity (more than 80%) against the wide range of <i>Candida spp.</i> , due to multimodal activity. | Phuna et al. (2020) |
| | Natamycin/methyl- β -cyclodextrin (N/ME- β -CD) | <i>A. Japonicus</i> , <i>Gibberella persicaria</i> , <i>Botrytis cinerea</i> , and <i>P. expansum</i> | Inhibits ergosterol biosynthesis in the plasma membrane resulting in mitochondrial dysfunction. | Kavitha et al. (2020); Yang et al. (2019) |
| <i>Origanum majorana</i> essential oil | Chitosan nanoemulsion | <i>A. flavus</i> , <i>A. parasiticus</i> , <i>F. graminearum</i> | Inhibits the production of ergosterol followed by release of cellular ions, inhibition of methylglyoxal, <i>in situ</i> inhibition of lipid peroxidation. | Chaudhari et al. (2020) |
| Clove essential oil | Eugenol (EG) incorporation into β -cyclodextrin (β -CD) | <i>Peromophthora litchii</i> | Causes hyphal and/or sporangiphore cell wall and plasma membrane damage leading to cell shrinkage, wrinkling, and partial distortion. | Gong et al. (2016) |

dimension of the polymeric mesh, polarity and release of incorporated bioactive molecules can easily be manipulated by using different types of cross-linkers and varying the degree of cross-linking (Sherje et al., 2017; Pawar et al., 2019; Ananya et al., 2020). There are three widely known isomers of cyclodextrins, viz., alpha (α), beta (β) and gamma (γ) (Haimhoffer et al., 2019) that form inclusion and non-inclusion with different drugs to improve their stability, permeability, solubility and cytotoxicity (Osmani et al., 2014; Kumar et al., 2018).

Nanosponges can be used to mitigate challenges within the agricultural industry. For example, they can be used as biofungicides to replace synthetic pesticides currently used in agriculture to protect plants against pests and diseases for increased yield and quality of the agricultural products (Asghari et al., 2016). Nanosponges can be encapsulated with herbs that can be used for phytotoxicity problems on agricultural products. Asghari et al. (2016) reported the effects of nano-encapsulated herbicides against parasitic weeds. The herbicide encapsulated configuration allowed for the use of herbicides in different modes to selected areas since they were not affected (i.e. degraded) by the crop. The herbicides were applied in low doses and accumulated in the parasitic weed via the sink effect (Asghari et al., 2016). The use of nanosponges in mitigating fungal invasion and mycotoxin contamination can be crucial. Despite the beneficial properties of nanosponges and their wide applicability, very limited studies have investigated their use in mycotoxicology. Appell and Jackson (2012) investigated OTA sorption ability and characteristics of a β -cyclodextrin-based nanosponge (cyclodextrin-polyurethane polymer) in aqueous solutions. The results revealed that cyclodextrin nanosponge materials were effective at reducing OTA levels (between 1–10 $\mu\text{g/L}$) in spiked aqueous solutions. In fact, it was possible to reduce OTA levels from up to 10 $\mu\text{g/L}$ in spiked red wine to within the recommended levels of 2 $\mu\text{g/L}$. Furthermore, inference from Langmuir isotherm of sorption data revealed that nanosponge has maximum binding capacity of 220 μg OTA per g of polymer. Elsewhere, solid phase extraction sorbent material based on a nanosponge was developed with β -cyclodextrin and methylene bis-diphenyl diisocyanate in the ratio of 1:5 (Appell et al., 2018). The novel polymer (i.e. β -cyclodextrin polyurethane polymer) was used to extract and clean-up OTA from wine and grape juice. Recoveries of OTA in the spiked beverages (0.5–20 ng/mL) as found, ranged from 69.1–86.5% in grape juice and 77.0–89.4% in wine (Appell et al., 2018). In another study, polyurethane- β -cyclodextrin polymers, in particular, polymer cross-linked with tolylene 2,4-diisocyanate were synthesized and the complex presented populations of binding sites which was able to extract patulin from apple juice (Appell and Jackson, 2010).

Researchers from Hungary investigated the pharmacoeactive effects of β -cyclodextrin against the toxic effects of the xenoestrogenic mycotoxin, ZEN, in Tg (vtg1:mCherry) zebrafish embryos and HeLa cells. The results revealed that ZEN was able to form stable complexes with methyl-, sulfoethyl-, and succinyl-methyl-substituted β -cyclodextrins at pH 7.4 ($K = 1.4\text{--}4.7 \times 10^4$ L/mol). The resultant complexes (i.e., modified cyclodextrins) strongly reduced or eliminated ZEN-induced mortality in zebrafish embryos and loss of cell viability in HeLa cells. Co-treatment with β -cyclodextrins further alleviated the sub-lethal effects of ZEN (Faisal et al., 2020). A study by Fliszar-Nyul et al. (2019) investigated association of alternariol toxin with γ - and β -CDs using molecular and fluorescence spectroscopy techniques. They also examined the extraction of the mycotoxin alternariol from water-based (aqueous) solutions using β -CD bead polymer (BBP) that is insoluble. The results demonstrated that natural γ -CD at pH 7.4 firmly increased the fluorescence of alternariol, which formed most stable complexes with natural γ -CD and quaternary ammonium derivatives under acidic/physiological pH and at pH of 10.0. Furthermore, BBP successfully extracted alternariol toxin from aqueous medium. The alternariol-binding capacity of BBP and β -CD polymers were strongly increased because of the β -CD content of the polymer. The effect of selective encapsulation of star anise essential oil (SAEO) by hydroxypropyl- β -cyclodextrin (HPCD) on its composition, stability, volatility, and antibacterial activity was investigated and results showed

that encapsulation significantly decreased the irritating odour of SAEO and also improved the inhibitory effect of SAEO on *Saccharomyces cerevisiae*, *Rhizopus stolonifer*, and *E. coli* and its antibacterial stability in 24 h (Zhang et al., 2018).

Studies have shown that natural products encapsulated in nanosponges yielded better results than with natural products alone (Ansari et al., 2011; Kumar et al., 2018). Nanosponges have been used as a stable carrier for the encapsulation of various therapeutic agents. Different types of phytochemical bioactive compounds such as resveratrol, curcumin, quercetin, rutin, oryzanol, chlorogenic acid, etc. have been encapsulated in cyclodextrin nanosponges (Kumar et al., 2018; Osmani et al., 2018; Pawar et al., 2019). A study by Ramírez-Ambrosi et al. (2014) evaluated the efficacy of β -cyclodextrin nanosponges for encapsulation of polyphenols phloridzin, rutin, and chlorogenic acid. The results showed that rutin had an encapsulation efficiency of 83.7 % (which was the highest) using 1,1'-carbonyldiimidazole as cross-linker in a 1:3 ratio of nanosponge/cross-linker, while phloridzin (87.2%) and chlorogenic acid (77.5 %), with best results seen with HMDI. Radic et al. (2020) evaluated the influence of olive pomace extracts matrix and cyclodextrins (CDs) on bio-accessibility and intestinal permeability of main olive pomace polyphenols. Their results demonstrated that olive pomace polyphenols were stable during gastrointestinal digestion and encapsulation of olive pomace polyphenol with cyclodextrins significantly increased bioaccessibility of tyrosol by forming inclusion complexes and preventing tyrosol adhesion to bile salts and other macromolecules present in reaction mixtures during simulation of olive pomace extract digestion. Despite the acclaimed pharmacological efficacy, there has been no investigation on cyclodextrin-based nanosponges encapsulation of phytochemicals for applications as biofungicides and nutraceuticals against mycotoxigenic fungi and mycotoxins. Nanosponges may play crucial role in the detoxification of mycotoxins in this era of the fourth industrial revolution (4IR) due to their binding and neutralization capacity (Fliszar-Nyul et al., 2019). This is because, these cyclodextrin-based nanosponges can reduce toxic effects associated with several microorganisms including mycotoxigenic fungi by binding and neutralising attendant harmful secondary metabolites in the body without leaving any harmful effects.

Cyclodextrins as nanosponges have been extensively used in a variety of applications, and a new class of them include integrated hydrogel cellulose nanosponges referred to as nanoemulgels that have recently attracted the attention of many scientists in the discovery and development of a variety of drugs. They are polymeric emulsion systems that provide synergistic complimentary properties from nanosponges or nanofibers and hydrogels, thereby reinforcing the overall antifungal activity in the food packaging sector (Oun et al., 2019; Munteanu and Vasile, 2020). Aldawasri et al. (2015) formulated lemongrass-loaded ethyl cellulose nanosponges with a topical hydrogel that forms a protective coating on the applied surface, whilst the encapsulated lemongrass oil had an enhanced antifungal activity against *C. albicans*. Bandyopadhyay et al. (2020) developed an effective green packaging technology for food safety and quality of gouda cheese. This technology was based extending the shelf life of gouda cheese using cinnamaldehyde and eugenol essential oils integrated to polyvinylpyrrolidone-carboxymethyl cellulose-bacterial cellulose-guar gum (PVP-CMC-BC-GG) hydrogel film (Munteanu and Vasile, 2020). The results revealed that this technology extended the shelf-life of cheese by more than 10–12 days, with a high antifungal activity against the tested *Aspergillus* and *Candida* species. Moreover, nanoemulgel formulations are ideal within the agricultural sector because they possessed repellent activities against insects. Lehtonen et al. (2020) demonstrated that addition of hexanal in aerogel extended the shelf life of plant-based products (*i.e.*, blueberries and cherry tomatoes). The use of nanoemulgels in agriculture is still in its infancy but widely exploited in the medical sector. However, the popularity of nanoemulgels may likely increase as alternative antifungal agents in the agricultural sector. However, before that happens, it is imperative to conduct cycle analysis of any

nanoformulation for agricultural applications so as to evaluate their overall ecotoxicology.

4. Conclusion and future perspectives

The search for alternative mycotoxin detoxification strategies is of paramount interest in food safety. From the literature reviewed herein, it was noted that the use of botanical extracts and their phytochemicals coupled with nanosponge encapsulation technology can reduce/detoxify mycotoxins with little or no negative consequences. Only a few studies have been carried out on the decontamination of mycotoxins using nano-encapsulated bioactive compounds. Cyclodextrin based-nanosponge encapsulated plant extracts or bioactive compounds can improve the efficacy of plant extracts or their phytochemicals for decontamination of mycotoxins as reviewed in this paper. This approach further increases the bioavailability of benign bioactive compounds utilized in agriculture as environmentally-friendly fungicides, highly-effective at low concentration with strong antifungal and mycotoxin-inhibiting activities.

Future work should seek to develop methods that may enable the use of these agents as food and feed additives for the same purpose. Detailed studies on the mechanisms of interaction between nano-encapsulated bioactive compounds and food components, together with their effects on human and animal health, should also be explored. Indeed, the potential of such nanosponges and plant bioactive compounds to play a dual pharmacological and nutraceutical function as biofungicides and detoxifying agents in mitigating the effects mycotoxins is very interesting.

Declarations

Author contribution statement

All authors listed have significantly contributed to the development and the writing of this article.

Funding statement

This work was supported by the National Research Foundation (DSI/NRF), the joint LEAP-AGRI and NRF funding, and the University of Johannesburg, South Africa.

Competing interest statement

The authors declare no conflict of interest.

Additional information

No additional information is available for this paper.

References

- Abdel-Fattah, S.M., Badr, A.N., Seif, F.A.-H., Ali, S.M., Hassan, A., 2018. Antifungal and antimycotoxigenic impact of eco-friendly extracts of wild stevia. *J. Biol. Sci.* 18, 488–499.
- Abdel-Fattah, S.M., Abosrea, Y.H., Shehata, F.E., Flourage, M.R., Helal, A.D., 2010. The efficacy of thyme oil as antitoxicant of aflatoxin(s) toxicity in sheep. *Journal of American Science* 6, 948–960.
- Adebisi, J.A., Kayitesi, E., Adebisi, O.A., Changwa, R., Njobeh, P.B., 2019. Food fermentation and mycotoxin detoxification: an African perspective. *Food Contr.* 106, 106731.
- Adebisi, O.A., Njobeh, P.B., Gbashi, S., Nwinyi, O.C., Mavumengwana, V., 2017. Review on microbial degradation of aflatoxins. *Crit. Rev. Food Sci. Nutr.* 57, 3208–3217.
- Adebisi, O.A., Gabriela Medina-Meza, I., 2020. Impact of fermentation on the phenolic compounds and antioxidant activity of whole cereal grains: a mini review. *Molecules* 25, 927.
- Adebisi, O.A., Molelekoa, T., Makhuvele, R., Adebisi, J.A., Oyedeji, A.B., Gbashi, S., Adefisoye, M.A., Ogundele, O.M., Njobeh, P.B., 2020. A review on novel non-thermal food processing techniques for mycotoxin reduction. *Int. J. Food Sci. Technol.*
- Adekoya, I., Njobeh, P., Obadina, A., Landschoot, S., Audenaert, K., Okoth, S., De Boever, M., De Saeger, S., 2019. Investigation of the metabolic profile and toxigenic

- variability of fungal species occurring in fermented foods and beverage from Nigeria and South Africa using UPLC-MS/MS. *Toxins* 11, 85.
- Alabi, O., Anokwuru, C., Ezekiel, C., Ajibaye, O., Nwadike, U., Fasasi, O., Abu, M., 2011. Anti-mutagenic and anti-genotoxic effect of Ethanolic extract of neem on dietary-aflatoxin induced Genotoxicity in mice. *J. Biol. Sci.* 11, 307–313.
- Alberts, J., Rheeder, J., Gelderblom, W., Shephard, G., Burger, H., 2019. Rural subsistence maize farming in South Africa: risk assessment and intervention models for reduction of exposure to Fumonisin mycotoxins. *Toxins* 11, 334.
- Aldawasri, H.M., Badr-Eldin, S.M., Labib, G.S., El-Kamel, A.H., 2015. Design and formulation of a topical hydrogel integrating lemon grass-loaded nanosponges with an enhanced antifungal effect: in vitro/in vivo evaluation. *Int. J. Nanomed.* 10, 893–902.
- Alotaibi, M.A., Tayel, A.A., Zidan, N.S., El Rabey, H.A., 2019. Bioactive coatings from nano-biopolymers/plant extract composites for complete protection from mycotoxigenic fungi in dates. *J. Sci. Food Agric.* 99, 4338–4343.
- Atyn, I., Twaruzek, M., 2020. Mycotoxin contamination concerns of herbs and medicinal plants. *Toxins* 12, 182.
- Ananya, K.V., Preethi, S., Amit, B.P., Gowda, D.V., 2020. Recent review on Nano sponge. *Int. J. Res. Pharm. Sci.* 11 (1), 1085–1096.
- Anjorin, T.S., Salako, E.A., Makun, H.A., 2013. Control of Toxicogenic Fungi and Mycotoxins with Phytochemicals: Potentials and Challenges. INTECH, p. 181. Chapter 8.
- Ansari, K.A., Vavia, P.R., Trotta, F., Cavalli, R., 2011. Cyclodextrin-based nanosponges for delivery of resveratrol: in vitro characterisation, stability, cytotoxicity and permeation study. *AAPS PharmSciTech* 12, 279–286.
- Appell, M., Jackson, M.A., 2010. Synthesis and evaluation of cyclodextrin-based polymers for patulin extraction from aqueous solutions. *J. Inclusion Phenom. Macrocycl. Chem.* 68, 117–122.
- Appell, M., Jackson, M.A., 2012. Sorption of Ochratoxin A from aqueous solutions using β -cyclodextrin-Polyurethane polymer. *Toxins* 4, 98–109.
- Appell, M., Evans, K.O., Jackson, M.A., Compton, D.L., 2018. Determination of ochratoxin A in grape juice and wine using nanosponge solid phase extraction clean-up and liquid chromatography with fluorescence detection. *J. Liq. Chromatogr. Relat. Technol.* 41, 949–954.
- Asghari, F., Jahanshahi, Z., Imani, M., Shams-Ghahfarokhi, M., Razzaghi-Abyaneh, M., 2016. Antifungal nanomaterials: synthesis, properties, and applications. In: *Nanobiomaterials in Antimicrobial Therapy*. Elsevier, pp. 343–383.
- Ashiq, S., Hussain, M., Ahmad, B., 2014. Natural occurrence of mycotoxins in medicinal plants: a review. *Fungal Genet. Biol.* 66, 1–10.
- Aydin, S., Palabiyik, S.S., Erkekoglu, P., Sahin, G., Başaran, N., Giray, B.K., 2013. The carotenoid lycopene protects rats against DNA damage induced by ochratoxin A. *Toxicol.* 73, 96–103.
- Ayele, T.T., 2018. A review on traditionally used medicinal plants/herbs for cancer therapy in Ethiopia: current status, challenge and future perspectives. *Org. Chem. Curr. Res.* 7 (2).
- Azzouz, M.A., 1981. The Inhibitory Effects of Herbs, Spices and Other Plant Materials on Mycotoxigenic Moulds. Doctoral dissertation. University of Nebraska, Lincoln, NE, USA.
- Bahrami, A., Delshadi, R., Assadpour, E., Jafari, S.M., Williams, L., 2020. Antimicrobial-loaded nanocarriers for food packaging applications. *Adv. Colloid. Interface. Sci.* 278.
- Balendres, M.A.O., Karlovsky, P., Cumagun, C.J.R., 2019. Mycotoxigenic fungi and mycotoxins in agricultural crop commodities in the Philippines: a review. *Foods* 8 (7), 249.
- Bandyopadhyay, S., Saha, N., Zandrea, O., Pummerova, M., Saha, P., 2020. Essential oil based PVP-CMC-BC-GG functional hydrogel sachet for 'cheese': its shelf life confirmed with anthocyanin (Isolated from red cabbage) bio stickers. *Foods* 9, 307.
- Behbehani, J.M., Irshad, M., Shreaz, S., Karched, M., 2019. Synergistic effects of tea polyphenol epigallocatechin 3-O-gallate and azole drugs against oral *Candida* isolates. *J. Mycol. Med.* 29 (2), 158–167.
- Bennett, J.W., Klich, M., 2003. Mycotoxins. *Clin. Microbiol. Rev.* 16, 497–516.
- Bhattacharya, R., Firozi, P., 1988. Effect of plant flavonoids on microsomal catalyzed reactions of aflatoxin B₁ leading to activation and DNA adduct formation. *Canc. Lett.* 39, 85–91.
- Bhattachar, S., 2011. Natural antimutagens: a review. *Res. J. Med. Plant* 5, 116–126.
- Bhowmik, H., Venkatesh, D.N., Kuila, A., Kumar, K.H., 2018. Nanosponges: a review. *Int. J. Appl. Pharm.* 10, 1.
- Castellanos, L.M., Olivás, N.A., Ayala-Soto, J., Contreras, C.M.D., Ortega, M.Z., Salas, F.S., Ochoa, L.H., 2020. In vitro and in vivo antifungal Activity of clove (*Eugenia caryophyllata*) and pepper (*Piper nigrum* L.) essential oils and functional extracts against *Fusarium oxysporum* and *Aspergillus niger* in tomato (*Solanum lycopersicum* L.). *International Journal of Microbiology*.
- Celik, K., 2020. The efficacy of mycotoxin-detoxifying and biotransforming agents in animal nutrition. In: *Nanomycotoxicology*. Elsevier, pp. 271–284.
- Chaudhari, A.K., Singh, V.K., Deepika, S.D., Prasad, J., Dwivedy, A.K., Dubey, N.K., 2020. Improvement of in vitro and in situ antifungal, AFB₁ inhibitory and antioxidant activity of *Origanum majorana* L. essential oil through nanoemulsion and recommending as novel food preservative. *Food Chem. Toxicol.* 111536.
- Chen, E., Benso, B., Seleem, D., Ferreira, L.E.N., Pasetto, S., Pardi, V., Murata, R.M., 2018. Fungal-host interaction: curcumin modulates proteolytic enzyme activity of *Candida albicans* and inflammatory host response in vitro. *International Journal of Dentistry* 2018.
- Choi, H., Lee, D.G., 2015. Lycopene induces apoptosis in *Candida albicans* through reactive oxygen species production and mitochondrial dysfunction. *Biochimie* 115, 108–115.
- Čolović, R., Puvača, N., Cheli, F., Avantaggiato, G., Greco, D., Đuragić, O., Kos, J., Pinotti, L., 2019. Decontamination of Mycotoxin-contaminated feedstuffs and compound feed. *Toxins* 11 (11), 617.
- da Cruz Cabral, L., Fernández Pinto, V., Patriarca, A., 2013. Application of plant derived compounds to control fungal spoilage and mycotoxin production in foods. *Int. J. Food Microbiol.* 166, 1–14.
- da Silva Bomfim, N., Kohiyama, C.Y., Nakasugi, L.P., Nerilo, S.B., Mossini, S.A.G., Romoli, J.C.Z., Mikcha, J.M.G., de Abreu Filho, B.A., Machinski Jr., M., 2020. Antifungal and antifungal activity of rosemary essential oil (*Rosmarinus officinalis* L.) against *Aspergillus flavus*. *Food Addit. Contam.* 37 (1), 153–161.
- Das, S., Chaudhari, A.K., Deepika, A.S., Singh, V.K., Dwivedy, A.K., Dubey, N.K., 2020. 6- Foodborne microbial toxins and their inhibition by plant-based chemicals. *Functional and Preservative Properties of Phytochemicals*. Academic Press, pp. 165–207.
- Dhakar, N.K., Matencio, A., Caldera, F., Argenziano, M., Cavalli, R., Dianzani, C., Zanetti, M., Lopez-Nicolas, J.M., Trotta, F., 2019. Comparative evaluation of solubility, cytotoxicity and photostability studies of resveratrol and oxysresveratrol loaded nanosponges. *Pharmaceutics* 11, 545.
- Dias, D.A., Urban, S., Roessner, U., 2012. A historical overview of natural products in drug discovery. *Metabolites* 2, 303–336.
- Dikhoba, P., Mongalo, N., Elgorashi, E., Makhafola, T., 2019. Antifungal and antimycotoxigenic activity of selected South African medicinal plants species. *Heliyon* 5, e02668.
- Do, K., An, T., Oh, S., Moon, Y., 2015. Nation-based occurrence and endogenous biological reduction of mycotoxins in medicinal herbs and spices. *Toxins* 7, 4111–4130.
- El-Desouky, T., Ammar, H., 2016. Honey mediated silver nanoparticles and their inhibitory effect on aflatoxins and ochratoxin A. *J. Appl. Pharmaceut. Sci.* 6, 83–90.
- El-Hamss, R., Analla, M., Campos-Sanchez, J., Alonso-Moraga, A., Muñoz-Serrano, A., Idaomar, M., 1999. A dose dependent anti-genotoxic effect of turmeric. *Mutat. Res. Genet. Toxicol. Environ. Mutagen* 446, 135–139.
- El-Sawi, N.M., Al-Seeni, M.N., 2009. Assessment of flavonoids as rutin for detoxification of T-2 toxin. *J. Appl. Anim. Res.* 35, 57–60.
- El-Shayeb, N.M., 1980. Inhibition of aflatoxin formation by some spices. *Z. Lebensm. Unters. Forsch.* 171, 344–347.
- Eskola, M., Kos, G., Elliott, C.T., Hajslová, J., Mayor, S., Krska, R., 2019. Worldwide contamination of food-crops with mycotoxins: validity of the widely cited 'FAO estimate' of 25%. *Crit. Rev. Food Sci. Nutr.* 1–17.
- Faisal, Z., Garai, E., Csepregi, R., Bakos, K., Fliszár-Nyúl, E., Szente, L., Balázs, A., Cserhádi, M., Kószegi, T., Urbányi, B., Csenki, Z., Poór, M., 2020. Protective effects of beta-cyclodextrins vs. zearalenone-induced toxicity in HeLa cells and Tg(vtg1:mCherry) zebrafish embryos. *Chemosphere* 240, 124948.
- Farombi, E.O., 2004. Diet-related cancer and prevention using anticarcinogens. *Afr. J. Biotechnol.* 3, 651–661.
- Fliszár-Nyúl, E., Lemli, B., Szente, L., Poor, M., 2019. Interactions of mycotoxins alternariol with cyclodextrins and its removal from aqueous solution by beta-cyclodextrin bead polymer. *Biomolecules* 9, 428.
- Gacem, M.A., Gacem, H., Telli, A., Khelil, A.O., 2020. Mycotoxins: decontamination and nanocontrol methods. In: *Nanomycotoxicology*. Elsevier, pp. 189–196.
- Galvano, F., Piva, A., Ritieni, A., Galvano, G., 2001. Dietary strategies to counteract the effects of mycotoxins: a review. *J. Food Protect.* 64, 120–131.
- Gbashi, S., Madala, N.E., De Saeger, S., De Boevre, M., Adekoya, I., Adebo, O.A., Njobeh, P.B., 2019. The socio-economic impact of mycotoxin contamination in Africa. In: Njobeh, P.B., Stepman, F. (Eds.), *Mycotoxins: Impact and Management Strategies*. InTech, Rijeka, Croatia, pp. 3–22.
- Gong, L., Li, T., Chen, F., Duan, X., Yuan, Y., Zhang, D., Jiang, Y., 2016. An inclusion complex of eugenol into β -cyclodextrin: preparation, and physicochemical and antifungal characterization. *Food Chem.* 196, 324–330.
- Gowda, N.K., Ledoux, D.R., Rottinghaus, G.E., Bermudez, A.J., Chent, Y.C., 2008. Efficacy of tumeric (*Curcuma longa*), containing a known level of curcumin, and a hydrated sodium calcium aluminosilicate to ameliorate the adverse effects of aflatoxin in broiler chicks. *Nutr. Metab.* 87, 1125–1130.
- Gross-Steinmeyer, K., Eaton, D.L., 2012. Dietary modulation of the biotransformation and genotoxicity of aflatoxin B₁. *Toxicology* 299, 69–79.
- Gyamfi, M.A., Aniya, Y., 1998. Medicinal herb, *Thonningia sanguinea* protects against aflatoxin B₁ acute hepatotoxicity in Fischer 344 rats. *Hum. Exp. Toxicol.* 17, 418–423.
- Haimhoffer, Á., Rusznayk, Á., Réti-Nagy, K., Vasvári, G., Váradi, J., Vecsernyés, M., Bácskay, I., Fehér, P., Ujhelyi, Z., Fenyvesi, F., 2019. Cyclodextrins in drug delivery systems and their effects on biological barriers. *Sci. Pharm.* 87, 33.
- Haq, M.A., Wang, Y., Shen, Z., Li, X., Saleemi, M.K., He, C., 2020. Mycotoxin contamination and control strategy in human, domestic animal and poultry: a review. *Microb. Pathog.* 104095.
- Hedayati, N., Naeini, M.B., Nezami, A., Hosseinzadeh, H., Wallace Hayes, A., Hosseini, S., Imenshahidi, M., Karimi, G., 2018. Protective effect of lycopene against chemical and natural toxins: a review. *Biofactors* 45, 5–23.
- Hitokoto, H., Morozumi, S., Wauke, T., Sakai, S., Kurata, H., 1980. Inhibitory effects of spices on growth and toxin production of toxigenic fungi. *Appl. Environ. Microbiol.* 39, 818–822.
- Hojnik, N., Cvelbar, U., Tavcar-Kalcher, G., Walsh, J., Krizaj, I., 2017. Mycotoxin decontamination of food: cold atmospheric pressure plasma versus "Classic" decontamination. *Toxins* 9, 151.
- Horn, R.C., Vargas, V.M., 2008. Mutagenicity and antimutagenicity of teas used in popular medicine in the Salmonella/microsome assay. *Toxicol. Vitro* 22, 1043–1049.

- Hosseini, A., Hosseinzadeh, H., 2018. Antidotal or protective effects of curcuma longa (turmeric) and its active ingredient, curcumin, against natural and chemical toxicities: a review. *Biomed. Pharmacother.* 99, 411–421.
- Hu, Y., Luo, J., Kong, W., Zhang, J., Logrieco, A.F., Wang, X., Yang, M., 2015. Uncovering the antifungal components from turmeric (*Curcuma longa* L.) essential oil as *Aspergillus flavus fumigants* by partial least squares. *RSC Adv.* 5 (52), 41967–41976.
- Hussain, A., Shafiqatullah, Ali, J., Zia-Ur-Rehman, 2012. Inhibition of aflatoxin producing fungus growth using chemical, herbal compounds/spices and plants. *Pure and Applied Biology* 1, 8–13.
- Inano, H., Onoda, M., 2002. Prevention of radiation induced mammary tumors. *International Journal of Radiation Oncology* 52, 212–223.
- Ip, S.P., Mak, D.H., Li, P.C., Poon, M.K., Ko, K.M., 1999. Effect of a lignan-enriched extract of Schisandra Chinensis aflatoxin B₁ and cadmium chloride-induced Hepatotoxicity in rats. *Pharmacol. Toxicol.* 78, 413–416.
- Iram, W., Anjum, T., Iqbal, M., Ghaffar, A., Abbas, M., 2015. Mass spectrometric identification and toxicity assessment of degraded products of aflatoxin B₁ and B₂ by *Corymbia citriodora* aqueous extracts. *Sci. Rep.* 5 (1).
- Iram, W., Anjum, T., Iqbal, M., Ghaffar, A., Abbas, M., 2016. Structural elucidation and toxicity assessment of degraded products of aflatoxin B₁ and B₂ by aqueous extracts of *Trachyspermum ammi*. *Front. Microbiol.* 7.
- Juodeikiene, G., Basinskiene, L., Bartkiene, E., Matusевичius, P., 2012. Mycotoxin decontamination aspects in food, feed and renewables using fermentation processes. *Structure and Function of Food Engineering* 171–204.
- Kagot, V., Okoth, S., De Boevre, M., De Saeger, S., 2019. Biocontrol of *Aspergillus* and *Fusarium* mycotoxins in Africa: benefits and limitations. *Toxins* 11 (2), 109.
- Karlovska, P., Suman, M., Berthiller, F., De Meester, J., Eisenbrand, G., Perrin, I., Oswald, I.P., Speijers, G., Chiodini, A., Recker, T., Dussort, P., 2016. Impact of food processing and detoxification treatments on mycotoxin contamination. *Mycotoxin Res.* 32, 179–205.
- Kavitha, K., Vijaya, N., Krishnaveni, A., Arthanareeswari, M., Rajendran, S., Al-Hashem, A., Subramania, A., 2020. Nanomaterials for antifungal applications. *Nanotoxicity* 385–398.
- Khan, H., Ullah, H., Nabavi, S.M., 2019. Mechanistic insights of hepatoprotective effects of curcumin: therapeutic updates and future prospects. *Food Chem. Toxicol.* 124, 182–191.
- Kocic-Tanackov, S., Dimic, G., Jaksic, S., Mojovic, L., Djukic-Vukovic, A., Mladenovic, D., Pejcin, J., 2019. Effects of caraway and juniper essential oils on aflatoxigenic fungi growth and aflatoxins secretion in polenta. *J. Food Process. Preserv.* 43.
- Kumar, S., Sihag, P., Trotta, F., Rao, R., 2018. Encapsulation of Babchi oil in cyclodextrin-based nanospheres: physicochemical characterization, photodegradation and *in vitro* cytotoxicity studies. *Pharmaceutics* 10, 169.
- Lahlou, Y., Rhandour, Z., Amraoui, B.E., Bamhaoud, T., 2019. Screening of antioxidant activity and the total polyphenolic contents of six medicinal Moroccan plants extracts. *J. Mater. Environ. Sci.* 10, 133–1348.
- Leal, M., Shimada, A., Ruíz, F., González, E., Mejía, D., 1999. Effect of lycopene on lipid peroxidation and glutathione-dependent enzymes induced by T-2 toxin *in vivo*. *Toxicol. Lett.* 109, 1–10.
- Lehtonen, M., Kekalainen, S., Nikkila, I., Kilpelainen, P., Tenkanen, M., Mikkonen, K.S., 2020. Active food packaging through controlled in situ production and release of hexanal. *Food Chem.* (5), 100074.
- Liew, W.-P., Mohd-Redzwan, S., 2018. Mycotoxin: its impact on gut health and microbiota. *Frontiers in cellular and infection microbiology* 8, 60.
- Limaye, A., Yu, R.-C., Chou, C.-C., Liu, J.-R., 2018. Protective and detoxifying effects conferred by dietary selenium and curcumin against AFB₁-mediated toxicity in livestock: a review. *Toxins* 10, 25.
- Loi, M., Paciolla, C., Logrieco, A., Mule, G., 2020. Plant bioactive compounds in pre- and post-harvest management for aflatoxins reduction. *Front. Microbiol.* 11.
- Luo, Y., Liu, X., Li, J., 2018. Updating techniques on controlling mycotoxins - a review. *Food Contr.* 89, 123–132.
- Lyagin, I., Efremenko, E., 2019. Enzymes for detoxification of various mycotoxins: origins and mechanisms of catalytic action. *Molecules* 24 (13), 2362.
- Madrigal-Santillan, E., Morales-Gonzalez, J.A., Vargas-Mendoza, N., Reyes-Ramirez, P., Cruz-Jaime, S., Sumaya-Martinez, T., Perez-Pasten, R., Madrigal-Bujaidar, E., 2010. Antigenotoxic studies of different substances to reduce the DNA damage induced by aflatoxin B₁ and ochratoxin A. *Toxins* 2, 738–757.
- Mahato, D.K., Lee, K.E., Kamle, M., Devi, S., Dewangan, K., Kumar, P., Kang, S.G., 2019. Aflatoxins in food and feed: an overview on prevalence, detection and control strategies. *Front. Microbiol.* 10.
- Makhafola, T.J., Elgorashi, E.E., McGaw, L.J., Awouafack, M.D., Verschaeve, L., Eloff, J.N., 2017. Isolation and characterization of the compounds responsible for the antimutagenic activity of Combretum microphyllum (Combretaceae) leaf extracts. *BMC Compl. Alternative Med.* 17.
- Makhuvele, R., Matshoga, R.G., Antonissen, R., Pieters, L., Verschaeve, L., Elgorashi, E.E., 2018a. Genotoxicity and antigenotoxicity of selected South African indigenous plants. *South Afr. J. Bot.* 114, 89–99.
- Makhuvele, R., Foubert, K., Apers, S., Pieters, L., Verschaeve, L., Elgorashi, E., 2018b. Antimutagenic constituents from *Monanthonotaxis caffra* (sond.) verdc. *J. Pharm. Pharmacol.* 79, 976–984.
- Marroquin-Cardona, A.G., Johnson, N.M., Phillips, T.D., Hayes, A.W., 2014. Mycotoxins in a changing global environment- A review. *Food Chem. Toxicol.* 69, 220–230.
- Mathuria, N., Verma, R.J., 2007. Aflatoxin induced hemolysis and its amelioration by tumeric extracts and curcumin *in vitro*. *Acta Poloniae Pharmaceutica Drug Research* 64, 165–168.
- Meng, D., Garba, B., Ren, Y., Yao, M., Xia, X., Li, M., Wang, Y., 2020. Antifungal activity of chitosan against *Aspergillus ochraceus* and its possible mechanisms of action. *Int. J. Biol. Macromol.* 158, 1063–1070.
- Milicevic, D.R., Skrinjar, M., Baltic, T., 2010. Real and perceived risks for mycotoxin contamination in foods and feeds: challenges for food safety control. *Toxins* 2, 572–592.
- Moghadamtousi, S.K., Kadir, H.A., Hassandarvish, P., Tajik, H., Abubakar, S., Zandi, K.A., 2014. Review on Antibacterial, Antiviral, and Antifungal Activity of Curcumin. *BioMed Research International*, p. 186864.
- Moura-Costa, G.F., Nocchi, S.R., Ceole, L.F., de Mello, J.C., Nakamura, C.V., Filho, B.P., Temponi, L.G., Ueda-Naamura, T., 2012. Antimicrobial activity of plants used as medicinals on an indigenous reserve in Rio das cobras, Parana, Brazil. *J. Ethnopharmacol.* 143, 631–638.
- Mulaudzi, R.B., 2019. Mycotoxins of medicinal plants and human health. *Elsevier* 524–530.
- Munteanu, S.B., Vasile, C., 2020. Vegetable additives in food packaging polymeric materials. *Polymers* 12, 28.
- Narayanan, V.S., Muddaiah, S., Shashidara, R., Sudheendra, U.S., Deepthi, N.C., Samaranayake, L., 2020. Variable antifungal activity of curcumin against planktonic and biofilm phase of different candida species. *Indian J. Dent. Res.* 31, 145–148.
- Nayak, S., Sashidhar, R.B., 2010. Metabolic intervention of aflatoxin B₁ toxicity by curcumin. *J. Ethnopharmacol.* 127, 641–644.
- Neelofar, K., Shreaz, S., Rimple, B., Muralidhar, S., Nikhat, M., Khan, L.A., 2011. Curcumin as a promising anticandidal of clinical interest. *Can. J. Microbiol.* 57 (3), 204–210.
- Negera, M., Washe, A.P., 2019. Use of natural dietary spices for reclamation of food quality impairment by aflatoxin. *Hindawi Journal of Food Quality* 1–10.
- Nerilo, S.B., Romoli, J.C.Z., Nakasugi, L.P., Zampieri, N.S., Mossini, S.A.G., Rocha, G.H.O., Micotti da, E.G., Abreu Filho, B.A., Machinski Jr., M., 2020. Antifungal activity and inhibition of aflatoxins production by Zingiber officinale Roscoe essential oil against *Aspergillus flavus* in stored maize grains. *Ciência Rural.* 50 (6), e20190779, 11.
- Nleya, N., Adetunji, M.C., Mwanza, M., 2018. Current status of mycotoxin contamination of food commodities in Zimbabwe. *Toxins* 10 (5), 89.
- Oliveira, R.C., Carvajal-Moreno, M., Mercado-Ruaro, P., Rojo-Callejas, F., Correa, B., 2020. Essential oils trigger an antifungal and anti-aflatoxigenic effect on *Aspergillus flavus* via the induction of apoptosis-like cell death and gene regulation. *Food Contr.* 110, 107038.
- Osmani, R.A., Thirumaleswar, S., Bhosale, R.R., Kulkarni, P.K., 2014. Nanospheres: the spanning accession in drug delivery- an updated comprehensive review. *Der Pharm. Lett.* 5, 7–21.
- Osmani, R.A., Kulkarni, P., Manjunatha, S., Vaghela, R., Bhosale, R., 2018. Cyclodextrin nanosphere-based systems in drug delivery and nanotherapeutics: current progress and future prospects. In: *Organic Materials as Smart Nanocarriers for Drug Delivery*, pp. 659–704.
- Oun, A.A., Shankar, S., Rhim, J.-W., 2019. Multifunctional nanocellulose/metal and metal oxide nanoparticle hybrid nanomaterials. *Crit. Rev. Food Sci. Nutr.* 60, 435–460.
- Palabiyik, S.S., Erkekoglu, P., Zeybek, N.D., Kizilgun, M., Baydar, D.E., Sahin, G., Giray, B.K., 2013. Protective effect of lycopene against ochratoxin A induced renal oxidative stress and apoptosis in rats. *Exp. Toxicol. Pathol.* 65, 853–861.
- Palombo, E.A., 2011. Traditional medicinal plant extracts and natural products with activity against oral bacteria: potential application in the prevention and treatment of oral diseases. *Evid. base Compl. Alternative Med.* 1–15.
- Pawar, A.Y., Naik, A.K., Jadhav, K.R., 2016. Nanospheres: a novel drug delivery system. *Asian J. Pharm. Pharmacol.* 10, S456–S463.
- Pawar, S., Shende, P., Trotta, F., 2019. Diversity of β -cyclodextrin-based nanospheres for transformation of actives. *Int. J. Pharm.* 565, 333–350.
- Perincherry, L., Lalak-Kanczugowska, J., Stepien, L., 2019. Fusarium-produced mycotoxins in plant-pathogen interactions. *Toxins* 11.
- Polasa, K., Naidu, A.N., Ravindranath, I., Krishnaswamy, K., 2004. Inhibition of B(a)P induced strand breaks in presence of curcumin. *Mutat. Res.* 557, 203–213.
- Powers, C.N., Satyal, P., Mayo, J.A., McFeeters, H., McFeeters, R.L., 2019. Bigger data approach to analysis of essential oils and their antifungal activity against *Aspergillus niger*, *Candida albicans*, and *Cryptococcus neoformans*. *Molecules* 24 (16).
- Phuna, Z.X., Yu, J.K.E., Tee, J.Y., Chuah, S.Q., Tan, N.W.H., Vijayabalan, S., Manap, A.S.A., Sisinthy, S.P., Madhavan, P., 2020. *In vitro* evaluation of nanoemulsions of curcumin, piperine, and tualang honey as antifungal agents for *Candida* species. *Journal of Applied Biotechnology Reports*.
- Ponzilacqua, B., Rottinghaus, G.E., Landers, B.R., Oliveira, C.F., 2019. Effects of medicinal herb and Brazilian traditional plant extracts on *in vitro* mycotoxin decontamination. *Food Contr.* 100, 24–27.
- Prakash, B., Kumar, A., Singh, P.P., Sogachan, L.S., 2020. Antimicrobial and antioxidant properties of phytochemicals: current status and future perspective. *Functional and Preservative Properties of Phytochemicals*. Elsevier, pp. 1–45.
- Prakash, B., Kedia, A., Mishra, P.K., Dubey, N.K., 2015. Plant essential oils as food preservatives to control moulds, mycotoxin contamination and oxidative deterioration of agri-food commodities- Potentials and challenges. *Food Contr.* 47, 381–391.
- Pushpalatha, R., Selvamuthukumar, S., Kilimozhi, D., 2018. Cross-linked, cyclodextrin-based nanospheres for curcumin delivery- Physicochemical characterization, drug release, stability and cytotoxicity. *J. Drug Deliv. Sci. Technol.* 45, 45–53.
- Radic, K., Dukovski, B.J., Cepo, D.V., 2020. Influence of pomace matrix and cyclodextrin encapsulation on olive pomace polyphenols' bioaccessibility and intestinal permeability. *Nutrients* 12, 669.

- Rai, M., Ingle, A.P., Pandit, R., Paralikar, P., Anasane, N., Santos, C.A.D., 2020. Curcumin and curcumin-loaded nanoparticles: antipathogenic and antiparasitic activities. *Expert Rev. Anti Infect. Ther.* 18 (4), 367–379.
- Ramírez-Ambrosi, M., Caldera, F., Trotta, F., Berrueta, L.Á., Gallo, B., 2014. Encapsulation of apple polyphenols in β -CD nanospheres. *J. Inclusion Phenom. Macrocycl. Chem.* 80, 85–92.
- Reddy, K.N., Nurdijati, S.B., Salleh, B., 2010. An overview of plant-derived products on control of mycotoxigenic fungi and mycotoxins. *Asian J. Plant Sci.* 9, 126–133.
- Redondo-Blanco, S., Fernandez, J., Lopez-Ibanez, S., Miguez, E.M., Villar, C.J., Lombo, F., 2019. Plant phytochemicals in food preservation: antifungal bioactivity: a review. *J. Food Protect.* 83 (1), 163–171.
- Sadeghi-Ghadi, Z., Vaezi, A., Ahangarkani, F., Ilkit, M., Ebrahimnejad, P., Badali, H., 2020. Potent *in vitro* activity of curcumin and quercetin co-encapsulated in nanovesicles without hyaluronan against *Aspergillus* and *Candida* isolates. *J. Mycol. Med.* 101014.
- Sadhasivam, S., Shapiro, O.H., Ziv, C., Barda, O., Zakin, V., Sionov, E., 2019. Synergistic inhibition of mycotoxigenic fungi and mycotoxin production by combination of pomegranate peel extract and azole fungicide. *Front. Microbiol.* 10, 1919.
- Sajid, M., Cameotra, S.S., Sajjad, M., Khan, A., Ahmad, I., 2019. Nanoparticle-based delivery of phytomedicines: challenges and opportunities. In: *New Look to Phytomedicine*. Elsevier, pp. 597–623.
- Sharma, M., Manoharlal, R., Puri, N., Prasad, R., 2010. Antifungal curcumin induces reactive oxygen species and triggers an early apoptosis but prevents hyphae development by targeting the global repressor TUP1 in *Candida albicans*. *Biosci. Rep.* 30 (6), 391–404.
- Shehata, M.G., Badr, A.N., Abdel-Razek, A.G., Hassanein, M.M., Amra, H.A., 2017. Oil-bioactive films as an antifungal application to save post-harvest food crops. *Annual Research and Review in Biology* 16, 1–16.
- Sherje, A.P., Dravyakar, B.R., Kadam, D., Jadhav, M., 2017. Cyclodextrin-based nanospheres: a critical review. *Carbohydr. Polym.* 173, 37–49.
- Shin, B., Park, W., 2018. Zoonotic diseases and phytochemical medicines for microbial infections in veterinary sciences: current state and future perspective. *Frontiers in Veterinary Science* 5.
- Singh, C., Prakash, C., Mishra, P., Tiwari, K.N., Mishra, S.K., More, R.S., Kumar, V., Singh, J., 2019. Hepatoprotective efficacy of *Premna integrifolia* L. leaves against aflatoxin B₁-induced toxicity in mice. *Toxicology* 166, 88–100.
- Smerak, P., Polivkova, Z., Sestakova, H., Stetina, R., Barta, I., Langova, M., Turek, B., Bartova, J., 2006a. Antimutagenic effect of curcumin and its effect on the immune response in mice. *Czech J. Food Sci.* 24, 72–83.
- Smerak, P., Sestakova, H., Polivkova, Z., Stetina, R., Langova, M., Barta, I., Turek, B., Bartova, J., 2006b. Antimutagenic effect of Epigallocatechin gallate and its effect on the immune response in mice. *Czech J. Food Sci.* 24, 180–192.
- Song, L., Zhang, F., Yu, J., Wei, C., Han, Q., Meng, X., 2020. Antifungal effect and possible mechanism of curcumin mediated photodynamic technology against *Penicillium expansum*. *Postharvest Biol. Technol.* 167 (2020), 111234.
- Soni, K.B., Lahiri, M., Chackradeo, P., Bhide, S.V., Kuttan, R., 1997. Protective effect of food additives on aflatoxin-induced mutagenicity and hepatocarcinogenicity. *Canc. Lett.* 115, 129–133.
- Sorrenti, V., Di Giacomo, C., Acquaviva, R., Bognanno, M., Grilli, E., D'Orazio, N., Galvano, F., 2012. Dimethylarginine dimethylaminohydrolase/nitric oxide synthase pathway in liver and kidney: protective effect of cyanidin 3-O- β -D-glucoside on ochratoxin-A toxicity. *Toxins* 4, 353–363.
- Sridhar, M., Suganthi, R.U., Thammiah, V., 2015. Effect of dietary resveratrol in ameliorating aflatoxin B₁-induced changes in broiler birds. *J. Anim. Physiol. Anim. Nutr.* 99, 1094–1104.
- Stoev, S.D., Njobeh, P., Zarkov, I., Mircheva, T., Zapryanova, D., Denev, S., Dimitrova, B., 2019. Selected herbal feed additives showing protective effects against ochratoxin A toxicosis in broiler chicks. *World Mycotoxin J.* 12, 257–268.
- Street, R.A., Prinsloo, G., 2013. Commercially important medicinal plants of South Africa: a review. *J. Chem.* 1–16.
- Tabeshpour, J., Mehri, S., Behbahani, F.S., Hosseinzadeh, H., 2018. Protective effects of *Vitis vinifera* (grapes) and one of its biologically active constituents, resveratrol, against natural and chemical toxicities: a comprehensive review. *Phytother. Res.* 32, 2164–2190.
- Thipe, V.C., Keyster, M., Katti, K.V., 2018. Sustainable nanotechnology: mycotoxin detection and protection. In: Abd-El Salam, K.A. (Ed.), *Nanobiotechnology Applications in Plant Protection, Nanotechnology in the Life Sciences*, pp. 323–349.
- Thipe, V.C., Panjtan Amiri, K., Bloebaum, P., Raphael Karikachery, A., Khoobchandani, M., Katti, K.K., Jurisson, S.S., Katti, K.V., 2019. Development of resveratrol-conjugated gold resveratrol corona on anti-tumor efficacy against nanoparticles: interrelationship of increased breast, pancreatic and prostate cancers. *Int. J. Nanomed.* 14, 4413–4428.
- Thipe, V.C., Bloebaum, P., Khoobchandani, M., Karikachery, A.R., Katti, K.K., Katti, K.V., 2020. Green nanotechnology: nanoformulations against toxigenic fungi to limit mycotoxin production. In: Rai, M., Abd-El Salam (Eds.), *Nanomycotoxicology Treating Mycotoxins in the Nano Way*. Elsevier, pp. 155–188.
- Tian, Y., Tan, Y., Liu, N., Liao, Y., Sun, C., Wang, S., Wu, A., 2016. Functional agents to biologically control deoxynivalenol contamination in cereal grains. *Front. Microbiol.* 7.
- Udomkun, P., Wiredu, A.N., Nagle, M., Muller, J., Vanlauwe, B., Bandyopadhyay, R., 2017. Innovative technologies to manage aflatoxins in foods and feeds and the profitability of application- A review. *Food Contr.* 76, 127–138.
- Udomkun, P., Njukwe, E., 2020. Nanotechnological methods for aflatoxin control. In: *Nanomycotoxicology*, pp. 385–396.
- Van Wyk, B.-E., Van Oudtshoorn, B., Gericke, B., 1997. *Medicinal Plants of South Africa*. Briza, Publications, Pretoria.
- Velu, G., Palanichamy, V., Rajan, A.P., 2018. Phytochemical and pharmacological importance of plant secondary metabolites in modern medicine. In: Roopan, S.M., Madhumitha, G. (Eds.), *Bioorganic Phase in Natural Food: an Overview*. Springer International Publishing AG, pp. 135–156.
- Venkatesh, N., Keller, N.P., 2019. Mycotoxins in conversation with bacteria and fungi. *Front. Microbiol.* 10.
- Verschaeve, L., Van Staden, J., 2008. Mutagenic and antimutagenic properties of extracts from South African traditional medicinal plants. *J. Ethnopharmacol.* 119, 575–587.
- Vila-Donat, P., Marín, S., Sanchis, V., Ramos, A., 2018. A review of the mycotoxin adsorbing agents, with an emphasis on their multi-binding capacity, for animal feed decontamination. *Food Chem. Toxicol.* 114, 246–259.
- Vipin, A.V., Raksha, R.K., Nawneet, K.K., Anu, A.K., Venkateswaran, G., 2017. Protective effects of phenolics rich extract of ginger against aflatoxin B₁-induced. *Biomed. Pharmacother.* 91, 415–424.
- World Health Organization (WHO), 2001. *Legal Status of Traditional Medicine and Complementary/Alternative Medicine: A World Review*. Geneva.
- Wu, Q., Wang, X., Nepovimova, E., Wang, Y., Yang, H., Li, L., Zhang, X., Kuca, K., 2017a. Antioxidant agents against trichothecenes: new hints for oxidative stress treatment. *Oncotarget* 8, 110708–110726.
- Wu, J.-C., Lai, C.-S., Tsai, M.-L., Ho, T., Wang, Y.-J., Pan, M.-H., 2017b. Chemopreventive effect of natural dietary compounds on xenobiotic-induced toxicity. *J. Food Drug Anal.* 25, 176–186.
- Xu, J.-L., Li, L.-W., Luo, Y.-X., Yuan, S.-H., Liu, N.-N., 2020. Antifungal nanomaterials: current progress and future directions. *Innovations in Digital Health, Diagnostics, and Biomarkers* 1.
- Yang, G., Zhong, L., Jiang, L., Geng, C., Cao, J., Sun, X., Liu, X., Chen, M., Ma, Y., 2011. 6-gingerol prevents patulin-induced genotoxicity in HepG2 cells. *Phytother. Res.* 25, 1480–1485.
- Yang, Y., Huan, C., Liang, X., Fang, S., Wang, J., Chen, J., 2019. Development of starch-based antifungal coatings by incorporation of natamycin/methyl- β -cyclodextrin inclusion complex for postharvest treatments on cherry tomato against *Botrytis cinerea*. *Molecules* 24 (21), 3962.
- Zain, M.E., 2011. Impact of mycotoxins on humans and animals. *Journal of Saudi Chemical Society* 15, 129–144.
- Zhang, G., Yuan, C., Sun, Y., 2018. Effect of selective encapsulation of hydroxypropyl- β -cyclodextrin on components and antibacterial properties of star anise essential oil. *Molecules* 23.
- Zhu, C., Lei, M., Andargie, M., Zeng, J., Li, J., 2019. Antifungal activity and mechanism of action of tannic acid against *Penicillium digitatum*. *Physiol. Mol. Plant Pathol.* 107, 46–50.