

ORIGINAL ARTICLE

Supplementation of fetal bovine serum increased the quality of *in vitro* fertilized porcine embryo

Dibyendu Biswas^{1,2} , Sang Hwan Hyun¹ 

¹Laboratory of Veterinary Embryology and Biotechnology, College of Veterinary Medicine, Chungbuk National University, Chungbuk 28644, Republic of Korea

²Department of Medicine, Surgery and Obstetrics, Faculty of Animal Science and Veterinary Medicine, Patuakhali Science and Technology University, Barisal campus, Bangladesh

ABSTRACT

Objective: The present study aimed to explain the effect of fetal bovine serum (FBS) on the *in vitro* production of porcine embryos and the molecular effects of FBS on the growing of porcine embryos.

Materials and Methods: Immature porcine oocytes were matured and fertilized *in vitro*. The resulting zygotes were cultured in porcine zygotic medium-3- until day 7 and FBS was added on day 4. Without FBS, it was treated as a control group. Quantitative real-time PCR and 2',7'-dichloro-dihydro-fluorescein diacetate (H₂DCFDA) molecular staining techniques were used to detect the expression patterns of apoptosis-associated genes and the accumulation of reactive oxygen species (ROS), respectively. Paired student's *t*-test was used by GraphPad Prism statistical software.

Results: FBS supplementation boosted blastocyst (BL) development and total cell count per BL substantially ($p < 0.05$). However, hatching and hatched BLs also increased in the FBS-treated group compared to the control. We also found that ROS accumulation in FBS-treated embryos was significantly reduced ($p < 0.05$) compared to the control group. The expression of the anti-apoptotic gene BCL-2 was significantly increased in FBS-treated BLs, but the pro-apoptotic gene, caspase-3 expression, was significantly reduced in FBS-treated BLs.

Conclusion: Our results suggest that FBS supplementation in porcine culture media could increase porcine embryo production by decreasing ROS accumulation and increasing the anti-apoptotic gene expression in developing BLs.

ARTICLE HISTORY

Received July 05, 2021
Revised September 26, 2021
Accepted October 07, 2021
Published November 01, 2021

KEYWORDS

FBS; BCL-2; ROS; PZM-3



© The authors. This is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 License (<http://creativecommons.org/licenses/by/4.0>)

Introduction

In vitro porcine embryo production is essential for biomedical research as pigs are used for xenotransplantation research concerning the new drug discovery process [1]. However, unlike *in vivo* embryo production, *in vitro* high-quality embryo production is very challenging. To beat this challenge, many attempts were introduced to develop gold standard protocols for embryo production. Supplementing different growth factors/hormones or other nutrients during the developmental stage can reduce the stress caused by *in vitro* conditions [2,3]. During *in*

vitro embryo production, some byproducts are generated inside embryos that are detrimental to their development. However, they are very unstable and highly reactive with other molecules that also donate or accept an electron. These free radicals include nitric oxide, superoxide anion, hydrogen peroxide, oxygen singlet, hydroxyl, hypochlorite, and peroxynitrite [4]. These free radicals mostly target the vulnerable macromolecules and cause cell damage, thus disrupting the embryo's homeostasis environment.

Several research groups have developed different embryo culture media, such as North Carolina State

Correspondence Dibyendu Biswas ✉ dipupstu2012@pstu.ac.bd 📧 Laboratory of Veterinary Embryology and Biotechnology, College of Veterinary Medicine, Chungbuk National University, Chungbuk 28644, Republic of Korea.

Sang Hwan Hyun ✉ shhyun@cbu.ac.kr 📧 Laboratory of Veterinary Embryology and Biotechnology, College of Veterinary Medicine, Chungbuk National University, Chungbuk 28644, Republic of Korea.

How to cite: Biswas D, Hyun SH. Supplementation of fetal bovine serum increased the quality of *in vitro* fertilized porcine embryo. J Adv Vet Anim Res 2021; 8(4):589–597.

University (NCSU) 23 and 37 media [5], Iowa State University medium [6], and porcine zygotic medium-3 (PZM-3) [7], to be studied for *in vitro* culture (IVC) of porcine embryos. However, NCSU23 and PZM-3 media have been widely used to culture porcine embryos. Porcine zygotes of various origins, either *in vitro* fertilization (IVF), parthenogenesis, or somatic cell nuclear transfer, have been reported to have higher development rates in PZM-3 media than in NCSU23 media [8]. It was also reported that the embryonic development rate was increased when the medium was supplemented with fetal bovine serum (FBS) [9] or fetal calf serum (FCS) [10]. It was proved that supplementation of animal serum acts as an energy source, osmoregulator, pH stabilizers, numerous antioxidants, and growth factors [11,12].

During the early growing phase, the embryo needs more energy due to the presence of more blastomeres in the embryo. Embryo development and hatching rates were increased when FBS or FCS was integrated into the embryo culture media during the mouse and rat model experiments [13]. However, the mechanism underlying how the integration of FBS into IVC medium enhances embryonic development in the laboratory is still unknown. On the other hand, during *in vitro* embryo culture, blockage of four to eight cells is a common phenomenon [14]. Many approaches have been adopted to overcome this obstacle by supplementing with hypotaurine, FBS, FCS, and bovine serum albumin (BSA) during IVC of the porcine embryo. Integrating serum into the embryo culture medium can improve embryo quality by preventing apoptosis and/or decreasing reactive oxygen species (ROS) accumulation. Therefore, this study was designed to investigate the role and the molecular mechanisms of porcine embryo development during the integration of FBS into culture media.

Materials and Methods

Chemicals

All the chemicals used in this experiment were bought from Sigma-Aldrich, St. Louis, MO, unless otherwise stated.

Ovary collection, recovery, and *in vitro* oocyte maturation

Pre-pubertal gilt ovaries were collected from an abattoir house in Chungbuk city, and the collected ovaries were transported to the laboratory within 2 h. The transport medium consisted of NaCl (0.9%), streptomycin sulfate (100 mg/ml), and penicillin-G (100 IU/ml). The contents of the follicular (3–7 mm in diameter) were aspirated with a 5 ml disposable syringe with a 20 gauge needle. A minimum of three layers of compact cumulus cells with homogenous ooplasmic cumulus–oocyte complexes (COC) was selected using a stereomicroscope. A selected group of 50–60 immature COCs were placed in 4-well culture dishes

(Nunc, Roskilde, Denmark) with the oocyte maturation medium. The media consisted of tissue culture medium 199 (Invitrogen, Grand Island, NY), cysteine (0.6 mM), pyruvate (0.91 mM), epidermal growth factor (15 ng/ml), kanamycin (75 µg/ml), insulin (1 µg/ml), and porcine follicular fluid (pFF) (10%, v/v). A four-well dish containing immature COCs was then placed at 39°C in an exceedingly humidified atmosphere containing 5% CO₂. These culture conditions were carried out for 20–22 h with 10 IU/ml equine chorionic gonadotropin (eCG; Intervet International BV, Boxmeer, Netherlands) and 10 IU/ml human chorionic gonadotropin (hCG; Intervet International BV, Boxmeer, Netherlands). After 20–22 h of culture, the oocytes were rinsed with fresh medium and cultured again for an additional 18 h without hormonal supplementation. The pFF was prepared according to Biswas et al. [15] and preserved at –20°C for further use. For maturation, each well of the 4-well Nunc plate contained 500 µl of maturation medium.

IVF of porcine oocytes and culture

IVF and IVC were carried out according to our previous study [16]. Briefly, on day 2, after *in vitro* maturation (IVM), de-cumulated M-II oocytes were co-incubated for 20 min with spermatozoa at a concentration of 5×10^5 /ml in 50 µl microdrop of modified Tris buffer medium (mTBM) at 39°C in a humidified atmosphere containing 5% CO₂ and 95% air. After 20 min of co-incubation, gentle pipetting was applied to remove loosely attached sperms from the zona pellucida. The zygotes were then washed thrice with mTBM and incubated with mTBM without sperm for 5–6 h in the same environment. After that, the zygotes were washed thrice with PZM-3 medium and cultured in 30 µl microdrops (10 zygotes per drop) of PZM-3 and covered with prewarmed mineral oil. The putative zygotes were incubated at 39°C for 168 h in a humidified atmosphere containing 5% O₂, 5% CO₂, and 90% N₂. The zygotes were randomly allocated into two groups after successful IVF. On day 4 post-IVF, 10% FBS was injected into the culture medium, and without FBS was treated as the control group. Each experiment was repeated at least thrice.

Embryo evaluation and total blastocyst (BL) count

On day 2 and day 7, zygotes were checked for cleavage and BL formation, respectively. To calculate the total number of blastomere in the BL at day 7, the BL was washed with 1% (wt/vol) phosphate-buffered saline BSA (PBS-BSA) for 5 min and stained with 5 mg/ml of Hoechst 33342 (bis-benzimide). Subsequently, the embryos were briefly fixed in 4% paraformaldehyde. Next, the BL was mounted on a glass slide in a drop of 100% glycerol and gently covered with a cover slip. The stained BLs were observed at 400x using a fluorescence microscope (Nikon Corporation, Japan) and counted the total number of blastomere.

Quantitative real-time reverse transcription polymerase chain reaction (RT-PCR) and the gene expression pattern

In each group, approximately 8–10 IVF BLs were stored at –80°C after washing with PBS until further use for the gene expression analysis. Quantitative real-time RT-PCR analyzed the expression of B cell lymphoma-2 (BCL-2) and caspase-3 mRNA in BL. TRIzol reagent (Invitrogen, Carlsbad, CA) was used to extract the total RNA from pooled BLs according to the manufacturer’s instructions. A spectrophotometer (Micro UV-Vis spectrophotometer, Hangzhou, 310023, China) was used to measure the total RNA concentration at a wavelength of 260 nm. A mixture of moloney murine leukemia virus reverse transcriptase (Invitrogen, Carlsbad), random primer (9-mers; TaKaRa Bio, Otsu, Shiga, Japan), and 1 µg of total RNA was used to synthesize the first-strand complementary DNA (cDNA). Real-time RT-PCR was carried out in a 20 µl PCR reaction volume consisting of 10 µl of 2 × SYBR premix Ex Taq (TaKaRa Bio Inc.), 1 µl cDNA, 1U of Taq polymerase (Intron Bio Technologies, Seongnam, Korea), and 10 pM of each specific primer in an MX3000P thermocycler engine (Stratagene-Agilent Technologies, Waldbronn, Germany).

The reaction was set for 40 cycles with the following parameters: denaturation at 95°C for 30 sec, followed by 30 sec of annealing at 57°C and 30 sec extension at 72°C. All oligonucleotide primer sequences are presented in Table 1. At the end of the extension, the fluorescence intensity of each gene was recorded from the PCR product. The fluorescence intensity threshold value of all samples was set manually. The number of reaction cycles where the fluorescence intensity of each PCR product crossed the threshold value was considered the cycle threshold (Ct) in

the exponential phase of PCR amplification. The relative comparative quantification of the expression of each target gene was based on the Ct values at constant fluorescence intensity. It was normalized to the RN18S as an internal control gene. The relative mRNA expression of each target gene (*R*) was calculated using the following equation, $R = 2^{-[\Delta Ct \text{ sample} - \Delta Ct \text{ control}]}$. The experiment was repeated at least thrice.

ROS accumulation assay in BLs

On day 7, BLs from both groups were used for intracellular ROS analysis. Briefly, the BLs were incubated with 10 mM of 2',7'-dichloro-dihydro-fluorescein diacetate (H₂DCFDA; Invitrogen) for 30 min in a dark place, and after incubation, the treated BLs were washed thrice in HEPES-buffered Tyrode’s medium-polyvinyl alcohol (TLH-PVA) solution. The stained BLs were mounted on a glass slide and visualized under an epifluorescence microscope (TE2000-U, Nikon, Japan). The final photos were captured with a digital camera and the fluorescence intensity of both BLs was measured with the ImageJ software (version 1.47; National Institutes of Health, USA). The fluorescence intensity was normalized with the control. This experiment was repeated at least four times.

Statistical analysis

The data percentage of the cleaved embryo and various types of BL formation was analyzed by paired student’s *t*-test using GraphPad Prism software. All data are presented as mean ± SEM. Differences at *p* < 0.05 were considered statistically significant

Table 1. Sequences of primer sets used for selective gene expression on day 7 porcine BL.

Symbol	Primer sequences (5'-3')	Product size	Accession number
BCL-2	F: AGGGCATTCACTGACCTGAC	193	NM_214285
	R: CGATCCGACTCACCAATACC		
Caspase-3	F: CGTGCTTCTAAGCCATGGTG	186	NM_214131
	R: GTCCCACTGTCCTCTCAAT		
RN18S	F: CGCGTTCTATTTTGTGGT	219	NR_046261.1
	R: AGTCGGCATCGTTTATGGTC		

Table 2. Effects of FBS supplementation on *in vitro* development of porcine IVF zygote.

Culture system	Total zygotes culture	Cleavage (%)	BL (%)	Average blastomere/BL
Control	573	310 (55.58 ± 8.05)	138 (23.85 ± 3.04) ^a	46.20 ± 0.84 ^a
+FBS	901	606 (68.19 ± 4.50)	353 (38.23 ± 3.98) ^b	102.2 ± 1.50 ^b

^{a,b} Values with different superscripts in the same column are significantly different (*p* < 0.05). Data are given as mean ± SEM from each replicate.




Figure 1. Effect of FBS on *in vitro* production of porcine embryo quality on day 7. The data are presented as mean \pm SEM. Bars with different letters in each endpoint differ from each other, which are statistically significant ($p < 0.05$). ErBL: early blastocyst; ExBL: expanded blastocyst; HgBL: hatching blastocyst; HdBL: hatched blastocyst.

Results

Effects of FBS on BL quality derived from IVF embryos in PZM-3 culture medium

As shown in Table 2, the embryo developmental potential at the BL stage and the total blastomere per BL were significantly higher ($p < 0.05$) in the FBS supplemented group than in the control group. On day 2, the total cleavage rate was not significantly different between the two groups. Furthermore, the group that received FBS during the culture period also showed significant ($p < 0.05$) hatching rates (5.82 ± 1.08) and hatched BL formation (12.22 ± 2.47) compared to the control group (1.51 ± 0.81 and 0.4 ± 0.26 , respectively). However, there was no significant difference in early BL and expanded BL formation between the groups (Fig. 1). Morphologically, embryos appeared to look good in the FBS-treated group compared to the control group (Fig. 2).

Effect of FBS on ROS production in day-7 porcine BLs

The H_2DCFDA molecular staining technique was applied to the determination of ROS levels on day-7 BLs. The present study showed that the ROS production was significantly reduced ($p < 0.05$) in BL cultured with FBS compared to the untreated control group (lower H_2DCFDA signal) (Fig. 3).

Effect of FBS supplementation on the mRNA expression of apoptosis-related genes in day-7 porcine BLs

The relative abundance of BCL-2 mRNA in day-7 BL was significantly higher ($p < 0.05$) in FBS-treated BLs compared to control, and caspase-3 mRNA level in day-7 BL

was also significantly lower ($p < 0.05$) in the FBS treatment group compared to the control group (Fig. 4).

Discussion

FBS is the most widely used cell culture and can also be used as an energy source for growing cells [17]. It supplies cocktail factors for cell growth, proliferation, and attachment [18]. Several studies have shown that supplementation of serum-like FBS in IVM or IVC medium increased the development of the embryo other than BSA or other macromolecules. In the present study, 10% FBS was supplemented on day 4 in the PZM-3 during the culture of *in vitro* fertilized porcine embryos. The percentages of BLs development and the total cell number per BL were significantly increased in the medium supplemented with FBS. Similar results were found in IVF porcine embryo development where 10% FBS was added in the chemically defined medium [19], and also 2.5% FBS was used for IVF bovine embryo production [20]. However, it was found that 5% FBS has been advantageous during the pre-cleavage stage, and 10%–20% FBS has been advantageous during the post-cleavage stage of the bovine embryos and also overcomes the deleterious effects of serum on early developing embryos by exploiting embryotropic effects during later stage [21].

Another study on bovine embryo culture indicated that adding FBS in the culture medium promoted development to the morula and BL stage [20]. Mostly, FBS acts as a protein source in the culture medium. Numerous embryonic growth-promoting factors and other unknown cell survival components are essential during embryonic growth [22]. Adding FBS on day 4 may improve the metabolic pathways when the IVF embryos are mostly at the morula stage [23],




Figure 2. On day 7, the different types of BL were found with or without FBS supplementation during *in vitro* porcine embryo production. Bar = 100 µm.




Figure 3. The fluorescence intensity of ROS levels was detected by H₂DCFDA staining of BL derived from IVF embryos cultured with FBS in the PZM-3 culture medium. The fluorescence intensity of BLs was analyzed by ImageJ software. The arbitrary data according to the control group was set as 1. The number of BLs examined in each group is shown in the bars. Values represent mean \pm SEM of the separate experiments. Bars with different letters in their respective endpoints differ from each other and are treated as statistically significant ($p < 0.05$).

and at that time, embryos need more nutrients and energy. It has also been reported that serum strongly inhibits the division of the first stage of the fertilized embryos and has no beneficial effect on the development from two cells to the morula stage [24]. This study found that the total cell number per BL on day 7 was increased twice when FBS was introduced into the embryo culture medium on day 4. A growing embryo needs more energy for replicating quickly and smoothly. The presence of FBS in the culture medium influences the embryonic cell division in the BL due to increasing intracellular lipid metabolism [23,25].

The increased metabolism leads to the formation of new cells in growing embryos. It could be suggested that the presence of one of the main factors in FBS is phosphate, and the concentration is higher as inorganic phosphorus can stimulate the development of eight-cell embryos in a rat model [26]. Moreover, it was also found that adding FBS on day 4 also improved the hatching and hatched BL formation compared to the control. It indicated that FBS had biphasic effects on developing embryos [24], but the stimulation from the serum at late-stage embryo development is unknown. However, FBS could trigger the cell cycle




Figure 4. Relative levels of BCL 2 and caspase 3 mRNA expression IVF-derived porcine BLs cultured with FBS and without FBS (control group) in the PZM-3 culture medium. The experiment was repeated at least thrice. The data are expressed as mean \pm SEM of this experiment ($p < 0.05$).

process and result in more BL and blastomere found in this experiment. Indeed, increased total blastomere dictating that developmental competence was excellent in accordance with trophectoderm cell number [27], and the total cell number is the suitable paramere for assessing *in vitro* produced embryos.

Our study found that with H_2DCFDA staining of day-7 BLs, ROS generation was significantly decreased in FBS-treated embryos. Several other studies have found that adding FBS to the IVC media often triggers ROS accumulation in mammalian cells [28]. In that case, bovine embryos were used and nicotinamide adenine dinucleotide phosphate (NADPH) oxidases enzymes were activated in the presence of FBS, which led to accelerating the cell cycle progression [29]. But this study showed the opposite result where ROS generation was decreased in the presence of FBS in culture medium. The variation of this result may be due to the different types of embryos used in this study. The porcine embryo contains high lipid contents in the ooplasm and has different metabolic activities during embryogenesis. Another study indicated that numerous antioxidant transcripts were upregulated after adding FBS in the culture medium. Excessive accumulation of ROS can damage the cellular component of growing embryos, and FBS inhibits the ROS production by activation of p-p38 mitogen-activated protein kinase (MAPK) and phospho-a serine/threonine protein kinase (p-AKT) signal pathways [19].

Programmed cell death or apoptosis is jointly involved in early embryonic development and differentiation. An unfavorable culture media sometimes induces unscheduled embryonic cells' apoptosis, leading to cessation or abnormal early developmental processes [30]. However, the balanced expression of pro- and anti-apoptotic genes controls the apoptotic pathway. The BCL-2 gene family is

known for its pro- and anti-apoptotic subgroups, and the BCL-2 gene is known to protect against apoptosis. In this observation, it was found that the addition of FBS in the porcine embryo culture media significantly affected the anti-apoptotic gene BCL-2 in FBS-treated embryos and significantly decreased the expression of the pro-apoptotic gene, caspase-3 compared to the control group. Another subgroup is highly conserved to the regulator of apoptosis gene expression like caspase-3. From these observations, it can be explained that FBS can reduce the apoptotic process in porcine BLs by preventing apoptosis either by preventing the activation of caspase-3 or by preventing the release of factors that induce apoptosis [31].

BCL-2 is a potent suppressor of cell death and its upstream expression can reduce the gene that promotes cell death, such as caspase-3 in maternal stores in developing embryos. This would be consistent with the idea that superior quality embryos contain higher mRNA levels from the anti-apoptotic gene and conversely lower levels of the pro-apoptotic transcriptome. However, FBS had some detrimental effects when used in porcine parthenote, and in that case, the total numbers of cells per BL were significantly reduced and apoptotic cells were significantly increased [20]. In fact, increased expression of the anti-apoptotic gene BCL-2 in embryos determined the presence of embryos with good morphological quality in IVC media [32].

Conclusion

In conclusion, these data indicated that the supplementation of the porcine culture media with FBS positively affects *in vitro* embryo development. The addition of FBS to the culture medium on day 4 significantly increased the hatched BL production *in vitro* and decreased ROS

accumulation. Expression of the anti-apoptotic BCL-2 gene was significantly increased in developing BLs.

List of Abbreviations

FBS, fetal bovine serum; PZM-3, porcine zygotic medium-3; ROS, reactive oxygen species; BL, blastocyst; NCSU, North Carolina State University; IVF, *in vitro* fertilization; FCS, fetal calf serum; COC, cumulus-oocyte complexes; pFF, porcine follicular fluid; IVC, *in vitro* culture; IVM, *in vitro* maturation; mTBM, modified Tris buffer medium; h, hours; min, minutes; sec, seconds; cDNA, complementary DNA; BCL-2, B cell lymphoma-2; eCG, equine chorionic gonadotropin; hCG, human chorionic gonadotropin; RT-PCR, reverse transcription polymerase chain reaction; TLH-PVA, HEPES-buffered Tyrode's medium-polyvinyl alcohol; NADPH, nicotinamide adenine dinucleotide phosphate; MAPK, mitogen-activated protein kinase; p-AKT, phospho-a serine/threonine protein kinase.

Acknowledgments

This research was funded, in part, by grants from the "National Research Foundation of Korea," funded by the Korean Government (2017K1A4A3014959, 2020R1A2C2008276), and the "Korea Institute of Planning and Evaluation for Technology in Food, Agriculture, Forestry, and Fisheries (IPET) (grant numbers:318016-5, 819029-2, and 320005-4)," Korea.

Conflict of interest

The authors declare no financial conflict of interests.

Authors' contributions

DB designed the study, interpreted the results, and arranged the publication draft and paper preparation. SH took part in funding acquisition and critically edited this manuscript. All authors reviewed the manuscript individually and approved the final manuscript.

References

- [1] Watanabe M, Kurome M, Matsunari H, Nakano K, Umeyama K, Shiota A, et al. The creation of transgenic pigs expressing human proteins using BAC-derived, full-length genes and intracytoplasmic sperm injection-mediated gene transfer. *Transgenic Res* 2012; 21(3):605-18; <https://doi.org/10.1007/s11248-011-9561-3>
- [2] Li J, Chen Q, Wang J, Huang G, Ye H. Does growth hormone supplementation improve oocyte competence and IVF outcomes in patients with poor embryonic development? A randomized controlled trial. *BMC Pregnancy Childbirth* 2020; 20(1):310; <https://doi.org/10.1186/s12884-020-03004-9>
- [3] Ou XH, Li S, Wang ZB, Li M, Quan S, Xing F, et al. Maternal insulin resistance causes oxidative stress and mitochondrial dysfunction in mouse oocytes. *Hum Reprod* 2012; 27(7):2130-45; <https://doi.org/10.1093/humrep/des137>
- [4] Lobo V, Patil A, Phatak A, Chandra N. Free radicals, antioxidants and functional foods: impact on human health. *Pharmacogn Rev* 2010; 4(8):118-26; <https://doi.org/10.4103/0973-7847.70902>
- [5] Petters RM, Wells KD. Culture of pig embryos. *J Reprod Fertil Supp* 1993; 48:61-73.
- [6] Youngs CR, Ford SP, McGinnis LK, Anderson LH. Investigations into the control of litter size in swine: I. Comparative studies on *in vitro* development of Meishan and Yorkshire preimplantation embryos. *J Anim Sci* 1993; 71:1561-5; <https://doi.org/10.2527/1993.7161561x>
- [7] Yoshioka K, Suzuki C, Tanaka A, Anas IM, Iwamura S. Birth of piglets derived from porcine zygotes cultured in a chemically defined medium. *Biol Reprod* 2002; 66:112-9; <https://doi.org/10.1095/biolreprod66.1.112>
- [8] Cao Z, Sui L, Li Y, Ji S, Zhang X, Zhang Y. Effects of chemically defined medium on early development of porcine embryos derived from parthenogenetic activation and cloning. *Zygote* 2012; 20(3):229-36; <https://doi.org/10.1017/S0967199411000153>
- [9] Okada K, Krylov V, Kren R, Fulka J Jr. Development of pig embryos after electro-activation and *in vitro* fertilization in PZM-3 or PZM supplemented with fetal bovine serum. *J Reprod Dev* 2006; 52(1):91-8; <https://doi.org/10.1262/jrd.17059>
- [10] Abdel-Wahab A, Abdel Aziz RL, Helmy NA, Ibrahim SS. The impact of using newborn bovine serum as fetal calf serum substitute in the *in vitro* bovine embryos production system. *Porto Biomed J* 2018; 3(2):e3; <https://doi.org/10.1016/j.pbj.0000000000000003>
- [11] Booth PJ, Holm P, Callesen H. The effect of oxygen tension on porcine embryonic development is dependent on embryo type. *Theriogenology* 2005; 63(7):2040-52; <https://doi.org/10.1016/j.theriogenology.2004.10.001>
- [12] van der Valk J, Brunner D, De Smet K, Fex Svenningsen A, Honegger P, Knudsen LE, et al. Optimization of chemically defined cell culture media replacing fetal bovine serum in mammalian *in vitro* methods. *Toxicol In Vitro* 2010; 24(4):1053-63; <https://doi.org/10.1016/j.tiv.2010.03.016>
- [13] Miyoshi K. Development of a culture medium for rat 1-cell embryos. *J Mamm Ova Res* 2016; 33(1):11-6. <https://doi.org/10.1274/jmor.33.11>
- [14] Oliveira CS, de Barros BAF, Monteiro CAS, Rosa PMS, Leal GR, Serapião RV, et al. Individual assessment of bovine embryo development using a homemade chamber reveals kinetic patterns of success and failure to reach blastocyst stage. *Syst Biol Reprod Med* 2019; 65(4):301-11; <https://doi.org/10.1080/19396368.2019.1589601>
- [15] Biswas D, Jung EM, Jeung EB, Hyun SH. Effects of vascular endothelial growth factor on porcine preimplantation embryos produced by *in vitro* fertilization and somatic cell nuclear transfer. *Theriogenology* 2011; 75(2):256-67; <https://doi.org/10.1016/j.theriogenology.2010.08.012>
- [16] Biswas D, So KH, Hwang SU, Yoon JD, Kim M, Kim DY, et al. Embryotropic effects of vascular endothelial growth factor on porcine embryos produced by *in vitro* fertilization. *Theriogenology* 2018; 120:147-56; <https://doi.org/10.1016/j.theriogenology.2018.07.024>
- [17] Arora M. Cell Culture Media: a review. *Mater Methods* 2013; 3:175; <https://doi.org/10.13070/mm.en.3.175>
- [18] Gstraunthaler G, Lindl T, van der Valk J. A plea to reduce or replace fetal bovine serum in cell culture media. *Cytotechnology* 2013; 65(5):791-3; <https://doi.org/10.1007/s10616-013-9633-8>
- [19] Mun SE, Sim BW, Yoon SB, Jeong PS, Yang HJ, Choi SA, et al. Dual effect of fetal bovine serum on early development depends on stage-specific reactive oxygen species demands in pigs. *PLoS One* 2017; 12(4):e0175427; <https://doi.org/10.1371/journal.pone.0175427>
- [20] Arnold DR, Corrêa CAP, Lorena LLG, Gaspar RC, Rossi GF, Ifran AM, et al. Supplementation of fetal bovine serum alters histone modification H3R26me2 during preimplantation development of *in vitro*

- produced bovine embryos. *Pesq Vet Bras* 2015; 35(7):605–12; <https://doi.org/10.1590/S0100-736X2015000700002>
- [21] Ferré LB, Kjelland ME, Strøbech LB, Hyttel P, Mermillod P, Ross PJ. Review: recent advances in bovine *in vitro* embryo production: reproductive biotechnology history and methods. *Animal* 2020; 14(5):991–1004; <https://doi.org/10.1017/S1751731119002775>
- [22] Chelladurai KS, Christyraj JDS, Rajagopalan K, Yesudhasan BV, Venkatachalam S, Mohan M, et al. Alternative to FBS in animal cell culture - an overview and future perspective. *Heliyon* 2021; 7(8):e07686; <https://doi.org/10.1016/j.heliyon.2021.e07686>
- [23] Guo S, Liu S, Bou G, Guo J, Jiang L, Chai Z, et al. Fetal bovine serum promotes the development of *in vitro* porcine blastocysts by activating the Rho-associated kinase signalling pathway. *Reprod Fertil Dev* 2019; 31(2):366–76; <https://doi.org/10.1071/RD18070>
- [24] Sena-Netto SB, Sprícigo JFW, Leme LO, Guimarães ALS, Caixeta FMC, Dode MAN, et al. The replacement of fetal bovine serum with bovine serum albumin during oocyte maturation and embryo culture does not improve blastocyst quality after slow freezing cryopreservation. *Biopreserv Biobank* 2020; 18(3):171–9; <https://doi.org/10.1089/bio.2019.0059>
- [25] Kim SH, Yoon JT. Effect of serum-containing and serum-free culture medium-mediated activation of matrix metalloproteinases on embryonic developmental competence. *Czech J Anim Sci* 2019; 64(12):473–82; <https://doi.org/10.17221/205/2019-CJAS>
- [26] Ma Y, Gu M, Chen L, Shen H, Pan Y, Pang Y, et al. Recent advances in critical nodes of embryo engineering technology. *Theranostics* 2021; 11(15):7391–424; <https://doi.org/10.7150/thno.58799>
- [27] Redel BK, Spate LD, Lee K, Mao J, Whitworth KM, Prather RS. Glycine supplementation *in vitro* enhances porcine preimplantation embryo cell number and decreases apoptosis but does not lead to live births. *Mol Reprod Dev*. 2016; 83(3):246–528; <https://doi.org/10.1002/mrd.22618>
- [28] Park SH, Cho HS, Yu IJ. Effect of bovine follicular fluid on reactive oxygen species and glutathione in oocytes, apoptosis and apoptosis-related gene expression of *in vitro*-produced blastocysts. *Reprod Domest Anim* 2014; 49(3):370–7; <https://doi.org/10.1111/rda.12281>
- [29] Vaquero EC, Edderkaoui M, Pandol SJ, Gukovsky I, Gukovskaya AS. Reactive oxygen species produced by NAD(P)H oxidase inhibit apoptosis in pancreatic cancer cells. *J Biol Chem* 2004; 279(33):34643–54; <https://doi.org/10.1074/jbc.M400078200>
- [30] Haouzi D, Hamamah S. Pertinence of apoptosis markers for the improvement of *in vitro* fertilization (IVF). *Curr Med Chem* 2009; 16(15):1905–16; <https://doi.org/10.2174/092986709788186075>
- [31] Guan M, Zhang Y, Huang Q, He L, Fang Q, Zhang J, et al. Fetal bovine serum inhibits neomycin-induced apoptosis of hair cell-like HEI-OC-1 cells by maintaining mitochondrial function. *Am J Transl Res* 2019; 11(3):1343–58.
- [32] Ramos-Ibeas P, Gimeno I, Cañón-Beltrán K, Cañón-Beltrán A, Rizo D, Gómez E. Senescence and apoptosis during *in vitro* embryo development in a bovine model. *Front Cell Dev* 2020; 8:619902; <https://doi.org/10.3389/fcell.2020.619902>