

# Associated risk factor analysis and the prognostic impact of positive resection margins after endoscopic resection in early esophageal squamous cell carcinoma

YONG FENG<sup>1</sup>, WEI WEI<sup>2</sup>, SHUO GUO<sup>3</sup> and BAO-QING LI<sup>1</sup>

Departments of <sup>1</sup>Thoracic Surgery, <sup>2</sup>Outpatients and <sup>3</sup>Department of Gastroenterology, Fourth Hospital of Hebei Medical University, Shijiazhuang, Hebei 050011, P.R. China

Received February 24, 2022; Accepted May 3, 2022

DOI: 10.3892/etm.2022.11384

**Abstract.** Endoscopic resection for early esophageal cancer has a risk of residual margins. The risk these residual margins pose have not been fully evaluated. The present study aimed to investigate the associated risk factors and prognosis of residual margins following the endoscopic resection of early esophageal squamous cell carcinoma. In total, 369 patients (381 lesions) with early esophageal squamous cell carcinoma treated in the Fourth Hospital of Hebei Medical University (Shijiazhuang, China) with endoscopic resection were retrospectively analyzed. Sex, age, location, tumor diameter, depth of tumor invasion, endoscopic treatment, endoscopic ultrasonography (EUS) before resection, work experience of endoscopists and the degree of tumor differentiation were all evaluated as potential risk factors. In addition, the prognosis of patients with positive margins were analyzed. A total of 73 patients (73/381, 19.2%) had positive margins after endoscopic resection. Amongst the 65 patients who were successfully followed up, five patients succumbed to cardiovascular and cerebrovascular diseases, one patient received radiotherapy, two patients received radiotherapy and chemotherapy whilst one patient received chemotherapy. By contrast, 12 patients received surgery and 20 patients received additional endoscopic mucosal resection or endoscopic submucosal dissection. The other 29 patients were followed up regularly and no recurrence could be found. Univariate analysis revealed that tumor diameter, endoscopic treatment, depth of invasion, EUS before resection, degree of tumor differentiation and direction of invasion were all associated with the positive margin. Multivariate logistic regression analysis then found that EUS before resection, degree of tumor differentiation and depth of tumor invasion are independent

risk factors for positive margins after endoscopic resection. These results suggest that poorly differentiated lesions and deeper invasion depth can increase the risk of positive margin after endoscopic resection. As a result, EUS evaluation before resection may reduce the risk of invasion depth. In addition, for poorly differentiated lesions, more aggressive treatment regimens may be recommended for preventing recurrence.

## Introduction

Esophageal cancer is a common type digestive system malignancy and currently ranks sixth in terms of the rate of cancer-associated mortality worldwide (1). Despite progress in the development of novel treatment strategies, the 5-year survival rate of esophageal cancer remains <20% (2). Esophageal cancer in the early stages is confined to the mucosa or superficial submucosa and accounts ~20% of all types of esophageal cancers (3). Surgical resection has long been the standard treatment method for early esophageal cancer (4). However, the high incidence of complications (such as gastroesophageal reflux and respiratory failure) as a result of this technique renders it unattractive, due to its severe negative effects on the quality of life of patients (4). Nevertheless, the development of gastroscopy has greatly improved the quality of life of patients with early esophageal cancer.

As advancements in endoscopic technology are made continuously, endoscopic resection is becoming the standard, minimally invasive treatment procedure for early esophageal cancer (5-7). It has been found to shorten the length of hospital stay and reduce the incidence of complications without affecting the quality of life, compared with those after esophagectomy (5-7).

However, similar to traditional esophagectomy, endoscopic resection also has the risk of a positive margin (positive margin refers to the presence of atypical cells at the lateral or deep resection margins). Positive margin after the endoscopic resection of early esophageal carcinoma has direct implications on the choice of treatment and disease prognosis (8). To date, previous studies of patients with early esophageal cancer (9-11) after undergoing endoscopic submucosal dissection (ESD) have produced sporadic results based on a small number of cases. Amongst the available reports on positive

---

*Correspondence to:* Dr Yong Feng, Department of Thoracic Surgery, Fourth Hospital of Hebei Medical University, 12 Jiankang Road, Shijiazhuang, Hebei 050011, P.R. China  
E-mail: 952453363@qq.com

**Key words:** early esophageal carcinoma, endoscopic treatment, risk factors, positive resection margins

margins, there are no reports of multicenter, large-sample trials with long-term follow-up periods.

Therefore, to explore methods of improving the prognosis of patients after the endoscopic resection of esophageal cancer, the present study investigated the potential risk factors for positive margins by comprehensively analyzing clinical and pathological data.

## Materials and methods

**Patients.** The present study retrospectively analyzed the clinical, endoscopic and pathological data of patients with esophageal mucosal lesions treated with endoscopic resection in the Fourth Hospital of Hebei Medical University (Shijiazhuang, China) from January 2011 to December 2020.

The inclusion criteria of the lesions were as follows: i) Complete resection of lesions; ii) pathological diagnosis after endoscopic treatment was atypical cells (cells that looked different and function differently than normal) [low-grade intraepithelial neoplasia (LIN), high-grade intraepithelial neoplasia (HIN) or invasive cancer]; and iii) patients provided informed consent. The exclusion criterion was patients who were pathologically diagnosed with adenocarcinoma or inflammation after endoscopic treatment. Of note, intraepithelial neoplasia was defined as the neoplastic change before epithelial invasion, which can be divided further into ‘low-grade’ and ‘high-grade’ according to whether the structural and cytological abnormalities observed occupy the upper part of the epithelium (12).

A total of 369 patients (381 lesions) received endoscopic resection due to early esophageal carcinoma. Of these, 236 were males and 133 were females with a mean age of  $63.5 \pm 9.9$  years. The main clinical symptoms were retropharyngeal asphyxia and retrosternal pain. No other symptoms or findings could be observed during physical examination. All patients and their relatives were aware of the risks and benefits of endoscopic resection, following which written informed consent was obtained from all participants. The present study was approved by the Ethics committee of the Fourth Hospital of Hebei Medical University (Shijiazhuang, China).

**Endoscopic treatment and pathological examination.** Under the guidance of endoscopy, two treatment methods were used to remove the tumor: Endoscopic mucosal resection (EMR) and ESD.

**Endoscopy-guided surgery.** The type of endoscopic treatment performed in the present study was by professionally trained endoscopists using a single-channel endoscope (Olympus H260; Olympus Corporation). Argon plasma coagulation (APC; APC probes for flexible endoscope; Erbe Elektromedizin GmbH) was used to mark ~5 mm outside the boundary of the lesion. After fully marking the border, an epinephrine-containing hypertonic saline solution was injected into the submucosa for submucosal lifting. A circumferential mucosal incision was then created around the marking spots. In cases where the EMR technique was used, asymmetrical polypectomy snare (MTW Endoskopie) resection was initiated. For ESD, the submucosal layer was dissected using the IT-knife2 (KD-611L; Olympus Corporation). After the

lesions were resected, the wounds were closed with APC, hot biopsy forceps coagulation or using titanium clips according to the wound conditions (Fig. 1A and B). The lesion was completely removed *en bloc* (Fig. 1C). The pathologist then evaluated tumor involvement in the lateral or deep resection margin (Fig. 1D).

**Post-operative pathological examination.** The resected specimens were processed and fixed immediately, before the lesion size was measured. After fixation with 10% formalin (20–25°C for 12–24 h), the specimens were sent for pathological examination. The samples were cut into continuous sections at 2-mm intervals from top to bottom and embedded in paraffin. In total, three sections were prepared from each tissue and then stained with hematoxylin and eosin using the Ventana HE 600 automatic staining machine (Roche Diagnostics; 20–25°C). An experienced pathologist (Dr Yao Liu, Department of Pathology, Fourth Hospital of Hebei Medical University) then used a light microscope (DM1000; Leica Microsystems GmbH; magnification, x200) to evaluate tumor involvement in the lateral or deep resection margin to avoid misjudgment. When examining the tissue sections for pathological evaluation, all sections were observed from the top of the tissue to identify the nature of the lesion, the degree of tumor differentiation, whether the vertical/transverse edge of the tumor was positive and whether there was vascular infiltration. A positive resection margin as defined as the presence of atypical cells (LIN, HIN or invasive cancer) at the lateral or deep resection margin.

**Statistical analysis.** Sex, age, lesion location, tumor diameter, depth of invasion, endoscopic treatment, endoscopic ultrasonography (EUS) before resection, working experience of the endoscopist and degree of tumor differentiation were all analyzed as potential risk factors.

Continuous data are presented as the mean  $\pm$  standard deviation.  $\chi^2$  and nonparametric Mann-Whitney U tests were used for statistical analysis. Univariate and multivariate logistic regression models were used to examine the association between the variables and positive margins risk factors.  $P < 0.05$  was considered to indicate a statistically significant significance. The SPSS v19.0 (IBM Corp.) was used for data processing.

## Results

**General data.** The present study was a retrospective analysis of the data collected from 369 patients with early esophageal cancer who met the inclusion criteria. Among all study participants, 236 were male and 133 were female with a ratio of 1.8:1. The average age was  $63.5 \pm 9.9$  (range, 31–85) years.

**Residues at the resection margin and follow-up data.** In total, 73 patients had positive margin, such that the positive rate was 19.2%, where 64 (16.8%) were positive for lateral margins and nine (2.4%) were positive for vertical margins. In addition, it was found that the residual rates of EMR and ESD were 26.9 and 16.7%, respectively. Only 7.5% (23/308) poorly differentiated lesions were diagnosed in negative margins. By contrast, the proportion of positive margin specimens was




Figure 1. Endoscopic resection procedure for early esophageal squamous cell cancer. (A) Upper gastrointestinal endoscopy revealed a lesion located at the middle esophagus (the lesion is indicated by arrows). (B) Upper gastrointestinal endoscopy image of the esophagus after endoscopic resection. (C) En bloc image of the resected specimen, 6x4 cm (the lesion was indicated by arrows). (D) High-grade intraepithelial neoplasia in the deep resection margin as shown by H&E staining. The lesion was indicated by arrows. Magnification, x200.

28.8% (21/73). There was no vascular invasion or submucosal lymphatic metastasis. After communicating with the patients and their families, 29 patients asked for regular follow-up, while others with positive margins received another ESD or EMR and were followed up 1 month after operation. During the follow-up period, endoscopy was performed 3, 6 and 12 months after endoscopic resection, before being performed every year thereafter. Chest CT was also performed 6 and 12 months after endoscopic resection followed by once a year thereafter as a precaution for distant metastasis. Follow up was halted 5 years after endoscopic resection. If tumor cells were found at the edge of the deep resection, additional esophagectomy or esophageal radiotherapy and chemotherapy would be performed. These treatment options were selected according to the situation of each patient and the wishes of patients and their families. In the present study, 65 cases of residual tumors were successfully followed up for 2-60 months, with an average follow-up time of  $28.1 \pm 10.1$  months. The follow-up rate was 89%. However, eight patients were lost to follow-up. During the follow-up, five patients succumbed to cardiovascular and cerebrovascular diseases, whilst the other 60 patients remained alive during the follow-up period.

*Comparison of groups with or without residual margins.* A number of associated factors can affect the residual margin, including sex, age, lesion location, tumor diameter, depth of invasion, endoscopic treatment, EUS before resection, working experience of the endoscopist and degree of tumor differentiation (12). Univariate and multivariate analyses were performed to determine if any of these factors can affect the residual margin after endoscopic resection. As shown in Table I, there was no significant difference in sex, age, lesion location or working experience of endoscopists in the univariate analysis of incision margins. By contrast, tumor diameter, endoscopic treatment, depth of invasion, EUS before resection and the degree of tumor differentiation were all risk factors for residual margins. To exclude any confounding factors, multivariate logistic regression analysis (Table II) was performed. Only the depth of tumor invasion, degree of differentiation and EUS before resection were regarded to be risk factors for residual margins.

## Discussion

Applications of EMR and ESD are becoming increasingly common for the treatment of early esophageal cancer (7). Endoscopic resection is able to not only preserve the integrity of the esophagus, but also circumvent the considerable risk of morbidity and mortality associated with esophageal resection (13). Previous studies have reported that although the effect of endoscopic resection on early esophageal cancer is equivalent to that of surgical resection, a lower incidence of complications and superior quality of life was associated with endoscopic resection (5,7). Several cohort studies (14-17) have recommended the use of EMR or ESD for T1a esophageal tumors, including highly atypical hyperplasia, adenocarcinoma or squamous cell carcinoma, which are limited to the superficial mucosa and do not extend to the muscular mucosa. In addition, Manner *et al* (18) have evaluated the efficacy and safety of endoscopic resection

Table I. Comparison of the groups with and without post-endoscopic resection residues at resection margins.

Parameter	Residues at resection margin (n=73)	No Residues at resection margin (n=308)	P-value
Age (years)	64.8±8.9	60.9±11.4	0.891
Sex			0.435
Female	23	112	
Male	50	196	
Location			0.136
Upper	3	23	
Middle	42	201	
Lower	28	84	
Tumor diameter			<0.001
≤1 cm	2	29	
1.1-3 cm	28	190	
>3 cm	43	89	
Endoscopic treatment methods			0.03
Endoscopic mucosal resection	25	68	
Endoscopic submucosal dissection	48	240	
Endoscopic ultrasonography evaluation before resection			<0.001
Yes	42	257	
No	31	51	
Working experience			0.673
≥5 years	45	198	
<5 years	28	110	
Depth of invasion			<0.001
Intramucosal cancer (M1)	30	208	
Lamina propria (M2)	11	33	
Muscularis mucosa (M3)	9	45	
Shallow and deep submucosal layer (SM1-2)	23	22	
Differentiation			<0.001
Well	23	233	
Moderate	29	52	
Poor	21	23	

Table II. Results of multivariate logistic regression analysis.

Parameter	P-value	Odds ratio	95% Confidence interval
Age	0.820	1.051	0.686-1.610
Sex	0.437	0.763	0.385-1.510
Maximum diameter of resected specimen	0.39	1.325	0.698-2.515
Location	0.663	1.187	0.549-2.564
Endoscopic resection procedures	0.526	0.774	0.350-1.710
Depth of tumor invasion	<0.001	2.182	1.704-2.795
Work experience	0.155	0.624	0.325-1.196
Degree of tumor differentiation	0.018	0.451	0.166-1.224
Endoscopic ultrasonography before resection	<0.001	35.826	7.400-173.454

in selected cases with mucosal myometrial infiltration and upper third submucosal involvement. These studies further

verified the application of endoscopic resection in early esophageal cancer.

After the endoscopic resection of early esophageal cancer, the positive rate of resection edge varies greatly according to the literature, ranging 1.7-22% (9-11). In the present study, observation of atypical cells (including LIN, HIN or invasive cancer) at the resection margin was used as the criteria, where the positive rate of the resection margin was 19.2%. If only invasive cancer was considered, the positive rate of the resection edge then decreases to 5.4%. Previous studies have found that ESD has a higher *en bloc* resection and complete resection rates compared with those following traditional EMR (19-21). This finding is consistent with results from the present study, which found that the residual rates of EMR and ESD were 26.9 and 16.7%, respectively. However, based on logistic regression analysis, endoscopic resection was not found to be an independent risk factor, which is consistent with the results of Sgourakis *et al* (22). This finding may be associated with the results of retrospective analysis and not to the results of randomized controlled trials.

Over recent years, an increasing number of gastroenterologists in China consider ESD to be the optimal choice for the treatment of early esophageal cancer (23,24). They believe that with continuous advancements in ESD, the incidence of serious complications, such as perforation and bleeding, can be controlled to negligible levels (23-25). These studies further confirm that ESD is safe and effective in the treatment of early esophageal cancer.

Isomoto *et al* (26) previously reported that tumor size had no significant association with curative resection. However, the tumor size was significantly associated with segmental resection, where the cure resection rate was significantly lower compared with that of whole resection (26,27). In the present study, univariate analysis revealed that the maximum diameter of the primary tumor was associated with residual margin, but not in the multivariate analysis, which may be due to the nonrandomized endoscopic treatment methods (EMR and ESD) in the present retrospective analysis. In addition, the present study found that preoperative EUS examination and depth of tumor invasion are independent risk factors for residual resection margin. Previous studies have shown that EUS is the optimal noninvasive tool for T1 esophageal cancer, with a sensitivity of 85% and a specificity of 87% (28,29). This finding was also illustrated in previous studies of gastrointestinal neuroendocrine tumors (30,31) and is consistent with results from the present study.

Preoperative evaluation of the depth of tumor invasion can minimize the residual edge of deep resection (32). Lugol iodine staining, NBI amplification and EUS can all be used to evaluate lesion size and depth of tumor invasion (33,34). Through the above methods, it is possible to reduce the residual margin in the process of endoscopic resection. Furthermore, unnecessary resection should be avoided to prevent esophageal stenosis.

The degree of tumor differentiation has also been previously established to be an independent risk factor (31,35-37). However, no similar reports exist regarding esophageal cancer. Poorly differentiated tumors are typically more likely to invade the vascular system and lymph nodes earlier or cause deep infiltration (38). In the present study, only 23 (7.5%) poorly differentiated lesions were diagnosed in 308 specimens with negative margins. By contrast, the proportion of positive margin specimens (21 poorly differentiated, representing

28.8% of 73 specimens) was significantly higher compared with that of negative margin specimens. In a previous study with gastric cancer, in the high-risk category, the benefit of surgery appears to be positive, since the cancer-specific survival rate in the salvage surgery group was higher compared that in the follow-up group (39). These data suggest that if biopsy indicates poorly differentiated tumors and if the patient's physical conditions allow, more aggressive treatment strategies should be recommended. Due to the limited follow-up time, no corresponding data were available and these patients require continuous close attention.

It remains unclear whether surgery and other medical interventions should be actively performed for patients with residual margins after endoscopic resection. For the designation of subsequent treatment plans, the patient's age, complications and willingness should also be considered. In this group of data, amongst the nine patients with positive deep margins, six patients received surgical treatment, two patients received radiotherapy and chemotherapy, whereas one patient was closely observed. No recurrence was found during the follow-up. Among the 64 patients with positive lateral margins, one patient received radiotherapy, one patient received chemotherapy, six patients received surgical treatment, 20 patients received additional EMR or ESD, eight patients were lost to follow-up and the remaining 28 patients were closely observed. During the follow-up period, five patients succumbed to cardiovascular and cerebrovascular diseases whereas no recurrence was found in other patients. In the present study, a preliminary prognostic analysis of patients with positive margins after endoscopic resection of early esophageal cancer was performed. A more detailed prognostic analysis needs to be verified by a multicenter study with a longer term of follow-up.

The present study has a number of limitations. The present study is only a retrospective single-center study, not a randomized controlled study. Only one pathologist participated in the analysis of the tissues. Therefore, there may be potential selection bias in the present study. Prospective clinical trials may be conducted in the future to further verify these results.

In conclusion, the present study analyzed the risk factors associated with the residual margin and the prognosis of patients. According to the results of data analysis, it is highly recommended to apply EUS for evaluating the depth of tumor invasion, determining the depth of lesion invasion and the indication of endoscopic resection before operation, all of which can effectively prevent a residual margin. If biopsy indicates a poorly differentiated tumor, more aggressive treatment strategies may be required to prevent recurrence after endoscopic resection.

#### Acknowledgements

Not applicable.

#### Funding

The present study was supported by The Key topics of medical science research of Hebei Provincial Health Commission (grant no. 20190765).



### Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

### Authors' contributions

YF and BQL designed the study. YF and SG confirm the authenticity of all the raw data. YF and WW collected clinical and pathological data of patients. SG and YF analyzed the data. BQL and WW contributed to the interpretation of results. All authors critically reviewed the manuscript, and read and approved the final manuscript.

### Ethics approval and consent to participate

All patients and their families agreed to participate in the present study and signed an informed consent form. The study was approved by the Ethics Committee of the Fourth Hospital of Hebei Medical University (Shijiazhuang, China).

### Patient consent for publication

Not applicable.

### Competing interests

The authors declare that they have no competing interests.

### References

- Zhang L, Zhou Y, Cheng C, Cui HY, Cheng L, Kong PZ, Wang JQ, Li Y, Chen WL, Song B, *et al*: Genomic analyses reveal mutational signatures and frequently altered genes in esophageal squamous cell carcinoma. *Am J Hum Genet* 96: 597-611, 2015.
- Njei B, McCarty TR and Birk JW: Trends in esophageal cancer survival in United States adults from 1973 to 2009: A SEER database analysis. *J Gastroenterol Hepatol* 31: 1141-1146, 2016.
- Naveed M and Kubiliun N: Endoscopic treatment of early-stage esophageal cancer. *Curr Oncol Rep* 20: 71, 2018.
- Wani S, Early D, Edmundowicz S and Sharma P: Management of high-grade dysplasia and intramucosal adenocarcinoma in Barrett's esophagus. *Clin Gastroenterol Hepatol* 10: 704-711, 2012.
- Neuhaus H: Endoscopic submucosal dissection in the upper gastrointestinal tract: Present and future view of Europe. *Dig Endosc* 21 (Suppl 1): S4-S6, 2009.
- Chennat J, Konda VJ, Ross AS, de Tejada AH, Noffsinger A, Hart J, Lin S, Ferguson MK, Posner MC and Waxman I: Complete Barrett's eradication endoscopic mucosal resection: an effective treatment modality for high-grade dysplasia and intramucosal carcinoma-an American single-center experience. *Am J Gastroenterol* 104: 2684-2692, 2009.
- Oyama T, Tomori A, Hotta K, Morita S, Kominato K, Tanaka M and Miyata Y: Endoscopic submucosal dissection of early esophageal cancer. *Clin Gastroenterol Hepatol* 3 (Suppl 1): S67-S70, 2005.
- Mariette C, Fabre S, Balon JM, Finzi L and Triboulet JP: Factors predictive of complete resection of operable esophageal cancer: review of 746 patients. *Gastroenterol Clin Biol* 26: 454-462, 2002.
- Fujishiro M, Yahagi N, Kakushima N, Kodashima S, Muraki Y, Ono S, Yamamichi N, Tateishi A, Shimizu Y, Oka M, *et al*: Endoscopic submucosal dissection of esophageal squamous cell neoplasms. *Clin Gastroenterol Hepatol* 4: 688-694, 2006.
- Repici A, Hassan C, Carlino A, Pagano N, Zullo A, Rando G, Strangio G, Romeo F, Nicita R, Rosati R and Malesci A: Endoscopic submucosal dissection in patients with early esophageal squamous cell carcinoma: Results from a prospective Western series. *Gastrointest Endosc* 71: 715-721, 2010.
- Ono S, Fujishiro M, Niimi K, Goto O, Kodashima S, Yamamichi N and Omata M: Long-term outcomes of endoscopic submucosal dissection for superficial esophageal squamous cell neoplasms. *Gastrointest Endosc* 70: 860-886, 2009.
- Hamilton SR and Aaltonen LA: World Health Organization classification of tumours: Pathology and genetics of tumours of the digestive system. IARC Press, Lyon, pp11-36, 2000.
- Birkmeyer JD, Siewers AE, Finlayson EV, Stukel TA, Lucas FL, Batista I, Welch HG and Wennberg DE: Hospital volume and surgical mortality in the United States. *N Engl J Med* 346: 1128-1137, 2002.
- Ell C, May A, Pech O, Gossner L, Guenter E, Behrens A, Nachbar L, Huijsmans J, Vieth M and Stolte M: Curative endoscopic resection of early esophageal adenocarcinomas (Barrett's cancer). *Gastrointest Endosc* 65: 3-10, 2007.
- Pech O, Behrens A, May A, Nachbar L, Gossner L, Rabenstein T, Manner H, Guenter E, Huijsmans J, Vieth M, *et al*: Long-term results and risk factor analysis for recurrence after curative endoscopic therapy in 349 patients with high-grade intraepithelial neoplasia and mucosal adenocarcinoma in Barrett's oesophagus. *Gut* 57: 1200-1206, 2008.
- Pech O, Gossner L, May A, Vieth M, Stolte M and Ell C: Endoscopic resection of superficial esophageal squamous-cell carcinomas: Western experience. *Am J Gastroenterol* 99: 1226-1232, 2004.
- Stahl M, Budach W, Meyer HJ and Cervantes A: Esophageal cancer: Clinical practice guidelines for diagnosis, treatment and follow-up. *Ann Oncol* 21 (Suppl 5): v46-v49, 2010.
- Manner H, May A, Pech O, Gossner L, Rabenstein T, Günter E, Vieth M, Stolte M and Ell C: Early Barrett's carcinoma with 'low-risk' submucosal invasion: Long-term results of endoscopic resection with a curative intent. *Am J Gastroenterol* 103: 2589-2597, 2008.
- Hazama H, Tanaka M, Kakushima N, Yabuuchi Y, Yoshida M, Kawata N, Takizawa K, Ito S, Imai K, Hotta K, *et al*: Predictors of technical difficulty during endoscopic submucosal dissection of superficial esophageal cancer. *Surg Endosc* 33: 2909-2915, 2019.
- Plum PS, Hölscher AH, Pacheco Godoy K, Schmidt H, Berth F, Chon SH, Alakus H and Bollschweiler E: Prognosis of patients with superficial T1 esophageal cancer who underwent endoscopic resection before esophagectomy-A propensity score-matched comparison. *Surg Endosc* 32: 3972-3980, 2018.
- Wang HY, Zeng X, Bai SY, Pu K, Zheng Y, Ji R, Guo QH, Guan QL, Wang YP and Zhou YN: The safety and efficacy of endoscopic submucosal dissection for treating early oesophageal carcinoma: A meta-analysis. *Ann R Coll Surg Engl* 102: 702-711, 2020.
- Sgourakis G, Gockel I and Lang H: Endoscopic and surgical resection of T1a/T1b esophageal neoplasms: A systematic review. *World J Gastroenterol* 19: 1424-1437, 2013.
- Qi ZP, Chen T, Li B, Ren Z, Yao LQ, Shi Q, Cai SL, Zhong YS and Zhou PH: Endoscopic submucosal dissection for early esophageal cancer in elderly patients with relative indications for endoscopic treatment. *Endoscopy* 50: 839-845, 2018.
- Wu Y, Zhang H, Zhou B, Han SY and Zhang YR: Clinical efficacy of endoscopic submucosal dissection in the treatment of early esophageal cancer and precancerous lesions. *J Cancer Res Ther* 14: 52-56, 2018.
- Yip HC and Chiu PW: Endoscopic diagnosis and management of early squamous cell carcinoma of esophagus. *J Thorac Dis* 9 (Suppl 8): S689-S696, 2017.
- Isomoto H, Shikuwa S, Yamaguchi N, Fukuda E, Ikeda K, Nishiyama H, Ohnita K, Mizuta Y, Shiozawa J and Kohno S: Endoscopic submucosal dissection for early gastric cancer: A large-scale feasibility study. *Gut* 58: 331-336, 2009.
- Chung IK, Lee JH, Lee SH, Kim SJ, Cho JY, Cho WY, Hwangbo Y, Keum BR, Park JJ, Chung HJ, *et al*: Therapeutic outcomes in 1000 cases of endoscopic submucosal dissection for early gastric neoplasms: Korean ESD study group multicenter study. *Gastrointest Endosc* 69: 1228-1235, 2009.
- Thosani N, Singh H, Kapadia A, Ochi N, Lee JH, Ajani J, Swisher SG, Hofstetter WL, Guha S and Bhutani MS: Diagnostic accuracy of EUS in differentiating mucosal versus submucosal invasion of superficial esophageal cancers: A systematic review and meta-analysis. *Gastrointest Endosc* 75: 242-253, 2012.

29. Yang J, Luo GY, Liang RB, Zeng TS, Long H, Fu LH, Xu GL, Yang MZ, Li S, Zhang LJ, *et al*: Efficacy of endoscopic ultrasonography for determining clinical T category for esophageal squamous cell carcinoma: Data from 1434 surgical cases. *Ann Surg Oncol* 25: 2075-2082, 2018.
30. Ishii N, Horiki N, Itoh T, Maruyama M, Matsuda M, Setoyama T, Suzuki S, Uchida S, Uemura M, Iizuka Y, *et al*: Endoscopic submucosal dissection and preoperative assessment with endoscopic ultrasonography for the treatment of rectal carcinoid tumors. *Surg Endosc* 24: 1413-1419, 2010.
31. Zhou FR, Huang LY and Wu CR: Endoscopic mucosal resection for rectal carcinoids under micro-probe ultrasound guidance. *World J Gastroenterol* 19: 2555-2559, 2013.
32. Lee TH, Cho JY, Chang YW, Kim JO, Lee JS, Cho WY, Kim HG, Kim WJ, Park YS and Jin SY: Appropriate indications for endoscopic submucosal dissection of early gastric cancer according to tumor size and histologic type. *Gastrointest Endosc* 71: 920-926, 2010.
33. Fujishiro M, Kodashima S, Goto O, Ono S, Niimi K, Yamamichi N, Oka M, Ichinose M and Omata M: Endoscopic submucosal dissection for esophageal squamous cell neoplasms. *Dig Endosc* 21: 109-115, 2009.
34. Yoshida T, Inoue H, Usui S, Satodate H, Fukami N and Kudo SE: Narrow-band imaging system with magnifying endoscopy for superficial esophageal lesions. *Gastrointest Endosc* 59: 288-295, 2004.
35. Kakushima N, Ono H, Tanaka M, Takizawa K, Yamaguchi Y and Matsubayashi H: Factors related to lateral margin positivity for cancer in gastric specimens of endoscopic submucosal dissection. *Dig Endosc* 23: 227-232, 2011.
36. Wen J, Linghu EQ, Yang YS, Liu QS, Yang J and Lu ZS: Associated risk factor analysis for positive resection margins after endoscopic submucosal dissection in early-stage gastric cancer. *J BUON* 20: 421-427, 2015.
37. Hwang JJ, Park KJ, Park YS, Lee HS, Yoon H, Shin CM, Kim N and Lee DH: A scoring system for patients with a tumor-positive lateral resection margin after endoscopic resection of early gastric cancer. *Surg Endosc* 30: 2751-2758, 2016.
38. Oyama T, Inoue H, Arima M, Momma K, Omori T, Ishihara R, Hirasawa D, Takeuchi M, Tomori A and Goda K: Prediction of the invasion depth of superficial squamous cell carcinoma based on microvessel morphology: Magnifying endoscopic classification of the Japan esophageal society. *Esophagus* 14: 105-112, 2017.
39. Hatta W, Gotoda T, Oyama T, Kawata N, Takahashi A, Yoshifuku Y, Hoteya S, Nakagawa M, Hirano M and Esaki M, *et al*: Is the eCura system useful for selecting patients who require radical surgery after noncurative endoscopic submucosal dissection for early gastric cancer? A comparative study. *Gastric Cancer* 21: 481-489, 2018.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.