BRIEF REPORT

ANIMAL GENETICS WILEY

A de novo start-lost variant in ANKRD28 in a Holstein calf with dwarfism

BACKGROUND

Osteochondrodysplasias comprise a broad spectrum of hereditary developmental disorders of bone and cartilage, but can also have significant effects on tendons, ligaments, and muscles (Warman et al., 2011). According to Online Mendelian Inheritance in Animals, there are six known dwarfism phenotypes in cattle, which are inherited both recessively and dominantly and are due to structural or signaling pathways disruptions (OMIA 000299-9913; Boegheim et al., 2017).

ANALYSIS

A 3.5-month-old, female, purebred Holstein calf was reported because of its short stature, protruding scapulohumeral joints, increased concavity of the frontal bone, and bossing of the intercornual prominence (Figure S1). A whole-genome sequencing (WGS) approach was performed as described before (Jacinto et al., 2022) using genomic DNA extracted from blood of the affected calf, her dam, and from semen of her sire. Reads were mapped to the ARS-UCD1.2 assembly (Rosen et al., 2020) resulting in an average read depth of 20.7× in the calf, 21.8× in the dam, and 22.1× in the sire, and then processed as reported earlier (Jacinto et al., 2021). We hypothesized that the affected calf had a novel form of dwarfism and investigated the genetic origin. The trio-based WGS approach identified no homozygous protein-changing variants present exclusively in the genome of the affected calf. Consequently, a recessive inheritance seems unlikely. Given that the reported calf was the only case and dwarfism had never before been reported in the Canadian Holstein population, we hypothesized that a dominant mode of inheritance due to a de novo mutation event was more likely. We found three heterozygous private protein-changing variants present in the calf and absent in both parental genomes and in 5365 controls (Table S1). Review of the sequencing data for each of these heterozygous variants revealed that all were indeed de novo variants affecting different genes. Only one of these variants concerns an interesting putative candidate gene (*ANKRD28*) for the observed phenotype. This heterozygous variant at chr1:152807533C>CA represents a 1-bp insertion in exon 2 of the *ANKRD28* gene that directly affect the translation initiation (start) codon (XM_024989836.1: c.2dupT; Figure S1). The predicted consequence of this start-lost variant is that amino acid Met1 is converted to Leu after activation of an upstream translation initiation site at cDNA position -20, resulting in N-terminal protein extension by insertion of six amino acids between Met1 and Gly2 (XP_024845604.1: p.Met1_Gly2insIleValGlyGlyLysAlaLeu).

COMMENTS

We propose the heterozygous c.2dupT variant as a candidate causative variant for the observed congenital disorder and *ANKRD28* as a novel candidate gene for dwarfism phenotypes. The protein-altering nature of this de novo mutation strongly suggests the causality of the variant. The Genome Aggregation Database showed that *ANKRD28* falls into the class of loss-of-function haploinsufficient genes (Karczewski et al., 2020). Therefore, two different situations could explain the perceived phenotype: either haploinsufficiency or co-expression of an N-terminally extended protein, leading to non- or abnormal expression of the mutant allele.

Nonetheless, since this is a single case study and we have no functional confirmation, this result must be considered preliminary and should be interpreted with caution. It must also be noted that the analysis of short-read genomic data is not ideal for identifying larger structural variants. Further individual cases of dwarfism in cattle or other animal species could be investigated for *ANKRD28* variants by DNA sequencing.

ACKNOWLEDGEMENTS

We thank the Interfaculty Bioinformatics Unit of the University of Bern for providing high-performance computational infrastructure. Irene M. Häfliger was funded

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2022 The Authors. Animal Genetics published by John Wiley & Sons Ltd on behalf of Stichting International Foundation for Animal Genetics.

by the Swiss National Science Foundation, grant number 172911. Wayne Dickieson of Birkentree Holsteins is gratefully acknowledged. Open access funding provided by University of Bern.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

DATA AVAILABILITY STATEMENT

The WGS data are available under the study accession no. PRJEB18113 at the European Nucleotide Archive (www.ebi.ac.uk/ena; calf SAMEA8565094, dam SAMEA8565093, sire SAMEA8565095).

Joana G. P. Jacinto^{1,2} De Irene M. Häfliger² Christine F. Baes^{2,3} De Hinayah R. de Oliveira Cord Drögemüller² De Irene M. Jacinto Jacoba P. Jacinto J. Jacinto Jacoba P. Jacinto J. Jacinto Jacoba P. Jacinto J. Jacoba P. Jac

¹Department of Veterinary Medical Sciences, University of Bologna, Ozzano Emilia (Bologna), Italy

²Vetsuisse Faculty, Institute of Genetics, University of Bern, Bern, Switzerland

³Department of Animal Biosciences, Centre for the Genetic Improvement of Livestock, University of Guelph, Guelph, Ontario, Canada

Correspondence

Cord Drögemüller, Vetsuisse Faculty, Institute of Genetics, University of Bern, 3001 Bern, Switzerland.

Email: cord.droegemueller@vetsuisse.unibe.ch

ORCID

Joana G. P. Jacinto https://orcid.org/0000-0002-6438-7975

Irene M. Häfliger https://orcid.org/0000-0002-5648-963X

Christine F. Baes https://orcid.org/0000-0001-6614-8890

Hinayah R. de Oliveira https://orcid.org/0000-0002-0355-8902

Cord Drögemüller thttps://orcid.org/0000-0001-9773-522X

REFERENCES

Boegheim, I.J.M., Leegwater, P.A.J., van Lith, H.A. & Back, W. (2017) Current insights into the molecular genetic basis of dwarfism in livestock. *The Veterinary Journal*, 224, 64–75. Available from: https://doi.org/10.1016/j.tvjl.2017.05.014

Jacinto, J.G.P., Häfliger, I.M., Akyürek, E.E., Sacchetto, R., Benazzi, C., Gentile, A. et al. (2021) KCNG1-related syndromic form of congenital neuromuscular channelopathy in a crossbred calf. Genes, 12(11), 1792. Available from: https://doi.org/10.3390/genes 12111792

Jacinto, J.G.P., Häfliger, I.M., Bernardini, M., Mandara, M.T., Bianchi, E., Bolcato, M. et al. (2022) A homozygous missense variant in laminin subunit beta 1 as candidate causal mutation of hemifacial microsomia in Romagnola cattle. *Journal of Veterinary Internal Medicine*, 36(1), 292–299. Available from: https://doi.org/10.1111/jvjm.16316

Karczewski, K.J., Francioli, L.C., Tiao, G., Cummings, B.B., Alföldi, J., Wang, Q. et al. (2020) The mutational constraint spectrum quantified from variation in 141,456 humans. *Nature*, 581(7809), 434–443. Available from: https://doi.org/10.1038/s4158 6-020-2308-7

Rosen, B.D., Bickhart, D.M., Schnabel, R.D., Koren, S., Elsik, C.G., Tseng, E. et al. (2020) De novo assembly of the cattle reference genome with single-molecule sequencing. *GigaScience*, 9(3), 1–9. Available from: https://doi.org/10.1093/gigascience/giaa021

Warman, M.L., Cormier-Daire, V., Hall, C., Krakow, D., Lachman, R., LeMerrer, M. et al. (2011) Nosology and classification of genetic skeletal disorders: 2010 revision. *American Journal of Medical Genetics Part A*, 155A(5), 943–968. Available from: https://doi.org/10.1002/ajmg.a.33909

SUPPORTING INFORMATION

Additional supporting information may be found in the online version of the article at the publisher's website.

How to cite this article: Jacinto, J.G.P., Häfliger, I.M., Baes, C.F., de Oliveira, H.R. & Drögemüller, C. (2022) A de novo start-lost variant in *ANKRD28* in a Holstein calf with dwarfism. *Animal Genetics*, 53, 470–471. Available from: https://doi.org/10.1111/age.13204