Theoretical Biology and Medical Modelling



Editorial Open Access

Editorial: hypotheses about protein folding - the proteomic code and wonderfolds

Paul S Agutter

Address: Theoretical Medicine and Biology Group, 26 Castle Hill, Glossop, Derbyshire, SK13 7RR, UK Email: Paul S Agutter - psa@tmedbiol.com

Published: 24 December 2009

Theoretical Biology and Medical Modelling 2009, 6:31 doi:10.1186/1742-4682-6-31

This article is available from: http://www.tbiomed.com/content/6/1/31

© 2009 Agutter; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/2.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Theoretical biology journals can contribute in many ways to the progress of knowledge. They are particularly well-placed to encourage dialogue and debate about hypotheses addressing problematical areas of research. An online journal provides an especially useful forum for such debate because of the option of posting comments within days of the publication of a contentious article.

Editorial

Theoretical biology' encompasses proposals ranging from new mathematical models of well-studied biological processes to speculative notions that inhabit the borderland between science and philosophy. *Theoretical Biology and Medical Modelling* can accommodate everything within that range subject to peer review and editorial approval. Novel hypotheses addressing phenomena that have defied satisfactory explanation are especially welcome, provided they meet the basic criterion of testability (at least in principle), because they can stimulate debate or excite controversy and are *ipso facto* healthy for science.

Theoretical biology has not enjoyed the status of, say, theoretical physics because biology is primarily a science of particular phenomena rather than general laws. 'Grand theories' in biology seldom prove useful or even tenable, as some widely-discussed instances have shown during the past decade. Nevertheless, there are problematic areas of the life sciences that invite theoretical exploration. Protein folding is an example. Most research in this field, as in others, is empirical and pertains *a fortiori* to only a limited range of polypeptides and/or species (e.g. [1,2]). Broad hypotheses about general mechanisms of protein

folding may therefore initiate significant contributions to knowledge.

Received: 16 December 2009 Accepted: 24 December 2009

The 'proteomic code' is one such hypothesis. Its basic claim is that while protein primary structure is encoded in the base sequence of mRNA, the rules for protein folding are encoded in other features of messenger structure. Jan Biró of the Homulus Foundation, Los Angeles, has developed the idea in a recent series of papers [3-6], some of them published in this journal, and has recently published a book that explains it in detail [7]. The proteomic code hypothesis is likely to find support among some workers in the protein structure field, but is equally likely to find powerful opponents.

Biró's point of departure is the well-known redundancy of the genetic code. Studying 81 messengers, he showed [3,4] that mRNA subsequences comprising 1st and/or 3rd codon residues have significantly higher free folding energies than subsequences containing only 2nd residues (p < 0.0001). No such periodically distributed differences in free folding energy were found in intron transcripts. This suggests selection for local secondary structures in RNA coding regions, and these structures resemble the folding

profiles of the encoded proteins. In particular, codons synonymous in respect of their encoded amino acids may nevertheless signify differences in protein secondary or tertiary structure. Thus, messengers not only direct the assembly of polypeptides with the correct primary sequence (the genetic code), they also direct the correct folding of those polypeptides (the proteomic code) [5,6].

This concept was first suggested a quarter of a century ago by Biró himself, and independently by Mekler, and was developed in studies by Blalock, Root-Bernstein, Siemion, Miller and others [7]. In 2003, Biró and colleagues published a common periodic table of codons and amino acids, which elaborates the proteomic code hypothesis in specific detail [8]. The idea is strikingly consistent with studies such as those of Chiusano et al. [9], who showed that the nucleotide frequencies in second codon positions are remarkably different among coding regions that correspond to different protein secondary structures and to amino acids with different physicochemical properties. It is also broadly compatible with the work of Ikehara and colleagues [10,11] and of Rodin and Rodin [12] on the origin and evolution of the genetic code.

However, some research conflicts with the proteomic code concept. A salient example is the work by Berezovsky and colleagues [13,14], whose emphasis is on polymer physics and on the selection for protein stability that causes preferred polypeptide structures to emerge. These authors have identified structural motifs that they dub 'wonderfolds', which arise repeatedly as native states of stable polypeptides resulting from the mutation and selection of random sequences. They reason that superfamilies with wonderfolds may have played an important part in early evolution. This approach to the study of protein folding has no connection at all with mRNA structure or the distinctive properties of codon bases. It seems likely that Berezovsky and his colleagues would dismiss the proteomic code hypothesis as speculative and unproductive, whereas proponents of the proteomic code may wish to relate 'wonderfolds' to particular recurrent combinations of mRNA codons (which would then, in turn, require explanation).

This is a potentially fruitful arena for continuing debate and discussion. Currently, the main questions seem to be (1) whether either hypothesis satisfactorily explains empirical results such as those in [1,2] and (2) whether the two hypotheses - which at present seem incompatible - can ultimately be reconciled. By fostering the further exploration of these and related questions, theoretical biology journals are in a position to make valuable contributions to knowledge. Theoretical Biology and Medical Modelling is particularly well placed in this regard because it provides the option of posting comments on contentious articles within days of their online publication.

References

- Preuss M, Miller AD: The affinity of the GroEL/GroES complex for peptides under conditions of protein folding. FEBS Lett 2000. 466:75-79.
- Pintar A, Pongor S: The "first in-last out" hypothesis on protein folding revisited. Proteins 2005, 60:584-590.
- Biró JC: Indications that "codon boundaries" are physicochemically defined and that protein-folding information is contained in the redundant exon bases. Theor Biol Med Model
- Biró JC: Protein folding information in nucleic acids which is not present in the genetic code. Ann N Y Acad Sci 2006, 1091:399-411.
- Biró JC: The Proteomic Code: a molecular recognition code for proteins. Theor Biol Med Model 4:45
- Biró JC: Discovery of proteomic code with mRNA assisted
- protein folding. Int J Mol Sci 2008, 9:2424-2446. Biró JC: Principia Bi®o-Informatica. Creative ideas in Molecular Biology & Bioinformatics Los Angeles: Homulus Foundation; 2009. ISBN 978-0-
- Biro JC, Benyó B, Sansom C, Szlávecz A, Fördös G, Micsik T, Benyó Z: A common periodic table of codons and amino acids. Biochem Biophys Res Commun 2003, 306:408-415.
- Chiusano ML, Alvarez-Valin F, Di Giulio M, D'Onofrio G, Ammirato G, Colonna G, Bernardi G: Second codon positions of genes and the secondary structures of proteins. Relationships and implications for the origin of the genetic code. Gene 2000, **261:**63-69.
- Ikehara K, Niihara Y: Origin and evolutionary process of the genetic code. Curr Med Chem 2007, 14:3221-3231.
- Ikehara K: Pseudo-Replication of [GADV]-proteins and origin of life. Int J Mol Sci 2009, 10:1525-1537
- Rodin SN, Rodin AS: On the origin of the genetic code: signatures of its primordial complementarity in tRNAs and aminoacyl-tRNA synthetases. Heredity 2008, 100:341-355
- 13. Berezovsky IN, Trifonov EN: Flowering buds of globular proteins: transpiring simplicity of protein organization. Comp Funct Genomics 2002, 3:525-534.
- Zeldovich KB, Berezovsky IN, Shakhnovich El: Physical origins of protein superfamilies. | Mol Biol 2006, 357:1335-1343.

Publish with **Bio Med Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- · yours you keep the copyright

Submit your manuscript here: http://www.biomedcentral.com/info/publishing adv.asp

