



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



## Vitamin D status can affect COVID-19 outcomes also in pediatric population

Michele Miraglia del Giudice<sup>a,\*</sup>, Cristiana Indolfi<sup>a</sup>, Giulio Dinardo<sup>a</sup>, Fabio Decimo<sup>a</sup>,  
Alberto Decimo<sup>b</sup>, Angela Klain<sup>a</sup>

<sup>a</sup> Department of Woman, Child and General and Specialized Surgery, University of Campania "Luigi Vanvitelli", Naples, Italy

<sup>b</sup> Department of Mental, Physical Health and Preventive Medicine, University of Campania "Luigi Vanvitelli", Naples, Italy

### ARTICLE INFO

#### Keywords:

Vitamin D  
COVID-19  
SARS-CoV-2 infection  
Children  
Immunity

### ABSTRACT

**Background:** vitamin D influences the immune system and the inflammatory response. It is known that vitamin D supplementation reduces the risk of acute respiratory tract infection. In the last two years, many researchers have investigated vitamin D's role in the pathophysiology of COVID-19 disease.

**Results:** the findings obtained from clinical trials and systematic reviews highlight that most patients with COVID-19 have decreased vitamin D levels and low levels of vitamin D increase the risk of severe disease. This evidence seems to be also confirmed in the pediatric population.

**Conclusions:** further studies (systematic review and meta-analysis) conducted on children are needed to confirm that vitamin D affects COVID-19 outcomes and to determine the effectiveness of supplementation and the appropriate dose, duration and mode of administration.

### 1. Exploring the immune link between vitamin D and SARS-CoV-2

Vitamin D is the collective name for cholecalciferol (vitamin D3) and ergocalciferol (vitamin D2); the active hormonal form is calcitriol or 1,25-dihydroxyvitamin D, obtained by hydroxylation in the liver and kidney [1]. Vitamin D plays a crucial role in the regulation of the metabolism of calcium and phosphates, regulating bone growth and the maintenance of mineral homeostasis, as well as acting on different peripheral organs and tissues such as kidneys, bones, parathyroids, skin, thymus, pituitary glands, uterus, placenta, and various cellular elements of the immune system. Vitamin D works on the adaptive and innate immune systems, with anti-inflammatory and immunomodulating activity [2]. Cells of the adaptive immune system are not only targets of vitamin D metabolites, but they are also able to activate and inactivate vitamin D metabolites. Both monocytes and dendritic cells (DCs) express vitamin D receptors (VDRs). Surface costimulatory molecules like CD40, CD80, and CD86 are decreased in monocytes cultured with 1,25(OH)<sub>2</sub>D [3]. 1,25(OH)<sub>2</sub>D inhibits the differentiation of monocytes into immature

DCs. Studies show that DCs matured in vitro in the presence of 1,25(OH)<sub>2</sub>D, have an altered cytokine and chemokine profile: the release of Th1 and Th17 inducing cytokines, interleukin (IL)-12 and IL-23, is inhibited, while the production of IL-10 (anti-inflammatory cytokine) and Treg cells (regulatory T cell) is enhanced [4,5]. Altogether these results show that vitamin D can block DC maturation, IL-12 production and enhance the production of IL-10 and Treg cells. Most in vitro studies on human cluster of differentiation (CD) 4 + T cells show that 1,25(OH)<sub>2</sub>D is able to reduce interferon-gamma (IFN-γ) secretion, inhibit Th1 cells and Th2 cell differentiation, and enhance the secretion of IL-4 by the already differentiated Th2 cells [2,6]. Vitamin D has been shown to interact with the innate immune system by activating Toll-like receptors (TLRs) or increasing the levels of cathelicidins and β-defensins. TLRs play an important role in recognizing viral particles and activating the innate immune system. Vitamin D increases TLR2 and TLR4 on the cord blood monocytes, TLR2/6 on the keratinocytes and TLR2, TLR3, TLR8 on the immune cells, strengthening the immune responses against microbial infections [7]. Different TLRs, like TLR2, TLR3, TLR4, TLR6, TLR7, TLR8, and TLR9 are potentially important in COVID-19 infection,

**Abbreviations:** DCs, dendritic cells; VDRs, vitamin D receptors; IL, interleukin; Treg cells, CD, regulatory T cell cluster of differentiation; IFN-γ, reduce interferon-gamma; TLRs, Toll-like receptors; RAS, renin-angiotensin system; ACE2, angiotensin-converting enzyme 2; IgE, immunoglobulin E; PAMPs, pathogen-associated molecular patterns; DAMPs, damage-associated molecular patterns; PRRs, pattern recognition receptors; TNF-α, tumor necrosis factor-α; ICU, intensive care unit; NK, natural killer; RCTs, randomized control trials; RR, risk ratio; CI, confidence interval; OR, odds ratio; HR, Hazard Risk; MD, mean difference.

\* Correspondence to: Department of Woman, Child and General and Specialized Surgery, University of Campania "Luigi Vanvitelli", Via Luigi de Crecchio, 4, 80138 Naples, Italy.

E-mail address: [michele.miraglia@unicampania.it](mailto:michele.miraglia@unicampania.it) (M. Miraglia del Giudice).

<https://doi.org/10.1016/j.phanu.2022.100319>

Received 3 May 2022; Received in revised form 12 October 2022; Accepted 12 October 2022

Available online 14 October 2022

2213-4344/© 2022 Published by Elsevier B.V.

representing a potential target in controlling the infection in the early stages of the disease [8]. Adequate vitamin D levels are also required in order to reduce renin-angiotensin system (RAS) activity and increase angiotensin-converting enzyme 2 (ACE2) concentrations in acute lung injury [9]. Furthermore, low circulating vitamin D levels have been associated with the risk of asthma, atopic dermatitis, and elevated total immunoglobulin E (IgE) [10–15].


In COVID-19 pathogenesis, SARS-CoV-2 is able to activate both the innate and adaptive immune response. The viral particle attaches to ACE2 receptor, which is present on lung cells, reduces ACE2 expression and then enters via endocytosis [16]. ACE2 regulates RAS, thus its downregulation breaks RAS homeostasis, affecting blood pressure, electrolyte balance, increasing inflammation and vascular permeability in the airways. Innate immune cells detect the pathogen-associated molecular patterns (PAMPs) and damage-associated molecular patterns (DAMPs) present on the virus particle using pattern recognition receptors (PRRs), which include TLRs, and generate local inflammation and releasing cytokines and chemokines including IFN- $\gamma$ , IL-6, MCP1, and IP-10. SARS-CoV-2-specific CD4 + T cells display IFN- $\gamma$ , tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ), IL-2 chemokines, enhancing the activation of Th1 immune response and cell-mediated immunity [9,17]. T cells, CD4 +, Th1 and Th17, and CD8 + T cells play a critical role in the adaptive immunity response. It consists in promoting the secretion of pathogen-specific antibodies by inducing T-dependent B cells and killing the virus-infected cells. Furthermore, virus-specific memory CD8 + T cells have a significant role in host protection from lethal SARS-CoV-2 infection by the production of IFN- $\gamma$ , TNF- $\alpha$ , IL-2, and cytolytic molecules, such as granzyme B [18] [Fig. 1]. In a case-control study by Bayraktar et al., the role and relationship between the cytokine profile and vitamin D were investigated. The serum cytokine levels IL-1, IL-6, IL-10, IL-21, TNF- $\alpha$  and vitamin D were measured in the COVID-19 patient group (intensive care unit (ICU) patients with severe illnesses) and the control group (individuals without a history of serious illness or infection): in the COVID-19 group cytokine levels were significantly higher than control groups, while serum vitamin D was significantly low [19].

The pathophysiological and clinical disparities between children and adults with COVID-19 may be due to different factors. Children have a more potent innate immune response with more natural killer (NK) cells, which serves as the initial line of defence against SARS-CoV-2 [20]. Another significant element is "trained immunity," which is the process by which innate immune cells undergo epigenetic reprogramming as a result of exposure to specific stimuli, such as infections and immunizations, and develop "memory." Children also have greater proportions of

lymphocytes and absolute numbers of T and B cells in connection to adaptive immunity, while ageing is linked to a decline in thymic activity and a decline in naive T cell numbers [21]. Children may also be protected by more frequent concurrent and recurrent infections, innate immunity to coronaviruses, variations in their microbiome, increased melatonin levels, and protective off-target effects of live vaccinations [21].

## 2. Does vitamin D deficiency increase the risk and severity of COVID-19?

Several studies have investigated a possible association between vitamin D serum levels and the risk and severity of SARS-CoV-2 infection. The most important evidence on the link between vitamin D and COVID-19 comes from systematic reviews and meta-analyses. In the meta-analysis by Rawat et al., 5 studies (3 randomized control trials (RCTs) and 2 Quasi-experimental) including 467 patients were included: vitamin D didn't reduce mortality (risk ratio (RR) 0.55, 95 % confidence interval (CI) 0.22–1.39,  $p = 0.21$ ), ICU admission rates (RR 0.20, 95 % CI 0.01–4.26,  $p = 0.3$ ) and need for invasive ventilation (RR 0.24, 95 % CI 0.01–7.89,  $p = 0.42$ ) [22]. In the systematic review by Pereira et al., vitamin D deficiency was not associated with a higher chance of infection by COVID-19 (OR = 1.35; 95 % CI = 0.80–1.88), but severe cases of COVID-19 present 64 % (OR = 1.64; 95 % CI = 1.30–2.09) more vitamin D deficiency compared with mild cases. Vitamin D concentration insufficiency increased hospitalization (OR = 1.81, 95% CI = 1.41–2.21) and mortality from COVID-19 (OR = 1.82, 95 % CI = 1.06–2.58) [23]. Twenty-three studies containing 11,901 participants entered into the meta-analysis by Ghasemian et al., 41 % of COVID-19 patients were suffering from vitamin D deficiency (95 % CI, 29–55 %), and in 42 % of patients, levels of vitamin D were insufficient (95 % CI, 24–63 %). The serum 25-hydroxyvitamin D concentration was 20.3 ng/ml among all COVID-19 patients (95 % CI, 12.1–19.8). The odds of getting infected with SARS-CoV-2 were 3.3 times higher among individuals with vitamin D deficiency (95 % CI, 2.5–4.3), while the chance of developing severe COVID-19 was about five times higher in patients with vitamin D deficiency (OR: 5.1, 95 % CI, 2.6–10.3). There was no significant association between vitamin D status and higher mortality rates (OR: 1.6, 95 % CI, 0.5–4.4) [24]. In the systematic review by Kazemi et al., thirty-nine studies were included: fifteen studies evaluated associations between vitamin D deficiency and composite severity. In the studies that were adjusted (OR: 2.57; 95 % CI: 1.65, 4.01; I<sup>2</sup> = 0.0 %) and nonadjusted for confounders (OR: 10.61; 95 % CI: 2.07, 54.23; I<sup>2</sup> = 90.8 %) there was a higher severity in the vitamin D deficiency group. Analysis of studies



**Fig. 1.** Comparison of the immune response induced by vitamin D administration and SARS-CoV-2 infection. Vitamin D increases the production of Toll-like receptors (TLRs), IL-10 (anti-inflammatory cytokine) and Treg cells (regulatory T cell), while the release of Th1 and Th17 inducing cytokines, interleukin (IL)–12 and IL-23, is inhibited. In the COVID-19 pathogenesis, pattern recognition receptors (PRRs), which include TLRs, are used by innate immune cells to recognize the pathogen-associated molecular patterns (PAMPs) and damage-associated molecular patterns (DAMPs) found on virus particles. This mechanism triggers local inflammation and the release of cytokines and chemokines such as IFN- $\gamma$  (interferon-gamma), IL-6, MCP1, and IP-10. IFN- $\gamma$ , tumor necrosis factor (TNF), and IL-2 chemokines are expressed on SARS-CoV-2-specific CD4 + T cells, which promote the activation of the Th1 immune response and cell-mediated immunity.

with crude OR (OR: 2.62; 95 % CI: 1.13, 6.05; I2: 47.9 %), and adjusted studies that used the Cox survival method (HR: 7.67; 95 % CI: 3.92, 15.03; I2: 0.0 %) indicated a significant association of vitamin D deficiency with mortality, while in adjusted studies that used logistic regression, no relation was observed (OR: 1.05; 95 % CI: 0.63, 1.75; I2: 76.6 %). The results of studies that examined relations between vitamin D deficiency and ICU admission, pulmonary complications, hospitalization, and inflammation were inconsistent [25]. In October 2021 Petrelli et al. assessed the association between vitamin D and risk, severity, and mortality for COVID-19 infection, through a review of 43 observational studies. Among subjects with deficient vitamin D values, risk of COVID-19 infection was higher compared to those with replete values (OR = 1.26; 95 % CI, 1.19–1.34;  $P < 0.01$ ). Vitamin D deficiency was also associated with worse severity and higher mortality than in non-deficient patients (OR = 2.6; 95 % CI, 1.84–3.67;  $P < 0.01$  and OR = 1.22; 95 % CI, 1.04–1.43;  $P < 0.01$ , respectively) [26]. According to the review by Pal et al., vitamin D use in COVID-19 was significantly associated with reduced ICU admission/mortality (OR 0.41, 95 % CI: 0.20, 0.81,  $p = 0.01$ , I2 = 66 %, random-effects model) and decreased the risk of adverse outcomes (pooled OR 0.27, 95 % CI: 0.08, 0.91,  $p = 0.03$ , I2 = 80 %, random-effects model). Subgroup analysis showed that vitamin D supplementation was associated with improved clinical outcomes only in patients receiving the drug post-COVID-19 diagnosis and not in those who had received vitamin D before diagnosis [27]. In 2022 the meta-analysis of Shah et al. showed that vitamin D supplementation reduces the risk of mortality (OR: 0.48, 95 % CI: 0.346–0.664;  $p < 0.001$ ), the need for ICU (OR: 0.35; 95 % CI: 0.28–0.44;  $p < 0.001$ ) and mechanical ventilation (OR: 0.54; 95 % CI: 0.411–0.708;  $p < 0.001$ ) requirement in COVID patients [28]. In an adult population-based study, the researchers observed that in comparison to non-supplemented 25-hydroxyvitamin D-deficient patients, those receiving cholecalciferol treatment and achieving vitamin D levels 30 ng/ml had a lower risk of SARS-CoV2 infection, a lower risk of severe COVID-19, and a lower risk of COVID-19 mortality (56/9474 [0.6 %] vs 96/7616 [1.3 %]; Hazard Risk (HR) 0.66 [CI 95 % 0.46–0.93],  $p = 0.018$ ). In the entire cohort ( $n = 134,703$ ), calcifediol use was not linked to a lower incidence of SARS-CoV2 infection or mortality. When compared to 25-hydroxyvitamin D-deficient patients not receiving vitamin D supplements, patients on calcifediol treatment who achieved serum vitamin D levels 30 ng/ml also had a lower risk of SARS-CoV2 infection, a lower risk of severe COVID-19, and a lower risk of COVID-19 mortality (88/16276 [0.5 %] vs 96/7616 [1.3 %]; HR 0.56 [CI 95 % 0.42–0.76],  $p < 0.001$ ) [29]. In the analysis by Wang et al., including 17 observational studies with 2756 individuals, in comparison to non-deficient vitamin D status, vitamin D deficiency was linked to significantly higher rates of mortality (OR: 2.47, 95 % CI 1.50–4.05; 12 studies; HR: 4.11, 95 % CI: 2.40–7.04; 3 studies), higher rates of hospital admissions (OR: 2.18, 95 % CI: 1.48–3.21; 3 studies), and longer hospital stays (0.52 days, 95 % CI: 0.25–0.80; 2 studies) [30]. In the review by Teshome et al., the pooled analysis showed that individuals with vitamin D deficiency were 80 % more likely to acquire COVID-19 infection as compared to those who have sufficient vitamin D levels (OR = 1.80; 95 % CI: 1.72, 1.88) [31]. In a systematic review conducted in Sri Lanka, 72 observational studies were included ( $n = 1\,976,099$ ). Vitamin D deficiency/insufficiency increased the odds of developing COVID-19 (OR=1.46; 95 % CI, 1.28–1.65;  $P < 0.0001$ ), severe disease (OR 1.90; 95 % CI, 1.52–2.38;  $P < 0.0001$ ), and death (OR 2.07; 95 % CI, 1.28–3.35;  $P = 0.003$ ). Patients with COVID-19 had lower 25-hydroxy vitamin D concentrations than controls (mean difference [MD]  $-3.85$  ng/ml; 95 % CI,  $-5.44$  to  $-2.26$ ;  $P = 0.0001$ ), patients with severe COVID-19 had lower 25-hydroxy vitamin D concentrations than controls with non-severe COVID-19 (MD  $-4.84$  ng/ml; 95 % CI,  $-7.32$  to  $-2.35$ ;  $P = 0.002$ ) [32]. In the systematic review by Hosseini et al., including 23 randomized and non-randomized control studies, vitamin D supplementation was significantly associated with a reduced risk of ICU admission (RR = 0.35, 95 % CI: 0.20, 0.62) and mortality (RR = 0.46, 95 % CI:

0.30, 0.70), but it had no significant impact on the risk of COVID-19 infection [33]. Also in the study by Tentolouris et al., including a total of 11 studies with 22,265 COVID-19 patients, the findings implied that vitamin D supplementation improved COVID-19 outcomes: vitamin D supplementation was associated with a reduction in intensive care unit admission rate (OR 0.27; 95 % CI: 0.09–0.76,  $p = 0.010$ , I2 = 70 %, random-effect modelling); reduction of the need for mechanical ventilation (OR 0.34; 95 % CI: 0.16–0.72,  $p = 0.005$ , I2 = 61 %, random-effect modelling) and reduction of mortality from COVID-19 (OR 0.37; 95 % CI: 0.21–0.66,  $p < 0.001$ , I2 = 50 %, random-effect modelling) [34]. In the systematic review by Akbar et al., consisting of 14 studies comprising of 999,179 participants, when compared to the control group, COVID-19 patients had lower serum vitamin D (OR = 2.71 [1.72, 4.29],  $p < 0.001$ ; I2: 92.6 %). Higher rate of severe COVID-19 was observed in patients with low serum 25-OHD (OR = 1.90 [1.24, 2.93],  $p = 0.003$ ; I2: 55.3 %). Decreased serum vitamin D was associated with higher mortality (OR = 3.08 [1.35, 7.00],  $p = 0.011$ ; I2: 80.3 %), with a sensitivity of 85 % and specificity of 35 %. Male gender and diabetes were found to be factors in the connection between low serum 25-OHD and mortality (OR = 1.22 [1.08, 1.39],  $p = 0.002$  and OR = 0.88 [0.79, 0.98],  $p = 0.019$ , respectively) [35].

### 2.1. Vitamin D and COVID-19 in children

Pandemic-related restrictions have caused a significant decrease in vitamin D levels in children, in particular school-aged children and adolescents [36]. Despite, in the pediatric population, most cases of SARS-CoV-2 infection are asymptomatic or paucisymptomatic [37–39], even five months after discharge and beyond, children who have long-COVID might experience debilitating symptoms [40–42]. Many studies show that allergy and asthma, if under control, did not represent risk factors for the susceptibility to SARS-CoV-2 [43–46]. The role of spirometry, in the COVID era, still remains fundamental in the diagnosis and follow-up of respiratory diseases, in compliance with current containment measures [47].

The role of vitamin D in SARS-CoV-2 infection was also investigated in the pediatric population. In the systematic review by Shah et al., eight eligible studies were included: in infected pediatric patients, low levels of vitamin D increased the risk of severe disease (OR  $-5.5$ ; 95 % CI: 1.560–19.515;  $P = 0.008$ ). It was also found that children and adolescents having vitamin D deficiency had a greater risk of COVID infection as compared to patients with normal vitamin D levels [48]. In a study by Yilmaz et al., 40 children hospitalized with COVID-19 and 45 healthy controls were divided into two groups; those with vitamin D levels  $< 20$  ng/ml were classified as Group 1, and those with  $\geq 20$  ng/ml as Group 2. Children with COVID-19 had significantly lower vitamin D levels than the controls ( $p < 0.001$ ). The symptom of fever was significantly higher in COVID-19 patients who had deficient and insufficient vitamin D levels than in patients who had sufficient vitamin D levels ( $p = 0.038$ ) [49]. In the retrospective study by Bayramoğlu et al., consisting of 103 pediatric cases with COVID-19, low 25OH vitamin D levels were linked to higher inflammatory indicators (C-reactive Protein, procalcitonin, fibrinogen, D-dimer) and more severe clinical course [50]. In a Turkish report, including seventy-five 1–18 years old patients, the mean serum vitamin D level was significantly lower in the COVID-19 group than the control group ( $21.5 \pm 10.0$  vs.  $28.0 \pm 11.0$  IU,  $P < 0.001$ ). In comparison to the control group, the COVID-19 group had a considerably higher percentage of patients who were vitamin D deficient (44 % vs. 17.5 %,  $P < 0.001$ ). Patients with low levels of vitamin D were older than those with normal levels of vitamin D ( $11.6 \pm 4.9$  vs.  $6.2 \pm 1.8$  years,  $P = 0.016$ ). There was a significant male preponderance in the normal vitamin D group compared with the low vitamin D group (91.7 % vs. 50.8 %,  $P = 0.03$ ). C-reactive protein level was higher in the low vitamin D group, although the difference did not reach statistical significance ( $9.6 \pm 2.2$  vs.  $4.5 \pm 1.6$  mg/l,  $P = 0.074$ ) [51]. [Table1].

**Table 1**

Comparison of current evidence on vitamin D and COVID-19 in pediatric population, based on retrospective studies and a systematic review.

| Reference | Study design         | Objectives  | Population  | Methods  | Results  |
|-----------|----------------------|---|---|--|--|
| [48]      | systematic review    | To study the relationship between vitamin D level, risk and severity of Coronavirus disease of 2019 (COVID-19) infection in pediatric population                        | 8 studies involving children (age $\leq 18$ ) reporting vitamin D status and COVID-19 infection in pediatric patients.  | –  | In infected pediatric patients, low levels of vitamin D increased the risk of severe disease (odds ratio-5.5; 95 % CI: 1.560–19.515; $P = 0.008$ ). It was also found that children and adolescents with vitamin D deficiency had a greater risk of COVID infection than patients with normal vitamin D levels.  |
| [49]      | Retrospective study. | Vitamin D deficiency prevalence and the association between vitamin D deficiency and clinical and inflammatory markers in patients hospitalized for COVID-19 infection. | 85 children between the ages of 1 month to 18 years<br>40 COVID-19 + (101.76 $\pm$ 27.91 months range, 3 months-18 years)   | Vitamin D status was assessed retrospectively using accredited laboratory methods. Patients with 25-OH vitamin D levels below 20 ng/ml ( $< 50$ nmol/l) were considered to have vitamin D deficiency; those with 25-OH vitamin D levels between 21 and 29 ng/ml (52.5 and 72.5 nmol/l) were considered to have vitamin D insufficiency; and those with 25-OH vitamin D levels above 30 ng/ml were considered to have a normal vitamin D level. | 72.5 % of cases were vitamin D deficient or insufficient, and 2 patients in need of treatment in the ICU had the vitamin D level of below 10 ng/ml, and had comorbid diseases, but there were no reported cases of mortality. In the study, the distribution of disease severity according to vitamin D levels was not found significantly different ( $p = 0.097$ )   |
| [50]      | Retrospective study. | To evaluate the relationship between vitamin D levels and clinical severity and inflammation markers in children and adolescents with COVID-19.                         | 103 children. The mean age was 12.2 $\pm$ 4.92 (range 1–17) years and 52.4 % ( $n = 54$ ) were male.  | Data were assessed retrospectively. Patients were grouped according to their clinical severity (asymptomatic, mild, and moderate-to-severe) and vitamin D levels (25 OH vitamin D serum levels sufficient ( $> 20$ ng/ml), deficient (12–20 ng/ml), and insufficient ( $< 12$ ng/ml)).   | The serum vitamin D level was associated with clinical severity and markers of inflammation in children and adolescents with COVID-19. Interestingly, these associations were observed especially when there was a deficiency (i.e., 25 OH vit D) vitamin D insufficiency was not found to be associated with disease severity or inflammation markers in COVID-19   |
| [51]      | Retrospective study. | To compare the vitamin D levels of paediatric patients with mild/moderate coronavirus disease 2019 (COVID-19) disease and a healthy control group                       | 75 COVID-19 patients and 80 healthy controls. The mean age of the COVID-19 patients was 10.7 $\pm$ 5.5 years (range 1–18 years); 43 (57.3 %) patients were male<br>The control group was composed of 43 (53.8 %) males and 37 (46.2 %) females and had a mean age of 9.9 $\pm$ 4.6 years (range 1–17 years) | Vitamin D status was assessed retrospectively using accredited laboratory methods. Patients with 25-OH vitamin D levels below 20 ng/ml ( $< 50$ nmol/l) were considered to have vitamin D  | The mean serum vitamin D level was significantly lower in the COVID-19 group than the control group (21.5 $\pm$ 10.0 vs. 28.0 $\pm$ 11.0 IU, $P < 0.001$ ). In comparison to the control group, the COVID-19 group had a considerably higher percentage of patients who were vitamin D deficient (44 % vs. 17.5 %, $P < 0.001$ ). Patients with low levels of vitamin D were older than those with normal levels of vitamin D (11.6 $\pm$ 4.9 vs. 6.2 $\pm$ 1.8 years, $P = 0.016$ ). There was a significant male preponderance in the normal vitamin D group compared with the low vitamin D group (91.7 % vs. 50.8 %, |

(continued on next page)



Table 1 (continued)

| Reference | Study design | Objectives | Population | Methods   | Results   |
|-----------|--------------|------------|------------|---|---|
|           |              |            |            | deficiency; those with 25-OH vitamin D levels between 21 and 29 ng/ml (52.5 and 72.5 nmol/l) were considered to have vitamin D insufficiency; and those with 25-OH vitamin D levels above 30 ng/ml were considered to have a normal vitamin D level | P = 0.03). C-reactive protein level was higher in the low vitamin D group, although the difference did not reach statistical significance (9.6 ± 2.2 vs. 4.5 ± 1.6 mg/l, P = 0.074) |

Current evidence suggests that taking a vitamin D supplement to maintain a serum concentration of 25-OH vitamin D of at least 30 ng/ml (preferred range 40–60 ng/ml), can help reduce the risk of COVID-19 and its severe outcomes, including mortality [52]. According to evidence on the paediatric population, although robust systematic and meta-analyses are lacking, it can be recommended in any case to maintain adequate levels of vitamin D (>30 ng/ml) in order to maintain a healthy immune system.

### 3. Conclusion

In conclusion, although the examined studies were different in methods and statistical approach, most of them indicated a significant relation between vitamin D serum levels and SARS-CoV-2 infection, clinical outcomes and mortality. Even in children with COVID-19, insufficient levels of vitamin D have been found and low levels of vitamin D would seem to be associated with worse outcomes. While there is currently insufficient evidence to support the use of vitamin D as a therapeutic treatment for SARS-CoV-2 infection, those who may be deficient should raise their levels through diet, lifestyle changes, and supplementation because it is affordable and generally safe, in order to achieve any potential prophylactic benefits, but more significantly because preserving appropriate vitamin D levels is crucial for overall wellness.

### Financial disclosure statement

No funding was received.

### CRedit authorship contribution statement

M.M.d.G. wrote and conceptualized the manuscript. C.I. revised the manuscript. G.D. contributed to drafting the introduction. F.D. revised the manuscript. A.D. contributed to the research in the literature. A.K. supervised the manuscript. All authors have read and agreed to the published version of the manuscript.

### Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

### Data availability

No data was used for the research described in the article.

### Acknowledgements

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

### References

- [1] G. Saggese, F. Vierucci, F. Prodam, et al., Vitamin D in pediatric age: consensus of the Italian Pediatric Society and the Italian Society of Preventive and Social Pediatrics, jointly with the Italian Federation of Pediatricians, Ital. J. Pediatr. 44 (1) (2018) 51, <https://doi.org/10.1186/s13052-018-0488-7>.
- [2] M. Miraglia Del Giudice, C. Indolfi, C. Strisciuglio, Vitamin D: immunomodulatory aspects, J. Clin. Gastroenterol. 52 (Suppl 1) (2018), <https://doi.org/10.1097/MCG.0000000000001112>.
- [3] C. Almerighi, A. Sinistro, A. Cavazza, C. Ciapri, G. Rocchi, A. Bergamini, 1Alpha,25-dihydroxyvitamin D3 inhibits CD40L-induced pro-inflammatory and immunomodulatory activity in human monocytes, Cytokine 45 (3) (2009) 190–197, <https://doi.org/10.1016/j.cyto.2008.12.009>.
- [4] G. Penna, L. Adorini, 1 Alpha,25-dihydroxyvitamin D3 inhibits differentiation, maturation, activation, and survival of dendritic cells leading to impaired alloreactive T cell activation, J. Immunol. 164 (5) (2000) 2405–2411, <https://doi.org/10.4049/jimmunol.164.5.2405>.
- [5] A.G. van Halteren, E. van Etten, E.C. de Jong, R. Bouillon, B.O. Roep, C. Mathieu, Redirection of human autoreactive T-cells Upon interaction with dendritic cells modulated by TX527, an analog of 1,25 dihydroxyvitamin D(3), Diabetes 51 (7) (2002) 2119–2125, <https://doi.org/10.2337/diabetes.51.7.2119>.
- [6] E. Peelen, S. Knippenberg, A.H. Muris, M. Thewissen, J. Smolders, J.W. Tervaert, R. Hupperts, J. Damoiseaux, Effects of vitamin D on the peripheral adaptive immune system: a review, Autoimmun. Rev. 10 (12) (2011) 733–743, <https://doi.org/10.1016/j.autrev.2011.05.002>.
- [7] M.K. Arababadi, R. Nosratabadi, G. Asadikaram, Vitamin D and toll like receptors, Life Sci. 203 (2018) 105–111, <https://doi.org/10.1016/j.lfs.2018.03.040>.
- [8] S. Khanmohammadi, N. Rezaei, Role of Toll-like receptors in the pathogenesis of COVID-19, J. Med Virol. 93 (5) (2021) 2735–2739, <https://doi.org/10.1002/jmv.26826>.
- [9] S.-K. Chiu, K.-W. Tsai, C.-C. Wu, C.-M. Zheng, C.-H. Yang, W.-C. Hu, Y.-C. Hou, K.-C. Lu, Y.-C. Chao, Putative role of vitamin D for COVID-19 vaccination, Int. J. Mol. Sci. 22 (2021) 8988, <https://doi.org/10.3390/ijms22168988>.
- [10] G. Heine, K. Anton, B.M. Henz, M. Worm, 1alpha,25-dihydroxyvitamin D3 inhibits anti-CD40 plus IL-4-mediated IgE production in vitro, Eur. J. Immunol. 32 (12) (2002) 3395–3404, [https://doi.org/10.1002/1521-4141\(200212\)32:12<3395::AID-IMMU3395>3.0.CO;2-#](https://doi.org/10.1002/1521-4141(200212)32:12<3395::AID-IMMU3395>3.0.CO;2-#).
- [11] D. Manousaki, L. Paternoster, M. Standl, M.F. Moffatt, M. Farrall, E. Bouzigon, D. P. Strachan, F. Demeis, M. Lathrop, W.O.C.M. Cookson, J.B. Richards, Vitamin D levels and susceptibility to asthma, elevated immunoglobulin E levels, and atopic dermatitis: a Mendelian randomization study, PLoS Med 14 (5) (2017), e1002294, <https://doi.org/10.1371/journal.pmed.1002294>.
- [12] M. Miraglia Del Giudice, A. Allegorico, The role of vitamin D in allergic diseases in children, J. Clin. Gastroenterol. 50 (Suppl 2) (2016), <https://doi.org/10.1097/MCG.0000000000000679>.
- [13] G. Saggese, F. Vierucci, F. Prodam, F. Cardinale, I. Cetin, E. Chiappini, G.L. De' Angelis, M. Massari, E. Miraglia Del Giudice, M. Miraglia Del Giudice, D. Peroni, L. Terracciano, R. Agostiniani, D. Careddu, D.G. Ghigloni, G. Bona, G. Di Mauro, G. Corsello, Vitamin D in pediatric age: consensus of the Italian Pediatric Society and the Italian Society of Preventive and Social Pediatrics, jointly with the Italian Federation of Pediatricians, Ital. J. Pediatr. 44 (1) (2018) 51, <https://doi.org/10.1186/s13052-018-0488-7>.
- [14] M. Miraglia Del Giudice, A. Allegorico, A. Grandone, F. Galdo, C. Cuppari, M. Capasso, N. Maiello, Vitamin D and bronchial inflammation in asthmatic children, J. Biol. Regul. Homeost. Agents 29 (2 Suppl 1) (2015) 125–129.

- [15] M. Miraglia Del Giudice, N. Maiello, A. Allegorico, L. Iavarazzo, M. Capasso, C. Capristo, G. Ciprandi, *Lactobacillus reuteri* DSM 17938 plus vitamin D3 as ancillary treatment in allergic children with asthma, *Ann. Allergy Asthma Immunol.* 117 (6) (2016) 710–712, <https://doi.org/10.1016/j.anai.2016.09.004>.
- [16] S. Beyerstedt, E.B. Casaro, É.B. Rangel, COVID-19: angiotensin-converting enzyme 2 (ACE2) expression and tissue susceptibility to SARS-CoV-2 infection, *Eur. J. Clin. Microbiol. Infect. Dis.* 40 (5) (2021) 905–919, <https://doi.org/10.1007/s10096-020-04138-6>.
- [17] R. Kumar, H. Rathi, A. Haq, S.J. Wimalawansa, A. Sharma, Putative roles of vitamin D in modulating immune response and immunopathology associated with COVID-19, *Virus Res.* 292 (2021), 198235, <https://doi.org/10.1016/j.virusres.2020.198235>.
- [18] A. Hosseini, V. Hashemi, N. Shomali, F. Asghari, T. Gharibi, M. Akbari, S. Gholizadeh, A. Jafari, Innate and adaptive immune responses against coronavirus, *Biomed. Pharmacother.* 132 (2020), 110859, <https://doi.org/10.1016/j.biopha.2020.110859>.
- [19] N. Bayraktar, H. Turan, M. Bayraktar, A. Ozturk, H. Erdogdu, Analysis of serum cytokine and protective vitamin D levels in severe cases of COVID-19, *J. Med. Virol.* 94 (1) (2022) 154–160, <https://doi.org/10.1002/jmv.27294>.
- [20] F. Filippatos, E.B. Tatsi, A. Michos, Immune response to SARS-CoV-2 in children: a review of the current knowledge, *Pediatr. Investig.* 5 (3) (2021), e12283, <https://doi.org/10.1002/ped4.12283>.
- [21] P. Zimmermann, N. Curtis, Why is COVID-19 less severe in children? A review of the proposed mechanisms underlying the age-related difference in severity of SARS-CoV-2 infections, *Arch. Dis. Child.* (2020), <https://doi.org/10.1136/archdischild-2020-320338>.
- [22] D. Rawat, A. Roy, S. Maitra, V. Shankar, P. Khanna, D.K. Baidya, "Vitamin D supplementation and COVID-19 treatment: a systematic review and meta-analysis", *Diabetes Metab. Syndr.* 15 (4) (2021), 102189, <https://doi.org/10.1016/j.dsx.2021.102189>.
- [23] M. Pereira, A. Dantas Damascena, L.M. Galvão Azevedo, T. de Almeida Oliveira, J. da Mota Santana, Vitamin D deficiency aggravates COVID-19: systematic review and meta-analysis, *Crit. Rev. Food Sci. Nutr.* 62 (5) (2022) 1308–1316, <https://doi.org/10.1080/10408398.2020.1841090>.
- [24] R. Ghasemian, A. Shamshirian, K. Heydari, M. Malekan, R. Alizadeh-Navaei, M. A. Ebrahimzadeh, M. Ebrahimi Warkiani, H. Jafarpour, S. Razavi Bazaz, A. Rezaei Shahmirzadi, M. Khodabandeh, B. Seyfari, A. Motamedzadeh, E. Dadgostar, M. Aalinezhad, M. Sedaghat, N. Razzaghi, B. Zarandi, A. Asadi, V. Yaghoubi Naei, R. Beheshti, A. Hessami, S. Azizi, A.R. Mohseni, D. Shamshirian, The role of vitamin D in the age of COVID-19: a systematic review and meta-analysis, *Int. J. Clin. Pract.* 75 (11) (2021), e14675, <https://doi.org/10.1111/ijcp.14675>.
- [25] A. Kazemi, V. Mohammadi, S.K. Aghababae, M. Golzarand, C.C.T. Clark, S. Babajafari, Association of Vitamin D Status with SARS-CoV-2 Infection or COVID-19 Severity: a Systematic Review and Meta-analysis [published correction appears in *Adv Nutr.* 2021 Oct 1;12(5):2040-2044], *Adv. Nutr.* 12 (5) (2021) 1636–1658, <https://doi.org/10.1093/advances/nmab012>.
- [26] F. Petrelli, A. Luciani, G. Perego, G. Dognini, P.L. Colombelli, A. Ghidini, Therapeutic and prognostic role of vitamin D for COVID-19 infection: a systematic review and meta-analysis of 43 observational studies, *J. Steroid Biochem. Mol. Biol.* 211 (2021), 105883, <https://doi.org/10.1016/j.jsbmb.2021.105883>.
- [27] R. Pal, M. Banerjee, S.K. Bhadada, A.J. Shetty, B. Singh, A. Vyas, Vitamin D supplementation and clinical outcomes in COVID-19: a systematic review and meta-analysis, *J. Endocrinol. Investig.* 45 (1) (2022) 53–68, <https://doi.org/10.1007/s40618-021-01614-4>.
- [28] K. Shah, P.V. V. U. Sharma, D. Mavalankar, Does vitamin D supplementation reduce COVID-19 severity? - a systematic review, *QJM* (2022) hcac040, <https://doi.org/10.1093/qjmed/hcac040>.
- [29] J. Oristrell, J.C. Oliva, E. Casado, et al., Vitamin D supplementation and COVID-19 risk: a population-based, cohort study, *J. Endocrinol. Investig.* 45 (1) (2022) 167–179, <https://doi.org/10.1007/s40618-021-01639-9>.
- [30] Z. Wang, A. Joshi, K. Leopold, et al., Association of vitamin D deficiency with COVID-19 infection severity: systematic review and meta-analysis, *Clin. Endocrinol.* 96 (3) (2022) 281–287, <https://doi.org/10.1111/cen.14540>.
- [31] A. Teshome, A. Adane, B. Girma, Z.A. Mekonnen, The impact of vitamin D level on COVID-19 infection: systematic review and meta-analysis, *Front Public Health* 9 (2021), 624559, <https://doi.org/10.3389/fpubh.2021.624559>.
- [32] H.A. Dissanayake, N.L. de Silva, M. Sumanatilleke, et al., Prognostic and therapeutic role of vitamin D in COVID-19: systematic review and meta-analysis, *J. Clin. Endocrinol. Metab.* 107 (5) (2022) 1484–1502, <https://doi.org/10.1210/clinem/dgab892>.
- [33] B. Hosseini, A. El Abd, F.M. Ducharme, Effects of vitamin D supplementation on COVID-19 related outcomes: a systematic review and meta-analysis, *Nutrients* 14 (10) (2022) 2134, <https://doi.org/10.3390/nu14102134>.
- [34] N. Tentolouris, G. Samakidou, I. Eleftheriadou, A. Tentolouris, E.B. Jude, The effect of vitamin D supplementation on mortality and intensive care unit admission of COVID-19 patients. A systematic review, meta-analysis and meta-regression, *Diabetes Metab. Res. Rev.* 38 (4) (2022), e3517, <https://doi.org/10.1002/dmrr.3517>.
- [35] M.R. Akbar, A. Wibowo, R. Pranata, B. Setiabudiawan, Low Serum 25-hydroxyvitamin D (Vitamin D) Level Is Associated With Susceptibility to COVID-19, Severity, and Mortality: a systematic review and meta-analysis [published correction appears in *Front Nutr.* 2021 Sep 27;8:754539], *Front Nutr.* 8 (2021), 660420, <https://doi.org/10.3389/fnut.2021.660420>.
- [36] G. Beyazgül, Ö. Bağ, İ. Yurtseven, F. Coşkunol, S. Başer, D. Çiçek, G.İ. Kanberoglu, F. Çelik, Ö. Nalbantoğlu, B. Özkan, How did vitamin D levels of children change during COVID-19 pandemic: a comparison of pre-pandemic/pandemic periods due to seasonal differences, *J. Clin. Res. Pediatr. Endocrinol.* (2022), <https://doi.org/10.4274/jcrpe.galenos.2022.2021-10-6>.
- [37] G.F. Parisi, C. Indolfi, F. Decimo, S. Leonardi, M. Miraglia Del Giudice, COVID-19 pneumonia in children: from etiology to management, *Front Pediatr.* 8 (2020), 616622, <https://doi.org/10.3389/fped.2020.616622>.
- [38] G.F. Parisi, G. Brindisi, C. Indolfi, L. Diaferio, G. Marchese, D.G. Ghiglioni, A. M. Zicari, M. Miraglia Del Giudice, Upper airway involvement in pediatric COVID-19, *Pediatr. Allergy Immunol.* 31 (Suppl 26) (2020) 85–88, <https://doi.org/10.1111/pai.13356>.
- [39] G.F. Parisi, G. Brindisi, C. Indolfi, L. Diaferio, G. Marchese, D.G. Ghiglioni, A. M. Zicari, M. Miraglia Del Giudice, COVID-19, anosmia, and ageusia in atopic children, *Pediatr. Allergy Immunol.* 33 (Suppl 27) (2022) 99–101, <https://doi.org/10.1111/pai.13644>.
- [40] G.F. Parisi, L. Diaferio, G. Brindisi, C. Indolfi, G.R. Umamo, A. Klain, G. Marchese, D.G. Ghiglioni, A.M. Zicari, G.L. Marseglia, M. Miraglia Del Giudice, Cross-sectional survey on long term sequelae of pediatric COVID-19 among Italian pediatricians, *Children* 8 (9) (2021) 769, <https://doi.org/10.3390/children8090769>.
- [41] R. Lordan, S. Prior, E. Hennessy, et al., Considerations for the safe operation of schools during the coronavirus pandemic, *Front. Public Health* 9 (2021), 751451, <https://doi.org/10.3389/fpubh.2021.751451>.
- [42] I.M. Osmanov, E. Spiridonova, P. Bobkova, A. Gamirova, A. Shikhaleva, M. Andreeva, et al., Risk factors for long covid in previously hospitalized children using the ISARIC global follow-up protocol: a prospective cohort study, *Eur. Respir. J.* 210 (2021) 1341, <https://doi.org/10.1183/13993003.01341-2021>.
- [43] G. Brindisi, A.M. Zicari, G.F. Parisi, L. Diaferio, C. Indolfi, G. Marchese, D. G. Ghiglioni, G.R. Umamo, A. Klain, G.L. Marseglia, M.M. Del Giudice, Prevalence of COVID-19 in children affected by allergic rhinoconjunctivitis and asthma: results from the second "SIAP rhinosinusitis and conjunctivitis committee" survey, *Ital. J. Pediatr.* 48 (1) (2022) 1, <https://doi.org/10.1186/s13052-021-01198-y>.
- [44] M. De Filippo, M. Votto, I. Brambilla, R. Castagnoli, L. Montagna, C. Caffarelli, F. Cardinale, M. Miraglia Del Giudice, M. Tosca, S. Caimmi, A. Licari, G. L. Marseglia, Allergy and COVID-19, *Acta Biomed.* 92 (S7) (2021), e2021522, <https://doi.org/10.23750/abm.v92iS7.12402>.
- [45] L. Indinnimeo, E. Chiappini, M. Miraglia Del Giudice, Italian Panel for the management of acute asthma attack in children Roberto Bernardini, Guideline on management of the acute asthma attack in children by Italian Society of Pediatrics, *Ital. J. Pediatr.* 44 (1) (2018) 46, <https://doi.org/10.1186/s13052-018-0481-1>.
- [46] M.E. Di Cicco, M. Leone, M. Scavone, M. Miraglia Del Giudice, A. Licari, M. Duse, I. Brambilla, G. Ciprandi, C. Caffarelli, M. Tosca, Intermittent and mild persistent asthma: how therapy has changed, *Acta Biomed.* 92 (S7) (2021), e2021523, <https://doi.org/10.23750/abm.v92iS7.12422>.
- [47] A. Klain, C. Indolfi, G. Dinardo, F. Decimo, M. Miraglia Del Giudice, Covid-19 and spirometry in this age, *Ital. J. Pediatr.* 48 (1) (2022) 11, <https://doi.org/10.1186/s13052-022-01199-5>.
- [48] K. Shah, V.P. Varna, A. Pandya, D. Saxena, Low vitamin D levels and prognosis in a COVID-19 pediatric population: a systematic review, *QJM* 114 (7) (2021) 447–453, <https://doi.org/10.1093/qjmed/hcab202>.
- [49] K. Yilmaz, V. Şen, Is vitamin D deficiency a risk factor for COVID-19 in children? *Pediatr. Pulmonol.* 55 (12) (2020) 3595–3601, <https://doi.org/10.1002/ppul.25106>.
- [50] E. Bayramoğlu, G. Akkoç, A. Ağbaş, et al., The association between vitamin D levels and the clinical severity and inflammation markers in pediatric COVID-19 patients: single-center experience from a pandemic hospital, *Eur. J. Pediatr.* 180 (8) (2021) 2699–2705, <https://doi.org/10.1007/s00431-021-04030-1>.
- [51] A. Alpcan, S. Tursun, Y. Kandur, Vitamin D levels in children with COVID-19: a report from Turkey, *Epidemiol. Infect.* 149 (2021), e180, <https://doi.org/10.1017/S0950268821001825>.
- [52] J.H. Bae, H.J. Choe, M.F. Holick, S. Lim, Association of vitamin D status with COVID-19 and its severity: Vitamin D and COVID-19: a narrative review, *Rev. Endocr. Metab. Disord.* 23 (3) (2022) 579–599, <https://doi.org/10.1007/s11154-021-09705-6>.