



POSTER PRESENTATION

Open Access

HERV-W/MRSV envelope transcripts detection in blood of multiple sclerosis patients after Natalizumab treatment

Luiz HS Nali¹, Guilherme S Olival², Augusto CP de Oliveira³, Jorge Casseb⁴, Jose E Vidal³, Lenira Moraes⁵, Maria CD Fink¹, Laura M Sumita¹, Camila M Romano^{1*}

From 16th International Conference on Human Retroviruses: HTLV and Related Viruses
Montreal, Canada. 26-30 June 2013

Multiple sclerosis (MS) is an inflammatory disorder of the central nervous system (CNS) and is the most common cause of neurological disability in young adults. MS-Associated Retrovirus (MSRV) is member of Human Endogenous Retroviruses W family, and their increased activity in MS patients is associated to the disease immunopathogenesis. Natalizumab, an antibody-based therapy, hinders migration of T cells into the CNS and is currently the most potent treatment for MS. Although Natalizumab interferes with gene expression relevant for function and differentiation of lymphocytes, its effects on genes involved in immunopathogenesis are unknown. Here, we report the effect of different treatments on the HERV-W/MRSV expression in patients with relapsing-remitting MS. MRSV transcripts were quantified by qRT-PCR in peripheral blood mononuclear cells of 9 patients receiving Natalizumab for at least 6 months (MSNat group) and 11 patients under immunosuppressive treatments (MSI group). The mean age was 28 years (18-35) for MSNat group and 44 (28-54) for MSI. The mean Expanded Disability Status Scale (EDSS) score was 4 (2-6) and 2.6 (1-6.5) for MSNat and MSI groups respectively. MSRV transcripts level was slightly higher in MSNat group, although not significantly, suggesting that Natalizumab does not interfere on HERV-W expression. Patients included in Natalizumab protocol usually do not respond to other treatments and present higher EDSSs. Possibly, EDSS and age have more impact in retroelements activity, as already demonstrated. This is the first comparison of HERV/MSRV expression between

different therapy groups in MS, and other studies are needed to confirm such findings.

Funding

This work was supported by FAPESP 2010/10619-0.

Authors' details

¹Departamento de Moléstias Infecciosas e Parasitárias (LIMHC), Instituto de Medicina Tropical de São Paulo, Universidade de São Paulo, São Paulo, SP, Brazil. ²Irmãmandade da Santa Casa de Misericórdia de São Paulo, São Paulo, SP, Brazil. ³Instituto de Infectologia Emílio Ribas, São Paulo, SP, Brazil.

⁴Laboratório de Imunodeficiências e Dermatologia, Instituto de Medicina Tropical de São Paulo e Faculdade de Medicina, Universidade de São Paulo, São Paulo, SP, Brazil. ⁵Hospital das Clínicas da Faculdade de Medicina, Universidade de São Paulo, São Paulo, SP, Brazil.

Published: 7 January 2014

doi:10.1186/1742-4690-11-S1-P131

Cite this article as: Nali et al.: HERV-W/MRSV envelope transcripts detection in blood of multiple sclerosis patients after Natalizumab treatment. *Retrovirology* 2014 **11**(Suppl 1):P131.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



* Correspondence: CMR-cmromano@usp.br

¹Departamento de Moléstias Infecciosas e Parasitárias (LIMHC), Instituto de Medicina Tropical de São Paulo, Universidade de São Paulo, São Paulo, SP, Brazil

Full list of author information is available at the end of the article