

Differential induction of total IgE by two *Salmonella enterica* serotypes

Zhanna A. Ktsoyan¹, Mkhitar S. Mkrtychyan¹, Magdalina K. Zakharyan¹,
Armine A. Mnatsakanyan², Karine A. Arakelova¹, Zaruhi U. Gevorgyan²,
Lusntag A. Ktsoyan³, Anahit İ. Sedrakyan¹, Alvard I. Hovhannisyan¹,
Karine A. Ghazaryan¹, Anna S. Boyajyan¹ and Rustam I. Aminov^{4*}

¹ Institute of Molecular Biology of National Academy of Sciences of Republic of Armenia, Yerevan, Armenia, ² "Nork" Clinical Hospital of Infectious Diseases, Ministry of Health of the Republic of Armenia, Yerevan, Armenia, ³ Yerevan State Medical University after Mkhitar Heratsi, Yerevan, Armenia, ⁴ National Veterinary Institute, Technical University of Denmark, Frederiksberg, Denmark

The main goal of this study was to establish how the inflammation caused by infection with two different *Salmonella enterica* serotypes, *S. Typhimurium* and *S. Enteritidis*, may lead to the predisposition to allergy as measured by total IgE level in the blood. Infection by *S. Typhimurium* did not affect the systemic IgE concentration while in *S. Enteritidis*-infected patients there was a significant 3.5-fold increase. This effect was especially profound in patients >4 years old, with up to the 8-fold increase above the norm. The degree of dysbiosis in these two infections measured with the comparative counts of cultivated bacteria showed an inverse relationship with the IgE concentration. Earlier we reported the elevated level of IL-17 in patients infected by *S. Enteritidis*. In the current study a significant correlation was found between the concentrations of IL-17 and IgE suggesting a possible role played by this cytokine in triggering the production of IgE in response to *S. Enteritidis* infection.

Keywords: *Salmonella*, salmonellosis, serotype, IgE, IL-17

OPEN ACCESS

Edited by:

Brice Rotureau,
Institut Pasteur, France

Reviewed by:

Olivier Disson,
Institut National de la Santé et de la
Recherche Médicale, France
Brice Rotureau,
Institut Pasteur, France

*Correspondence:

Rustam I. Aminov,
National Veterinary Institute, Technical
University of Denmark, Bülowsvej 27,
1870 Frederiksberg C, Denmark
rusam@vet.dtu.dk

Received: 18 December 2014

Accepted: 02 May 2015

Published: 26 May 2015

Citation:

Ktsoyan ZA, Mkrtychyan MS,
Zakharyan MK, Mnatsakanyan AA,
Arakelova KA, Gevorgyan ZU, Ktsoyan
LA, Sedrakyan AI, Hovhannisyan AI,
Ghazaryan KA, Boyajyan AS and
Aminov RI (2015) Differential induction
of total IgE by two *Salmonella enterica*
serotypes.
Front. Cell. Infect. Microbiol. 5:43.
doi: 10.3389/fcimb.2015.00043

Introduction

Intestinal infectious diseases remain a serious problem for the health care systems. In many countries the most frequent intestinal infection is salmonellosis, which is caused by various serotypes of non-typhoidal *Salmonella* (Rabsch et al., 2001). Except for norovirus infections, *Salmonella* is a single most common causative agent among food-borne diseases (Centers for Disease Control and Prevention, 2013). Gastroenteritis is most frequently caused by *Salmonella enterica* serovar Typhimurium (*S. Typhimurium*) and *S. enterica* serovar Enteritidis (*S. Enteritidis*), which are also prevalent in Armenia (Ktsoyan et al., 2013). The former serovar has been predominantly reported before the 1980's but beginning from the 1990's the latter becomes the predominant serovar isolated from the salmonellosis cases (Angulo and Swerdlow, 1999; Munro et al., 1999; Tschape et al., 1999).

At the genomic level, *S. enterica* serovars are very close, with a large and stable core genome, while the accessory genome is dominated by mobile genetic elements such as phages, prophages, genomic islands, transposons and plasmids (Thomson et al., 2008; Jacobsen et al., 2011). Interestingly, a particular gene content often correlates with a serotype of *S. enterica* thus providing a genotypic background for the earlier circumstantial observations on the dependence of clinical picture of salmonellosis on the serotype (Jones et al., 2008).

The pairwise comparison of *S. Typhimurium* and *S. Enteritidis* similarly emphasizes the role of mobile genetic elements in genomic differences between the two. In particular, this is the presence of genes encoded by bacteriophages Fels-1, Fels-2, Gifsy-1, and Gifsy-2, as well as by two other phage-related regions in the chromosome of *S. Enteritidis* (Olson et al., 2007). Other differences include the lack SG11 in *S. Enteritidis* as well as the carriage of dissimilar virulence plasmids by these two serovars. Despite a better understanding of genomic differences between the two serotypes, it remains unclear what factors contributed to the emergence of *S. Enteritidis* as a highly successful pathogen in the 1990's (Van Immerseel, 2010).

The normal microbiota plays a critical role in the establishment and maintenance of colonization resistance and proper immune responses. The impact on the intestinal microbiota in childhood is believed to be a major risk factor for chronic immune diseases such as allergy and autoimmune diseases in later life (Licciardi et al., 2010). During infection, *Salmonella* induces a strong pro-inflammatory response, which affects the normal intestinal microbiota and allows *Salmonella* to gain a competitive advantage (Stecher et al., 2007). The clearance of *Salmonella* infection, however, is not the end of the story. Gastroenteritis caused by *S. Enteritidis* during childhood, for example, is a risk factor for irritable bowel syndrome in adulthood (Cremon et al., 2014).

The etiology of allergic diseases remains uncertain, and many factors have been implicated as contributing to their development. One of the pioneering works in this area has been the hygiene hypothesis proposed by David Strachan (Strachan, 1989). The hypothesis implicated an insufficient microbial exposure in modern life that may result in the increasing number of allergic diseases. More recent works also suggested a major role played by commensal microbiota in the development of atopic disorders (Penders et al., 2007a,b). If the structure-and-function of commensal microbiota is compromised, this in turn may affect the proper functioning of the immune system and colonization resistance (Prakash et al., 2011). Because of the complexity of commensal microbiota, the molecular mechanisms, by which the microbiota as a whole exerts beneficial or detrimental effects, remain largely unknown (Sekirov et al., 2010).

The link between allergic disorders and infectious diseases, bacterial, and viral, remain contradictory. There are indications that early viral or bacterial exposure could be beneficial to inhibit excessive allergic Th2 reactions by inclining the immune system toward Th1 responses (Herz et al., 2000). It is known, for example, that attenuated *Salmonella* may reduce the induced airway inflammation and Th2 responses (Wu et al., 2006). At the same time, however, infections may worsen atopic conditions. For a long time the cause of allergic diseases has been predominantly discussed within the Th1/Th2 framework. Recently, however, a number of studies have implicated the Th17 pathway in asthma and allergic diseases (Oboki et al., 2008; Cosmi et al., 2011; Newcomb and Peebles, 2013; Manni et al., 2014; Naji et al., 2014). In summary, the prevailing views are largely within the context of alterations in the microbiome that interfere with immune system maturation (Feehley et al., 2012). The compounding factors for alterations are multiple and

include decreased exposure to infections due to vaccination and sanitation, mode of delivery, antibiotic use, infant formulae, and diet.

Previously, we have reported that infection by two serotypes of *S. enterica* results in a serotype-specific host response, with the induction of a specific set of cytokines and antimicrobial protein calprotectin (Ktsoyan et al., 2013). In particular, the concentration of IL-1 β was significantly higher in patients infected with *S. Typhimurium* while that of IL-17 was higher in patients infected with *S. Enteritidis*. On the basis of our previous findings, the main aim of this study was to identify factors that may play a role in the predisposition to allergy following infection by the two *S. enterica* serotypes.

The hallmark of allergic inflammatory response is the binding of IgE to an allergen (De Amici and Ciprandi, 2013). The level of systemic IgE is considered as a typical biomarker for allergies. Moreover, the level of total IgE may indicate the likelihood of sensitization even in patients with negative specific allergy tests (Kerkhof et al., 2003).

Materials and Methods

The study cohorts included patients with salmonellosis admitted to the infectious disease hospital Nork in Yerevan, RA. A total of 56 patients with acute salmonellosis caused by *S. Typhimurium* ($n = 21$) or *S. Enteritidis* ($n = 35$) and 18 healthy subjects were enrolled in this study. The effect of age was investigated by separating data from the above three cohorts into two age groups: (i) less than 4 years old (14 patients infected with *S. Typhimurium*, 15—with *S. Enteritidis* and 7 healthy subjects) and (ii) more than 4 years old (7 patients infected with *S. Typhimurium*, 20 with *S. Enteritidis* and 11 healthy subjects). Diagnosis was based on clinical presentation and laboratory analyses. Clinical picture consistent with gastroenteritis were: diarrhea, fever, nausea, vomiting, and abdominal cramps. Anamneses of the food consumed, water sources, social gatherings, any contact with a similar illness, and recent travels were also recorded. The patients or guardians reported no previous allergies or autoimmune diseases.

The patients selected for the study were not taking any type of medication, including antibiotics, before the hospital admission. Blood and fecal samples were taken on the first or second day of admission to the hospital. At the time of discharge from the hospital, no presence of *Salmonella* has been detected in the fecal samples or blood of any of the patients. For detoxification and rehydration, all patients were receiving the standard infusion therapy.

All study subjects (or parents or guardians if a child) gave their written consent to give fecal and blood samples for the study. The study protocol was approved by the Ethics Committee of the Institute of Molecular Biology NAS RA (IORG number 0003427, Assurance number FWA00015042, and IRB number 00004079).

The composition of intestinal microbiota was evaluated by standard microbiological methods and selective media using the fecal samples collected from patients with acute salmonellosis caused by *S. Typhimurium* or *S. Enteritidis*. The term “dysbiosis”

is not an universally accepted clinical definition but is rather broadly defined as "...any change to the composition of resident commensal communities relative to the community found in healthy individuals" (Petersen and Round, 2014). We examined the degree of dysbiosis by counting the increase or decrease of certain fecal bacterial groups in salmonellosis compared to healthy control subjects. The degree of dysbiosis was determined on a 0–4 scale, ranging from 0 (normal) to 4 (severe dysbiosis) according to the previously proposed criteria (Mitrokhin et al., 1998; Tabolin et al., 1998).

Biochemical assays for identification of *Salmonella* were: fermentation of glucose, negative urease reaction, lysine decarboxylase, negative indole test, H₂S production and fermentation of galactitol (dulcitol). Serotypes of *Salmonella* were determined using the standard Kauffman-White scheme with the use of commercially available polyvalent antisera for flagellar (H) and lipopolysaccharide (O) antigens.

The concentration of IgE in the sera of patients with acute salmonellosis due to *S. Typhimurium* or *S. Enteritidis* infection was used to estimate the predisposition to allergy. The IgE concentration was measured using the ECL technology. The corresponding concentrations were determined with the use of the immunoassay analyzer cobas e 411 (Roche, USA) according to the manufacturer's protocols. The concentration of IL-17 was measured as described before (Ktsoyan et al., 2013).

GraphPad Prism 5 (GraphPad Software, USA) was used to perform the Mann–Whitney *U*-test to determine the statistical significance of differences among the groups studied. The *p*-values < 0.05 were considered statistically significant. Discriminant function analysis was carried out with IBM SPSS Statistics 19 (IBM, USA).

Results

Systemic IgE in *S. Typhimurium* and *S. Enteritidis* Infections

We found a significant 3.5-fold increase of systemic IgE in patients of both age groups infected by *S. Enteritidis* compared to control ($p < 0.0001$), while in patients of both age groups infected by *S. Typhimurium* the level was approximately the same as in the control group ($p = 0.12$) (Figure 1). The level of IgE was 6-fold higher in patients of both age groups infected with *S. Enteritidis* compared to patients of both age groups infected with *S. Typhimurium* ($p < 0.0001$) (Figure 1). Compared to the healthy cohorts, *S. Enteritidis*-infected patients of the younger age had the IgE concentration 2-fold higher ($p = 0.2$, statistically not significant), while the elder patients with *S. Enteritidis* infection—8-fold higher ($p < 0.0001$). We observed a higher variability of IgE concentration in patients less than 4 years old infected by *S. Enteritidis*, which possibly compromised the statistical significance (Figure 1). In the elder age group the difference was consistent and significant, with the 86% of cases having the concentration of IgE higher than in healthy controls.

Dysbiosis Due to *S. Typhimurium* and *S. Enteritidis* Infections

In the search for factors contributing to the elevated production of IgE due to *S. Enteritidis* infection we investigated a possible role of dysbiosis induced by these two serotypes of *S. enterica* (Table 1). Changes in the qualitative and quantitative composition of the fecal microbiota during the acute disease stage were investigated using the standard cultivation-based techniques with selective media. In general,




FIGURE 1 | Systemic IgE concentration in patients of different age and infected by different serotypes of *S. enterica* (median with interquartile range). Total: all patients infected by *S. Typhimurium* or *S. Enteritidis*, irrespective of age; control and either *S. Enteritidis*- or *S. Typhimurium*-infected. The second and

third columns: patients are separated according to age and *S. enterica* serotypes. *Significant difference compared to control ($p \leq 0.0002$, Mann–Whitney *U*-test). **Significant difference between *S. Enteritidis*- and *S. Typhimurium*-infected patients ($p \leq 0.0002$, Mann–Whitney *U*-test).

TABLE 1 | Dysbiotic changes due to *S. Typhimurium* and *S. Enteritidis* infections.

Microbial groups	<i>S. Typhimurium</i> (% of patients ^a)	<i>S. Enteritidis</i> (% of patients)
1 Total <i>Salmonella</i> spp. count	↑ ^x 83.0*	↑ ^x 37.5*
2 Commensal <i>E. coli</i>	↓ ^y 55.0*	↓ ^y 25.5*
3 <i>E. coli</i> with low β-galactosidase activity	0	↑ ^x 10.5
4 Lactose-negative <i>Enterobacteriaceae</i>	↑ ^x 76.6*	↑ ^x 22.5*
5 Hemolytic <i>E. coli</i>	↑ ^z 30	↑ ^z 22.5*
6 Cocci among the total bacterial count (%)	↑ ^x 43.3	↑ ^x 47.5
7 Hemolytic <i>Staphylococcus</i> spp.	0	0
8 <i>Bifidobacterium</i> spp.	↓ ^w 80.7*	↓ ^w 52.5*
9 <i>Lactobacillus</i> spp.	↓ ^w 81.7*	↓ ^w 70.0*
10 <i>Enterococcus</i> spp.	↑ ^w 30.7	↑ ^w 28.0
11 <i>Proteus</i> spp.	0	0
12 <i>Staphylococcus aureus</i>	0	0
13 <i>Candida</i> spp.	↑ ^w 43.3*	↑ ^w 30*
14 Anaerobic cocci	0	0

^aPercentage of patients that have the increased (↑) or decreased (↓) count of the corresponding microbial groups compared to control.

*Statistically significant difference between the infected and control groups ($p < 0.05$).

^yDifference between the infected and control groups is ≥ 2 -fold.

^xAbsence in the infected group.

^zPresence in the infected group.

^wDifference between the infected and control groups is ≥ 100 -fold.

the *Salmonella*-induced dysbiosis was characterized by a sharp decline in the numbers of *Bifidobacterium* spp., *Lactobacillus* spp., and *Escherichia coli* (Table 1). Also a quantitative increase in numbers of lactose-negative bacteria, *Enterococcus* spp., *Candida* spp., and enteric bacteria with pathogenic potential was observed in fecal samples of *Salmonella*-infected patients compared to control. *E. coli* strains with low β-galactosidase activity were detected only in *S. Enteritidis*-infected patients while the increase in the lactose-negative *Enterobacteriaceae* was more pronounced in patients infected with *S. Typhimurium*. The departure from the normal composition of gut microbiota was more pronounced in patients infected with *S. Typhimurium* compared to *S. Enteritidis*-infected. The degree of dysbiosis in the former group was significantly higher than in the latter (2.75 vs. 1.0, $p = 0.005$; Figure 2).

Further intergroup and intragroup comparative analyses revealed that the degree of dysbiosis was most pronounced in the age group < 4 years old, with a higher average degree of dysbiosis in patients infected by *S. Typhimurium* compared to *S. Enteritidis*-infected (3.5 vs. 1.5, $p = 0.03$). In the age group >4 years old the difference was not significant (Figure 2). The intra-group analysis revealed a significant difference in the degree of dysbiosis between the <4 and >4 years old groups in the case of *S. Enteritidis* infections (1.5 vs. 1.0, $p = 0.01$) while in the case of *S. Typhimurium* infections the difference was not statistically significant (3.5 vs. 1.75, $p = 0.22$) (Figure 2). Thus, the comparative analyses demonstrated significant differences

in the dysbiotic changes depending on the age of patients and the serotype of *S. enterica* that caused salmonellosis. The degree of dysbiosis in *S. Enteritidis*-infected patients, however, was significantly lower compared to *S. Typhimurium* infection. The cumulative dysbiotic score, therefore, could not be considered as a main factor contributing to the increased production of IgE in *S. Enteritidis*-infected patients.

Multivariate Discriminant Function Analyses

To get into the finer level of resolution and establish the role of other possible factors involved, the raw data were further subjected to multivariate analyses to identify the relationships between more variables and their relevance to the correct classification of subjects in the study. In particular, we used discriminant function analyses (DA) with the aim to determine the distribution of patterns from the complex datasets obtained for each group studied. Various DA models have been tested, with the division into groups based on the age (patients and healthy subjects < or > than 4 years old) as well as on the *S. enterica* serotypes. The variable predictors used in modeling were the titers of certain microbial groups, which demonstrated significant changes compared to control (Table 1). Additional variable predictors included the degree of dysbiosis and age of the subjects thus making the number of variable predictors 9 (Figure 3A). Another predictor, concentration of IgE, was added to set of DA models to estimate its role in discrimination and correct classification (Figure 3B). The most successful models generated are summarized in Figure 3.


The DA analyses revealed a significant dependence of the dysbiosis on the serotype of *S. enterica* in the younger age group (Figure 3.1A), while in the older age group this dependence was less evident (Figure 3.2A). In *S. Typhimurium* infections there was a significant dependence of dysbiosis from the age (Figure 3.4A), while this was less evident for the *S. Enteritidis*-infected patients (Figure 3.3A).

The results demonstrated that the DA models with the inclusion of a IgE concentration variable had a better predictive accuracy allowing a highly confident classification of subjects in the study (Figures 3.1B–3.4B). These results also suggest that the serotype of *S. enterica* plays a role not only in the induction of a specific pro-inflammatory response (Ktsoyan et al., 2013) but is also involved in the formation of immunoglobulin responses in the adaptive arm of immunity that includes IgE production by B cells.

Correlation Between IL-17 and IgE in *S. Enteritidis*-Infected Patients

Previously we have revealed the serotype-specific induction of cytokines upon infection by *S. Typhimurium* and *S. Enteritidis* (Ktsoyan et al., 2013). In particular, acute infection by the latter serovar resulted in a significant increase of systemic IL-17 (6-fold increase compared to control and 2-fold increase compared to *S. Typhimurium*). Importantly, the elevated IL-17 level persisted in convalescent patients postinfection with *S. Enteritidis* (Ktsoyan et al., 2013).

Besides the involvement in a variety of host defenses and autoimmune diseases this cytokine has been recently implicated



in a number of allergic diseases and asthma as well (Oboki et al., 2008; Cosmi et al., 2011; Newcomb and Peebles, 2013; Manni et al., 2014; Naji et al., 2014). In particular, it has been shown that IL-17 directly promotes IgE production in human B cells (Milovanovic et al., 2010). Thus we measured the systemic concentrations of IL-17 and IgE in nine patients infected with *S. Enteritidis* and performed statistical dependence analysis between the two variables (Figure 4). The Spearman's rho, 0.683, suggested a statistically significant dependence between the systemic concentrations of IL-17 and IgE in these patients. Need to reiterate here that the concentration of IgE is total, not specific, and it was measured, together with IL-17, on the first or second day of hospital admission. Thus this is not a textbook case of the conventional specific antibody response that needs the development of a sufficient Th cell response to promote the class switch, which requires about 8 days (Murray et al., 2012).

Discussion

In this work we attempted to establish the impact of salmonellosis on the possible development of predisposition to allergies. For this, we monitored the systemic level of IgE, which is widely used as a marker for screening and detection of allergic conditions, in salmonellosis patients and control subjects. We found a significantly elevated level of IgE in the sera of patients infected by *S. Enteritidis* while infections caused by *S. Typhimurium* had no such effect. The impact of the *S. Enteritidis* infection on the elevated IgE concentration was more pronounced in the group


of patients more than 4 years old while the younger patients demonstrated insignificant departure from the control levels, possibly due to the extreme variability of IgE concentrations in this age group.

The main function of IgEs is providing immune defense against parasites such as helminthes and protozoa (Erb, 2007; Duarte et al., 2007). It was highly unexpected that infection by *S. Enteritidis* may result in the surge of systemic IgE, which production is primarily implemented via the Th2 signaling pathway (Shakib et al., 2008). On the contrary, it is generally accepted that bacteria such as *S. Enteritidis* induce mainly Th1-cell-dependent cellular and humoral responses, with a negligible contribution from the Th2 signaling pathway (Lehmann et al., 2006). This unusual finding led us to investigate the possible factors contributing to a significant induction of IgE in our patients infected by *S. Enteritidis*.

As noticed before, the host microbiota plays a substantial role in the development and maintenance of normal immune responses. Starting from the pioneering work of Strachan (1989), allergic conditions are mainly considered as a consequence of insufficient exposure of the immune system to the environmental microbiota in the contemporary lifestyle. There is a substantial body of evidence suggesting a direct link between allergic diseases and dysbiotic changes in gut microbiota (Prakash et al., 2011; Penders et al., 2007a,b).

Although the time-scale for the development of predisposition to allergic diseases is substantially longer compared to the time-course of *S. enterica* infection, we attempted to clarify a temporal dysbiosis effect during salmonellosis on the systemic


1A. Intergroup DA. All predictors, without IgE (n = 9).



1. *S. Enteritidis*-infected patients < 4 years old
 2. *S. Typhimurium*-infected patients < 4 years old
 3. Control < 4 years old
- 84.2% of the original group cases were correctly classified;
Wilks' Lambda = 0.1

Serotype/control	Predicted group membership			Total (N)
	SE < 4	ST < 4	Control < 4	
SE < 4	5	2	0	7
ST < 4	1	6	0	7
Control < 4	0	0	5	5
%				
SE < 4	71.4	28.6	0	100.0
ST < 4	14.3	85.7	0	100.0
Control < 4	0	0	100	100.0


1B. Intergroup DA. All predictors, with IgE (n = 10).



1. *S. Enteritidis*-infected patients < 4 years old
 2. *S. Typhimurium*-infected patients < 4 years old
 3. Control < 4 years old
- 94.7% of the original group cases were correctly classified;
Wilks' Lambda = 0.063


Serotype/control	Predicted group membership			Total (N)
	SE < 4	ST < 4	Control < 4	
SE < 4	6	1	0	7
ST < 4	0	7	0	7
Control < 4	0	0	5	5
%				
SE < 4	85.7	14.3	0	100.0
ST < 4	0	100	0	100.0
Control < 4	0	0	100	100.0

2A. Intergroup DA. All predictors, with IgE (n = 9).



1. *S. Enteritidis*-infected patients > 4 years old
 2. *S. Typhimurium*-infected patients > 4 years old
 3. Control > 4 years old
- 61.5% of the original group cases were correctly classified;
Wilks' Lambda = 0.4

2B. Intergroup DA. All predictors, with IgE (n = 10).




1. *S. Enteritidis*-infected patients > 4 years old
 2. *S. Typhimurium*-infected patients > 4 years old
 3. Control > 4 years old
- 84.6% of the original group cases were correctly classified;
Wilks' Lambda = 0.091

FIGURE 3 | Continued

Serotype/control	Predicted group membership			Total (N)
	SE > 4	ST > 4	Control > 4	
SE > 4	7	1	3	11
ST > 4	2	1	3	6
Control	1	0	8	9
%				
SE > 4	63.6	9.1	27.3	100.0
ST > 4	33.3	16.7	50.0	100.0
Control > 4	11.1	0	88.9	100.0

3A. Intragroup DA. All predictors, without IgE (n = 9).




1. *S. Enteritidis*-infected patients > 4 years old
 2. *S. Enteritidis*-infected patients < 4 years old
 3. Control > 4 years old
 4. Control < 4 years old
- 81.8% of the original group cases were correctly classified;
Wilks' Lambda = 0.15

Serotype/ control	Predicted group membership				Total (N)
	SE > 4	SE < 4	Control > 4	Control < 4	
SE > 4	7	1	3	0	11
SE < 4	0	6	0	2	8
Control > 4	0	0	9	0	9
Control < 4	0	0	0	5	5
%					
SE > 4	63.6	9.1	27.3	0	100.0
SE < 4	0	75.0	0	25.0	100.0
Control > 4	0	0	100	0	100.0
Control < 4	0	0	0	100	100.0

FIGURE 3 | Continued

Serotype/control	Predicted group membership			Total (N)
	SE > 4	ST > 4	Control > 4	
SE > 4	10	1	0	11
ST > 4	0	3	3	6
Control	0	0	9	9
%				
SE > 4	90.9	9.1	0	100.0
ST > 4	0	50.0	50.0	100.0
Control > 4	0	0	100.0	100.0


3B. Intragroup DA. All predictors, with IgE (n = 10).



1. *S. Enteritidis*-infected patients > 4 years old
 2. *S. Enteritidis*-infected patients < 4 years old
 3. Control > 4 years old
 4. Control < 4 years old
- 93.9% of the original group cases were correctly classified;
Wilks' Lambda = 0.048

Serotype/ control	Predicted group membership				Total (N)
	SE > 4	SE < 4	Control > 4	Control < 4	
SE > 4	11	0	0	0	11
SE < 4	0	6	0	2	8
Control > 4	0	0	9	0	9
Control < 4	0	0	0	5	5
%					
SE > 4	100	0	0	0	100.0
SE < 4	0	75.0	0	25.0	100.0
Control > 4	0	0	100	0	100.0
Control < 4	0	0	0	100	100.0


4A. Intragroup DA. All predictors, without IgE (n = 9).



1. *S. Typhimurium*-infected patients > 4 years old
 2. *S. Typhimurium*-infected patients < 4 years old
 3. Control > 4 years old
 4. Control < 4 years old
- 89.3% of the original group cases were correctly classified;
Wilks's Lambda = 0.031

Serotype/ control	Predicted group membership				Total (N)
	ST > 4	ST < 4	Control > 4	Control < 4	
ST > 4	4	0	2	0	6
ST < 4	0	7	0	1	8
Control > 4	0	0	9	0	9
Control < 4	0	0	0	5	5
%					
ST > 4	66.7	0	33.3	0	100.0
ST < 4	0	87.5	0	12.5	100.0
Control > 4	0	0	100	0	100.0
Control < 4	0	0	0	100	100.0

4B. Intragroup DA. All predictors, with IgE (n = 10).



1. *S. Typhimurium*-infected patients > 4 years old
 2. *S. Typhimurium*-infected patients < 4 years old
 3. Control > 4 years old
 4. Control < 4 years old
- 92.9% of the original group cases were correctly classified;
Wilks's Lambda = 0.028

Serotype/ control	Predicted group membership				Total (N)
	ST > 4	ST < 4	Control > 4	Control < 4	
ST > 4	4	0	2	0	6
ST < 4	0	8	0	0	8
Control > 4	0	0	9	0	9
Control < 4	0	0	0	5	5
%					
ST > 4	66.7	0	33.3	0	100.0
ST < 4	0	100	0	0	100.0
Control > 4	0	0	100	0	100.0
Control < 4	0	0	0	100	100.0

FIGURE 3 | DA models generated with predictor variables: counts of (i) *Salmonella*, (ii) *E. coli*, (iii) lactose-negative *Enterobacteriaceae*, (iv) *Bifidobacterium* spp., (v) *Lactobacillus* spp., (vi) *Enterococcus* spp.,

and (vii) *Candida* spp.; as well as (viii) the degree of dysbiosis, (ix) age of subjects, and (x) IgE concentration. IgE concentration is omitted in models in column A and included in column B models.

IgE concentration. It is well known that infection by *Salmonella* causes inflammation and dysbiotic changes in the intestinal microbiota. In particular, the growth of the Bacteroidetes and Firmicutes is suppressed, which allows the overgrowth of *Salmonella* and *Enterococcus* (Deatherage Kaiser et al., 2013). To estimate the role of the dysbiosis factor, we investigated the composition of fecal microbiota in salmonellosis caused by the two serotypes of *S. enterica* using cultivation-based techniques. Our results indicated a significantly higher degree of dysbiosis in the case of *S. Typhimurium* infection compared to *S. Enteritidis*. The level of dysbiosis was higher in *S. Typhimurium*-infected patients of the younger (<4 years old) age compared to the elder (>4 years old) patients, which is not surprising since the comprehensive colonization resistance mechanisms may be


not completely functional in the younger age group (Lawley and Walker, 2013). In general, there was a negative association between the degree of dysbiosis and IgE concentration. Although the importance of a balanced gut microbiota for the proper development and functioning of the immune system is well established, in particular for the prevention of allergic disorders, the salmonellosis-induced dysbiosis may not have an immediate effect on the production of this immunoglobulin isotype. Thus the temporal dysbiosis during salmonellosis can be ruled out as an imminent factor contributing to the excessive production of systemic IgE during *S. Enteritidis* infection.

To identify factors that may contribute to this phenomenon, we performed DA analyses with the involvement of almost all variables collected during this study. These results demonstrated

that the main factor affecting the correct classification is the serotype of *S. enterica*. In our previous work we established that *S. Enteritidis* infection induces a significantly higher level of IL-17 compared to *S. Typhimurium* (Ktsoyan et al., 2013). Moreover, even after the clearance of *S. Enteritidis* infection, the post-infection convalescent patients still maintained an elevated systemic level of IL-17 (3-fold higher compared to control). It becomes increasingly clear that the Th17 lineage plays an important role in the host defense against a variety of pathogens and fills a crucial gap that could not be attributed to the previously defined functions of the Th1 and Th2 lineages (Curtis and Way, 2009). Besides the participation in the host response against pathogens, IL-17 is also involved in a number of other pathologies that include allergic and autoimmune diseases (Oboki et al., 2008; Cosmi et al., 2011; Newcomb and Peebles, 2013; Zhao et al., 2013; Manni et al., 2014; Naji et al., 2014). The Th17 pathway, therefore, has been proposed as a potential target for the treatment of autoimmune and allergic disorders (Robinson et al., 2013).

We hypothesized, therefore, that the excessive production of IL-17 during *S. Enteritidis* infection might contribute to the increased level of IgE found in these patients. Molecular mechanisms of how IL-17 promotes the IgE production in human B cells have been recently revealed by Milovanovic et al. (2010). They established that IL-17 directly promotes IgE production and increases the number of IgE-producing B cells. In the rodent models of murine atopic dermatitis, it has been shown that IL-17 serves as an inducer for Th2 immune responses, including the induction of IgE (Nakajima et al., 2014). On the other hand, infection by *S. Typhimurium* is characterized by a lesser level of IL-17 but a higher level of IL-1 β compared to *S. Enteritidis* infection (Ktsoyan et al., 2013). IL-1 β , together with IL-18, is a central target for the activation of inflammatory response by various inflammasomes (Sollberger et al., 2014). This interleukin appeared to be crucial for the early Th17 cell differentiation (Chung et al., 2009), and indeed we observed the elevated level of IL-17 in *S. Typhimurium*-infected patients compared to control (Ktsoyan et al., 2013). This level, however, remained lower than during *S. Enteritidis* infection and did not affect the IgE level. In a murine model, the attenuated *S. Typhimurium* may reduce the allergen-induced inflammation and Th2 response (Wu et al., 2006) thus demonstrating a protective effect of this *S. enterica* serotype against allergic conditions.

The pioneering work of Mosmann et al. (1986) has laid a foundation for the functional division of T helper cells into the two main populations, Th1 and Th2. With the accumulating knowledge, however, it becomes increasingly difficult to fit the experimental observations into this simple dichotomy. Later the list has been extended to include Treg cells (Asseman et al., 1999), Th17 cells (Steinman, 2007), Th9 cells (Veldhoen et al., 2008), and Th22 cells (Duhon et al., 2009; Trifari et al., 2009). These divisions are conditional to a certain extent, and the differentiation and stability of the phenotypes of T helper cells depend on a variety of external and internal cues to respond appropriately toward them (Murphy and Stockinger, 2010). The



present view on allergic sensitization suggests the involvement of all T cell types including Th9, Th17, Th22, and Treg, in addition to Th1 and Th2 (van Ree et al., 2014). The resulting cytokine profile, with the dominance of IL-4 and/or IL-13, switch B cells to IgE production. In this work we provided direct experimental evidence suggesting a significant correlation between the elevated systemic concentrations of IL-17 and IgE in *S. Enteritidis* infection.

But what are the mechanisms specific for *S. Enteritidis* but absent in *S. Typhimurium*, which make the former more virulent and possibly contribute to the enhanced immunogenicity leading to the induction of IL-17 and IgE? There are two lines of evidence implicating the fimbriae of *S. Enteritidis* in conferring such traits compared to *S. Typhimurium*. First, whole-genome comparisons of *S. Enteritidis* and *S. Typhimurium* revealed an extensive core genome with >90% of coding sequences shared (Thomson et al., 2008). Coding sequences present in *S. Enteritidis* but absent in *S. Typhimurium* LT2 are dominated by prophage-related elements, which however, are related and harbor the same genes as prophage found previously in other *S. enterica*. Among the 13 fimbrial clusters in *S. Enteritidis*, there is a novel cluster not found in *S. Typhimurium*, which have been termed *peg* (Thomson et al., 2008). The role of the Peg fimbriae, in colonization and virulence has been established in a number of model experiments. In a recent study, a library of 54,000 transposon mutants of *S. Enteritidis* was screened for mutations attenuating the colonization ability *in vivo* using the BALB/c mouse model and a microarray-based negative-selection screening (Silva et al., 2012). Genes and genomic islands that are not present in *S. Typhimurium* or in most other *Salmonella* serovars but contribute to the pathogenicity of *S. Enteritidis* included: a type I restriction/modification system (SEN4290 to SEN4292), the *peg* fimbrial operon (SEN2144A to SEN2145B), a putative pathogenicity island (SEN1970 to

SEN1999), and a type VI secretion system remnant SEN1001 (Silva et al., 2012). In another study of *S. Enteritidis*-specific virulence genes the transposon mutants have been screened for the attenuated invasiveness in human and chicken cells (Shah et al., 2012). The *S. Enteritidis* genes that are absent in *S. Typhimurium* or in most other *Salmonella* serovars included *pegD*, SEN1152, SEN1393, and SEN1966. In a later study of the same group the *pegD* mutant was found to be defective in intestinal colonization of chickens (Addwebi et al., 2014). Thus the consistent involvement of the *peg* fimbrial proteins in the virulence of *S. Enteritidis* makes them the prime candidates for the possible involvement in induction of IL-17 and IgE by this serovar.

Recent statistical data suggest an extremely high number of conditions with an allergic component, which include allergies

to food, drugs and insects, as well as general and skin allergies, allergic rhinitis and sinusitis (<http://www.aaaai.org/about-the-aaaai/newsroom/allergy-statistics.aspx>). The worldwide rise in the number of these diseases is continuing for more than 50 years now (World Allergy Organization, 2013). In our work we established that certain bacterial infections could make a contribution to this general trend of ever growing sensitization by stimulating total IgE production. What is interesting, though, is that even closely related bacteria within the same species may induce differential IgE levels. Thus the exposure to bacterial infections may have differential outcomes for the development and severity of allergic disorders (Herz et al., 2000). More research is needed to understand the properties *S. enterica* serotypes that are closely related but, nevertheless, induce differential cytokine and IgE responses.

References

- Addwebi, T. M., Call, D. R., and Shah, D. H. (2014). Contribution of *Salmonella* Enteritidis virulence factors to intestinal colonization and systemic dissemination in 1-day-old chickens. *Poult. Sci.* 93, 871–881. doi: 10.3382/ps.2013-03710
- Angulo, F. J., and Swerdlow, D. L. (1999). "Epidemiology of human *Salmonella enterica* serovar Enteritidis infections in the United States," in *Salmonella enterica Serovar Enteritidis in Humans and Animals*, eds A. M. Saeed, R. K. Gast, M. E. Potter, and P. G. Wall (Ames, Iowa State University Press), 33–41.
- Asseman, C., Mauze, S., Leach, M. W., Coffman, R. L., and Powrie, F. (1999). An essential role for interleukin 10 in the function of regulatory T cells that inhibit intestinal inflammation. *J. Exp. Med.* 190, 995–1004. doi: 10.1084/jem.190.7.995
- Centers for Disease Control and Prevention (CDC). (2013). Surveillance for foodborne disease outbreaks—United States, 2009–2010. *MMWR Morb. Mortal. Wkly. Rep.* 62, 41–47. Available online at: <http://www.cdc.gov/mmwr/preview/mmwrhtml/mm6203a1.htm>
- Chung, Y., Chang, S. H., Martinez, G. J., Yang, X. O., Nurieva, R., Kang, H. S., et al. (2009). Critical regulation of early Th17 cell differentiation by interleukin-1 signaling. *Immunity* 30, 576–587. doi: 10.1016/j.immuni.2009.02.007
- Cosmi, L., Liotta, F., Maggi, E., Romagnani, S., and Annunziato, F. (2011). Th17 cells: new players in asthma pathogenesis. *Allergy* 66, 989–998. doi: 10.1111/j.1398-9995.2011.02576.x
- Cremon, C., Stanghellini, V., Pallotti, F., Fogacci, E., Bellacosa, L., Morselli-Labate, A. M., et al. (2014). *Salmonella* gastroenteritis during childhood is a risk factor for irritable bowel syndrome in adulthood. *Gastroenterology* 147, 69–77. doi: 10.1053/j.gastro.2014.03.013
- Curtis, M. M., and Way, S. S. (2009). Interleukin-17 in host defence against bacterial, mycobacterial and fungal pathogens. *Immunology* 126, 177–185. doi: 10.1111/j.1365-2567.2008.03017.x
- De Amici, M., and Ciprandi, G. (2013). The age impact on serum total and allergen-specific IgE. *Allergy Asthma Immunol. Res.* 5, 170–174. doi: 10.4168/aaair.2013.5.3.170
- Deatherage Kaiser, B. L., Li, J., Sanford, J. A., Kim, Y. M., Kronewitter, S. R., Jones, M. B., et al. (2013). A multi-omic view of host-pathogen-commensal interplay in *Salmonella*-mediated intestinal infection. *PLoS ONE* 8:e67155. doi: 10.1371/journal.pone.0067155
- Duarte, J., Deshpande, P., Guiyedi, V., Mécheri, S., Fesel, C., Cazenave, P., et al. (2007). Total and functional parasite specific IgE responses in *Plasmodium falciparum*-infected patients exhibiting different clinical status. *Malar. J.* 6:1. doi: 10.1186/1475-2875-6-1
- Duhen, T., Geiger, R., Jarrossay, D., Lanzavecchia, A., and Sallusto, F. (2009). Production of interleukin 22 but not interleukin 17 by a subset of human skin-homing memory T cells. *Nat. Immunol.* 10, 857–863. doi: 10.1038/ni.1767
- Erb, K. J. (2007). Helminths, allergic disorders and IgE-mediated immune responses: where do we stand? *Eur. J. Immunol.* 37, 1170–1173. doi: 10.1002/eji.200737314
- Feehley, T., Stefka, A. T., Cao, S., and Nagler, C. R. (2012). Microbial regulation of allergic responses to food. *Semin. Immunopathol.* 34, 671–688. doi: 10.1007/s00281-012-0337-5
- Herz, U., Lacy, P., Renz, H., and Erb, K. (2000). The influence of infections on the development and severity of allergic disorders. *Curr. Opin. Immunol.* 12, 632–640. doi: 10.1016/S0952-7915(00)00155-2
- Jacobsen, A., Hendriksen, R. S., Aaresturp, F. M., Ussery, D. W., and Friis, C. (2011). The *Salmonella enterica* pan-genome. *Microb. Ecol.* 62, 487–504. doi: 10.1007/s00248-011-9880-1
- Jones, T. F., Ingram, L. A., Cieslak, P. R., Vugia, D. J., Tobin-D'Angelo, M., Hurd, S., et al. (2008). Salmonellosis Outcomes Differ Substantially by Serotype. *J. Infect. Dis.* 198, 109–114. doi: 10.1086/588823
- Kerkhof, M., Dubois, A. E. J., Postma, D. S., Schouten, J. P., and de Monchy, J. G. R. (2003). Role and interpretation of total serum IgE measurements in the diagnosis of allergic airway disease in adults. *Allergy* 58, 905–911. doi: 10.1034/j.1398-9995.2003.00230.x
- Ktsoyan, Z., Ghazaryan, K., Manukyan, G., Martirosyan, A., Mnatsakanyan, A., Arakelova, K., et al. (2013). Inflammatory responses to *Salmonella* infections are serotype-specific. *Int. J. Bacteriol.* 2013:168179. doi: 10.1155/2013/168179
- Lawley, T. D., and Walker, A. W. (2013). Intestinal colonization resistance. *Immunology* 138, 1–11. doi: 10.1111/j.1365-2567.2012.03616.x
- Lehmann, J., Springer, S., Werner, C. E., Lindner, T., Bellmann, S., Straubinger, R. K., et al. (2006). Immunity induced with a *Salmonella enterica* serovar Enteritidis live vaccine is regulated by Th1-cell-dependent cellular and humoral effector mechanisms in susceptible BALB/c mice. *Vaccine* 24, 4779–4793. doi: 10.1016/j.vaccine.2006.03.024
- Licciardi, P. V., Wong, S., Tang, L. K., and Karagiannis, T. C. (2010). Epigenome targeting by probiotic metabolites. *Gut Pathog.* 2:24. doi: 10.1186/1757-4749-2-24
- Manni, M. L., Robinson, K. M., and Alcorn, J. F. (2014). A tale of two cytokines: IL-17 and IL-22 in asthma and infection. *Expert Rev. Respir. Med.* 8, 25–42. doi: 10.1586/17476348.2014.854167
- Milovanovic, M., Drozdenko, G., Weise, C., Babina, M., and Worm, M. (2010). Interleukin-17A promotes IgE production in human B cells. *J. Invest. Dermatol.* 130, 2621–2628. doi: 10.1038/jid.2010.175
- Mitrokhin, S. D., Nikushkin, E. V., Minayeff, V. I., Vorobyov, A. A., and Kryuchkov, M. I. (1998). *Complex Laboratory Diagnostics - a Basis of Monitoring of Microbial ecology of the Human Gut*. Moscow: Kremlevskaya Medicina; Klinicheskiy Vestnik.
- Mosmann, T. R., Cherwinski, H., Bond, M. W., Giedlin, M. A., and Coffman, R. L. (1986). Two types of murine helper T cell clones. I. Definition according to profiles of lymphokine activities and secreted proteins. *J. Immunol.* 136, 2348–2357.
- Munro, D. S., Girdwood, R. W. A., and Reilly, W. J. (1999). "Salmonella enterica serovar Enteritidis in Scotland," in *Salmonella enterica Serovar Enteritidis in Humans and Animals*, eds A. M. Saeed, R. K. Gast, M. E. Potter, and P. G. Wall (Ames, Iowa State University Press), 27–31.

- Murphy, K. M., and Stockinger, B. (2010). Effector T cell plasticity: flexibility in the face of changing circumstances. *Nat. Immunol.* 11, 674–680. doi: 10.1038/ni.1899
- Murray, P. R., Rosenthal, K. S., and Pfaller, M. A. (2012). *Medical Microbiology, 7th Edn.* Amsterdam: Elsevier Health Sciences.
- Naji, N., Smith, S. G., Gauvreau, G. M., and O'Byrne, P. M. (2014). T Helper 17 cells and related cytokines after allergen inhalation challenge in allergic asthmatics. *Int. Arch. Allergy Immunol.* 165, 27–34. doi: 10.1159/000367789
- Nakajima, S., Kitoh, A., Egawa, G., Natsuaki, Y., Nakamizo, S., Moniaga, C. S., et al. (2014). IL-17A as an inducer for Th2 immune responses in murine atopic dermatitis models. *J. Invest. Dermatol.* 134, 2122–2130. doi: 10.1038/jid.2014.51
- Newcomb, D. C., and Peebles, R. S. Jr. (2013). Th17-mediated inflammation in asthma. *Curr. Opin. Immunol.* 25, 755–760. doi: 10.1016/j.coi.2013.08.002
- Oboki, K., Ohno, T., Saito, H., and Naka, S. (2008). Th17 and Allergy. *Allergol. Int.* 57, 121–134. doi: 10.2332/allergolint.R-07-160
- Olson, A. B., Andrysiak, A. K., Tracz, D. M., Guard-Bouldin, J., Demczuk, W., Ng, L. K., et al. (2007). Limited genetic diversity in *Salmonella enterica* serovar Enteritidis PT13. *BMC Microbiol.* 7:87. doi: 10.1186/1471-2180-7-87
- Penders, J., Stobberingh, E. E., van den Brandt, P. A., and Thijs, C. (2007a). The role of the intestinal microbiota in the development of atopic disorders. *Allergy* 62, 1223–1236. doi: 10.1111/j.1398-9995.2007.01462.x
- Penders, J., Thijs, C., van den Brandt, P. A., Kummeling, I., Snijders, B., Stelma, F., et al. (2007b). Gut microbiota composition and development of atopic manifestations in infancy: the KOALA Birth Cohort Study. *Gut* 56, 661–667. doi: 10.1136/gut.2006.100164
- Petersen, C., and Round, J. L. (2014). Defining dysbiosis and its influence on host immunity and disease. *Cell. Microbiol.* 16, 1024–1033. doi: 10.1111/cmi.12308
- Prakash, S., Rodes, L., Coussa-Charley, M., and Tomaro-Duchesneau, C. (2011). Gut microbiota: next frontier in understanding human health and development of biotherapeutics. *Biologics* 5, 71–86. doi: 10.2147/BTT.S19099
- Rabsch, W., Tschape, H., and Baumler, A. J. (2001). Non-typhoidal salmonellosis: emerging problems. *Microbes Infect.* 3, 237–247. doi: 10.1016/S1286-4579(01)01375-2
- Robinson, K. M., Manni, M. L., Biswas, S. P., and Alcorn, J. F. (2013). Clinical consequences of targeting IL-17 and TH17 in autoimmune and allergic disorders. *Curr. Allergy Asthma Rep.* 13, 587–595. doi: 10.1007/s11882-013-0361-0
- Sekirov, I., Russell, S. L., Antunes, L. C. M., and Finlay, B. B. (2010). Gut microbiota in health and disease. *Physiol. Rev.* 90, 859–904. doi: 10.1152/physrev.00045.2009
- Shah, D. H., Zhou, X., Kim, H. Y., Call, D. R., and Guard, J. (2012). Transposon mutagenesis of *Salmonella enterica* serovar Enteritidis identifies genes that contribute to invasiveness in human and chicken cells and survival in egg albumen. *Infect. Immun.* 80, 4203–4215. doi: 10.1128/IAI.00790-12
- Shakib, F., Ghaemmaghami, A. M., and Sewell, H. F. (2008). The molecular basis of allergenicity. *Trends Immunol.* 29, 633–642. doi: 10.1016/j.it.2008.08.007
- Silva, C. A., Blondel, C. J., Quezada, C. P., Porwollik, S., Andrews-Polymenis, H. L., Toro, C. S., et al. (2012). Infection of mice by *Salmonella enterica* serovar Enteritidis involves additional genes that are absent in the genome of serovar Typhimurium. *Infect. Immun.* 80, 839–849. doi: 10.1128/IAI.05497-11
- Sollberger, G., Strittmatter, G. E., Garstkiewicz, M., Sand, J., and Beer, H. D. (2014). Caspase-1: the inflammasome and beyond. *Innate Immun.* 20, 115–125. doi: 10.1177/1753425913484374
- Stecher, B., Robbiani, R., Walker, A. W., Westendorf, A. M., Barthel, M., Kremer, M., et al. (2007). *Salmonella enterica* serovar Typhimurium exploits inflammation to compete with the intestinal microbiota. *PLoS Biol.* 5:e244. doi: 10.1371/journal.pbio.0050244
- Steinman, L. (2007). A brief history of T(H)17, the first major revision in the T(H)1/T(H)2 hypothesis of T cell-mediated tissue damage. *Nat. Med.* 13, 139–145. doi: 10.1038/nm1551
- Strachan, D. P. (1989). Hay fever, hygiene, and household size. *BMJ* 299, 1259–1260. doi: 10.1136/bmj.299.6710.1259
- Tabolin, V. A., Belmer, S. V., Gasilina, T. V., Muhina, U. G., and Korneva, T. I. (1998). *Rational Therapy of Intestinal Dysbacteriosis in Children.* Moscow: Medicina. Available online at: <http://www.rusmedserv.com/childgastro/dysb.htm>
- Thomson, N. R., Clayton, D. J., Windhorst, D., Vernikos, G., Davidson, S., Churcher, C., et al. (2008). Comparative genome analysis of *Salmonella* Enteritidis PT4 and *Salmonella* Gallinarum 287/91 provides insights into evolutionary and host adaptation pathways. *Genome Res.* 18, 1624–1637. doi: 10.1101/gr.077404.108
- Tschape, H., Liesegang, A., Gericke, B., Prager, R., Rabsch, W., and Helmuth, R. (1999). “Ups and downs of *Salmonella enterica* Serovar Enteritidis in Germany,” in *Salmonella enterica Serovar Enteritidis in Humans and Animals*, eds A. M. Saeed, R. K. Gast, M. E. Potter, and P. G. Wall (Ames, Iowa State University Press), 51–61.
- Trifari, S., Kaplan, C. D., Tran, E. H., Crellin, N. K., and Spits, H. (2009). Identification of a human helper T cell population that has abundant production of interleukin 22 and is distinct from T(H)-17, T(H)1 and T(H)2 cells. *Nat. Immunol.* 10, 864–871. doi: 10.1038/ni.1770
- Van Immerseel, F. (2010). Stress-induced survival strategies enable *Salmonella* Enteritidis to persistently colonize the chicken oviduct tissue and cope with antimicrobial factors in egg white: a hypothesis to explain a pandemic. *Gut Pathog.* 2:23. doi: 10.1186/1757-4749-2-23
- van Ree, R., Hummelshøj, L., Plantinga, M., Poulsen, L. K., and Swindle, E. (2014). Allergic sensitization: host-immune factors. *Clin. Transl. Allergy* 4:12. doi: 10.1186/2045-7022-4-12
- Veldhoen, M., Uyttenhove, C., van Snick, J., Helmby, H., Westendorf, A., Buer, J., et al. (2008). Transforming growth factor- β ‘reprograms’ the differentiation of T helper 2 cells and promotes an interleukin 9-producing subset. *Nat. Immunol.* 9, 1341–1346. doi: 10.1038/ni.1659
- World Allergy Organization (WAO). (2013). *White Book on Allergy 2013 Update.* Milwaukee: World Allergy Organization.
- Wu, C. J., Chen, L. C., and Kuo, M. L. (2006). Attenuated *Salmonella typhimurium* reduces ovalbumin-induced airway inflammation and T-helper type 2 responses in mice. *Clin. Exp. Immunol.* 145, 116–122. doi: 10.1111/j.1365-2249.2006.03099.x
- Zhao, J., Lloyd, C. M., and Noble, A. (2013). Th17 responses in chronic allergic airway inflammation abrogate regulatory T-cell-mediated tolerance and contribute to airway remodeling. *Mucosal Immunol.* 6, 335–346. doi: 10.1038/mi.2012.76

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2015 Ktsoyan, Mkrtchyan, Zakharyan, Mnatsakanyan, Arakelova, Gevorgyan, Ktsoyan, Sedrakyan, Hovhannisyanyan, Ghazaryan, Boyajyan and Aminov. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.