

CORRECTION

# Correction: Aerobic Exercise Training in Post-Polio Syndrome: Process Evaluation of a Randomized Controlled Trial

Eric L. Voorn, Fieke S. Koopman, Merel A. Brehm, Anita Beelen, Arnold de Haan, Karin H. L. Gerrits, Frans Nollet

There is an error in [Table 1](#). The values in the “ET” column are swapped with values in the “UC” column. Please see the corrected [Table 1](#) here.

**Table 1. Baseline characteristics of participants.**

	ET (N = 22)	UC (N = 22)
<b>Demographic data</b>		
Age (yr), mean (SD)	56.7 (8.9)	60.1 (7.4)
Gender, n female (%)	13 (59%)	11 (50%)
Body mass index ( $kg/m^2$ ), mean (SD)	25.6 (4.3)	26.5 (3.4)
<b>Polio characteristics</b>		
Age at acute polio (yr), median (range)	2 (0–16)	3 (1–40)
Time since new symptoms (yr), mean (SD)	14.6 (9.5)	13.2 (7.7)
Present walking distance, n(%) <sup>a</sup>		
Around the house	6 (27%)	9 (41%)
Seldom further than 1 km	11 (50%)	8 (36%)
Regularly further than 1 km	5 (23%)	5 (23%)
Manual muscle testing sum score legs (0 to 80), median (range) <sup>b</sup>	69.6 (43.0–80.0)	67.8 (41.8–80.0)
Manual muscle testing sum score arms (0 to 80), median (range) <sup>c</sup>	50.0 (16.8–50.0)	50.0 (32.0–50.0)

Abbreviations: ET, exercise therapy; UC, usual care.

<sup>a</sup> Walking distance was classified into 3 categories: around the house, seldom further than 1 km, and regularly further than 1 km.


<sup>b</sup> Sum score for muscle strength of the legs was calculated by adding 16 muscle groups. Each muscle group had a score between 0 and 5, sum score ranged from 0 to 80 [19].

<sup>c</sup> Sum score for muscle strength of the arms was calculated by adding 10 muscle groups. Each muscle group had a score between 0 and 5, sum score ranged from 0 to 50 [19].

<https://doi.org/10.1371/journal.pone.0192338.t001>

## Reference

1. Voorn EL, Koopman FS, Brehm MA, Beelen A, de Haan A, Gerrits KHL, et al. (2016) Aerobic Exercise Training in Post-Polio Syndrome: Process Evaluation of a Randomized Controlled Trial. PLoS ONE 11 (7): e0159280. <https://doi.org/10.1371/journal.pone.0159280> PMID: 27419388



## OPEN ACCESS

**Citation:** Voorn EL, Koopman FS, Brehm MA, Beelen A, de Haan A, Gerrits KHL, et al. (2018) Correction: Aerobic Exercise Training in Post-Polio Syndrome: Process Evaluation of a Randomized Controlled Trial. PLoS ONE 13(1): e0192338. <https://doi.org/10.1371/journal.pone.0192338>

**Published:** January 30, 2018

**Copyright:** © 2018 Voorn et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.