



## RESEARCH ARTICLE

**REVIS** **Typing complex meningococcal vaccines to understand diversity and population structure of key vaccine antigens**  
[version 2; referees: 2 approved]

Charlene M. C. Rodrigues <sup>1</sup>, Hannah Chan<sup>2</sup>, Caroline Vipond<sup>2</sup>, Keith Jolley <sup>1</sup>, Odile B. Harrison<sup>1</sup>, Jun Wheeler<sup>2</sup>, Gail Whiting <sup>2</sup>, Ian M. Feavers<sup>2</sup>, Martin C. J. Maiden <sup>1</sup>

<sup>1</sup>Department of Zoology, University of Oxford, Oxford, OX1 3SY, UK

<sup>2</sup>Division of Bacteriology, National Institute for Biological Standards and Control, Potters Bar, EN6 3QG, UK

**v2** **First published:** 29 Nov 2018, 3:151 (<https://doi.org/10.12688/wellcomeopenres.14859.1>)  
**Latest published:** 19 Mar 2019, 3:151 (<https://doi.org/10.12688/wellcomeopenres.14859.2>)

**Abstract**

**Background:** Protein-conjugate capsular polysaccharide vaccines can potentially control invasive meningococcal disease (IMD) caused by five (A, C, W, X, Y) of the six IMD-associated serogroups. Concerns raised by immunological similarity of the serogroup B capsule to human neural cell carbohydrates, meant that ‘serogroup B substitute’ vaccines target more variable subcapsular protein antigens. A successful approach using outer membrane vesicles (OMVs) as major vaccine components had limited strain coverage. In 4CMenB (Bexsero<sup>®</sup>), recombinant proteins have been added to ameliorate this problem.

**Methods:** Scalable, portable, genomic techniques were used to investigate the Bexsero<sup>®</sup> OMV protein diversity in meningococcal populations. Shotgun proteomics identified 461 proteins in the OMV, defining a complex proteome. Amino acid sequences for the 24 proteins most likely to be involved in cross-protective immune responses were catalogued within the [PubMLST.org/neisseria](http://PubMLST.org/neisseria) database using a novel OMV peptide Typing (OMVT) scheme.

**Results:** Among these proteins there was variation in the extent of diversity and association with meningococcal lineages, identified as clonal complexes (ccs), ranging from the most conserved peptides (FbpA, NEISp0578, and putative periplasmic protein, NEISp1063) to the most diverse (TbpA, NEISp1690). There were 1752 unique OMVTs identified amongst 2492/3506 isolates examined by whole-genome sequencing (WGS). These OMVTs were grouped into clusters (sharing  $\geq 18$  identical OMVT peptides), with 45.3% of isolates assigned to one of 27 OMVT clusters. OMVTs and OMVT clusters were strongly associated with cc, genogroup, and Bexsero<sup>®</sup> antigen variants, demonstrating that combinations of OMV proteins exist in discrete, non-overlapping combinations associated with genogroup and Bexsero<sup>®</sup> Antigen Sequence Type. This highly structured population of IMD-associated meningococci is consistent with strain structure models invoking host immune and/or metabolic selection.

**Open Peer Review**

**Referee Status:**

Invited Referees

1 2

**REVIS**

**version 2**

published  
19 Mar 2019

**version 1**

published  
29 Nov 2018



report



report

1 **Mariagrazia Pizza**, GlaxoSmithKline (GSK), Italy

**Margherita Bodini**, GlaxoSmithKline (GSK), Italy

2 **Peter van der Ley**, Institute for Translational Vaccinology (Intravacc), The Netherlands

Any reports and responses or comments on the article can be found at the end of the article.

**Conclusions:** The OMVT scheme facilitates region-specific WGS investigation of meningococcal diversity and is an open-access, portable tool with applications for vaccine development, especially in the choice of antigen combinations, assessment and implementation.

### Keywords

Outer membrane vesicle, OMV, *Neisseria meningitidis*, meningococcal, vaccines, diversity, proteomics, typing, invasive meningococcal disease

**Corresponding author:** Charlene M. C. Rodrigues ([charlene.rodrigues@gtc.ox.ac.uk](mailto:charlene.rodrigues@gtc.ox.ac.uk))

**Author roles:** **Rodrigues CMC:** Conceptualization, Data Curation, Formal Analysis, Funding Acquisition, Investigation, Methodology, Project Administration, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Chan H:** Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Project Administration, Visualization, Writing – Review & Editing; **Vipond C:** Conceptualization, Formal Analysis, Investigation, Methodology, Resources, Visualization, Writing – Review & Editing; **Jolley K:** Resources, Software, Writing – Review & Editing; **Harrison OB:** Methodology, Software, Writing – Review & Editing; **Wheeler J:** Formal Analysis, Methodology, Resources, Writing – Review & Editing; **Whiting G:** Formal Analysis, Methodology, Resources, Writing – Review & Editing; **Feavers IM:** Conceptualization, Formal Analysis, Methodology, Supervision, Writing – Review & Editing; **Maiden MCJ:** Conceptualization, Formal Analysis, Methodology, Project Administration, Resources, Software, Supervision, Writing – Original Draft Preparation, Writing – Review & Editing

**Competing interests:** No competing interests were disclosed.

**Grant information:** This study was funded by the Wellcome Trust (109031 and 087622).

*The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.*

**Copyright:** © 2019 Rodrigues CMC *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution Licence](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**How to cite this article:** Rodrigues CMC, Chan H, Vipond C *et al.* **Typing complex meningococcal vaccines to understand diversity and population structure of key vaccine antigens [version 2; referees: 2 approved]** Wellcome Open Research 2019, 3:151 (<https://doi.org/10.12688/wellcomeopenres.14859.2>)

**First published:** 29 Nov 2018, 3:151 (<https://doi.org/10.12688/wellcomeopenres.14859.1>)

**REVISED Amendments from Version 1**

We are grateful to the reviewers for their insightful and constructive comments. We have revised the manuscript to reflect these comments in particular clarifying: the proteomic methodology; decisions for choosing proteins to include in the typing scheme; the uncertainty regarding cross-protection of certain outer membrane proteins; and the limitations of "designer" OMV vaccine.

**See referee reports**

## Introduction

Ideal vaccines comprise a single antigenically conserved component of the target pathogen, preferably against a virulence determinant. The highly successful tetanus and diphtheria toxoid vaccines epitomise this approach, with a major impact on disease globally as part of the World Health Organisation Expanded Programme on Immunisation (WHO EPI)<sup>1</sup>. Similarly, the conjugate polysaccharide vaccines against *Haemophilus influenzae* type b and certain capsular variants of *Neisseria meningitidis*, the meningococcus, and *Streptococcus pneumoniae*, have been highly successful in eliminating the diseases they cause, largely through herd immunity effects<sup>2</sup>. A number of more complex vaccines have also been successful, but several of these, including the widely used tuberculosis vaccine BCG<sup>3</sup>, or those comprising killed whole cells, such as whole cell pertussis and certain typhoid and cholera vaccines, are more difficult to define and control, often achieving imperfect disease control. Antigenically diverse pathogens continue to challenge vaccine design, and it is necessary either: (i) to identify protective antigens that are invariant, as has been attempted with the RTS,S malaria vaccine<sup>4</sup>, or (ii) to include multiple components, perhaps varying composition over time, as has been done with influenza vaccines over many years.

There is a limited repertoire of meningococcal polysaccharide capsules associated with invasive meningococcal disease (IMD), identified as serogroups A, B, C, W, X, and Y<sup>5</sup>, potentially simplifying vaccine formulation. While vaccines based on one or a few capsular polysaccharides have been deployed successfully since the 1960s, with notable success achieved by protein-polysaccharide conjugate vaccines<sup>2</sup>, a comprehensive anti-capsular vaccine has been precluded by concerns about the safety and efficacy of formulations that include the serogroup B polysaccharide. Attempts to generate 'serogroup B substitute' vaccines have consequently focussed on subcapsular antigens, especially proteins, but in most cases these are highly variable. Meningococcal diversity extends beyond antigen genes, with a very wide range of distinct genotypes defined by variation in 'housekeeping genes' that encode essential cytoplasmic metabolic functions. This diversity is captured by multilocus sequence typing (MLST) approaches, including conventional MLST<sup>6</sup>, ribosomal MLST (rMLST), and core genome MLST (cgMLST)<sup>7</sup>. These approaches have shown that both genetic and antigenic diversity of this organism is structured into clusters of related organisms, which share a common ancestor (lineages). These clusters are recognised by conventional seven-locus MLST as clonal complexes (ccs)<sup>8</sup>. The ccs are correlated with clinical phenotypes, including the propensity to cause disease and its severity<sup>9</sup>, and antigenic characteristics including capsular and subcapsular antigens<sup>10</sup>. Comprehensive genomic

typing schemes have been developed that catalogue serogroups, ccs and principle vaccine antigens<sup>11</sup>.

The multiple variants of the subcapsular antigens included in serogroup B substitute vaccines do not necessarily generate cross-reactive protective immune responses against a wide range of distinct meningococci<sup>12</sup>. Two main approaches have been adopted to overcome this problem: (i) the generation of strain-specific vaccines that target epidemics caused by particular meningococci; (ii) the identification of conserved antigens that offer broad protection. The most widely used strain-specific vaccines have been based on meningococcal outer membrane vesicles (OMVs), which contain outer membrane proteins (OMPs) as principle antigens, especially the PorA porin<sup>5</sup>. The investigational vaccine 'Nonamen' included nine different PorA proteins in a single OMV background to increase population coverage<sup>13</sup>. 'Second generation' meningococcal B substitute vaccine development, using both conventional<sup>14</sup> and 'reverse vaccinology'<sup>15</sup> approaches, identified Factor H binding protein (fHbp) as a leading contender for a single, conserved vaccine antigen, leading to licensure of two vaccines: Bexsero<sup>®</sup> (4CMenB)<sup>16</sup>; and Trumenba<sup>®</sup> (rLP2086)<sup>14</sup>. The fHbp protein is, however, highly variable in meningococcal populations<sup>17</sup>, so the effectiveness of the protection offered, not fully established at the time of writing, is dependent on the generation of cross-protective immune responses<sup>12</sup>.

The bivalent Trumenba<sup>®</sup> vaccine, containing two lipidated fHbp variants was licenced for use in the US in 10–15-year-olds in 2014 and for those aged 10 years or over in Europe in 2017; however, as of summer 2018, Bexsero<sup>®</sup> vaccine was the only serogroup B substitute vaccine licenced, in 2013, for use in infants. This vaccine was introduced into the UK infant immunisation programme in 2015<sup>18</sup>, undergoing post implementation assessment<sup>19</sup> and comprising three recombinant meningococcal proteins: fHbp, neisserial heparin binding antigen (NHBA); and *Neisseria* adhesin A (NadA); together with the MeNZB<sup>TM</sup> OMV that contained PorA as a principal antigen<sup>20</sup>. The meningococcal antigen typing system (MATS) immunological assay had been used to assess potential cross-reactivity of toddler antibodies to the principal Bexsero<sup>®</sup> components<sup>21,22</sup> and a Bexsero<sup>®</sup> Antigen Sequence Typing scheme (BAST) was developed to assess the sequence variability of these vaccine components from meningococcal whole-genome sequence (WGS) data<sup>23,24</sup>. At the time of writing, however, no means were available to systematically assess the other protein components of the OMV, which could potentially contribute to cross-immunity generated by this vaccine. Shotgun proteomics was performed to identify the protein content of the Bexsero<sup>®</sup>-derived OMVs, and alongside published data<sup>25</sup>, informed the selection of proteins for use in the development of an Outer Membrane Vesicle peptide Typing (OMVT) scheme. Here we describe the OMVT scheme and investigate its value in supporting the assessment of vaccine impact and the improvement of vaccine formulation.

## Methods

### Proteomic OMV sample preparation

Two batches of meningococcal NZ98/254 derived OMVs, from GlaxoSmithKline, Siena Italy, were analysed in triplicate. Each replicate contained 100 µg total protein in 0.5% SDS and 200 mM

triethylammonium bicarbonate, (TEAB), the pH 8.0 was reduced with tris(2-carboxyethyl) phosphine and alkylated with iodoacetamide before an overnight acetone precipitation. A 200 mM TEAB solution containing 2.5 µg trypsin (Promega) was directly added to the protein pellet and digestion was performed at 37°C overnight. The resulting peptides from each sample in a set were labelled with a different isobaric tag (TMTs 126–131, ThermoFisher) before being mixed into a single sixplex sample.

A sample of labelled peptide mixture was injected on to a XBridge C18 column, (5 µm, 4.6 mm id and 25 cm long, Waters) for the first-dimension high pH RP-HPLC separation under a linear gradient consisting mobile phase A (10 mM ammonium formate, pH 10.0) and up to 70% B (90% acetonitrile in mobile phase A) for 2 hours at flow rate of 0.5 ml/min, using a Jasco system consisting an autosampler, semi-micro HPLC pumps and UV detector. Eluted fractions were collected and concatenated into eight tubes and vacuum dried. As OMPs are not all equally susceptible to enzymatic digestion, we evaluated different sample preparation methods. The method described provided an increased number of OMPs to other studies in our laboratory and the contribution of membrane to cytoplasmic proteins was comparable to other studies<sup>25–27</sup>.

#### Nano-LC-MS/MS

Nano-LC and tandem mass spectrometry (MS/MS) was performed using a U3000 direct nano system coupled with nano-electrospray and LTQ-Orbitrap Discovery mass spectrometer (Thermo). Resuspending in 0.1% formic acid, the HPLC fractions containing a mixture of sixplex labelled peptides were separated on a PepMap C18 reversed phase nano column (3 µm, 100 Å, 50 cm length, Thermo) under a column flow rate of 0.3 µl/min using linear gradient of 5–25% for 180 min, 25–32% for 20 min and 32–90% for 10 min of 95% acetonitrile and 0.1% formic acid. MS scan

and MS/MS fragmentation were carried out in Orbitrap and LTQ mass analysers, respectively, using 2 cycles of top 3 data-dependent acquisition with dynamic exclusion mode enabled and total cycle time at approximately 30 milliseconds. The first cycle used collision-induced dissociation (CID) fragmentation generating spectra for peptide sequencing, and the second High energy CID (HCD).

#### Proteomic data analysis

Mass spectra processing, database searching and quantitation were carried out using Thermo Proteome Discoverer 1.4 with built-in Sequest against the Uniprot *N. meningitidis* MC58 FASTA database, release 2014\_03. Spectra from the 8 fractions were added together as one sample during searching. Initial mass tolerances by MS were set to 10 ppm. Up to two missed tryptic cleavages were considered. Methionine oxidation was set as dynamic modification whereas carboxymethylation on cysteine and TMT6plex labels on N-terminal amino acid and lysine side chain were set as static modifications. Positive protein identification was considered when matched to minimum of two peptides sequenced at rank 1 with high confidence. Protein FASTA sequences identified were subsequently submitted to a web server for protein subCELlular LOcalisation prediction (<http://cello.life.nctu.edu.tw/cello2go/>) and PSORTb; Supplementary Table 1)<sup>28–30</sup>.

#### OMVT scheme development and nomenclature

A total of 25 proteins were chosen, primarily for pragmatic reasons as this was considered manageable for detailed curation within the typing scheme. The proteins represented the 20 most abundant outer membrane or periplasmic proteins identified in by Nano-LC-MS/MS with an additional five proteins chosen from the published core proteome<sup>25</sup>, also identified by Nano-LC-MS/MS (Table 1). All proteins chosen for the scheme were predicted to be either outer membrane or periplasmic and therefore more likely to be OMV antigens than the cytoplasmic proteins, which were

**Table 1. Details of the outer membrane vesicle (OMV) proteins included in the Outer Membrane Vesicle peptide Typing (OMVT) scheme including the corresponding NEISp number assigned with respect to the NEIS nucleotide sequence on PubMLST.org.**

| NEISp annotation | NEIS no. (Gene number) | UniProtKB identifier | Cellular location | Protein description                                       | Isolates with peptide variant present n (%) | Isolates with null peptide variant (0) n (%) |
|------------------|------------------------|----------------------|-------------------|---|---|--|
| 0073             | NEIS0073 (NMB0088)     | A0A0E0TMZ0           | OM                | Outer membrane protein, OmpP1 (OmpP1/FadL/TodX family)    | 3438 (98.06)                                | 64 (1.83)                                    |
| 0101             | NEIS0101 (NMB0109b)    | A0A0E0TRC4           | Unknown           | LysM domain/M23 peptidase domain protein                  | 3506 (100.00)                               | 0 (0.00)                                     |
| 0173             | NEIS0173 (NMB0182)     | A0A0E0TN68           | OM                | Outer membrane protein assembly factor BamA (Omp85)       | 3501 (99.86)                                | 0 (0.00)                                     |
| 0196             | NEIS0196 (NMB0204)     | A0A0E0TN82           | OM                | Outer membrane protein assembly factor BamE (Lipoprotein) | 3504 (99.94)                                | 0 (0.00)                                     |

| NEISp annotation | NEIS no. (Gene number)      | UniProtKB identifier | Cellular location | Protein description   | Isolates with peptide variant present n (%) | Isolates with null peptide variant (0) n (%) |
|------------------|-----------------------------|----------------------|-------------------|---|---|--|
| 0275             | NEIS 0275 (NMB0280)         | F0MIT7 (H44/76)      | OM                | LPS-assembly protein LptD / Putative organic solvent tolerance protein                        | 3442 (98.17)                                | 4 (0.11)                                     |
| 0408             | NEIS0408 (NMB1812)          | A0A0E0TNY5           | OM                | Type IV pilus secretin PilQ (PilQ)  | 3429 (97.80)                                | 36 (1.03)                                    |
| 0578             | NEIS0578 (NMB0634)          | A0A0E0TPP5           | PP                | Iron (III) ABC transporter, periplasmic iron (III)-binding protein (FbpA)                     | 3501 (99.86)                                | 0 (0.00)                                     |
| 0612             | NEIS0612 (NMB0663)          | A0A0E0TQI3           | OM                | Surface protein A (NspA)  | 3498 (99.77)                                | 0 (0.00)                                     |
| 0944             | NEIS0944 (NMB0964b)         | A0A0E0TQ95           | OM                | Iron receptor protein (Irp, TonB-dependent receptor)  | 3203 (91.36)                                | 64 (1.83)                                    |
| 1063             | NEIS1063 (NMB1124/ NMB1162) | A0A0E0TQ69           | OM                | Putative lipoprotein / Putative periplasmic protein   | 3451 (98.43)                                | 6 (0.17)                                     |
| 1066             | NEIS1066 (NMB1126/1164)     | A0A0E0TQU6           | PP                | CsgG family protein / Putative lipoprotein  | 3459 (98.66)                                | 40 (1.14)                                    |
| 1271             | NEIS1271 (NMB1333)          | A0A0E0TRB5           | OM                | M23 peptidase domain protein  | 3471 (99.00)                                | 3 (0.09)                                     |
| 1364             | NEIS1364 (NMB1429)          | A0A0E0TSG1           | OM                | PorA (Outer membrane protein IA)  | 3400 (96.98)                                | 52 (1.48)                                    |
| 1428             | NEIS1428 (NMB1497)          | A0A0E0TR48           | OM                | Heme-utilization protein (Hup, TonB-dependent receptor)                                       | 3436 (98.00)                                | 6 (0.17)                                     |
| 1468             | NEIS1468 (NMB1540)          | A0A0E0TRH9           | OM                | Lactoferrin-binding protein A (LbpA)  | 3240 (92.41)                                | 43 (1.23)                                    |
| 1487             | NEIS1487 (NMB1567)          | A0A0E0TRL0           | OM                | Macrophage infectivity potentiator (Peptidyl-prolyl cis-trans isomerasesomerase, Mip/FkpA)    | 3504 (99.94)                                | 0 (0.00)                                     |
| 1632             | NEIS1632 (NMB1714)          | A0A0E0TRR1           | OM                | Multiple transferable resistance system protein (Multidrug efflux pump channel protein, MtrE) | 3501 (99.86)                                | 4 (0.11)                                     |
| 1687             | NEIS1687 (NMB0464)          | A0A0E0TTD6           | OM                | Phospholipase A1 (PldA)   | 2732 (77.92)                                | 729 (20.79)                                  |
| 1690             | NEIS1690 (NMB0461)          | A0A0E0TSK9           | OM                | Transferrin-binding protein 1 (TbpA)  | 3268 (93.21)                                | 2 (0.06)                                     |
| 1783             | NEIS1783 (NMB0382)          | A0A0E0TTL2           | OM                | Outer membrane protein class 4 (OmpA family protein, RmpM)                                    | 3505 (99.97)                                | 0 (0.00)                                     |
| 1963             | NEIS1963 (NMB1988)          | A0A0E0TNL5           | OM                | Enterobactin receptor FetA (FetA)   | 3384 (96.52)                                | 52 (1.48)                                    |
| 2020             | NEIS2020 (NMB2039)          | A0A0E0TSH4           | OM                | Major outer membrane protein PIB (PorB)   | 3468 (98.92)                                | 1 (0.03)                                     |
| 2083             | NEIS2083 (NMB0375)          | A0A0E0TU30           | OM                | Adhesin MafA family (Putative lipoprotein MafA)   | 3046 (86.88)                                | 194 (5.53)                                   |
| 2198             | NEIS2198 (NMB1053)          | A0A0E0QB2            | OM                | Class 5 outer membrane protein (OpcA)   | 2217 (63.23)                                | 1282 (36.57)                                 |

EC, extracellular; IM, inner membrane; PP, periplasmic.



used to make the MeNZB™ vaccine. Every unique amino acid sequence variant subsequently identified was assigned a unique variant number, in order of discovery.

### Peptide variant designation and caveats

For each component of the OMVT scheme, peptide variants were assigned from deduced peptide sequences of the respective NEIS locus. Where the NEIS locus was a complete coding sequence, i.e. containing a start and terminal stop codons with the number of base pairs (bp) being a multiple of three and no internal stop codons, each unique peptide sequence was assigned a unique arbitrary variant number in order of identification. Where the gene contained a mutation generating a frameshift, indel, or an internal stop codon more than 50 bp from the consensus stop codon, then variant 0 (null) was assigned, following the deduction that no functional protein would be produced from such loci. Where internal stop codons occurred fewer than 50 bp from the consensus stop codon, the peptide variants were assigned. Where no NEIS or NEISp variant could be deduced, due to incomplete sequencing or genome assembly, no variant was assigned.

### OMVTs

For OMVT profiles with all 24 loci present, each unique combination was assigned an integer, the OMVT. For example, OMVT-1 corresponded to the OMV peptide variants present in the NZ05/33 genome. All subsequent OMVTs were numbered in order of identification. Although OMVTs provided a rapid means of identifying isolates with identical profiles, there were related isolates that differed at  $\leq 23$  loci. To analyse these relationships among OMVTs, a clustering method was applied using eBURSTv3, which determined non-overlapping groups of closely-related strains, i.e. no OMVT was allocated to more than one OMVT cluster, and entry into a cluster required matching at  $\geq 18$  loci to the central OMVT<sup>34</sup>. To ensure stability of OMVT clusters, a central OMVT was defined as the OMVT that differed from the largest number of other OMVTs at only a single peptide sequence. This was also used to name the cluster, for example OMVT-449 was the central OMVT of the OMVT-449 cluster. Bootstrapping for each group was assessed using the 23-peptide cut-off.

The OMVT scheme was used to catalogue the diversity of IMD isolates from the Meningitis Research Foundation Meningococcus Genome Library (MRF-MGL) and related isolate collections. This included 3506 WGS of all culture-confirmed meningococcal isolates in epidemiological years 2010/11 to 2016/17 from England, Wales, Scotland, and Northern Ireland, representing approximately 55% of all laboratory-confirmed IMD in these regions<sup>24,33</sup>. All isolates in the collection were automatically annotated for the OMVT variants, with manual curation of new variants. OMVTs and OMVT clusters were assigned accordingly.

### Cluster analysis

GrapeTree software<sup>35</sup> was used to cluster allelic profiles from large collections of WGS and visualise these relationships. It provides an interactive, web-based interface associated with relevant metadata including: year, serogroup, cc, and BAST. GrapeTree was run through the PubMLST.org plug-in.

### Statistical analysis

All statistical analyses were performed using R version 3.2.4. Cramer's V coefficient was used to assess the association of peptide loci with cc and was calculated using the 'Cramer's V' function in the 'lsr' package (v0.5) in R. The diversity of each OMVT protein was assessed using Shannon's index of diversity (H) and the "adjusted diversity" calculated from the natural log of the number of peptide variants/amino acid/isolate. H was calculated using the 'Vegan' package (v2.4-v2.6) in R, H represents the uncertainty in predicting the peptide variant of a new isolate, given the number of peptide variants and the evenness in abundance of isolates possessing each variant. H increases as richness (number of variants) and evenness (distribution) in the population increases.

### Results


#### The proteome of the OMV from Bexsero®

A total of 461 proteins, representing the proteome of the NZ98/254 OMV, were unambiguously identified in this study (Supplementary Table 1). The predicted subcellular location of each protein revealed 282 cytoplasmic proteins, 60 inner membrane or periplasmic proteins, 36 outer membrane or extracellular proteins and 83 that could not be definitively assigned to any location. Other proteomic analysis of OMVs have found between 100-300 proteins<sup>25-27</sup>, with an increasing number of proteins identified in the OMVs presumably due to technical advances improving the sensitivity of mass spectrometers. All studies share the finding that membrane and periplasmic proteins dominate the proteome in terms of abundance. The peak areas of the top three most abundant peptides were used to provide a relative quantity of each of the proteins. By summing the peak areas of all the proteins from each location, the protein abundance in each sub-cellular location was determined with 58% of the total protein from the outer membrane, 6% from either the inner membrane or periplasm, 32% cytoplasmic proteins, and 3% without a location defined. The 24 proteins chosen for the OMVT scheme encompass 60% of the total protein found in the OMVs (Figure 2).

#### Peptide variants in OMVT scheme

Peptide variants (including null, 0) were annotated in  $\geq 98\%$  of the WGS data of the 3506 UK isolates. As a consequence of incomplete gene assembly, annotation was lower for: Irp (NEISp0944), 3267/3506 isolates (93.2%); LbpA (NEISp1468), 3283/3506 (93.6%); TbpA (NEISp1690), 3270/3506 (93.3%); and MafA (NEISp2083), 3240/3506 (92.4%), (Table 1). H was used to assess the richness and evenness of variants, with TbpA the most diverse antigen (H=3.94) and putative periplasmic protein NEISp1063 the least diverse (H=1.04) (Figure 3, Table 2). The OMPs were more diverse than cytoplasmic proteins ( $p < 0.00001$ ). The adjusted diversity for each locus ranged from -10.3 for FbpA (NEISp0578) to -8.1 for TbpA (NEISp1690) and PorB (NEISp2020) (Figure 3, Table 2).

The distribution of OMVT protein variants within the UK IMD isolates was non-random. A cumulative frequency distribution of the peptide variants for each locus, for the 3506 IMD isolates, demonstrated the loci for which four variants are found in  $>90\%$  of isolates including: FbpA (NEISp0578); putative periplasmic



**Figure 2. Relative amount of protein as determined from peak area of tandem mass spectrometry data.** Proteins were assigned to a group using CELLO<sup>29,30</sup> and PSORTb<sup>28</sup> informatic tools, where the predictions differed proteins were analysed manually using UNIPROT and available published data to define a location. Where a location could not be predicted proteins were assigned an unknown status.


protein (NEISp1063); Mip/FkpA (NEISp1487); and PldA (NEISp1687) (Figure 4). Other loci, including TbpA, LbpA, FetA, PorA, and PorB, showed more diversity and variants were distributed more evenly across the 3506 isolates. The distribution of the variation within each protein was also non-random, i.e. the variation was not distributed evenly across the amino acid sequence. In many OMVT proteins there were regions of high diversity concentrated in the so-called “variable regions”, representing surface-exposed loops under immune selection, previously well-characterised for TbpA, PorB, and PorA<sup>36–38</sup> (Figure 5).

#### Distribution of OMVTs in IMD and association with cc

The OMVT scheme was applied to the WGS data from the 3506 UK meningococcal isolates. A total of 1752 OMVTs were assigned to 2492/3506 (71.1%) isolates. The remaining 1014 (28.9%) isolates had one or more incomplete OMVT protein-encoding loci and it was not possible to assign an OMVT. The most frequently occurring type was OMVT-1149 in 246/2492 (9.9%) isolates, which was associated with cc11 genogroup W isolates (W:cc11). The most frequent OMVTs represented by 10 or more

isolates were: OMVT-21, 64/2492 (2.6%); OMVT-50, 32/2492 (1.3%); OMVT-1275, 18/2492 (0.7%); OMVT-1161, 16/2492 (0.6%); OMVT-1199 16/2492 (0.6%); and OMVT-1158, 12/2492 (0.5%). There were: 1547 OMVTs that were represented by only one isolate (1547/2492 (62.1%); 133 represented by 2 isolates (266/2492, 10.7%); and 65 represented by 3–10 isolates (275/2492, 11.0%). Analysis of all isolates for which an OMVT was available (71.1%, 2492/3506), demonstrated that OMVT was strongly associated with cc: Cramer’s V=1, with values closer to 1 representing strong association (Figure 6)<sup>35</sup>. To determine whether a different subset of OMV proteins could alter the clustering of isolates, the same 3506 meningococcal genomes were analysed using 50 OMV proteins (OMPs, periplasmic and cytoplasmic), showing similarly strong association with clonal complex (data not shown). To determine the extent of association of individual OMVT proteins with cc, Cramer’s V was calculated for each protein. The values ranged from 0.92 for two TonB-dependent receptors (NEISp0944 and NEISp1428) to 0.34 for the putative periplasmic protein (NEISp1063) (Table 2), suggesting that OMVT proteins with lower Cramer’s V were found across ccs.





**Figure 3. Diversity of each protein in the Outer Membrane Vesicle peptide Typing scheme.** Diversity is measured by Shannon's index of diversity (H) for each peptide locus (indicated by the dark blue dots and left axis) and adjusted diversity (AD), the natural log transformation of the number of variants per amino acid per isolate (indicated by yellow dots and right axis). All data were deduced from a dataset of 3506 UK invasive meningococcal disease isolates. Four cytoplasmic proteins are shown on the left of the x axis for reference, three of which are ribosomal proteins RpsM, RplN, RpsG known to be highly conserved in bacteria. The recombinant proteins in the Bexsero<sup>®</sup> vaccine are marked with an \*, factor-H binding proteins (fHbp), Neisserial Heparin binding antigen (NHBA), *Neisseria* adhesin A (NadA) and Porin A variable region 2 (PorA VR2).

### OMVT clusters and association with BAST

Given the large number of possible OMVTs, they were grouped into OMVT clusters, which differed by up to six loci from a defined central OMVT, i.e. isolates in OMVT-1149 cluster were identical to OMV-1149 for  $\geq 18$  loci. There were 1587/3506 (45.3%) isolates assigned to 27 OMVT clusters. Each OMVT cluster was predominantly associated with a cc and a genogroup (Table 3). The recombinant Bexsero<sup>®</sup> antigens had a strong association with the OMVTs, as measured by Cramer's V (fHbp=0.98; NHBA=0.99; NadA=0.96; PorAVR1=1; PorAVR2=1) and OMVT clusters were also associated with BAST (Cramer's V = 0.96) (Table 3)<sup>23</sup>. These associations were visualised with the GrapeTree clustering algorithm for OMVT clusters and BAST for cc11, cc269, and cc23 (Figure 7a-c)<sup>35</sup>. Some large ccs exhibited a degree of subtracting; for example, cc11 exhibited three OMVT clusters (OMVT-1149 cluster, OMVT-218 cluster and OMVT-1158 cluster). Each OMVT cluster exhibited a non-overlapping repertoire of BASTs.

### Distribution of OMVT variants by OMV vaccine

OMV vaccines can be uniquely identified using OMVTs; for example MeNZB<sup>TM</sup> and Bexsero<sup>®</sup> vaccines are OMVT-1. Each OMVT deduced protein from all 3506 isolates were


compared to the variants present in OMVT-1, a perfect match giving a score of 24. Only 10/3506 (0.3%) of isolates did not share any peptide variants with MeNZB<sup>TM</sup>, these belonged to cc23, cc213, and cc116. The only isolates with a defined cc that had 11 or more shared variants with MeNZB<sup>TM</sup> belonged to cc41/44 (Figure 8a). There were 34 isolates with undefined cc with  $\geq 11$  shared variants with MeNZB<sup>TM</sup>, corresponding to STs that have not been assigned to a cc or isolates that had a partial MLST profile related to cc41/44. The cc41/44 isolates comprised 169 STs. Among these cc41/44 isolates, there was a bimodal distribution of OMVT variants shared with MeNZB<sup>TM</sup>, suggesting two distinct groups: ST-41, ST-1194, ST-485, ST-340, ST-154, and ST-2314 isolates had  $\geq 12$  shared variants, whereas ST-1097, ST-409, ST-6508, and ST-571 isolates had  $\leq 10$  shared variants with MeNZB<sup>TM</sup>. There were 2772/3506 (79%) isolates that shared 1–5 variants with MeNZB<sup>TM</sup>, this was due to the presence of identical variants of: FbpA (NEISp0578); NspA (NEISp0612); putative periplasmic protein (NEISp1063); CsgG family protein (NEISp1066); and Mip/FkpA (NEISp1487). These were found in a high proportion of isolates, independent of cc: NEISp0578 2524/3506 (72.0%); NEISp0612, 1275/3506 (36.4%); NEISp1063 2932/3506 (83.6%); NEISp1066 2115/3506 (60.3%); and NEISp1487 1637/3506 (46.7%).

**Table 2. Outer Membrane Vesicle peptide Typing (OMVT) scheme association with clonal complex and peptide diversity.** Association between individual outer membrane vesicle proteins included in the OMVT scheme and clonal complex (cc) as measured by the Cramer's V statistic, where values approaching one support an association. Measures of diversity of each locus are given as Shannon's index of diversity and adjusted diversity.


| OMVT proteins                            | Cramer's V | Shannon's index of diversity | Adjusted diversity |
|--|------------|------------------------------|--------------------|
| Irp (NEISp0944)                          | 0.92       | 2.84                         | -9.62              |
| Hup (NEISp1428)                          | 0.92       | 2.62                         | -9.55              |
| TbpA (NEISp1690)                         | 0.90       | 3.94                         | -8.12              |
| FetA (NEISp1963)                         | 0.89       | 3.01                         | -8.80              |
| LbpA (NEISp1468)                         | 0.87       | 3.40                         | -8.72              |
| PorA (NEISp1364)                         | 0.83       | 3.19                         | -8.37              |
| LptD (NEISp0275)                         | 0.80       | 3.11                         | -9.43              |
| PorB (NEISp2020)                         | 0.77       | 2.86                         | -8.12              |
| OmpP1 (NEISp0073)                        | 0.74       | 2.33                         | -9.74              |
| MtrE (NEISp1632)                         | 0.73       | 2.46                         | -9.51              |
| OMP85 (NEISp0173)                        | 0.69       | 2.71                         | -9.89              |
| PilQ (NEISp0408)                         | 0.67       | 3.11                         | -9.43              |
| MafA (NEISp2083)                         | 0.63       | 2.16                         | -9.41              |
| M23 peptidase domain protein (NEISp1271) | 0.60       | 2.22                         | -10.08             |
| OpcA (NEISp2198)                         | 0.58       | 1.77                         | -9.81              |
| NspA (NEISp0612)                         | 0.57       | 2.17                         | -9.21              |
| Lipoprotein (NEISp0196)                  | 0.52       | 2.16                         | -9.02              |
| FbpA (NEISp0578)                         | 0.44       | 1.23                         | -10.28             |
| Mip/FkpA (NEISp1487)                     | 0.41       | 1.65                         | -10.03             |
| PldA (NEISp1687)                         | 0.41       | 1.95                         | -9.59              |
| RmpM (NEISp1783)                         | 0.41       | 1.87                         | -9.57              |
| LysM domain protein (NEISp0101)          | 0.41       | 1.92                         | -10.16             |
| CsgG family protein (NEISp1066)          | 0.41       | 1.62                         | -9.46              |
| Putative periplasmic protein (NEISp1063) | 0.34       | 1.04                         | -9.85              |

The same approach was used to analyse other OMV vaccines: MenBvac<sup>®</sup>; MENGOCOC-VA<sup>®</sup>; and the Chilean OMV vaccine, all of which were derived from cc32 isolates and optimised OMV formulations (Figure 8b, c). By identifying variants that occurred at high frequency and across the most prevalent OMVT clusters, the composition of the 24 OMVT proteins was manipulated to improve the number of antigenic matches amongst the set of UK disease isolates. Devising a formulation comprising the protein variants found in the most frequently occurring ccs, which differed at 14/24 loci from OMVT-1, exhibited the highest degree of shared variants to an OMV formulation (Figure 8b). The maximum number of shared variants was 14 (n=40), and all isolates


shared  $\geq 1$  variant. The lower overall number of variant matches was because most isolates shared the loci that are independent of cc, and additionally only a small number of cc-specific variants. Another way to broaden potential vaccine coverage would be to create a vaccine containing multiple OMVTs, for example a formulation representing the five most prevalent hyperinvasive ccs (OMVTs: 160, 368, 1152, 27, and 21). With this formulation all isolates in the reference set shared  $\geq 2$  variants with the proposed vaccine, with 3310/3506 (94.4%) isolates having  $\geq 8$  variant matches, across all ccs, except cc11 (Figure 8c). All cc11 isolates had  $\leq 8$  shared variants with the multiple OMVT vaccine; however, they are highly uniform and the development of a



**Figure 4. Distribution of Outer Membrane Vesicle peptide Typing (OMVT) variants in UK meningococcal disease isolates.** The cumulative frequency distribution is shown for each locus in the OMVT scheme. For each peptide, it shows the proportion of the 3506 isolates that were annotated with each successive peptide variant. The grey dashed line indicates the proportion of isolates within the dataset which possess the four most frequent variants. The distribution is highly skewed, as for FbpA (NEISp0578), Putative periplasmic protein (NEISp1063), Mip/FkpA (NEISp1487) and PldA (NEISp1687) four variants annotated >90% of isolates, contrasting with TbpA for which only 35.6% of isolates possessed the four most frequent peptide variants.



**Figure 5. Variation in the diversity across the amino acid sequence for outer membrane proteins.** The number of allelic variants for each position of the protein are shown for the three most diverse proteins on the left (a) TbpA (NEISp1690), (b) PorB (NEISp2020), (c) PorA (NEISp1364) and the three most conserved (d) M23 peptidase domain protein (NEISp1271), (e) Putative peptidoglycan-binding periplasmic protein, LysM domain-containing protein (NEISp0101), (f) FbpA (NEISp0578). Variable regions are shown in yellow for TbpA, PorB and PorA, as previously described<sup>35–37</sup>. There were other proteins that had potential variable regions including well-described protein FetA (NEISp1963), LbpA (NEISp1468) and Ton-B dependent receptor (NEISp1428).




**Figure 6. UK meningococcal disease isolates clustered by the Outer Membrane Vesicle peptide Typing (OMVT) scheme.** Invasive meningococcal disease isolates from epidemiological years 2010 to 2017 (n=3506) analysed using the 24 loci OMVT scheme and visualised using GrapeTree software. The OMVT cluster by clonal complex (cc), which are shown by different colours, with unfilled nodes representing isolates from the other ccs. The size of the circles is proportional to the number of isolates.


**Table 3. OMVT clusters were associated with clonal complex, genogroup and Bexsero® Antigen Sequence Type (BAST), comprised of the other recombinant proteins found in the 4CMenB vaccine Bexsero®.**

| OMVT cluster      | No. of isolates | Clonal complex | Genogroup | % with genogroup | BASTs   |
|-------------------|-----------------|----------------|-----------|------------------|---|
| OMVT 104 Cluster  | 10              | 32             | Y         | 100              | 234, 539  |
| OMVT 1142 Cluster | 14              | 23             | B         | 100              | 4, 19   |
| OMVT 1149 Cluster | 502             | 11             | W         | 99.8             | 2, 3, 50, 59, 68, 116, 153, 179, 787, 820, 898, 913, 915, 917, 922, 928, 1402, 1491, 1762, 1780, 1880 |
| OMVT 1152 Cluster | 26              | 32             | B         | 100              | 5, 61, 62, 108, 110, 138, 155, 391, 535, 925, 1397, 1419, 1701, 1878                                  |
| OMVT 1158 Cluster | 42              | 11             | C         | 97.5             | 3, 8, 38  |
| OMVT 1229 Cluster | 4               | 32             | B         | 100              | 309   |
| OMVT 1247 Cluster | 23              | 461            | B         | 100              | 230, 296, 307, 407, 520, 525, 543, 571, 587, 1009, 1800   |

| OMVT cluster      | No. of isolates | Clonal complex | Genogroup | % with genogroup | BASTs   |
|-------------------|-----------------|----------------|-----------|------------------|---|
| OMVT 1256 Cluster | 12              | 32             | B         | 100              | 56, 88, 160, 195, 618, 918, 926, 1773, 2138   |
| OMVT 1316 Cluster | 29              | 162            | B         | 100              | 244, 246, 286, 326, 374, 502, 568, 807, 873   |
| OMVT 1341 Cluster | 16              | 32             | B         | 100              | 10, 178, 1764   |
| OMVT 151 Cluster  | 9               | 35             | B         | 100              | 290, 445  |
| OMVT 160 Cluster  | 67              | 41/44          | B         | 100              | 220, 223, 226, 229, 231, 236, 239, 280, 287, 312, 316, 347, 438, 448, 515   |
| OMVT 186 Cluster  | 10              | 269            | B         | 100              | 238, 268, 313, 319, 591   |
| OMVT 21 Cluster   | 240             | 23             | Y         | 99.1             | 221, 225, 228, 233, 357, 368, 387, 427, 444, 499, 579, 628, 964, 1473, 1792, 1824, 1954, 2498   |
| OMVT 211 Cluster  | 5               | 35             | B         | 100              | 257, 293, 320   |
| OMVT 218 Cluster  | 53              | 11             | C         | 84.3             | 126, 142, 235, 242, 521, 522, 546   |
| OMVT 257 Cluster  | 4               | 41/44          | B         | 100              | 220   |
| OMVT 27 Cluster   | 91              | 213            | B         | 100              | 224, 245, 258, 264, 304, 310, 324, 328, 333, 335, 336, 346, 356, 369, 383, 395, 400, 403, 417, 471, 504, 509, 517, 518, 537, 552, 556, 608, 617, 623, 634, 2879, 2882 |
| OMVT 368 Cluster  | 91              | 269            | B         | 100              | 222, 238, 249, 251, 252, 277, 358, 379, 455, 468, 480, 501, 550, 595, 602, 668, 973, 1413, 1505, 1766   |
| OMVT 3782 Cluster | 196             | 269            | B         | 100              | 219, 254, 267, 273, 291, 295, 431, 446, 452, 460, 473, 524, 536, 583, 598, 611, 616, 624, 633, 942, 946, 947, 953, 1273, 1424, 1816                                   |
| OMVT 414 Cluster  | 34              | 41/44          | B         | 100              | 220, 223, 226, 240, 260, 278, 562, 589, 2874  |
| OMVT 449 Cluster  | 47              | 22             | B         | 95.5             | 265, 300, 301, 330, 349, 386, 429, 440, 466, 1765, 1776   |
| OMVT 627 Cluster  | 17              | 269            | B         | 62.5             | 222, 249, 360, 405, 423, 485, 491, 510, 604, 1414, 1821   |
| OMVT 654 Cluster  | 11              | 103            | C         | 72.7             | 288, 311, 541, 544, 553, 566  |
| OMVT 905 Cluster  | 13              | 167            | Y         | 100              | 261, 461, 577, 1406, 1805   |
| OMVT 912 Cluster  | 5               | 23             | Y         | 100              | 228   |
| OMVT 932 Cluster  | 16              | 174            | Y         | 80               | 11, 14, 72, 117, 127, 135, 146, 177   |



**Figure 7. Outer Membrane Vesicle peptide Typing (OMVT) clusters and association with Bexsero® Antigen Sequence Types (BASTs).** (a) Clonal complex 11 (cc11) isolates (n=777) were analysed using the OMVT scheme. (i) Clusters of OMVTs that match 18 loci to the central OMVT are shown. Cc11 has 3 OMVT clusters, 1149, 218 and 1158. OMVT 1149 cluster represents lineage 11.2, OMVT clusters 218 (yellow) and 1158 (red) represent lineage 11.1. (ii) OMVT clusters are coloured by BASTs, demonstrating the strong non-overlapping association of BASTs with OMVT clusters. Unfilled nodes represent incomplete data, so a profile could not be assigned. (b) Cc269 isolates (n=500) were analysed using the OMVT scheme. (i) Cc269 has 4 OMVT clusters. (ii) OMVT clusters are coloured by BAST. (c) Cc23 isolates (n=459) were analysed using the OMVT scheme. (i). Cc23 has 3 OMVT clusters. (ii) OMVT clusters are coloured by BAST.



**Figure 8. The distribution of UK meningococcal disease isolates that match the peptide variants in different vaccine formulations.** Isolates were scored at each of the 24 loci that shared peptide sequence variants with real (a) and hypothetical vaccine formulations (b–d). (a) OMV vaccine MeNZB™ clonal complex 41/44 (cc41/44), (b) selected OMVT protein variants that occur at high frequency in this dataset, (c) engineered OMV including multiple OMVTs (-160, -368, -1152, -27, -21), (d) OMVT-1149 based vaccine (cc11). The disease isolates (n=3506) were stratified by cc (n shown after each cc) and then the number of matches, 1–24, are displayed on the right (n shown after the number of matches). The thickness of the lines is proportional to the number of isolates. Undefined cc represents STs that have not been assigned to a cc or isolates that had a partial multilocus sequence typing profile.

single-strain cc11 OMV vaccine, using for example OMVT-1149, would provide good coverage (Figure 8d), similar to that demonstrated by previously developed OMV vaccines in Cuba, Norway, and New Zealand.

## Discussion

The diversity of the meningococcus and the complexity of its infection biology<sup>39</sup>, in combination with fears of the possible

harmful effects of vaccines containing the serogroup B capsular antigen<sup>40</sup>, have hindered the development of a universal meningococcal childhood vaccine that comprises one or a few components<sup>12</sup>. The meningococcus is an ‘accidental’ pathogen, as disease plays no part in onward transmission and, indeed, is inimical to it<sup>39</sup>. While disease rates are relatively low, asymptomatic carriage rates can be very high, especially in young adults in high-income settings<sup>41</sup> and interruption of transmission is an

important element of the efficacy of conjugate polysaccharide vaccines<sup>42</sup>. OMV vaccines have been used a number of times, with evidence for efficacy against disease caused by the outbreak strain in Cuba<sup>43</sup>, Chile<sup>44</sup>, Norway<sup>45</sup>, New Zealand<sup>46</sup> and France<sup>47</sup>. Evidence for the efficacy of OMV vaccines against carriage is less compelling. Although Bexsero<sup>®</sup> was designed as a four-component vaccine<sup>20</sup>, the inclusion of the MeNZB<sup>™</sup> OMV<sup>48</sup> complicates assessment of its impact. At the time of writing, the vaccine was undergoing post-implementation investigation in the UK<sup>18,19,49</sup> and evaluation of its impact on carriage in Australia<sup>50</sup>. The OMVT scheme was devised and evaluated to assist in the assessment of the diversity of OMV vaccine components.

The majority of the OMVT scheme components were OMPs, constituting approximately 60% of the total protein content<sup>51</sup> known to be present in the meningococcal outer membrane, and therefore potentially exposed to diversifying selection imposed by immune responses, although the most diverse components Pil and Opa were too variable to be included in the scheme<sup>52</sup>. Consistent with this view, the cytoplasmic proteins present in the OMV formulation, three of which were ribosomal subunit proteins, were the least diverse; however, among the OMPs, there was a wide range of diversity. The least diverse proteins, a putative periplasmic protein (NEISp1063) and FbpA, (NEISp0578), are likely not to have surface-exposed regions and might also be influenced by functional constraints. In the most diverse proteins, TbpA (NEISSp1690), PorA (NEISSp1364) and PorB (NEISSp2020), peptide sequence variability was localised in particular 'variable regions' (VRs), shown in a number of studies to represent surface-exposed parts of the sequence subject to immune attack by bactericidal antibodies<sup>36–38</sup>. As our knowledge and understanding of the OMV proteome grows, in terms of understanding protein function, location and immunological relevance to disease and cross-protection, the OMVT scheme could be amended to remove or add relevant antigens.

As reported previously for a number of meningococcal protein antigens, the variable components of the OMVT were non-randomly distributed among isolates, with the most diverse antigens exhibiting a non-overlapping structure<sup>53,54</sup>. The persistence of particular non-random, non-overlapping combinations of PorA and FetA antigens, a few of which attain high frequency in the population<sup>55</sup>, has been used as evidence to support the existence of stable stain types in the meningococcal population driven by immune and/or metabolic selection<sup>53,54</sup>. This structuring was especially exhibited by the more diverse proteins in the OMVT scheme, with the exception of TbpA (NEISSp1690) and LbpA (NEISSp1468). Thus, although there were a very large number of variants of individual proteins and combinations of these variants, they grouped into a relatively small number of clusters, similar to those observed with FetA and PorA<sup>55</sup> and the BAST scheme<sup>23</sup>. In addition, as with PorA:FetA types<sup>9</sup> and BASTs<sup>24</sup>, OMVT clusters were strongly associated with ccs and capsular group. Ccs correspond to lineages within the meningococcal population, several of which have an increased propensity to cause invasive disease. These are referred to as the hyperinvasive lineages<sup>39</sup> and they dominate collections of isolates from IMD, although they are rarer in isolates obtained from asymptomatic carriage<sup>9</sup>. A number of mechanisms have been

proposed to explain the existence of persistence of these population structures in the highly recombinogenic meningococcus<sup>56</sup>, but whatever the mechanism that generates them, these associations can be exploited in vaccine formulation, potentially simplifying it<sup>10</sup>.

Indexing of the variation in the OMVTs in the context of representative isolate collections, such as the MRF-MGL<sup>33</sup>, along with the analysis tools embedded within the [PubMLST.org](http://PubMLST.org) website, enabled the *in silico* assessment of the likely performance of different OMV formulations. The OMVT from a given isolate could be rapidly determined and the likely content of an OMV vaccine made from the isolate deduced. The number of exact protein matches present in a given meningococcal population to this vaccine could then be readily established, indicating the likely degree of coverage that vaccine might attain. This analysis only considered exact matches to vaccine OMP variants as cross-protection is not well characterised for antigens beyond PorA and FetA. Analysis of the UK reference dataset showed that the MeNZB<sup>™</sup> OMV exhibited a similar number of exact matches to a putative cc11 OMV, with higher levels of exact matches attainable by alternative vaccine formulations or the inclusion of multiple OMVs in a single vaccine formulation. In the absence of evidence for very broad cross-protection, these comparisons suggest that multi-component or 'designer' OMVs that contained artificial repertoires of OMPs could have substantial advantages in terms of vaccine coverage, though this requires an understanding of regional meningococcal epidemiology with ongoing surveillance and the additional lipoproteins present in native OMV preparations.

Using a similar approach as the previously described MLST<sup>6</sup> and BAST<sup>23</sup> schemes, the OMVT scheme represents a portable and scalable means of cataloguing variation in the OMV components of meningococcal vaccines<sup>12</sup>. This provides a stable basis for the comparison of the extensive variation of these molecules. The OMVT scheme has the potential to be enhanced with: (i) bioinformatic approaches that predict B- and T-cell epitopes<sup>57</sup>; and (ii) phenotypic expression and immunological data, to provide a more sophisticated prediction of likely cross-protection provided by these vaccines<sup>58</sup>; however, while a peptide sequence catalogue can be comprehensive, this is unlikely to be the case for phenotypes and some deduction from sequence data is likely to remain essential for the foreseeable future. Correlation of the OMVT with immune responses in infants and those that affect carriage in older individuals is of particular importance in assessing the cost-effectiveness of novel meningococcal vaccines<sup>18</sup>. The embedding of the OMVT scheme within the [PubMLST.org/nesseria](http://PubMLST.org/nesseria) database<sup>59</sup> enables the scheme to be implemented genus-wide, permitting inter-species as well as intra-species diversity to be explored. This is of particular importance given the renewed interest in the development of novel vaccines against *Neisseria gonorrhoeae*, the gonococcus, and the possible impact of Bexsero<sup>®</sup> on gonococcal infection<sup>60</sup>. Intra-species diversity is also important when considering the possible impact of such vaccines on the oropharyngeal microbiota. Finally, this approach exemplifies how WGS data can be used to support the development and evaluation of complex multicomponent vaccines against highly variable pathogens.



## Data availability

Peptide sequences generated using Nano-LC-MS/MS have been deposited to the ProteomeXchange Consortium via the PRIDE partner repository, accession number PXD011622. DOI: <https://doi.org/10.6019/PXD011622>.

## Grant information

This study was funded by the Wellcome Trust (109031 and 087622).

*The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.*

## Acknowledgements

This study used the Meningitis Research Foundation Meningococcus Genome Library (<http://www.meningitis.org/research/genome>), which was developed in a collaborative effort by Public Health England, the Wellcome Trust Sanger Institute, and the University of Oxford. We acknowledge Sunil Maharjan and Oliver Lam of NIBSC for their technical assistance in the proteomics study. We thank the staff at the national reference laboratories that allow this endeavour to continue, and Glaxo-SmithKline (Siena, Italy) for providing vaccine grade material for the proteomic analysis. This study used the Neisseria PubMLST website (<http://pubmlst.org/neisseria/>), developed by Keith Jolley and sited at the University of Oxford.

## Supplementary material

**Supplementary Table 1. List of proteins identified from the NZ98/254 OMV component of Bexsero<sup>®</sup> identified using multi-dimensional LC-MS/MS peptide sequencing and database searching (*N. meningitidis* MC58, Swiss-Pro FASTA release 2014\_02).**

Following identification, protein FASTA sequences were submitted to a web server for cellular location prediction using two tools, subCELular Localisation prediction (<http://cello.life.nctu.edu.tw/cello2go/>) and PSORTb<sup>28</sup>. Peak area of the top three most abundant peptides were used to represent the quantity of the protein identified. OM=Outer-membrane, EC=Extracellular, PP=Periplasmic, CP=Cytoplasmic, IM=Inner-membrane. Protein number marked with \* were selected as part of the OMV peptide Typing (OMVT) scheme in this study. The OMV components of both NZ98/254 (MeNZB<sup>TM</sup>) and Bexsero<sup>®</sup> are detergent-extracted with a subsequent loss of fHbp when compared to a native OMV composition. Analysis of the MC58 and NZ98/254 meningococcal reference genomes on PubMLST.org revealed that Opa genes are absent in these strains.

[Click here to access the data](#)

## References

- Okwo-Bele JM, Cherian T: **The expanded programme on immunization: a lasting legacy of smallpox eradication.** *Vaccine*. 2011; **29** Suppl 4: D74–D9. [PubMed Abstract](#) | [Publisher Full Text](#)
- Maiden MC: **The impact of protein-conjugate polysaccharide vaccines: an endgame for meningitis?** *Philos Trans R Soc Lond B Biol Sci*. 2013; **368**(1623): 20120147. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Davenny T, McShane H: **Why don't we have an effective tuberculosis vaccine yet?** *Expert Rev Vaccines*. 2016; **15**(8): 1009–13. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Gordon DM, McGovern TW, Krzych U, *et al.*: **Safety, immunogenicity, and efficacy of a recombinantly produced *Plasmodium falciparum* circumsporozoite protein-hepatitis B surface antigen subunit vaccine.** *J Infect Dis*. 1995; **171**(6): 1576–85. [PubMed Abstract](#) | [Publisher Full Text](#)
- Stephens DS, Greenwood B, Brandtzaeg P: **Epidemic meningitis, meningococcaemia, and *Neisseria meningitidis*.** *Lancet*. 2007; **369**(9580): 2196–210. [PubMed Abstract](#) | [Publisher Full Text](#)
- Maiden MC, Bygraves JA, Feil E, *et al.*: **Multilocus sequence typing: a portable approach to the identification of clones within populations of pathogenic microorganisms.** *Proc Natl Acad Sci U S A*. 1998; **95**(6): 3140–5. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Maiden MC, Jansen van Rensburg MJ, Bray JE, *et al.*: **MLST revisited: the gene-by-gene approach to bacterial genomics.** *Nat Rev Microbiol*. 2013; **11**(10): 728–36. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Bratcher HB, Corton C, Jolley KA, *et al.*: **A gene-by-gene population genomics platform: de novo assembly, annotation and genealogical analysis of 108 representative *Neisseria meningitidis* genomes.** *BMC Genomics*. 2014; **15**: 1138. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Yazdankhah SP, Kriz P, Tzanakaki G, *et al.*: **Distribution of serogroups and genotypes among disease-associated and carried isolates of *Neisseria meningitidis* from the Czech Republic, Greece, and Norway.** *J Clin Microbiol*. 2004; **42**(11): 5146–53. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Urwin R, Russell JE, Thompson EA, *et al.*: **Distribution of surface protein variants among hyperinvasive meningococci: implications for vaccine design.** *Infect Immun*. 2004; **72**(10): 5955–62. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Jolley KA, Brehony C, Maiden MC: **Molecular typing of meningococci: recommendations for target choice and nomenclature.** *FEMS Microbiol Rev*. 2007; **31**(1): 89–96. [PubMed Abstract](#) | [Publisher Full Text](#)
- Feavers IM, Maiden MCJ: **Recent Progress in the Prevention of Serogroup B Meningococcal Disease.** *Clin Vaccine Immunol*. 2017; **24**(5): pii: e00566-16. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kaaijk P, van Straaten I, van de Waterbeemd B, *et al.*: **Preclinical safety and immunogenicity evaluation of a nonavalent PorA native outer membrane vesicle vaccine against serogroup B meningococcal disease.** *Vaccine*. 2013; **31**(7): 1065–71. [PubMed Abstract](#) | [Publisher Full Text](#)
- Zlotnick GW, Jones TR, Liberator P, *et al.*: **The discovery and development of a novel vaccine to protect against *Neisseria meningitidis* Serogroup B Disease.** *Hum Vaccin Immunother*. 2015; **11**(1): 5–13. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

15. Vernikos G, Medini D: **Bexsero® chronicle**. *Pathog Glob Health*. 2014; **108**(7): 305–16.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
16. Giuliani MM, Adu-Bobie J, Comanducci M, *et al.*: **A universal vaccine for serogroup B meningococcus**. *Proc Natl Acad Sci U S A*. 2006; **103**(29): 10834–9.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
17. Brehony C, Wilson DJ, Maiden MC: **Variation of the factor H-binding protein of *Neisseria meningitidis***. *Microbiology*. 2009; **155**(Pt 12): 4155–69.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
18. Ladhani SN, Ramsay M, Borrow R, *et al.*: **Enter B and W: two new meningococcal vaccine programmes launched**. *Arch Dis Child*. 2016; **101**(1): 91–5.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
19. Parikh SR, Andrews NJ, Beebejaun K, *et al.*: **Effectiveness and impact of a reduced infant schedule of 4CMenB vaccine against group B meningococcal disease in England: a national observational cohort study**. *Lancet*. 2016; **388**(10061): 2775–82.  
[PubMed Abstract](#) | [Publisher Full Text](#)
20. Serruto D, Bottomley MJ, Ram S, *et al.*: **The new multicomponent vaccine against meningococcal serogroup B, 4CMenB: immunological, functional and structural characterization of the antigens**. *Vaccine*. 2012; **30** Suppl 2: B87–B97.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
21. Donnelly J, Medini D, Boccadifuoco G, *et al.*: **Qualitative and quantitative assessment of meningococcal antigens to evaluate the potential strain coverage of protein-based vaccines**. *Proc Natl Acad Sci U S A*. 2010; **107**(45): 19490–5.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
22. Vogel U, Taha MK, Vazquez JA, *et al.*: **Predicted strain coverage of a meningococcal multicomponent vaccine (4CMenB) in Europe: a qualitative and quantitative assessment**. *Lancet Infect Dis*. 2013; **13**(5): 416–25.  
[PubMed Abstract](#) | [Publisher Full Text](#)
23. Brehony C, Rodrigues CMC, Borrow R, *et al.*: **Distribution of Bexsero® Antigen Sequence Types (BASTs) in invasive meningococcal disease isolates: Implications for immunisation**. *Vaccine*. 2016; **34**(39): 4690–7.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
24. Rodrigues CMC, Lucidarme J, Borrow R, *et al.*: **Genomic Surveillance of 4CMenB Vaccine Antigenic Variants among Disease-Causing *Neisseria meningitidis* Isolates, United Kingdom, 2010–2016**. *Emerg Infect Dis*. 2018; **24**(4): 673–82.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
25. Tani C, Stella M, Donnarumma D, *et al.*: **Quantification by LC-MS<sup>E</sup> of outer membrane vesicle proteins of the Bexsero® vaccine**. *Vaccine*. 2014; **32**(11): 1273–9.  
[PubMed Abstract](#) | [Publisher Full Text](#)
26. Lappann M, Otto A, Becher D, *et al.*: **Comparative proteome analysis of spontaneous outer membrane vesicles and purified outer membranes of *Neisseria meningitidis***. *J Bacteriol*. 2013; **195**(19): 4425–35.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
27. Masforro Y, Gil J, García D, *et al.*: **A deeper mining on the protein composition of VA-MENGOC-BC®: An OMV-based vaccine against *N. meningitidis* serogroup B and C**. *Hum Vaccin Immunother*. 2017; **13**(11): 2548–2560.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
28. Yu NY, Wagner JR, Laird MR, *et al.*: **PSORTb 3.0: improved protein subcellular localization prediction with refined localization subcategories and predictive capabilities for all prokaryotes**. *Bioinformatics*. 2010; **26**(13): 1608–15.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
29. Yu CS, Lin CJ, Hwang JK: **Predicting subcellular localization of proteins for Gram-negative bacteria by support vector machines based on n-peptide compositions**. *Protein Sci*. 2004; **13**(5): 1402–6.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
30. Yu CS, Chen YC, Lu CH, *et al.*: **Prediction of protein subcellular localization**. *Proteins*. 2006; **64**(3): 643–51.  
[PubMed Abstract](#) | [Publisher Full Text](#)
31. Ferrari G, Garaguso I, Adu-Bobie J, *et al.*: **Outer membrane vesicles from group B *Neisseria meningitidis* delta gna33 mutant: proteomic and immunological comparison with detergent-derived outer membrane vesicles**. *Proteomics*. 2006; **6**(6): 1856–66.  
[PubMed Abstract](#) | [Publisher Full Text](#)
32. Petkau A, Stuart-Edwards M, Stothard P, *et al.*: **Interactive microbial genome visualization with GViewer**. *Bioinformatics*. 2010; **26**(24): 3125–6.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
33. Hill DM, Lucidarme J, Gray SJ, *et al.*: **Genomic epidemiology of age-associated meningococcal lineages in national surveillance: an observational cohort study**. *Lancet Infect Dis*. 2015; **15**(12): 1420–8.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
34. Feil EJ, Li BC, Aanensen DM, *et al.*: **eBURST: inferring patterns of evolutionary descent among clusters of related bacterial genotypes from multilocus sequence typing data**. *J Bacteriol*. 2004; **186**(5): 1518–30.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
35. Zhou Z, Alikhan NF, Sergeant MJ, *et al.*: **GrapeTree: Visualization of core genomic relationships among 100,000 bacterial pathogens**. *bioRxiv*. 2017; 216788.  
[Publisher Full Text](#)
36. Pajón R, China G, Marrero E, *et al.*: **Sequence analysis of the structural *tbpA* gene: protein topology and variable regions within neisserial receptors for transferrin iron acquisition**. *Microb Pathog*. 1997; **23**(2): 71–84.  
[PubMed Abstract](#) | [Publisher Full Text](#)
37. van der Ley P, Heckels JE, Virji M, *et al.*: **Topology of outer membrane porins in pathogenic *Neisseria* spp**. *Infect Immun*. 1991; **59**(9): 2963–71.  
[PubMed Abstract](#) | [Free Full Text](#)
38. Russell JE, Jolley KA, Feavers IM, *et al.*: **PorA variable regions of *Neisseria meningitidis***. *Emerg Infect Dis*. 2004; **10**(4): 674–8.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
39. Caugant DA, Maiden MC: **Meningococcal carriage and disease—population biology and evolution**. *Vaccine*. 2009; **27** Suppl 2: B64–70.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
40. Finne J, Leinonen M, Makela PH: **Antigenic similarities between brain components and bacteria causing meningitis. Implications for vaccine development and pathogenesis**. *Lancet*. 1983; **2**(8346): 355–7.  
[PubMed Abstract](#) | [Publisher Full Text](#)
41. Christensen H, May M, Bowen L, *et al.*: **Meningococcal carriage by age: a systematic review and meta-analysis**. *Lancet Infect Dis*. 2010; **10**(12): 853–61.  
[PubMed Abstract](#) | [Publisher Full Text](#)
42. Trotter CL, Maiden MC: **Meningococcal vaccines and herd immunity: lessons learned from serogroup C conjugate vaccination programs**. *Expert Rev Vaccines*. 2009; **8**(7): 851–61.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
43. Sierra GVG, Campa CH, Garcia LI, *et al.*: **Efficacy evaluation of the Cuban vaccine VA-MENGOC-BC against disease caused by serogroup B *Neisseria meningitidis***. In: Achtman Mea, editor. *Neisseria 1990*. Berlin: Walter de Gruyter; 1991; 129–34.  
[Reference Source](#)
44. Zollinger WD, Boslego J, Moran E, *et al.*: **Meningococcal serogroup B vaccine protection trial and follow-up studies in Chile. The Chilean National Committee for Meningococcal Disease**. *NIPH Ann*. 1991; **14**(2): 211–2; discussion 213.  
[PubMed Abstract](#)
45. Bjune G, Høiby EA, Grønnesby JK, *et al.*: **Effect of outer membrane vesicle vaccine against group B meningococcal disease in Norway**. *Lancet*. 1991; **338**(8775): 1093–6.  
[PubMed Abstract](#) | [Publisher Full Text](#)
46. Galloway Y, Stehr-Green P, McNicholas A, *et al.*: **Use of an observational cohort study to estimate the effectiveness of the New Zealand group B meningococcal vaccine in children aged under 5 years**. *Int J Epidemiol*. 2009; **38**(2): 413–8.  
[PubMed Abstract](#) | [Publisher Full Text](#)
47. Caron F, du Châtelet IP, Leroy JP, *et al.*: **From tailor-made to ready-to-wear meningococcal B vaccines: longitudinal study of a clonal meningococcal B outbreak**. *Lancet Infect Dis*. 2011; **11**(6): 455–63.  
[PubMed Abstract](#) | [Publisher Full Text](#)
48. Oster P, Lennon D, O'Hallaigh J, *et al.*: **MeNZB: a safe and highly immunogenic tailor-made vaccine against the New Zealand *Neisseria meningitidis* serogroup B disease epidemic strain**. *Vaccine*. 2005; **23**(17–18): 2191–6.  
[PubMed Abstract](#) | [Publisher Full Text](#)
49. Parikh SR, Newbold L, Slater S, *et al.*: **Meningococcal serogroup B strain coverage of the multicomponent 4CMenB vaccine with corresponding regional distribution and clinical characteristics in England, Wales, and Northern Ireland, 2007–08 and 2014–15: a qualitative and quantitative assessment**. *Lancet Infect Dis*. 2017; **17**(7): 754–62.  
[PubMed Abstract](#) | [Publisher Full Text](#)
50. Marshall HS, McMillan M, Koehler A, *et al.*: **B Part of It protocol: a cluster randomised controlled trial to assess the impact of 4CMenB vaccine on pharyngeal carriage of *Neisseria meningitidis* in adolescents**. *BMJ Open*. 2018; **8**(7): e020988.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
51. Tani C, Stella M, Donnarumma D, *et al.*: **Quantification by LC-MS<sup>E</sup> of outer membrane vesicle proteins of the Bexsero® vaccine**. *Vaccine*. 2014; **32**(11): 1273–9.  
[PubMed Abstract](#) | [Publisher Full Text](#)
52. Wachter J, Hill S: **Positive Selection Pressure Drives Variation on the Surface-Exposed Variable Proteins of the Pathogenic *Neisseria***. *PLoS One*. 2016; **11**(8): e0161348.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
53. Buckee CO, Jolley K, Recker M, *et al.*: **Role of selection in the emergence of lineages and the evolution of virulence in *Neisseria meningitidis***. *Proc Natl Acad Sci U S A*. 2008; **105**(39): 15082–7.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
54. Gupta S, Maiden MC, Feavers IM, *et al.*: **The maintenance of strain structure in populations of recombining infectious agents**. *Nat Med*. 1996; **2**(4): 437–42.  
[PubMed Abstract](#) | [Publisher Full Text](#)
55. Watkins ER, Maiden MC: **Persistence of hyperinvasive meningococcal strain types during global spread as recorded in the PubMLST database**. *PLoS One*. 2012; **7**(9): e45349.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
56. Watkins ER, Maiden MC, Gupta S: **Metabolic competition as a driver of bacterial population structure**. *Future Microbiol*. 2016; **11**: 1339–57.  
[PubMed Abstract](#) | [Publisher Full Text](#)

57. Christodoulides M, Heckels J: **Novel approaches to *Neisseria meningitidis* vaccine design.** *Pathog Dis.* 2017; **75**(3): ftx033.  
[PubMed Abstract](#) | [Publisher Full Text](#)
58. Ladhani SN, Flood JS, Ramsay ME, *et al.*: **Invasive meningococcal disease in England and Wales: implications for the introduction of new vaccines.** *Vaccine.* 2012; **30**(24): 3710–6.  
[PubMed Abstract](#) | [Publisher Full Text](#)
59. Jolley KA, Maiden MC: **Automated extraction of typing information for bacterial pathogens from whole genome sequence data: *Neisseria meningitidis* as an exemplar.** *Euro Surveill.* 2013; **18**(4): 20379.  
[PubMed Abstract](#) | [Free Full Text](#)
60. Petousis-Harris H, Paynter J, Morgan J, *et al.*: **Effectiveness of a group B outer membrane vesicle meningococcal vaccine against gonorrhoea in New Zealand: a retrospective case-control study.** *Lancet.* 2017; **390**(10102): 1603–1610.  
[PubMed Abstract](#) | [Publisher Full Text](#)

# Open Peer Review

Current Referee Status:  

---

## Version 1

Referee Report 14 January 2019

<https://doi.org/10.21956/wellcomeopenres.16194.r34347>



**Peter van der Ley**

Institute for Translational Vaccinology (Intravacc), Bilthoven, The Netherlands

The paper describes a novel typing scheme to address the population diversity of meningococcal vaccine antigens. This constitutes an important step forward, as it allows a more "holistic" approach to estimate vaccine coverage. Instead of focusing on one or a few individual antigens, using this method one can assess the overall potential contributions of multiple antigens in OMV-based vaccines.

### Minor comments

1. Considering that the total OMV proteome listed in Supplementary Table 1 has 461 proteins, the selection of just 24 candidates for the scheme seems somewhat arbitrary. It would be helpful to give more discussion on the reasons for this particular selection, and how the results could be influenced by a different selection.
2. In our experience, not all outer membrane proteins are equally susceptible to the trypsin digestion step used during sample preparation for MS analysis, and therefore the details of the method used can influence the quantification results that are obtained. Have the authors compared different methods ?
3. Some well-known vaccine antigens are missing in Supplementary Table 1, in particular the Opa proteins and fHbp. Were these left out deliberately ?
4. Figure 4 contains very interesting information on variation of individual antigens. However, it is very difficult to identify the antigens corresponding to the different lines.
5. The NZ98/254 OMV used is obtained with detergent extraction. However, other methods for OMV preparation have also been described, which avoid the use of detergent extraction. Such native OMVs have different antigen composition, especially with regards to surface-exposed lipoproteins such as the well-known vaccine antigen factor H binding protein. They would therefore require different selections of OMVT antigens. It would be helpful to discuss this in the context of hypothetical novel OMV formulations (Figure 8).

**Is the work clearly and accurately presented and does it cite the current literature?**

Yes

**Is the study design appropriate and is the work technically sound?**

Yes

**Are sufficient details of methods and analysis provided to allow replication by others?**

Yes

**If applicable, is the statistical analysis and its interpretation appropriate?**

Yes

**Are all the source data underlying the results available to ensure full reproducibility?**

Yes

**Are the conclusions drawn adequately supported by the results?**

Yes

**Competing Interests:** No competing interests were disclosed.

**Reviewer Expertise:** Molecular Microbiology

**I have read this submission. I believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.**

Author Response 19 Mar 2019

**Charlene Rodrigues**, University of Oxford, UK

We are grateful for your constructive review and we have addressed all the following points:

Minor comments

1. Considering that the total OMV proteome listed in Supplementary Table 1 has 461 proteins, the selection of just 24 candidates for the scheme seems somewhat arbitrary. It would be helpful to give more discussion on the reasons for this particular selection, and how the results could be influenced by a different selection.

**We have clarified this in the “OMVT scheme development and nomenclature” section:**

*A total of 25 proteins were chosen, primarily for pragmatic reasons as this was considered manageable for detailed curation within the typing scheme. The proteins represented the 20 most abundant outer membrane or periplasmic proteins identified in by Nano-LC-MS/MS with an additional five proteins chosen from the published core proteome 25, also identified by Nano-LC-MS/MS ( Table 1). All proteins chosen for the scheme were predicted to be either outer membrane or periplasmic and therefore more likely to be OMV antigens than the cytoplasmic proteins, which were more abundant in the proteomic analysis. The cytoplasmic proteins are unlikely to be constituents of the meningococcal outer membrane and their presence in OMV samples is a probable consequence of cell lysis during detergent extraction causing cytosolic proteins to be released into the preparations.*

**We have additionally noted in the “Distribution of OMVTs in IMD and association with cc” section:**

*To determine whether a different subset of OMV proteins could alter the clustering of isolates, the same 3506 meningococcal genomes were analysed using 50 OMV proteins (OMPs, periplasmic and cytoplasmic), showing similarly strong association with clonal complex (data not shown).*

2. In our experience, not all outer membrane proteins are equally susceptible to the trypsin digestion step used during sample preparation for MS analysis, and therefore the details of the method used can influence the quantification results that are obtained. Have the authors compared different methods ?

**We have added this to the “Proteomic OMV sample preparation” section:**

*As OMPs are not all equally susceptible to enzymatic digestion, we evaluated different sample preparation methods. The method described provided an increased number of OMPs to other studies in our laboratory and the contribution of membrane to cytoplasmic proteins was comparable to other studies.*

3. Some well-known vaccine antigens are missing in Supplementary Table 1, in particular the Opa proteins and fHbp. Were these left out deliberately ?

**We have added this in the figure legend of Supplementary Table 1:**

*The OMV components of both NZ98/254 (MeNZB™) and Bexsero® are detergent-extracted with a subsequent loss of fHbp when compared to a native OMV composition. Analysis of the MC58 and NZ98/254 meningococcal reference genomes on PubMLST.org revealed that Opa genes are absent in these strains.*

4. Figure 4 contains very interesting information on variation of individual antigens. However, it is very difficult to identify the antigens corresponding to the different lines.

**We have made the markers on the graph larger as well as on the legend, so it will be easier to identify which colour represents the different antigens.**

5. The NZ98/254 OMV used is obtained with detergent extraction. However, other methods for OMV preparation have also been described, which avoid the use of detergent extraction. Such native OMVs have different antigen composition, especially with regards to surface-exposed lipoproteins such as the well-known vaccine antigen factor H binding protein. They would therefore require different selections of OMVT antigens. It would be helpful to discuss this in the context of hypothetical novel OMV formulations (Figure 8).

**We have clarified this in the figure legend of Supplementary Table 1:**

*The OMV components of both NZ98/254 (MeNZB™) and Bexsero® are detergent-extracted with a subsequent loss of fHbp when compared to a native OMV composition.*

**Additionally, the assessment of other OMV proteins has been added to the “Distribution of OMVTs in IMD and association with cc” section:**

*To determine whether a different subset of OMV proteins could alter the clustering of isolates, the same 3506 meningococcal genomes were analysed using 50 OMV proteins (OMPs, periplasmic and cytoplasmic), showing similarly strong association with clonal complex (data not shown).*

**Competing Interests:** No competing interests were disclosed.

Referee Report 18 December 2018

<https://doi.org/10.21956/wellcomeopenres.16194.r34348>



**Mariagrazia Pizza , Margherita Bodini**

GlaxoSmithKline (GSK), Siena, Italy

This manuscript describes the development of an Outer Membrane Vesicles Peptide Typing (OMVT) scheme based on a subset of selected proteins in 3506 UK invasive isolates. The paper is of great interest, well-written, and the methodology used to define the OMVT scheme and the definition of the relative clusters are clearly described. Moreover, the paper represents a milestone in vaccine development, because it provides important insight in the assessment of vaccine impact, improvement of vaccine formulations, and evaluation of inter-species and intra-species diversity.

Minor comments:

- page 4 **OMVT scheme development and nomenclature** paragraph: The criteria used for the selection of five additional proteins from the core proteome, and the basis for the selection of only five should be further clarified.

- page 7 **The proteome of the OMV from Bexsero** paragraph: It would be worth describing the differences between the number of proteins identified from this analysis, compared to what is already published.

- **Figure 4:** The figure would be clearer by increasing the size of the dots to enable the reader to easily identify the distribution for each single antigen. A possibility could be also to zoom in the region of the dashed line, related to the four most common variants.

- **Figure 5:** Only results related to distribution of variation for proteins known to contain variable regions and for three most conserved proteins are shown in Figure 5. However, it would be interesting to add a comment to what happens in-between: are there other peptides containing variation patterns compatible with variable regions?

- **Discussion** at page 16 and **Abstract:** The sentence “[...]has been used as evidence to support the existence of stable strain types in the meningococcal population driven by immune and metabolic selection 50,51]” should read “immune and/or metabolic selection”. The same applies to the abstract: ..... “is consistent with strain structure models invoking metabolic and/or host immune selection”.

- **Discussion:** The comparison between potential coverage of real and hypothetical vaccines is very interesting. However, it could be more appropriate to clarify a few points in the discussion section:

- The differences between antigens in which a perfect match is needed to confer protection (e.g. PorA) and antigens for which the influence that sequence variability could have on the cross-protective ability is unknown, and should be further emphasized.
- The ability of additional antigens to confer cross-protection and not included in the OMV analysis, should be mentioned and further discussed. It would be interesting to evaluate in future how much these antigens cluster with OMVT scheme.
- The sentence on page 16 : “In the absence of evidence for a very broad cross-protection[...]”, highlights the importance of this study and raises a very interesting concept related to a multi-component or “designer” OMV vaccine. It would be interesting to briefly introduce also the possible limitations that such an approach may have.

**Is the work clearly and accurately presented and does it cite the current literature?**

Yes

**Is the study design appropriate and is the work technically sound?**

Yes

**Are sufficient details of methods and analysis provided to allow replication by others?**

Yes

**If applicable, is the statistical analysis and its interpretation appropriate?**

Yes

**Are all the source data underlying the results available to ensure full reproducibility?**

Yes

**Are the conclusions drawn adequately supported by the results?**

Yes

**Competing Interests:** MP and MB are full time employee of GSK Vaccines.

**Reviewer Expertise:** Microbiology, Vaccinology

**We have read this submission. We believe that we have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.**

Author Response 19 Mar 2019

**Charlene Rodrigues**, University of Oxford, UK

We welcome your constructive review and have addressed all the points as follows:

Minor comments:

- page 4 OMVT scheme development and nomenclature paragraph: The criteria used for the selection of five additional proteins from the core proteome, and the basis for the selection of only five should be further clarified.

**We have clarified this in the "OMVT scheme development and nomenclature" section:**

*A total of 25 proteins were chosen, primarily for pragmatic reasons as this was considered manageable for detailed curation within the typing scheme. The proteins represented the 20 most abundant outer membrane or periplasmic proteins identified in by Nano-LC-MS/MS with an additional five proteins chosen from the published core proteome 25, also identified by Nano-LC-MS/MS ( Table 1). All proteins chosen for the scheme were predicted to be either outer membrane or periplasmic and therefore more likely to be OMV antigens than the cytoplasmic proteins, which were more abundant in the proteomic analysis. The cytoplasmic proteins are unlikely to be constituents of the meningococcal outer membrane and their presence in OMV samples is a probable consequence of cell lysis during detergent extraction causing cytosolic proteins to be released into the preparations.*

- page 7 The proteome of the OMV from Bexsero paragraph: It would be worth describing the differences between the number of proteins identified from this analysis, compared to what is already published.

**We have added the following to the "Proteome of the OMV of Bexsero®" Results section:**

*Other proteomic analysis of OMVs have found between 100-300 proteins (1-3), with an increasing number of proteins identified in the OMVs presumably due to technical advances improving the sensitivity of mass spectrometers. All studies share the finding that membrane and periplasmic proteins dominate the proteome in terms of abundance.*

1. Lappann M, Otto A, Becher D, Vogel U. 2013. Comparative proteome analysis of spontaneous outer membrane vesicles and purified outer membranes of *Neisseria meningitidis*. *Journal of Bacteriology* 195:4425-35.



2. Masforrol Y, Gil J, Garcia D, Noda J, Ramos Y, Betancourt L, Guirola O, Gonzalez S, Acevedo B, Besada V, Reyes O, Gonzalez LJ. 2017. A deeper mining on the protein composition of VA-MENGOC-BC(R): An OMV-based vaccine against *N. meningitidis* serogroup B and C. *Hum Vaccin Immunother* 13:2548-2560.

3. Tani C, Stella M, Donnarumma D, Biagini M, Parente P, Vadi A, Magagnoli C, Costantino P, Rigat F, Norais N. 2014. Quantification by LC-MS(E) of outer membrane vesicle proteins of the Bexsero® vaccine. *Vaccine* 32:1273-9.

- Figure 4: The figure would be clearer by increasing the size of the dots to enable the reader to easily identify the distribution for each single antigen. A possibility could be also to zoom in the region of the dashed line, related to the four most common variants.

**We have made the markers on the graph larger as well as on the legend, so it will be easier to identify which colour represents the different antigens.**

- Figure 5: Only results related to distribution of variation for proteins known to contain variable regions and for three most conserved proteins are shown in Figure 5. However, it would be interesting to add a comment to what happens in-between: are there other peptides containing variation patterns compatible with variable regions?

**We have added the following text to the figure 5 legend:**

*There were other proteins that had potential variable regions including well-described protein NEISp1963 (FetA), NEISp 1468 (LbpA) and NEISp 1428 (Ton-B dependent receptor).*

- Discussion at page 16 and Abstract: The sentence “[...]has been used as evidence to support the existence of stable stain types in the meningococcal population driven by immune and metabolic selection 50,51]” should read “immune and/or metabolic selection”. The same applies to the abstract: ..... “is consistent with strain structure models invoking metabolic and/or host immune selection”.

**Thank you, this has been changed in the updated manuscript.**

- Discussion: The comparison between potential coverage of real and hypothetical vaccines is very interesting. However, it could be more appropriate to clarify a few points in the discussion section:

- The differences between antigens in which a perfect match is needed to confer protection (e.g. PorA) and antigens for which the influence that sequence variability could have on the cross-protective ability is unknown, and should be further emphasized.

**We have added the following to the discussion:**

*This analysis only considered exact matches to vaccine OMP variants as cross-protection is not well characterised for antigens beyond PorA and FetA.*

- The ability of additional antigens to confer cross-protection and not included in the OMV analysis, should be mentioned and further discussed. It would be interesting to evaluate in future how much these antigens cluster with OMVT scheme.

**We have amended the following to include:**

*As our knowledge and understanding of the OMV proteome grows, in terms of understanding protein function, location, and immunological relevance to disease and cross-protection, the OMVT*

*scheme could be amended to remove or add relevant antigens.*

- The sentence on page 16 : "In the absence of evidence for a very broad cross-protection[...]", highlights the importance of this study and raises a very interesting concept related to a multi-component or "designer" OMV vaccine. It would be interesting to briefly introduce also the possible limitations that such an approach may have

**We have added the following limitations of this approach:**

*In the absence of evidence for very broad cross-protection, these comparisons suggest that multi-component or 'designer' OMVs that contained artificial repertoires of OMPs could have substantial advantages in terms of vaccine coverage, though this requires an understanding of regional meningococcal epidemiology with ongoing surveillance and the additional lipoproteins present in native OMV preparations.*

**Competing Interests:** No competing interests were disclosed.