



King Saud University

Saudi Journal of Biological Sciences

www.ksu.edu.sa  
www.sciencedirect.com



الجمعية السعودية لعلم الحياتة  
SAUDI BIOLOGICAL SOCIETY

ORIGINAL ARTICLE

# Phenolic composition and antiparasitic activity of plants from the Brazilian Northeast “Cerrado”



João Tavares Calixto Júnior<sup>a</sup>, Selene Maia de Moraes<sup>a</sup>, Celeste Vega Gomez<sup>b</sup>, Cathia Coronel Molas<sup>b</sup>, Miriam Rolon<sup>b</sup>, Aline Augusti Boligon<sup>d</sup>, Margareth Linde Athayde<sup>d</sup>, Cícera Datiane de Moraes Oliveira<sup>c</sup>, Saulo Relison Tintino<sup>c</sup>, Henrique Douglas Melo Coutinho<sup>c,\*</sup>

<sup>a</sup> Post Graduation Biotechnological Programme – RENORBIO, Laboratory of Natural Products, State University of Ceará, Itaperi Campus, Fortaleza, Ceará, Brazil

<sup>b</sup> Centro para el Desarrollo de la Investigación Científica (CEDIC), Fundación Moisés Bertoní/Laboratorios Díaz Gill, Asunción, Paraguay

<sup>c</sup> Department of Biological Chemistry, Regional University of Cariri, Crato, Ceará State, Brazil

<sup>d</sup> Department of Industrial Pharmacy, Federal University of Santa Maria, Santa Maria, Rio Grande do Sul State, Brazil

Received 11 July 2015; revised 11 October 2015; accepted 13 October 2015

Available online 23 October 2015

## KEYWORDS

*Guazuma ulmifolia*;  
Leishmanicidal activity;  
*Luehea paniculata*;  
*Prockia crucis*;  
Trypanocidal activity

**Abstract** This work describes the antiparasitic and cytotoxic activities of three plant species from the Cerrado biome, Northeastern Brazil. Significant antiparasitic inhibition was observed against *Trypanosoma cruzi* (63.86%), *Leishmania brasiliensis* (92.20%) and *Leishmania infantum* (95.23%) when using ethanol extract from leaves of *Guazuma ulmifolia* Lam. (Malvaceae), at a concentration of 500 µg/mL. However, low levels of inhibition were observed when assessing leishmanicidal and trypanocidal (Clone CL-B5) activities of crude ethanol extracts from leaves and bast tissue of *Luehea paniculata* (Malvaceae) and leaves and bark of *Prockia crucis* (Salicaceae) at a concentration of 500 µg/mL. The extracts revealed the presence of phenolic acids such as gallic acid, chlorogenic acid, caffeic acid and rosmarinic acid, as well as flavonoids such as rutin, luteolin, apigenin and quercetin – the latter detected only in *G. ulmifolia*. *G. ulmifolia* extract displayed higher leishmanicidal activity probably due to the presence of quercetin, a potent known leishmanicidal compound. A cytotoxicity test indicated values over 50% at the highest concentration (1000 µg/mL) for all natural products,

\* Corresponding author at: Laboratório de Microbiologia e Biologia Molecular, Departamento de Química Biológica, Universidade Regional do Cariri – URCA, Rua Cel. Antonio Luis 1161, Pimenta, 63105-000 Crato, Ceará, Brazil. Tel.: +55 (88) 31021212; fax: +55 (88) 31021291. E-mail address: [hdmcoutinho@gmail.com](mailto:hdmcoutinho@gmail.com) (M.C. Henrique Douglas).

Peer review under responsibility of King Saud University.



Production and hosting by Elsevier

which were considered cytotoxic. This points out the need for further tests to enable future *in vivo* trials, including antineoplastic activity on human tumor cells.

© 2015 The Authors. Production and hosting by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

Cerrado vegetation occurs particularly in Brazil's Central Highlands, especially in the states of Mato Grosso, Goiás and Minas Gerais. Disjunct areas were later identified in the states of São Paulo and Minas Gerais (Ferri, 1955) and in Ceará, where this biome was contained almost entirely within the semiarid domain, around the southern part of the state (Figueiredo and Fernandes, 1987). Cerrado features heterogeneous vegetation, with taxonomic biodiversity that is actually higher than that of the Amazon rainforest, including over 7000 native vascular plant species (Hiruma-Lima et al., 2006). Due to this notable floristic diversification, there has been a growing interest in investigating medicinal plants from the Cerrado as sources of bioactive compounds (Guarin Neto and Morais, 2003).

Leishmaniasis is a prevalent disease in 88 countries, located in 4 continents, with an estimated 1.6 million new cases annually, of which 500,000 are visceral and 1.1 million, cutaneous or mucocutaneous (WHO, 2010). In Brazil, leishmaniasis occurs in 19 states, with over 90% of human cases occurring in the Northeast region (Maia-Elkhoury et al., 2008). Rangel and Lainson (2009) state that American cutaneous leishmaniasis in Brazil is caused by a dermatropic variety of *Leishmania*, and the transmission of the causal agents involves different species of phlebotomines (*Diptera: Psychodimilian: Phlebotominae*) in strict relationship with mammal hosts of the parasites, resulting in a number of different transmission cycles nationwide.

Several substances have been tested as anti-*Leishmania* drugs. The main ones are directed toward inhibition of the vital and specific metabolic pathways of the parasite (Figueiredo et al., 2014). To Bezerra et al. (2004), the first-line drugs to treat leishmaniasis have been pentavalent antimonials (Sb5+). Natural products constitute an important source of new medications, because their derivatives are extremely useful as structures for synthetic modification and bioactivity optimization. Flavonoids have also been investigated in this field (Sülsen et al., 2007).

To Chai et al. (2005), the mechanism of action of pentavalent antimonials against *Leishmania* is still disputed and little understood. Liposomal amphotericin B, pentamidine, paromomycin and miltefosine are drugs of interest as they represent new therapeutic alternatives, but feature several problems such as side effects, product price and formula production (Pereira, 2011).

Chagas disease is another widespread parasitic disease occurring worldwide, caused by *Trypanosoma cruzi* and affecting about 10 million people in the Americas (WHO, 2010; Figueiredo et al., 2014). The most epidemiologically important forms of transmission are vectors – through hematophagous insects (triatomines), transfusion, congenital and oral (Coura et al., 2007). *T. cruzi* has a complex biological cycle, involving three evolutive forms (trypomastigote, epimastigote and

amastigote) and several species of triatomines and mammals, wild and domestic, which act respectively as vectors and reservoirs of the parasite (Lana and Tafuri, 2005).

Given the importance and the need to seek new antiparasitic drugs to combat leishmaniasis and Chagas disease, the aim of this study was to evaluate the leishmanicidal and trypanocidal potential as well as the cytotoxicity of crude ethanol extracts from three species of plants with few or almost no reports in the literature about antiparasitic activities – *Luehea paniculata* Mart. and Zucc. (Malvaceae), *Prockia crucis* P. Browne ex. L. (Salicaceae) and *Guazuma ulmifolia* Lam. (Malvaceae), medicinal trees of multiple effects – found in a Cerrado fragment located in the Northeast of Brazil – an area of interest relevant to the understanding of the biological potential of its flora.

## 2. Materials and methods

### 2.1. Place of collection of the plant material and obtaining of extracts

Leaves and bast tissue of *L. paniculata*, leaves and bark of *P. crucis* and leaves of *G. ulmifolia* were collected on November 19, 2012 (dry season), around 17:30 h, from the Cerrado located in the center of Caatinga area, Lavras da Mangabeira city, central mesoregion in South of Ceará, Northeastern Brazil. The area is characterized as a vegetation relic (Figueiredo and Fernandes, 1987), presenting itself in deep soil on a tabular relief on long dissected surfaces with penetration of the savannah flora. The patch of savannah in question is located in hills of 280–600 m in height, at the following coordinates: (06° 72' 2432' S and 38° 97' 7396' W). Botanical material was collected and exsiccates were obtained, which were deposited at the Prisco Bezerra Herbarium, Federal University of Ceará (UFC), Fortaleza, Ceará, Brazil, as per the numbering described in Table 1. The plant material was dried at 60 °C, weighed (Table 2) and subjected to maceration with ethanol PA for seven days. After this period, filtration and concentration of the extract were made distilling all the solvent on a rotary evaporator under reduced pressure (40 rpm at 60 °C) and then in a water bath at 60 °C, obtaining the crude extracts.

**Table 1** Plant species and parts used in the antiparasitic and cytotoxicity tests.

Species	Family	Plant part	Exsiccate
<i>Luehea paniculata</i> Mart. and Zucc.	Malvaceae	Leaves and bast tissue	54.641
<i>Prockia crucis</i> P. Browne ex. L.	Salicaceae	Leaves and bark	54.741
<i>Guazuma ulmifolia</i> Lam.	Malvaceae	Leaves	54.743

**Table 2** Dry weight and yield of the crude ethanol extracts.

Natural product	Solvent	Dry matter (g)	Crude extract (g)	Extract yields (%)
EELPL	Ethanol	595	61.649	10.36
EELPB	Ethanol	195	36.621	18.78
EEPCL	Ethanol	481	49.786	10.35
EEPC	Ethanol	247	23.604	9.55
EEGUL	Ethanol	558	54.457	9.76

EELPL = Ethanolic extract of *L. paniculata* leaves; EELPB = Ethanolic extract of *L. paniculata* bast; EEPCL = Ethanolic extract of *P. crucis* leaves; EEPC = Ethanolic extract of *P. crucis* bark; EEGUL = Ethanolic extract of *G. ulmifolia* leaves; AP(Li) = Antipromastigote activity of *L. infantum*; AP(Lb) = Antipromastigote activity of *L. brasiliensis*; AE = Antiepipromastigote activity (*T. cruzi*); DS = Standard deviation.

## 2.2. Chemical, apparatus and general procedures

All chemical were of analytical grade. Methanol, formic acid, gallic acid, chlorogenic acid, caffeic acid and rosmarinic acid were purchased from Merck (Darmstadt, Germany). Catechin, quercetin, quercitrin, coumarin, vitexin, luteolin, rutin, apigenin and kaempferol were acquired from Sigma Chemical Co. (St. Louis, MO, USA). High performance liquid chromatography (HPLC–DAD) was performed with a Shimadzu Prominence Auto Sampler (SIL-20A) HPLC system (Shimadzu, Kyoto, Japan), equipped with Shimadzu LC-20AT reciprocating pumps connected to a DGU 20A5 degasser with a CBM 20A integrator, SPD-M20A diode array detector and LC solution 1.22 SP1 software. The quantification of compounds by HPLC–DAD way performed by reverse phase chromatographic analyses was carried out under gradient conditions using C<sub>18</sub> column (4.6×150 mm) packed with 5 µm diameter particles; the mobile phase was water containing 1% formic acid (A) and methanol (B), and the composition gradient was: 12% of B until 10 min and changed to obtain 20%, 30%, 50%, 60%, 70%, 20% and 10% B at 20, 30, 40, 50, 60, 70 and 80 min, respectively, following the method described by Pereira et al. (2014) to *L. paniculata* extracts, with slight modifications and Klimaczewski et al. (2014) with slight modifications, to *P. crucis* and *G. ulmifolia* extracts. All chromatography operations were carried out at ambient temperature and in triplicate.

## 2.3. Cell lines used

For *in vitro* studies of *T. cruzi*, the clone CL-B5 was used (Le-Senne et al., 2002). Parasites stably transfected with the *Escherichia coli* b-galactosidase gene (lacZ), were provided by Dr. F. Buckner through Instituto Conmemorativo Gorgas (Panama). Epimastigotes were grown at 28 °C in liver infusion tryptose broth (Difco, Detroit, MI) with 10% fetal bovine serum (FBS) (Gibco, Carlsbad, CA), penicillin (Ern, S.A., Barcelona, Spain) and streptomycin (Reig Jofré S.A., Barcelona, Spain), as described previously (Roldos et al., 2008), and harvested during the exponential growth phase. Culture of *Leishmania brasiliensis* and *Leishmania infantum* was obtained from the Instituto de Investigaciones en Ciencias de la Salud, Asunción, Paraguay – IICS. The maintenance of strain, the form of cultivation and isolation shape promastigote followed

the procedures described by Roldos et al. (2008). The inhibition assays of promastigotes were performed using the strain of *L. brasiliensis* (MHOM/BR/75/M2903), grown at 22 °C in Schneider's Drosophila medium supplemented with 20% FBS. For the cytotoxic assays, was used the fibroblast cell line NCTC 929 grown in Minimal Essential Medium (Sigma). The culture medium was supplemented with heat-inactivated FBS (10%), penicillin G (100 U/ml) and streptomycin (100 µg/mL). Cultures were maintained at 37 °C in humid atmosphere with 5% CO<sub>2</sub>. The viability of these strains was assessed according to Roldos et al. (2008), through the use of resazurin as a colorimetric method.

## 2.4. Reagents

Resazurin sodium salt was obtained from Sigma–Aldrich (St. Louis, MO) and stored at 4 °C protected from light. A solution of resazurin was prepared in 1% phosphate buffer, pH 7, and filter sterilized prior to use. Chlorophenol red-β-D-galactopyranoside (CPRG; Roche, Indianapolis, IN) was dissolved in 0.9% Triton X-100 (pH 7.4). Penicillin G (Ern, S.A., Barcelona, Spain), streptomycin (Reig Jofré S.A., Barcelona, Spain) and dimethylsulfate were also used.

## 2.5. In vitro epimastigote susceptibility assay

The screening assay was performed in 96-well microplates with cultures that had not reached the stationary phase (Vega et al., 2005). Briefly, epimastigotes were seeded at  $1 \times 10^5$  mL<sup>-1</sup> in 200 µL of liver tryptose broth medium. The plates were then incubated with the drugs (0.1–50 µg/mL) at 28 °C for 72 h, at which time 50 µL of CPRG solution was added to give a final concentration of 200 µM. The plates were incubated at 37 °C for an additional 6 h and were then read at 595 nm. Each experiment was performed twice and independently, each concentration was tested in triplicate in each experiment. The efficacy of each compound was estimated by calculating the anti epimastigotes percentual (AE%).

## 2.6. In vitro leishmanicidal assay

The assay was performed using a modification of a previous method. Cultures of promastigotes of *L. brasiliensis* and *L. infantum* were grown to a concentration of 10<sup>6</sup> cells/mL and then transferred to the test. The compounds were dissolved in DMSO to the concentrations to be tested and were transferred to microplates. Each test was performed in triplicate. The activity of compounds was evaluated after 72 h by direct counting of cells after serial dilutions and compared with an untreated control.

## 2.7. Cytotoxic assays

NCTC929 fibroblasts were plated in 96-well microplates at a final concentration of  $3 \times 10^4$  cells/well. The cells were grown at 37 °C in an atmosphere of 5% CO<sub>2</sub>. After that, the culture medium was removed and the compounds were added to 200 µL, and performed a new culture for 24 h. After this incubation, 20 µL of a 2 mM solution of resazurin was added to each well. The plates were incubated for 3 h and the reduction of resazurin was measured using dual absorbance at

wavelengths of 490 and 595 nm. The value of the control (blank) was subtracted. Each concentration was tested in triplicate.

### 3. Results and discussion

The extracts evaluated in this work showed the yields demonstrated in Table 2, where we observed a higher yield of *L. paniculata* extracts compared to *P. crucis* and *G. ulmifolia*.

This work evaluated cytotoxicity using murine fibroblasts, as well as anti-parasite bioactivity against the epimastigote forms of *T. cruzi* and promastigote forms of *L. brasiliensis* and *L. infantum* of ethanol extracts from three species of plants occurring in a fragment of Cerrado in the midst of the Caatinga scrublands of northeastern Brazil.

Fig. 1 features the results for cytotoxic activity at concentrations of 1000–125 µg/mL for the natural products from the three species studied, as well as the LC<sub>50</sub> values (median lethal concentration capable of killing 50% of tested cells – murine fibroblasts). The extracts from leaves (EELLP) and bast tissue (EEBTLP) of *L. paniculata*, as well as the ethanol extract from the bark of *P. crucis* (EEBPC) showed low toxicity at the concentration of 500 µg/mL, with LC<sub>50</sub> values above 730 µg/mL for both. For its part, the leaf extract from *P. crucis* (EELPC) and leaf extract from *G. ulmifolia* (EELGU) displayed high cytotoxicity at the same concentration.

Fibroblasts are cells found in the conjunctive tissue of mammals. These cells are usually chosen for cytotoxicity tests because they are easy to maintain and produce results that show high correlation with biological results; also, they are present in wounds and constitute the main cell type present in regeneration (Ratner et al., 2004). Toxicity tests are devised in order to evaluate or predict toxic effects on biological systems and measure the relative toxicity of the substances (Forbes and Forbes, 1994). In that regard, the results can provide valuable information for the triage of natural products that can be regarded as probable candidates to become medications (Morais-Braga et al., 2013).

While evaluating anti-*Leishmania* activity, the best result was observed in EELGU, tested at four concentrations, with mortality of 95.23% (500 µg/mL) and 28.27% (250 µg/mL)


for *L. infantum* cells, and 92.20% (500 µg/mL) and 44.22% (250 µg/mL) for *L. brasiliensis* cells. EC<sub>50</sub> values of the tests with the respective parasites were 316.54 µg/mL and 282.63 µg/mL. The other extracts showed results of low to moderate anti-promastigote activity, with no cell mortality from EEBTLP at concentrations of 500 and 250 µg/mL. All extracts except EELGU showed EC<sub>50</sub> values over 850 µg/mL in anti-*Leishmania* tests.

With regard to the evaluation of anti-*Trypanossoma* activity, the best result was also observed for EELGU, with 68.93% inhibition of the protozoan at the concentration of 500 µg/mL and EC<sub>50</sub> = 454.68 µg/mL. The results of anti-parasite action from the studied extracts are shown in Table 3.

Plant extract triage is a valid strategy given that these natural products are explored in order to discover trypanocidal agents (Luize et al., 2006; Pizzolatti et al., 2008; Figueiredo et al., 2014). Several secondary metabolites with various structural patterns have proven active against *T. cruzi* (Saúde-Guimarães and Faria, 2007). For leishmaniasis, several studies have recommended the use of plants for treatment, both in ulcerated lesions from cutaneous leishmaniasis and in *in vitro* models (Paula-Junior et al., 2006; Bezerra et al., 2004; Santos et al., 2013).

Studies involving *Leishmania* spp. have focused on the extracellular form of the parasite, known as promastigote, instead of the amastigote form, due to easier *in vitro* growth and not requiring another cell culture such as macrophages (Castilhos et al., 2011).

The results of this work demonstrate that the extracts of *L. paniculata* and *P. crucis* were not clinically relevant for any of the tested protozoa, as they did not show values equal to or higher than 50% inhibition at a concentration of 500 µg/mL (EC<sub>50</sub> values above 850 µg/mL). To Rosas et al. (2007), this level of inhibition (50%) at a concentration of 500 µg/mL is considered to be clinically relevant. For its part, the leaf extract of *G. ulmifolia* showed considerable leishmanicidal potential, inhibiting 95.23% (*L. infantum*) and 92.20% (*L. brasiliensis*) of the growth of tested strains at a concentration of 500 µg/mL, with EC<sub>50</sub> values of 316.54 and 282.63 µg/mL, respectively. A significant result was also observed when evaluating anti-*Trypanossoma* potential, as the test using this extract




**Figure 1** Graph showing the percent cytotoxicity of extracts under study on murine fibroblasts. In which: (\*) LC<sub>50</sub> values (median lethal concentration capable of killing 50% of tested cells); (\*\*) LC<sub>50</sub> value over concentration of 1000 µg/mL.

**Table 3** Antipromastigote activities of *L. brasiliensis* and *L. infantum* and antiepipastigote activity (*T. cruzi*) of ethanol extracts from three species native to a fragment of Cerrado in Northeastern Brazil.

Natural product	Conc. ( $\mu\text{g/mL}$ )	%AP(Li)	$\pm$ %DS	%AP(Lb)	$\pm$ %DS	%AE	$\pm$ %DS
EELPL	1000	35.32	1.19	39.00	0.22	28.30	1.98
	500	30.56	1.05	27.64	0.22	23.18	0.57
	250	17.13	0.61	19.18	0.59	0.58	3.39
EELPB	1000	1.15	0.39	29.39	1.58	0.00	0.42
	500	0.00	1.99	25.69	0.71	0.00	0.49
	250	0.00	1.61	19.57	1.58	0.00	3.46
EEFPC	1000	49.98	1.07	45.65	0.15	55.82	1.56
	500	25.34	0.41	29.22	0.12	25.96	1.56
	250	18.24	0.14	21.12	1.17	14.46	0.92
EECPC	1000	23.25	0.07	23.26	1.37	25.28	1.12
	500	13.59	0.75	14.42	0.23	10.68	0.26
	250	10.39	0.57	11.18	0.26	0.00	1.24
EEFGU	1000	95.45	0.72	92.36	0.17	63.86	1.20
	500	95.23	0.39	92.20	0.04	61.15	0.99
	250	28.27	0.45	42.22	0.01	24.32	1.91
	125	5.80	0.44	0.00	2.14	4.03	0.99

EELPL = Ethanol extract of *L. paniculata* leaves; EELPB = Ethanol extract of *L. paniculata* bast; EEPCL = Ethanol extract of *P. crucis* leaves; EEPB = Ethanol extract of *P. crucis* bark; EEGUL = Ethanol extract of *G. ulmifolia* leaves; AP(Li) = Antipromastigote activity of *L. infantum*; AP(Lb) = Antipromastigote activity of *L. brasiliensis*; AE = Antiepipastigote activity (*T. cruzi*); DS = Standard deviation.



**Figure 2** Representative high performance liquid chromatography profile for *L. paniculata* leaves extract (A); *L. paniculata* sapwood extract (B); *P. crucis* leaves extract (C); *P. crucis* bark extract (D); *G. ulmifolia* leaves extract (E). Detection UV was at 325 nm, where: (A and B): gallic acid (peak 1), chlorogenic acid (peak 2), vitexin (peak 3), rosmarinic acid (peak 4), rutin (peak 5) and luteolin (peak 6); (C and D): gallic acid (peak 1), chlorogenic acid (peak 2), caffeic acid (peak 3), coumarin (peak 4), rutin (peak 5), quercitrin (peak 6), kaempferol (peak 7), luteolin (peak 8) and apigenin (peak 9); (E): catechin (peak 1), chlorogenic acid (peak 2), caffeic acid (peak 3), rutin (peak 4), quercitrin (peak 5), quercetin (peak 6) and luteolin (peak 7).

showed 68.93% inhibition in the growth of the strain of *T. cruzi* at a concentration of 500  $\mu\text{g/mL}$ .

One important criterion when searching for active compounds with leishmanicidal and trypanocidal activity is the evaluation of their toxicity in host cells. The cytotoxic activity of other plants has been evaluated in different human cell models, such as: peritoneal macrophages (Houghton et al., 2007), human lymphocytes (Reyes-Chilpa et al., 2008), MRC-5 cells (Cabral et al., 2010) and murine fibroblasts (Santos et al., 2013; Figueiredo et al., 2014).

The results of the cytotoxicity test in this study indicated values over 50% at the highest concentration (1000  $\mu\text{g/mL}$ ) for all natural products, meaning they were regarded as toxic. At concentrations in which cytotoxicity was low (extracts of *L. paniculata* and *P. crucis*), antiepipastigote and antipromastigote effects were irrelevant. Ethanol extract from the leaves of *G. ulmifolia*, which had considerable antiparasitic effect in this study, showed high cytotoxic activity at the concentrations of 1000  $\mu\text{g/mL}$  (88.41%) and 500  $\mu\text{g/mL}$  (68.93%), with  $\text{LC}_{50}$  value of 454.68  $\mu\text{g/mL}$ . This aspect becomes irrelevant, as

the cytotoxic potential of *G. ulmifolia* – in which the reactivity of the extract in mammal cells was verified – represents an essential step in natural products research that precedes *in vivo* tests, which confirm or negate the viability and safety of their therapeutic use in humans. The cytotoxicity presented can further be useful in studies on antineoplastic activity in tumor cells.

To Simões et al. (2010), toxic substances present in the plant may be limited to a given season of the year or to certain environmental conditions, or yet to specific varieties. Block et al. (2005), emphasize that the mechanism of action of cytotoxicity is related to the ability by plants to induce cell apoptosis.

The presence of phenolic content in the species was detected and quantified using the analytical method of high-performance liquid chromatography (HPLC). The extracts revealed the presence of phenolic acids such as gallic acid, chlorogenic acid, caffeic acid and rosmarinic acid, as well as flavonoids such as rutin, luteolin, apigenin and quercetin – the latter compound detected only in *G. ulmifolia* (Fig. 2).

Research results indicate that some of the phenolic compounds found in the extracts of the plants studied can show anti-kinetoplastidae activity (Morais-Braga et al., 2013). In *L. paniculata*, although low, the level of phenolic compounds found in the leaf extract is higher than in the bast tissue extract, with the most abundant compounds in the leaf extract being rosmarinic acid (1%) and luteolin (0.79%), whereas in the bast tissue vitexin (0.49%) and rosmarinic acid (0.41%) are most abundant. In *P. crucis*, the leaf extract contains kaempferol (1.57%) and coumarin (1.07%) at the highest rates, while quercitrin (1.89%) and gallic acid (1.5%) are most common in bark extract. In the leaf extract of *G. ulmifolia*, a natural product with significant antiparasitic results, chlorogenic acid (2.35%) and quercetin (2.15%) were found to be major compounds.

In a study on anti-*Leishmania* and anti-*Trypanosoma* activity by compounds isolated from *Croton lobatos* L., Lagnika et al. (2009) investigated, among other compounds, the action of chlorogenic acid, and found irrelevant activity. Tasdemir et al. (2006), in an *in vitro* study, indicated quercetin and its derivatives (7,8-dihydroxyflavone) as having potent and efficient antiparasitic activity against *Leishmania donovani*, with IC<sub>50</sub> of 1.0 µg/mL, although without significant activity for *T. cruzi*. Those authors further observed that the majority of most active compounds with trypanocidal and leishmanicidal effects have a typical flavone structure (2, 3 and C-4 of keto function) and/or a catechol substructure without replacements in the B-ring, in the benzochromone skeleton and in 7,8-dihydroxyflavone. Mittra et al. (2000) demonstrated a significant inhibitory effect *in vitro* by luteolin and quercetin on the growth of promastigote and amastigote forms of *L. donovani*. Muzitano et al. (2006), observed quercetin, isolated from *Kalanchoe pinnata*, to have leishmanicidal activity against *Leishmania amazonensis*. Matsuda et al. (2004) demonstrated that quercetin is a potent immunomodulator, capable of directly inducing death by apoptosis of *Trypanosoma brucei* gambiense, without affecting the viability of normal cells. These results highlight the use of quercetin as a potential antimicrobial and anti-inflammatory agent for the treatment of trypanosomiasis.

An extract, however, is a mixture of several compounds, which in their chemical interactions can combine synergistically,

antagonistically or indifferently, and thus alter the effect that each would have by itself (Morais-Braga et al., 2013). Furthermore, the low concentrations of phenolic compounds may have been responsible for the weak bioactivity of products from *L. paniculata* and *P. crucis* against *L. brasiliensis*, *L. infantum* and *T. cruzi*.

Although cited in several works for its use by local populations to clean internal ulcers and wounds (Brandão, 1991; Ritter et al., 2002; Carvalho, 2006; Tanaka et al., 2005), no records were found of works investigating the anti-*Leishmania* potential of *L. paniculata*, or its anti-*Trypanosoma* effect. *G. ulmifolia*, another medicinal tree with multiple uses, is also hailed in natural medicine for cleaning ulcers and healing open wounds (Teske and Trentini, 1997), and this is the first report on these activities for that specie, as well as for *P. crucis*.

#### 4. Conclusions

The ethanol leaf extract from *G. ulmifolia* was considered efficient against the tested parasite strains, emerging as a potential alternative source of natural products with activity against *T. cruzi*, *L. brasiliensis* and *L. infantum*, possibly due the presence of quercetin, a potent known leishmanicidal compound. The cytotoxicity seen in the extract from *G. ulmifolia* indicates the need for further tests, enabling future *in vivo* trials, including on antineoplastic activity in tumor cells.

#### References

- Bezerra, R.J.S., Leon, L., Genestra, M., 2004. Recentes avanços da quimioterapia das leishmanioses: moléculas intracelulares como alvo de fármacos. Rev. Bras. Ciênc. Farm. 40, 141–148.
- Block, S., Gerkens, P., Peulen, O., et al, 2005. Induction of apoptosis in human promyelocytic leukemia cells by a natural trachylobane diterpene. Anticancer Res. 25, 363–368.
- Brandão, M., 1991. Plantas Medicamentosas do Cerrado Mineiro, 15. Informe Agropecuário, Belo Horizonte, pp. 15–21.
- Cabral, M.M.O., Barbosa-Filho, J.M., Maia, G.L.A., et al, 2010. Neolignans from plants in Northeastern Brazil (Lauraceae) with activity against *Trypanosoma cruzi*. Exp. Parasitol. 124, 319–324.
- Carvalho, P.E.R., 2006. Espécies Arbóreas Brasileiras, 2. Embrapa Informações Tecnológicas, Embrapa Florestas, Brasília, Colombo, p. 627.
- Castilhos, P., Pereira, C.G., Silva, A.L.N., et al, 2011. Effects of *Bothrops moojeni* venom on *Leishmania amazonensis* promastigote forms. J. Venomous Anim. Toxins Incl. Trop. Dis. 17, 150–158.
- Chai, Y., Yan, S., Wong, I.L.K., Chow, L.M.C., Sun, H., 2005. Complexation of antimony (Sb-V) with guanosine 5'-monophosphate and guanosine 5'-diphospho-D-mannose: formation of both mono- and bis- adducts. J. Inorg. Biochem. 99, 2257–2263.
- Coura, J.R., Junqueira, A.C.V., Carvalho-Moreira, C.J., Borges-Pereira, J., Albajar, P.V., 2007. Uma visão sistêmica da endemia chagásica. In: Silveira, A.C. (Ed.), La Enfermedad de Chagas a la Puerta de los 100 Años del Conocimiento de Una Endemia Americana Ancestral. Organización Panamericana de la Salud/ Fundación Mundo Sano, Buenos Aires.
- Ferri, M.G., 1955. Contribuição ao Conhecimento da Ecologia do Cerrado e da Caatinga, 2. Boletim da Faculdade de Filosofia, Ciências e Letras, São Paulo, pp. 5–170.
- Figueredo, F.G., Tintino, S.R., Brito, D.I.V., et al, 2014. Avaliação das potenciais atividades tripanocida e antileishmaniana do extrato de folhas de *Piper arboreum* (Piperaceae) e de suas frações. Rev. Ciênc. Farm. Básica Apl. 35, 149–154.

- Figueiredo, M.A., Fernandes, A., 1987. Encraves de cerrado no interior do Ceará. *Rev. Ciênc. Agron.* 18, 103–106.
- Forbes, V.E., Forbes, T.L., 1994. *Ecotoxicology in Theory and Practice*. Chapman and Hall, London.
- Guarin Neto, G., Morais, R.G., 2003. Recursos medicinais de espécies do cerrado de Mato Grosso: um estudo bibliográfico. *Acta Bot. Bras.* 17, 561–584.
- Hiruma-Lima, C.A., Santos, L.C., Kushima, H., et al., 2006. Brazilian “cerrado” medicinal plant presents an important antiulcer activity. *J. Ethnopharmacol.* 104, 207–214.
- Houghton, P.J., Howes, M.J., Lee, C.C., Steventon, G., 2007. Uses and abuses of *in vitro* tests in ethnopharmacology: visualizing an elephant. *J. Ethnopharmacol.* 110, 391–400.
- Lana, M., Tafuri, W.L., 2005. *Trypanosoma cruzi* e a doença de chagas. In: Neves, D.P. (Ed.), *Parasitologia Humana*. Atheneu, São Paulo.
- Le-Senne, A., Muelas-Serrano, S., Fernandez-Portillo, C., Escario, J. A., Gomez-Barrio, A., 2002. Biological characterization of a betagalactosidase expressing clone of *Trypanosoma cruzi* CL strain. *Mem. Inst. Oswaldo Cruz* 97, 1101–1105.
- Luize, P.S., Ueda-Nakamura, T., Dias-Filho, B.P., Cortez, D.A.G., Nakamura, C.V., 2006. Activity of neolignans isolated from *Piper regnellii* (MIQ.) C. DC. var. *pallidum* (C. DC.) Yunck against *Trypanosoma cruzi*. *Biol. Pharm. Bull.* 29, 2126–2130.
- Maia-Elkhoury, A.N.S., Alves, W.A., Sousa-Gomes, M.L., Sena, J. M., Luna, E.A., 2008. Visceral leishmaniasis in Brazil: trends and challenges/Leishmaniose visceral no Brasil: evolução e desafios, 24. *Cadernos de Saúde Pública*, pp. 2941–2947.
- Matsuda, M.M., Rambert, J., Malvy, D., et al., 2004. Quercetin induces apoptosis of *Trypanosoma brucei* gambiense and decreases the proinflammatory response of human macrophages. *Antimicrob. Agents Chemother.* 48, 924–929.
- Mitra, B., Saha, A., Chowdhury, A.R., et al., 2000. Luteolin, an abundant dietary component is a potent anti-leishmanial agent that acts by inducing topoisomerase II-mediated kinetoplast DNA cleavage leading to apoptosis. *Mol. Med.* 6, 527–541.
- Morais-Braga, M.F.B., Souza, T.M., Santos, K.K.A., et al., 2013. Citotoxicidade e atividade antiparasitária de *Lygodium venustum* SW. *Acta Toxicol. Argent.* 21, 50–56.
- Muzitano, M.F., Tinoco, L.W., Guette, C., et al., 2006. The antileishmanicidal activity assessment of usual flavonoids from *Kalanchoe pinnata*. *Phytochemistry* 67, 2071–2077.
- Paula-Junior, W., Rocha, F.H., Donatti, L., Fadel-Picheth, C.M.T., Weffort-Santos, A.M., 2006. Leishmanicidal, antibacterial, and antioxidant activities of *Caryocar brasiliense* Cambess leaves hydroethanolic extract. *Rev. Bras. Farmacogn.* 16, 625–630.
- Pereira, I.O., Sacramento, L.V.S., Marques, M.J., 2011. Leishmanioses: “o Estado da Arte”, 9. *Revista da Universidade do Vale do Rio Verde*, pp. 220–238.
- Pizzolatti, M.G., Mendes, B.G., Cunha-Júnior, A., et al., 2008. Trypanocidal activity of coumarins and styryl-2-pyrone from *Polygala sabulosa* A.W. Bennet (Polygalaceae). *Rev. Bras. Farmacogn.* 18, 177–182.
- Rangel, E.F., Lainson, R., 2009. Proven and putative vectors of American cutaneous leishmaniasis in Brazil: aspects of their biology and vectorial competence. *Mem. Inst. Oswaldo Cruz* 104, 937–954.
- Ratner, B., Hoffman, A.S., Schoen, F.J., Lemons, J.E., 2004. In: *Biomaterials Science: An Introduction to Materials in Medicine*, second ed. Elsevier Academic Press, New York.
- Reyes-Chilpa, R., Estrada-Muñiz, E., Veja-Avila, E., et al., 2008. Trypanocidal constituents in plants. 7. Mamea-type coumarins. *Mem. Inst. Oswaldo Cruz* 103, 431–436.
- Ritter, M.R., Sobierajski, G.R., Schenkel, E.P., Mentz, L.A., 2002. Plantas usadas como medicinais no município de Ipê, RS, Brasil. *Rev. Bras. Farmacogn.* 12, 51–62.
- Roldos, V., Nakayama, H., Rolon, M., et al., 2008. Activity of a hydroxybiphenyl bryophyte constituent against *Leishmania* spp. and *Trypanosoma cruzi*: *in silico*, *in vitro* and *in vivo* activity studies. *Eur. J. Med. Chem.* 43, 1797–1807.
- Rosas, L.V., Cordeiro, M.S.C., Campos, F.R., et al., 2007. *In vitro* evaluation of the cytotoxic and trypanocidal activities of *Ampelozizyphus amazonicus* (Rhamnaceae). *Braz. J. Med. Biol. Res.* 40, 663–670.
- Santos, K.K.A., Rolón, M., Vega, C., et al., 2013. Atividade leishmanicida *in vitro* de *Eugenia uniflora* e *Momordica charantia*. *Rev. Ciênc. Farm. Básica Apl.* 34, 47–50.
- Saúde-Guimarães, D.A., Faria, A.R., 2007. Substâncias da natureza com atividade anti-*Trypanosoma cruzi*. *Rev. Bras. Farmacogn.* 17, 455–465.
- Simões, C.M.O., Schenkel, E.P., Gosmann, G., et al., 2010. In: *Farmacognosia: Da Planta ao Medicamento*, sixth ed. Editora da UFRGS/Editora da UFSC, Porto Alegre/Florianópolis.
- Sülsen, V.P., Cazorla, S.I., Frank, F.M., et al., 2007. Trypanocidal and leishmanicidal activities of flavonoids from argentine medicinal plants. *Am. J. Trop. Med. Hyg.* 77, 654–659.
- Tanaka, J.C.A., Silva, C.C., Dias Filho, B.P., et al., 2005. Chemical constituents of *Luehea divaricata* Mart. (Tiliaceae). *Quím. Nova* 28, 834–837.
- Tasdemir, D., Kaiser, M., Brun, R., et al., 2006. Antitrypanosomal and antileishmanial activities of flavonoids and their analogues. *In vitro*, *In vivo*, structure–activity relationship, and quantitative structure–activity relationship studies. *Antimicrob. Agents Chemother.* 50, 1352–1364.
- Teske, M., Trentini, A.M.M., 1997. In: *Herbarium: Compêndio de Fitoterapia*, third ed. Ingra, Curitiba.
- Vega, C., Rolon, M., Martinez-Fernandez, A.R., Escario, J.A., Gomez-Barrio, A., 2005. A new pharmacological screening assay with *Trypanosoma cruzi* epimastigotes expressing beta-galactosidase. *Parasitol. Res.* 95, 296–298.
- World Health Organization, 2010. Working to overcome the global impact of neglected tropical diseases: first WHO report on neglected tropical diseases. 2010 Dezembro [consulta em 03 de setembro de 2014]. Disponível em <[http://www.who.int/neglected\\_diseases/2010report/NTD\\_2010report\\_embargoed.pdf](http://www.who.int/neglected_diseases/2010report/NTD_2010report_embargoed.pdf)> .