

# pH homeostasis links the nutrient sensing PKA/TORC1/Sch9 ménage-à-trois to stress tolerance and longevity

Marie-Anne Deprez<sup>1,°</sup>, Elja Eskes<sup>1,°</sup>, Tobias Wilms<sup>1</sup>, Paula Ludovico<sup>2</sup>, Joris Winderickx<sup>1,\*</sup>

<sup>1</sup> Functional Biology, KU Leuven, Leuven, Belgium.

<sup>2</sup> Life and Health Sciences Research Institute (ICVS), School of Medicine, University of Minho, Braga, Portugal; ICVS/3B's - PT Government Associate Laboratory, Braga/Guimarães, Portugal.

<sup>°</sup> Contributed equally, in alphabetical order.

\* Corresponding Author:

Joris Winderickx, Functional Biology, KU Leuven, Kasteelpark Arenberg 31 box 2433; Tel: +32 (0)16 321516; E-mail: joris.winderickx@kuleuven.be

**ABSTRACT** The plasma membrane H<sup>+</sup>-ATPase Pma1 and the vacuolar V-ATPase act in close harmony to tightly control pH homeostasis, which is essential for a vast number of physiological processes. As these main two regulators of pH are responsive to the nutritional status of the cell, it seems evident that pH homeostasis acts in conjunction with nutrient-induced signalling pathways. Indeed, both PKA and the TORC1-Sch9 axis influence the proton pumping activity of the V-ATPase and possibly also of Pma1. In addition, it recently became clear that the proton acts as a second messenger to signal glucose availability via the V-ATPase to PKA and TORC1-Sch9. Given the prominent role of nutrient signalling in longevity, it is not surprising that pH homeostasis has been linked to ageing and longevity as well. A first indication is provided by acetic acid, whose uptake by the cell induces toxicity and affects longevity. Secondly, vacuolar acidity has been linked to autophagic processes, including mitophagy. In agreement with this, a decline in vacuolar acidity was shown to induce mitochondrial dysfunction and shorten lifespan. In addition, the asymmetric inheritance of Pma1 has been associated with replicative ageing and this again links to repercussions on vacuolar pH. Taken together, accumulating evidence indicates that pH homeostasis plays a prominent role in the determination of ageing and longevity, thereby providing new perspectives and avenues to explore the underlying molecular mechanisms.

doi: 10.15698/mic2018.03.618

Received originally: 29.09.2017;

in revised form: 24.12.2017,

Accepted 28.12.2017,

Published 12.01.2018.

**Keywords:** yeast, ageing, longevity, pH, V-ATPase, Pma1, PKA, TORC1, Sch9.

**Abbreviations:**

CLS – chronological lifespan

EGOC – EGO complex,

ER – endoplasmic reticulum,

pHc – cytosolic pH,

pHe – extracellular pH,

pHv – vacuolar luminal pH,

RLS – replicative lifespan,

WT – wild type.

## INTRODUCTION

pH homeostasis is of crucial importance for many molecular and physiological processes. In eukaryotic cells, intracellular pH affects protein folding and enzyme activity, is required for vesicle trafficking and impacts organelle function and integrity. As our knowledge of pH homeostasis progresses, it becomes increasingly evident that the proton not only acts as a facilitator for cellular functions but also as a potent second messenger for the regulation of growth and ageing. Dysregulation of pH and lysosomal dysfunction are being linked to numerous human diseases [1], including neurodegenerative disorders [2, 3]. Moreover, acidity-dependent probes are used as lysosomal storage disorder-

associated markers [4] and acidification of the extracellular space is known to promote cancer metastasis [5]. These observations led to a growing interest in pH-related research and in this domain the unicellular organism *Saccharomyces cerevisiae* proves to be a valuable model to unravel the fundamental molecular mechanisms underlying pH homeostasis and the relationship between pH, cell growth, stress resistance and longevity.

## THE MAIN PLAYERS IN pH HOMEOSTASIS

Cytosolic and organellar pH are tightly controlled in all eukaryotic cells. The two main players of pH homeostasis in yeast are the V-ATPase and Pma1. The V-ATPase is a pro-

ton translocating ATPase that pumps protons from the cytosol into the vacuole, endosomes and Golgi compartments. Pma1 is a P-type ATPase that pumps protons over the plasma membrane to acidify the extracellular space. As accurate control of pH is required for pH homeostasis and optimal functionality of the cell and its organelles, it is not surprising that the V-ATPase and Pma1 act in close harmony. Loss of V-ATPase activity leads to a partial mislocalization of Pma1 to the vacuole and other compartments, probably reflecting a compensatory mechanism [6]. In addition to the V-ATPase and Pma1, numerous other proton pumps and exchangers have been identified in yeast, and they probably fine-tune pH control of the cytoplasm and each of the organelles [7]. However, given the importance of pH for organelle-specific functions, surprisingly little is known about their precise regulation and contribution to pH homeostasis.

### Regulation of the vacuolar V-ATPase

The V-ATPase is a multisubunit enzyme composed of a peripheral  $V_1$  sector responsible for ATP hydrolysis, and a membrane-embedded  $V_0$  sector responsible for proton translocation [8]. In contrast to mammalian cells, which often exhibit tissue- and/or organelle-specific expression of multiple isoforms of one subunit, yeast cells only have one organelle-specific  $V_0$  isoform. Indeed, Vph1-containing V-ATPase complexes are localized at the vacuolar membrane, while Stv1-containing V-ATPase complexes cycle between the Golgi apparatus and endosomes [9]. The activity of Vph1-containing V-ATPases is mainly regulated by reversible disassembly of the  $V_0$  and  $V_1$  sectors, in which carbon source availability plays a major role. In the presence of glucose, the sectors are assembled at the vacuolar membrane and the V-ATPase is highly active, acidifying the vacuolar lumen. When glucose is scarce, the vacuolar  $V_1$  sector dissociates from the  $V_0$  sector and the vacuolar luminal pH (pH<sub>v</sub>) becomes more alkaline [10]. However, this is in no case an all-or-nothing event, as intermediate levels of assembly have been observed with varying nutrient conditions. The exact mechanism by which glucose impacts on vacuolar V-ATPase assembly remains poorly understood though several links with glucose-induced signalling events and glycolysis have been reported as further explained below. Interestingly, the vacuolar V-ATPase is also responsive to changes in cytosolic (pH<sub>c</sub>) and extracellular pH (pH<sub>e</sub>) [11-13]. For instance, V-ATPase disassembly upon glucose-starvation is significantly reduced when cells are grown at pH 7 and accordingly, enhanced V-ATPase activity is observed in isolated vacuoles from cells grown under these conditions as compared to cells grown at pH 5 [11, 12]. Hence, the V-ATPase appears to act as pH sensor and both in yeast and mammalian cells the vacuole-specific  $V_0$  subunit 'a', encoded by *VPH1* in yeast, has been proposed as pH-sensing protein [11, 14, 15]. Interestingly, this subunit also mediates the dissociation of the  $V_1$  and  $V_0$  complexes upon glucose-starvation [11]. In line with this, V-ATPases that reside in the Golgi compartment, containing the *STV1*-encoded subunit 'a' instead of Vph1, do not dissociate upon glucose depletion [16]. Besides glucose and

pH, lipids also impact on V-ATPase regulation as the signalling phosphoinositides PI[3,5]P<sub>2</sub> and PI[4]P promote V-ATPase activity at the vacuole and Golgi, respectively, through interaction with the appropriate 'a' subunits Vph1 and Stv1 [17-19]. Additional levels of V-ATPase regulation include the adjustment of coupling efficiency between ATP hydrolysis and proton translocation, and the mechanisms for assembly of the  $V_0$  complex in the endoplasmic reticulum (ER) and the subsequent export and delivery of the vacuolar V-ATPase to the vacuolar membrane [20-24].

### Regulation of the plasma membrane embedded Pma1

The Pma1 P-type ATPase is considered as the major determinant of pH<sub>c</sub> and plasma membrane potential in yeast [25, 26]. In contrast to genes encoding subunits of the V-ATPase, *PMA1* is an essential gene, making the protein a difficult target to study. Different environmental and nutritional factors control Pma1 activity, with glucose availability being the best studied. The transcription of *PMA1* is enhanced during growth on glucose [27]. In addition, the proton pump is controlled by reversible phosphorylation, which modulates an inhibitory interaction of the C-terminus with the active site of the H<sup>+</sup>-ATPase. The latter reflects a complex interplay of several signalling events, albeit not all players have yet been identified. Evidence obtained so far points to an involvement of the casein kinase 1 homologues Yck1 and Yck2, the protein kinase Ptk2 and the protein phosphatase Glc7 [28-30]. Calcium signalling was also shown to modulate the plasma membrane H<sup>+</sup>-ATPase activation in response to glucose [31]. Furthermore, the intra- and extracellular pH, as well as the plasma membrane potential affect Pma1 activity. Here, an important contribution has been ascribed to the pH-dependent regulation of the potassium transporter Trk1 and the compensatory roles of K<sup>+</sup> transport and H<sup>+</sup> efflux to maintain the electrochemical gradient [6, 32]. As mentioned before, a reduction in V-ATPase activity, for instance by glucose starvation, triggers sorting of Pma1 from the Golgi to the vacuole [33].

Note that the yeast genome encodes for another plasma membrane H<sup>+</sup>-ATPase, i.e. Pma2, but the expression of this pump is very low and, consistently, it only has a minor impact on cellular pH [34, 35].

The proton gradients established by the V-ATPase, Pma1 and other proton transport systems are of absolute importance for several processes. For instance, the driving force created by these gradients is crucial to maintain phosphate homeostasis [36] and the cation balance of alkali metals (Na<sup>+</sup> and K<sup>+</sup>), divalent cations (Ca<sup>2+</sup> and Mg<sup>2+</sup>) and trace metals (Fe<sup>2+</sup>, Zn<sup>2+</sup>, Cu<sup>2+</sup> and Mn<sup>2+</sup>). Their symport/antiport along with H<sup>+</sup> helps to maintain their physiological and non-toxic levels, so as to provide a suitable environment for various biochemical reactions. The exact regulation of this cation balance has been the subject of an excellent review [7]. Several membrane proteins rely on the proton driving force. Examples are the proton-coupled phosphate symporters and the different amino acid permeases that reside in the plasma membrane [36, 37] or the yeast AVT1-7 family members that mediate bidirectional

transport of amino acids across the vacuolar membrane [38].


**A MÉNAGE-À-TROIS FOR NUTRIENT SIGNALLING AND pH REGULATION**

As both the V-ATPase and Pma1 react closely to nutrient availability, it seems evident that proton pumping activity and pH homeostasis must be regulated by an interplay of diverse nutrient-induced signalling networks. Indeed, the protein kinases PKA and Sch9 and the protein kinase complex TORC1 play a central role. This so-called ménage-à-trois integrates input from several nutrient sensing systems in order to regulate metabolism, intracellular trafficking, proteome integrity, autophagy, stress resistance, cell size, cell cycle progression, growth and sporulation [39-42]. The activity of each of these kinases is tightly regulated in response to different nutritional cues.

**Nutrient-controlled regulation of the ménage-à-trois**

As shown in Fig. 1, the activity of PKA is regulated by the Ras-cAMP pathway and activation of adenylate cyclase, the

latter depending on extracellular glucose sensing via the Gpr1/Gpa2 GPCR system as well as on intracellular glucose sensing via activation of the small G-proteins Ras1 and Ras2 [43]. The protein kinase complex TORC1 is regulated by intracellular amino acid signalling and this occurs by different mechanisms depending on the quality of the amino acid as a nitrogen source [44]. Leucine and probably other neutral amino acids signal to TORC1 through the EGO complex (EGOC), the orthologue of the mammalian Rag-Ragulator complex. EGOC consists of Ego1, Ego2 and Ego3 that form a scaffold at the vacuolar membrane for the Rag GTPases Gtr1-2. The latter function as a heterodimer that receives input about the cytosolic amino acid content. When amino-acids are available in the cytoplasm, an active EGOC formation is triggered in which Gtr1 is in its GTP-bound form and Gtr2 in its GDP-bound form, leading to TORC1 activation. Amino acid starvation induces the opposite GTP/GDP-loading status of the GTPases and the consequent inhibition of TORC1 [45, 46]. Recent evidence obtained in mammalian systems suggests a role for the Rag-Ragulator in signalling lysosomal amino acid content and



**FIGURE 1: The nutrient signalling ménage-à-trois.** Glucose stimulates PKA activity through the GPCR system of Gpr1 and Gpa2 on the one hand, and through activation of Ras1/2 on the other. Both induce cAMP production by Cyr1, activating PKA. At the vacuolar membrane, the EGOC senses amino acids from the cytosol and presumably the vacuole, and activates TORC1 in return. TORC1 negatively regulates the energy sensor SNF1 and activates Sch9 through phosphorylation. Besides TORC1, also SNF1 influences Sch9 activity as well as the sphingolipid effectors Pkh1-3. The PKA and TORC1/Sch9 branches converge on multiple players, perhaps the most prominent being Rim15, whose import into the nucleus is inhibited by both branches and is also controlled by phosphate availability through Pho85. When the ménage-à-trois is inactive, Rim15 can enter the nucleus where it induces transcription of PDS genes by Gis1 and STRE genes by the Msn2/4 transcription factors.

transport in addition to cytosolic amino acid content [47, 48], but whether EGOc has a similar role in sensing the vacuolar amino acid load in yeast remains to be studied. Preferred nitrogen sources like glutamine can signal independently of EGOc to trigger a more sustained TORC1 activation required to support vigorous growth [44]. At present, the underlying mechanism for this Gtr1/2-independent TORC1 activation in yeast remains poorly understood, though one study identified Ypt1, a GTPase involved in ER-to-Golgi vesicular trafficking, as an alternative regulator of TORC1 [49]. Other studies suggested a role for the vacuolar membrane-associated phosphatidylinositol (PI) 3-phosphate binding protein Pib2 and the Vps15/34 PI 3-kinase complex [50, 51]. The third member of the ménage-à-trois, the AGC kinase Sch9, is the yeast orthologue of mammalian PKB/Akt and S6 kinase. It is a well-known target of TORC1 and as such it is involved in amino acid and nitrogen source signalling [52]. However, Sch9 has been implicated in the sensing and signalling of other nutrients as well. It plays a major role in lipid signalling as a regulator of the so-called sphingolipid rheostat [53] and, besides its phosphorylation by TORC1, Sch9 needs to be phosphorylated by sphingolipid-activated kinases Pkh1-3 at its PDK1 site in the activation loop to obtain full activity [52, 54, 55]. Moreover, Sch9 also responds to glucose availability, thereby acting in conjunction with PKA [56], and receives input of the protein kinase complex SNF1, a central regulator of cellular energy homeostasis [57]. Thus, Sch9 appears to play an integratory role in nutrient sensing that allows for coordination and fine-tuning of diverse signalling cascades. As such, it shares several effectors with the PKA and TORC1 signalling routes but for several of these targets the effect inflicted by Sch9 can either be additive or opposite depending on the activity status of the other kinases [41, 58-60].

An example of the interplay between the signalling pathways controlled by the ménage-a-trois is their convergence to regulate the activity and nuclear localization of Rim15, a protein kinase required for metabolic adaptation and the general stress response (Fig. 1) [61]. Here, also phosphate availability comes into the picture as the nuclear exit of Rim15 is controlled by the phosphate-responsive PHO-pathway, particularly Pho85 [41, 62].

### The nutrient status regulates pH homeostasis


The most straightforward evidence for the role of nutrient signalling in pH regulation are the alterations of extra- and intracellular pH in response to nutrient availability and growth rate. Albeit quite complex, we can summarize this as follows. When cells grow fermentatively, they produce organic acids that acidify the medium, such as acetic acid. In its protonated form, acetic acid is able to move through the plasma membrane back into the cell [63]. Once inside the cytosol, whose pH is maintained around neutrality during exponential growth, protons dissociate from the weak acid causing intracellular acidification. This acidification is counteracted by Pma1-mediated proton efflux and V-ATPase-mediated vacuolar acidification in order to maintain pH<sub>c</sub> homeostasis [9, 64-66]. During the diauxic shift or

upon glucose starvation, the activity of Pma1 reduces, the cytosol acidifies and the vacuolar V-ATPase disassembles [11, 67-70]. Re-addition of glucose or sucrose to glucose-starved cells initially results in a rapid drop in pH<sub>c</sub>, probably due to the resumption of glycolysis, but this is quickly followed by intracellular alkalization and extracellular acidification as a result of Pma1 and vacuolar V-ATPases regaining full activity [11, 70-72]. In contrast to the clear impact of carbon source availability, re-addition of amino acids or another nitrogen source to starved cells does not affect pH<sub>c</sub> [70, 73]. This demonstrates that adjusting pH<sub>c</sub> is not simply a matter of growth resumption, but that it is specifically controlled by carbon source availability.

### The ménage-à-trois is crucial for pH regulation

The ménage-à-trois plays an important role to integrate nutrient availability and pH homeostasis as shown in Fig. 2. Alterations that enhance PKA activity, like deletion of the Ras2 GTPase *IRA2* or the PKA regulatory subunit *BCY1*, were found to inhibit vacuolar V-ATPase disassembly upon glucose deprivation. This suggests that PKA is involved in glucose-dependent V-ATPase regulation [74]. Although PKA-dependent phosphorylation has been described for the V<sub>1</sub> subunit 'C' in insect cells [75] and for the V<sub>1</sub> subunit 'A' in human HEK-293T cells [76], no such event has been reported in yeast so far. Nonetheless, the Ras-cAMP pathway and PKA are known to regulate key enzymes of glycolysis in yeast [41, 77] and two glycolytic enzymes, i.e. aldolase and phosphofruktokinase, were reported to associate with the V-ATPase, placing them in pole position to mediate the glucose signal [67, 78-80]. Especially the phosphofruktokinase Pfk2 might be a good candidate as it is a direct target of PKA [81]. Pfk2 is indeed required to maintain vacuolar acidification and optimal RAVE-mediated reassembly of the V<sub>0</sub> and V<sub>1</sub> subunits upon re-addition of glucose to cells that were briefly deprived of the sugar [67, 78]. RAVE is the acronym for 'Regulator of H<sup>+</sup>-ATPase of Vacuolar and Endosomal membranes' and is a complex known to function as a scaffold that binds the V<sub>0</sub> and V<sub>1</sub> sectors in a glucose-dependent manner [82, 83]. Earlier studies also suggested a possible role for the Ras-cAMP pathway in the glucose-induced activation of Pma1 [84, 85]. However, subsequent detailed analysis contradicted such a role and showed that the plasma membrane H<sup>+</sup>-ATPase is still activated in mutant strains deficient for glucose-induced cAMP increase and that similar activation levels are obtained in strains with normal or attenuated PKA activity [86, 87].

Although starvation and resupplementation of nitrogen and amino acids does not influence pH<sub>c</sub> [70, 73], both TORC1 and Sch9 are involved in the regulation of pH homeostasis in function of glucose availability. When cells are grown on glucose both players are localized at the vacuolar membrane, but once cells enter the diauxic shift, Sch9 is displaced into the cytosol. The latter occurs concomitantly with the disassembly of the vacuolar V-ATPase [70]. In fact, Sch9 has a modulatory role in controlling the V-ATPase assembly state as evidenced by the observation that cells lacking Sch9 display an enhanced association of the V<sub>0</sub> and



**FIGURE 2: The interplay between signalling and pH control.** Pma1 and the V-ATPase, the two main players in pH control, both impact and are regulated by the ménage-à-trois. When stimulated by glucose or cytosolic protons, the vacuolar Vph1-containing V-ATPase stimulates TORC1 via the EGO-complex components Gtr1/2. The Golgi-specific Stv1-containing V-ATPase stimulates PKA via Ira1/2 inhibition by Arf1, pushing Ras1/2 towards the GTP-bound state and in turn activating Cyr1 to produce cAMP. Active TORC1 and PKA each regulate the vacuolar V-ATPase assembly state. TORC1 stimulates disassembly via Sch9, while PKA aids V-ATPase assembly presumably via Pfk2 which impacts on the V-ATPase scaffolding complex RAVE. Additionally, the vacuolar V-ATPase subunit Vph1 acts as a proton sensor to induce V-ATPase activity, and this may also be the case for the Stv1-containing V-ATPase. Additionally, active TORC1 also stimulates Pma1 activity.

V<sub>1</sub> sectors in the presence of glucose and a delayed disassembly of the proton pump upon glucose starvation. Similar effects were also seen upon rapamycin-induced TORC1 inhibition or upon replacement of the wild type Sch9 allele by the Sch9<sup>5A</sup> version that can no longer be activated by TORC1 [70]. Accordingly, *sch9Δ* cells have more acidic vacuoles both in the presence or absence of glucose, and while *sch9Δ* mutant cells are able to maintain their pH<sub>c</sub> within the same range as wild type (WT) cells during fermentative growth, once they traverse the diauxic shift a hyperacidification of the cytosol is observed [70]. Whether the latter points to a role of Sch9 in the regulation of Pma1 or other ion pumps and exchangers remains to be investigated in more detail, but at least two observations suggest that this may well be the case. Indeed, *sch9Δ* cells fail to rapidly acidify the extracellular medium when glucose is refed to glucose-starved cells and the additional deletion of *SCH9* in strains lacking a functional V-ATPase triggers a

further alkalization of the vacuolar lumen [70]. Interestingly, a most recent paper reported that TORC1 is required to obtain full Pma1 activity and although a possible role of Sch9 was not examined, the study demonstrated an involvement of the protein phosphatase Sit4, another TORC1 effector, in the turnover but not the phosphorylation status of Pma1. The same study also showed that both *sit4Δ* cells and cells lacking the TORC1 subunit Tco89 display reduced K<sup>+</sup> uptake and a more acidic intracellular pH as compared to WT cells, but solely in the *sit4Δ* mutant this was associated with a reduced Pma1 activity [88].

#### Additional players link nutrient signalling with pH homeostasis

Apart from the effects mediated by the PKA and TORC1-Sch9 axis on V-ATPase assembly/disassembly or Pma1 activity, other players involved in nutrient signalling also affect pH homeostasis. As mentioned, the signalling lipid

PI[3,5]P<sub>2</sub> was shown to be important for pH<sub>v</sub> regulation. Accordingly, the PI[3,5]P<sub>2</sub>-deficient mutants *vac7*, *vac14* and *fab1* all display vacuolar acidification defects [89-91]. Although PI[3,5]P<sub>2</sub> may directly affect V-ATPase activity [18], the phosphoinositide may also act indirectly via TORC1/Sch9 signalling, as TORC1 activity as well as the recruitment of Sch9 and its TORC1-dependent phosphorylation at the vacuolar membrane are dependent on the presence of PI[3,5]P<sub>2</sub> [92-94]. Moreover, increased PI[3,5]P<sub>2</sub> levels are associated with vacuolar fragmentation, which involves TORC1 and its downstream effectors [95, 96]. This relationship is further highlighted by the significant overlap in targets between a genome-wide screening aiming to identify genes conferring a synthetic sick/lethal phenotype when combined with the *SCH9* deletion and a screening for genes mediating inositol auxotrophy [70, 97]. Similar genome-wide screenings were performed to find proteins involved in the control of pH<sub>c</sub> or pH<sub>v</sub> during fermentative growth on glucose [24, 25] and at least with the screening on pH<sub>c</sub> a significant overlap is seen when compared to the data of the synthetic *SCH9* screening [70]. Besides proteins associated with vesicular transport, lipid metabolism or amino acid biosynthesis, both pH screenings retrieved proteins required for mitochondrial functions. This is intriguing because it indicates that mitochondria are essential to maintain pH homeostasis even during fermentative growth, where they are not required for energy supply. This underscores the importance of cross-talk between organelles, and the notable lack of research performed to date in this area.

#### pH as a second messenger for nutrient signalling

Changes in intracellular pH affect cell functioning at different levels as it impacts on protein folding, enzyme activities and the protonation of biological macromolecules, lipids and other metabolites [66]. The crucial question, however, is whether alterations in pH are sensed and perceived as signals. Perhaps the best way to answer this is to describe the recent advances in how pH affects the activities of the PKA/TORC1/Sch9 ménage-à-trois.

Already decades ago, it was reported that the treatment of fungi with depolarizing agents inflict a rapid increase in the level of cAMP [98-100]. Mechanistically, different scenarios were described as intracellular acidification was found to enhance both the affinity of the adenylate cyclase *Cyr1* for its substrate ATP [98] as well as Ras-GTP loading by inhibition of the GTPase-activating proteins, *Ira1* and *Ira2* [101]. Notably, a mild treatment of cells with a protonophore allows to bypass the requirement of hexose transport for glucose-induced activation of the Ras-cAMP pathway [102], which is consistent with the observation that the protonophore treatment of glucose-starved cells triggers a rapid drop in pH<sub>c</sub> similar as that seen immediately after addition of glucose to these starved cells [98]. A more recent study correlated the glucose-induced changes in pH<sub>c</sub> to V-ATPase assembly/disassembly and subsequent activation of PKA [11]. As mentioned, this study also proposed the V-ATPase to act as pH-sensor with Vph1, the V<sub>0</sub> subunit 'a', as putative pH-sensing protein.


Subsequently, the role of the V-ATPase for activation of the Ras-cAMP pathway was further defined, since in the presence of glucose the pump was shown to signal to the Ras proteins via Arf1, a GTPase that interacts with the Golgi-specific V<sub>0</sub> subunit Stv1. Moreover, the activity of the plasma membrane ATPase Pma1 was shown to stimulate growth through Ras activation by increasing pH<sub>c</sub> [73].

Additionally, pH<sub>c</sub> and the V-ATPase also confer signals for glucose availability to TORC1 and Sch9. To this end, the V-ATPase controls the GTP-load of Gtr1 and interacts with this EGOC Rag GTPase via the vacuole-specific V<sub>0</sub> subunit Vph1. This finding is rather remarkable as Gtr1 plays a pivotal role in amino acid sensing [45, 46], but amino acid availability itself does not affect pH<sub>c</sub> nor the V-ATPase assembly state [70, 73]. As the V-ATPase apparently acts as activator of both PKA and the TORC1-Sch9 axis, while each of the kinases has the ability to provide feedback by affecting the V-ATPase assembly/disassembly, a very balanced system based on feedback loops is established (Fig. 2). The necessity of this tight control relates undoubtedly to the crucial roles of the PKA/TORC1/Sch9 ménage-à-trois for the overall cellular functioning and the coordination between growth and the cell cycle [41]. The latter is further evidenced by the finding that the TORC1-Sch9 axis transmits signals from the vacuole that are required for cell cycle progression [94].

A signalling role of pH<sub>c</sub> also became apparent from data obtained from a genome-wide screening that investigated the correlation between aberrant intracellular pH and reduced growth rate of mutants. This analysis confirmed a tight connection between both, suggesting that pH<sub>c</sub> dictates the growth rate. For 19 out of the 173 mutants, however, the causal relationship between pH<sub>c</sub> and growth rate was completely abrogated, indicative that these mutants fail to properly sense the pH signal. Among them were mutants affected in mitochondrial translation, inositol phosphate biosynthesis and lipid biosynthesis [25]. Importantly, a similar study found no causal correlation between pH<sub>v</sub> and growth rate [24].

#### pH CONTROL, STRESS TOLERANCE AND LONGEVITY

It is well established that the nutrient pathways controlled by the PKA/TORC1/Sch9 ménage-à-trois have a significant impact on cellular ageing, whether being monitored as replicative lifespan (RLS) by assessing the number of divisions a mother cell can undergo before dying, or as chronological lifespan (CLS) by assessing the time span a non-dividing cell remains viable [103]. Both modes of ageing rely on partially overlapping cellular and molecular determinants and these have been the topic of excellent reviews to which we refer for more details [40, 104-112]. As nutrient availability and the PKA/TORC1/Sch9 ménage-à-trois affect pH control, it is not surprising that pH is being linked to the regulation of ageing/longevity in yeast. Hence, a number of groups have explored the interplay between extracellular, cytosolic and organelle pH and longevity, the topic of the sections below and summarized in Fig. 3.



**FIGURE 3: The interplay between pH control and ageing. (A) Young daughter cell.** Fermentatively growing cells produce acetic acid, which is extruded from the cell. When protonated in the acidic medium, acetic acid passes the plasma membrane causing cytosolic acidification and acetic acid stress. CR and medium buffering counteract this phenomenon by reducing fermentation and protonation of acetic acid respectively. The optimally active Pma1 and V-ATPase retain the cytosolic pH around neutrality by counteracting this proton influx. The resulting acidic vacuole allows for optimal amino acid uptake by the proton/amino acid antiporter Avt1 and efficient autophagy. Acetic acid stress induces the environmental stress response aided by transcription factors Msn2/4. Albeit mitochondria are not required for energy production on fermentative medium, they do contribute to the maintenance of pH homeostasis. The retrograde response, mediated by Rtg1/3, assures mitochondrial integrity. In the event of protein misfolding, the UPR is activated in the ER. Damaged and aggregated proteins are efficiently cleared by autophagy and the ubiquitin-proteasome system (UPS; not shown). **(B) Old mother cell.** Due to asymmetrical inheritance, mother cells possess an abundance of Pma1 proteins, presumably raising the cytosolic pH, which likely also causes an unavailability of protons to be traversed to the vacuole. The resulting vacuolar alkalization causes insufficient amino acid import by Avt1, leading to mitochondrial dysfunction by a yet unknown mechanism. The retrograde response is activated in an attempt to counteract mitochondrial dysfunction, though it may not be sufficient to maintain mitochondrial integrity. Misfolded proteins accumulate in the ER thereby triggering ER stress, and damaged proteins accumulate in the cytosol due to impaired autophagy and an overwhelmed UPS (not shown). Both reduced mitochondrial integrity and protein misfolding lead to the formation of reactive oxygen species (ROS) and they affect lipid homeostasis thereby further impairing vacuolar functioning. **(C) Chronological ageing of young daughter cell.** When traversing the diauxic shift, cells experience a disassembly of the V-ATPase and a reduction in Pma1 activity, aided by its endocytosis. As a result, the cytosol acidifies and the vacuole alkalinizes. This presumably, similar as in replicatively ageing cells, leads to an impairment of vacuolar amino acid uptake, autophagy, mitophagy and endocytosis. In post-diauxic shift cells, Msn2/4 and Gis1 are activated to induce stress responses and the production of trehalose, which helps to protect cells from acetic acid stress and toxicity of damaged proteins. Although stress responses are optimal in young daughter cells traversing post-diauxic shift, intracellular damage will accumulate over time, eventually surpassing the capacity of the protective machinery, leading to cell demise. **(D) Hypothetical model of chronological ageing in old mother cell.** In this model we assume a cumulative deregulation of the already decontrolled pH homeostasis in old mother cells following post-diauxic shift. This should lead to an additional impairment of vacuolar amino acid uptake, autophagy, mitophagy and endocytosis and cause a quick accumulation of intracellular damage, leading to more rapid cell demise.

### Acetic acid shortens longevity and induces programmed cell death

A nice example that illustrates the interconnections between lifespan, nutrient availability and their dependence on pH is the observation that chronologically aged cells display a reduced subsequent RLS, but that this can be attenuated when the chronological ageing occurs in buffered medium or in calorie restriction (CR) conditions [113, 114]. As mentioned, during fermentative growth yeast cells produce and secrete different organic acids. While some of these acids accumulate in the growth medium, acetic acid levels decline after post-diauxic shift, indicating that this acid is used as a carbon source. The drawback is, however, that acetic acid uptake by the cell comes with a degree of toxicity that affects longevity (Fig. 3). Consistently, growing cells under acidic conditions was found to shorten CLS and RLS, while CR significantly reduces the production of organic acids and extends CLS and RLS [114-116]. Buffering the medium also has a positive effect on lifespan as it lowers the difference between the extra- and intracellular pH values, thereby reducing the driving force for inwards diffusion of acetic acid. As such, buffering presumably prevents stress by reducing the amount of energy that needs to be consumed to maintain the intracellular pH [117].

Indeed, as a first line of defense against acetic-acid induced stress, cells rely on energy-consuming proton pumps to counteract cytosolic acidification caused by dissociation of protons from the acetic acid once it has entered the cell, and to maintain pH optima in all different cellular compartments. Here, the plasma membrane-embedded Pma1 fulfils an important role by extruding these protons from the cytosol, allowing healthy exponential WT cells to maintain pH<sub>c</sub> around neutrality independently of pH<sub>e</sub> (Fig. 3A) [71]. During ageing however, yeast cells struggle to maintain proper pH homeostasis. As mother cells undergo divisions, the pH<sub>c</sub> increases whereas daughter cells retain a more acidic cytosol (Fig. 3B). This phenomenon is attributed to asymmetrical distribution of Pma1, which predominantly remains in the plasma membrane of the mother cell [118]. This mother-specific increase in Pma1 activity and subsequent reduction of cytosolic proton content is believed to trigger the decline of vacuolar acidity during replicative ageing, as protons are unavailable for the V-ATPase to be pumped into the vacuolar lumen [118]. Conversely, when cells have traversed the diauxic shift or when encountering glucose depletion, a condition typically used to study chronological ageing, the activity of Pma1 declines as mentioned previously. Moreover, also the V-ATPase is disassembled under these conditions [11, 67-70] and this loss of V-ATPase activity signals ubiquitination and endocytosis of Pma1 [119]. In consequence, the pH<sub>c</sub> is significantly reduced both in post-diauxic cells and in chronologically aged cells and due to the lower activity of the V-ATPase, the vacuole presumably becomes more alkaline (Fig. 3C).

Secondly, mild stress induces the so-called general or environmental stress response pathway, an adaptive response that allows to acquire a form of stress resistance that protects cells from subsequent stress triggered by the same or another stressor [120, 121]. This adaptive re-

sponse applies to acetic acid-induced stress as well as to physiological stresses affecting lifespan [122] and it is the basis of hormesis effects that play in ageing [123, 124]. In the environmental stress response pathway, the transcription factors Msn2/4 play a central role. Msn2/4 are well known targets of the PKA/TORC1/Sch9 ménage-à-trois required to induce expression of several stress-responsive genes needed to protect cells from adverse conditions (Fig. 1; Fig. 3) [41]. Notably, these transcription factors are also induced when cells encounter alkaline stress [125, 126], suggesting that the actual stress factor might be a change in pH, the proton gradient or membrane potential. A study that aimed to identify genes essential for the acquisition of tolerance to different weak acids implicated Msn2/4 for acetic acid tolerance [127]. Interestingly, apart from vacuolar acidification, intracellular trafficking and ergosterol biosynthesis, acetic acid tolerance was found to specifically depend on a small set of genes and this included those encoding Ras2, the trehalose-6P synthase, different cytosolic and mitochondrial ribosomal proteins and two well-known players of the retrograde response pathway, i.e. Rtg2 and Rtg3 [127]. These data nicely complement observations connecting metabolism, cell death and longevity. The finding of mitochondrial ribosomal proteins and components of the retrograde response pathway is consistent with the requirement of functional mitochondria to maintain pH homeostasis [24, 25], and there is ample evidence that mitochondrial dysfunction accompanies acetic acid-induced programmed cell death (PCD) [128, 129] and the reduction of CLS and RLS (Fig. 3B-D) [130-135]. Upon mitochondrial dysfunction, the retrograde response pathway is triggered to transmit signals to the nucleus in order to make adjustments in cellular metabolic and biosynthetic activities. The retrograde response pathway is positively controlled by Ras2, explaining why this small GTPase is essential for the acquisition of acetic acid tolerance. In fact, PKA, similar as TORC1, negatively influences the retrograde response pathway [136]. Among other genes, the retrograde response pathway targets several cytosolic and mitochondrial ribosomal protein genes [137], explaining their involvement in acetic acid tolerance.

Some evidence suggests that the expression of genes involved in trehalose biosynthesis is also influenced by the retrograde response pathway [137]. Although the role of trehalose in conveying acetic acid tolerance is not fully understood, the long-lived *tor1Δ* and *sch9Δ* mutants appear to make optimal use of the protective properties of trehalose as these strains were found to switch their metabolism in the quiescent phase and use the acetic acid that was secreted during the pre-diauxic phase to produce trehalose [138]. A recent study reported that Tps1 can decelerate chronological ageing independently of its known trehalose-6P catalytic activity [139, 140]. Consistently, the *tps1Δ* mutant is short-lived. However, another study demonstrated that this phenotype is shared by mutants lacking the trehalose-6P phosphatase Tps2, or the neutral trehalases Nth1 and Nth2. In contrast, mutants lacking the regulatory subunits of the trehalose synthase



complex Tps3 or Tsl1, or the periplasmic acid trehalase Ath1, were found to be long-lived [141]. Analysis of these different mutants revealed that trehalose reduces the amount of oxidative carbonylated proteins during post-diauxic phase and that it lowers the level of protein aggregation in the quiescent state [141]. Thus, trehalose seems to protect cells by preventing proteotoxicity (Fig. 3C-D).

### pH and proteotoxicity

Cells are equipped with a complex network that ensures proteome integrity. Proteins that can no longer fulfil their function due to misfolding, damage or aggregation become substrates for the protein degradation machinery and they are cleared either via the proteasome, or via autophagy and vacuolar targeting (Fig. 3). Here the link to pH homeostasis is obvious, as the final step of autophagy, i.e. the disintegration of autophagic bodies, is linked to vacuolar membrane integrity and acidification of the vacuolar lumen by the V-ATPase [142-144]. Next to its detoxifying function, however, autophagy also has a role in nutrient recycling. In that regard, it seems logical that autophagy is upregulated in nutrient limiting conditions and that the nutrient signalling ménage-à-trois plays an important role in the regulation of autophagy [40, 41, 60]. Interestingly, the cellular capacity for autophagic degradation declines with age, which will itself also contribute to the accumulation of cellular and molecular damage. Accordingly, induction of autophagy extends lifespan, and this seems to be accompanied by vacuolar acidification (Fig. 3) [143, 145].

Recent evidence indicates that misfolded and damaged proteins are first partitioned in specific inclusions in the cell. In JunQ and INQ misfolded proteins are refolded, while those deposited in IPOD probably await clearance via autophagy [146, 147]. This partitioning requires Hsp104 and functional actin cables [148-150], the latter being another essential lifespan determinant [151, 152]. Partitioning of misfolded proteins is beneficial during chronological ageing [153] and assures the asymmetric inheritance of damaged and non-functional proteins during replicative ageing, thereby producing rejuvenated daughter cells [148-150]. A recent study connected the V-ATPase, vesicular trafficking and components involved in actin cable-dependent vacuole inheritance to this asymmetric inheritance [154], implying a pH dependency. Notably, apart from the vacuole also the ER, mitochondria and even mRNA are subject to asymmetric inheritance during the division of yeast cells (reviewed in [155]).

### Mitochondrial dysfunction and its interplay with the ER and the vacuole

Several molecular chaperones assist in the correct folding of proteins or the refolding in case protein misfolding occurs. These chaperone proteins can be found in the cytoplasm, the ER and mitochondria. In fact, a recent study revealed that yeast cells possess a cell-wide proteostasis system where proteotoxicity in one cellular compartment triggers a response in other compartments. This response, termed cross-organellar stress response (CORE) has a protective role and extends both CLS and RLS [156]. Aberrant

chaperone activity in each of the compartments leads to fragmented mitochondria, a loss of respiratory activity and an increase in cytosolic NADPH reducing power. This effect is associated with inactivation of TORC1, which acts as protein folding sensor, and the subsequent activation of Snf1 [156, 157]. The existence of a cell-wide proteostasis system is also inferred by the observation that yeast cells use a common system to monitor and ensure protein quality control in the ER and mitochondria. This system involves Cdc48/p97, an AAA ATPase known from the ubiquitin-proteasome system that is recruited by stressed ER or mitochondria to extract ubiquitinated proteins presented at the membranes of the organelles and to direct these to the proteasome [158, 159]. Interestingly, several physical organelle contact sites exist in yeast, although their involvement in CORE and the cell-wide proteostasis system remains to be elucidated. Here we mention ERMES, the ER-mitochondrial contact site [160], and vCLAMP, the vacuole-mitochondrion contact site [161]. Several nutrient and ion transporters have been shown to be enriched at these contact sites, indicating a role as hubs for the exchange of nutrients and ions between organelles [161] and pointing towards the idea that pH could be an essential regulator. In addition, ERMES and vCLAMP have been proposed to serve as dynamic metabolic signalling hubs [162]. Notably, they are co-regulated in response to nutrients and appear to fulfil partially overlapping functions as their simultaneous disruption is lethal [161].

It is obvious that mitochondria play an important role in the determination of lifespan that surpasses their function as energy supplying factory. It is essential for viability that dysfunctional mitochondria are removed, which occurs via mitophagy [40]. Efficient mitophagy requires ubiquitination of the ERMES components Mdm34 and Mdm12 [163]. Moreover, ERMES colocalizes with the site of mitophagosome generation and the ER was proposed to deliver the necessary lipids for membrane engulfment of the mitochondrion [164, 165]. Whether vCLAMP has a role in mitophagy has not been investigated yet. Nonetheless, it is known that it suffices to enhance vacuolar proton pumping to significantly reduce mitochondrial dysfunction, indicating a communication between these organelles which could rely on vCLAMP. Consistently, cells lacking functional V-ATPases display significant increased levels of markers for dysfunctional mitochondria and a dramatically shortened RLS and CLS (Fig. 3B-D) [70, 134]. Intriguingly, this appears not to be related to a reduced capacity of vacuolar protein degradation, but rather to the impaired ability of the vacuole to efficiently store amino acids. This is evidenced by the observation that enhanced expression of the neutral amino acid transporter Avt1 attenuates mitochondrial dysfunction in replicative ageing cells without preventing vacuolar alkalization [134]. How this relates to the regulation of autophagic processes by the PKA/TORC1/Sch9 ménage-à-trois [40, 166, 167] remains to be investigated.

### pH, lipid synthesis and liponecrosis

The synthesis of major membrane lipids is spatially organized and involves different organelles [168]. Moreover, many lipid synthesizing enzymes are enriched at the contact sites between the ER and other organelles and these contacts mediate non-vesicular selective lipid transport [169-175]. As organelles rely on pH for optimal functionality, it is expected that compartmental cross-talk for lipid metabolism is closely connected to pH regulation. This is supported by the observation that membrane contact sites constitute domains important for ion transport as well [161, 176], which in case of the ER-plasma membrane contact is linked to proton pumping by Pma1 [177]. Moreover, a study that used a systems biology approach to investigate the interdependence of pH control and CLS by comparison of young and old cells grown in buffered or non-buffered medium revealed that pH has a main impact on the reorganization of lipid metabolism. This reorganization has a beneficial effect on CLS by preserving mitochondrial and vacuolar health, the latter being dependent on V-ATPase activity [178].

Lipid homeostasis is of utmost importance to maintain longevity of yeast cells and persistent deviations thereof can lead to cell demise and a phenomenon described as liponecrosis [105, 179-181]. Several observations indicate that alterations in lipid homeostasis induce mechanisms involved in protein quality control. Conditions that perturb lipid biosynthesis, that alter the lipid compositions of the plasma membrane and endomembranes or that affect the lipid droplet content of cells were all reported to trigger ER stress and activation of the unfolded protein response (UPR) [182-188]. In contrast to acute ER stress, which induces PCD [189, 190], the lipid-associated induction of ER stress appears to be linked to compensatory mechanisms directed to reinstate lipid biosynthesis and lipid metabolism and to promote cell survival [191-195]. For instance, the activation of the UPR was shown to restore normal ceramide levels when sphingolipid biosynthesis was compromised [192] and to enhance synthesis of triacylglycerols and sterol esters in order to stimulate the formation of lipid droplets [196]. Besides their role for energy storage, these lipid droplets are essential for the regulation of autophagic processes and the clearance of damaged and aggregated proteins from the ER and mitochondria [197-201]. Interventions that affect lipid homeostasis are commonly associated with the appearance of fragmented vacuoles and V-ATPase dysfunction. This includes alterations in the biosynthesis of ergosterols, sphingolipids and ceramides or the availability of essential precursors like inositol [202-205]. However, preventing lipid droplet formation by blocking the synthesis of di- and triacylglycerol through deletion of the *PAH1*-encoded phosphatidic acid phosphatase appears to be an exception since *pah1Δ* cells are characterized by vacuolar fragmentation and enhanced lipid toxicity while still displaying improved acidification of the vacuolar lumen due to elevated expression of V-ATPase subunits [206, 207]. The latter relates to a negative effect of Pah1 on the transcription of several V-ATPase subunit

genes, which all contain an UAS<sub>INO</sub> element in their promoter [206]. This is interesting since this promoter element also links the expression of these V-ATPase genes to phospholipid biosynthesis and the availability of inositol, choline and phosphate [208].

### CONCLUDING REMARKS

Taken together, control of pH homeostasis is emerging as a key factor determining longevity and alterations culminate in many hallmarks of ageing. Although we are only beginning to uncover the importance of pH homeostasis, it is already amazing how many aspects of cell functioning are influenced by intracellular pH and the reciprocal regulation of mainly two players, the V-ATPase and Pma1. When thriving in an environment with plentiful nutrients, Pma1 and the V-ATPase maintain an ideal pH to support vigorous cell growth. During ageing, the activity of these two players changes, and while increased Pma1 activity may provide enhanced tolerance of a mother cell to the weak acids that were produced, the drawback is a reduction of vacuolar acidity. Indeed, vacuolar acidity is of absolute importance for endocytosis and vesicular trafficking, as well as cellular damage control, the latter via the clearance of malfunctioning proteins and organelles, important for both CLS and RLS (Fig. 3). In addition, damage control also occurs via asymmetrical inheritance during replicative ageing and also here links with vacuolar functioning and pH<sub>v</sub> are emerging. Undoubtedly, this field of research opens a promising path towards the understanding of intrinsic mechanisms of ageing and longevity, which could be of critical value for our insight into lysosomal-related human diseases and proteopathies, including neurodegenerative disorders, different types of cancer and lysosomal storage diseases.

### ACKNOWLEDGMENTS

We gratefully thank FWO Vlaanderen for a fellowship to EE and for support via the research grants G.0694.13, G.0A63.15 and SBO-S006617N. We also thank KU Leuven for support by granting the C14/17/063 project.

### CONFLICT OF INTEREST

The authors report no conflict of interest. J.W. declares that he is co-founder and shareholder of the KU Leuven spin-off companies ReMYND nv (Leuven, Belgium) and ADxNeuroSciences nv (Ghent, Belgium), but this did not influence in any way the content of this manuscript, nor is there a link between the studies reported and the activities of the aforementioned companies.

### COPYRIGHT

© 2018 Deprez *et al.* This is an open-access article released under the terms of the Creative Commons Attribution (CC BY) license, which allows the unrestricted use, distribution, and reproduction in any medium, provided the original author and source are acknowledged.

Please cite this article as: Marie-Anne Deprez, Elja Eskes, Tobias Wilms, Paula Ludovico, Joris Winderickx (2018). pH homeostasis links the nutrient sensing PKA/TORC1/Sch9 ménage-à-trois to stress tolerance and longevity. *Microbial Cell* 5(3): 119-136. doi: 10.15698/mic2018.03.618

## REFERENCES

- Platt FM (2014). Sphingolipid lysosomal storage disorders. *Nature* 510(7503): 68-75. doi: 10.1038/nature13476
- Bagh MB, Peng S, Chandra G, Zhang Z, Singh SP, Pattabiraman N, Liu A, Mukherjee AB (2017). Misrouting of v-ATPase subunit V0a1 dysregulates lysosomal acidification in a neurodegenerative lysosomal storage disease model. *Nat Commun* 8: 14612. doi: 10.1038/ncomms14612
- Onyenwoke RU, Brenman JE (2015). Lysosomal Storage Diseases—Regulating Neurodegeneration. *J Exp Neurosci* 9(Suppl 2): 81-91. doi: 10.4137/jen.s25475
- te Vrugte D, Speak AO, Wallom KL, Al Eisa N, Smith DA, Hendriksz CJ, Simmons L, Lachmann RH, Cousins A, Hartung R, Mengel E, Runz H, Beck M, Amraoui Y, Imrie J, Jacklin E, Riddick K, Yanjanin NM, Wassif CA, Rolfs A, Rimmele F, Wright N, Taylor C, Ramaswami U, Cox TM, Hastings C, Jiang X, Sidhu R, Ory DS, Arias B, et al. (2014). Relative acidic compartment volume as a lysosomal storage disorder-associated biomarker. *J Clin Invest* 124(3): 1320-1328. doi: 10.1172/jci72835
- Kato Y, Ozawa S, Miyamoto C, Maehata Y, Suzuki A, Maeda T, Baba Y (2013). Acidic extracellular microenvironment and cancer. *Cancer Cell Int* 13(1): 89. doi: 10.1186/1475-2867-13-89
- Martinez-Munoz GA, Kane P (2008). Vacuolar and plasma membrane proton pumps collaborate to achieve cytosolic pH homeostasis in yeast. *J Biol Chem* 283(29): 20309-20319. doi: 10.1074/jbc.M710470200
- Cyert MS, Philpott CC (2013). Regulation of cation balance in *Saccharomyces cerevisiae*. *Genetics* 193(3): 677-713. doi: 10.1534/genetics.112.147207
- Nishi T, Forgac M (2002). The vacuolar (H<sup>+</sup>)-ATPases—nature's most versatile proton pumps. *Nat Rev Mol Cell Biol* 3(2): 94-103. doi: 10.1038/nrm729
- Kane PM (2016). Proton Transport and pH Control in Fungi. In: Ramos J, Sychrova H, Kschischo M, editors. *Yeast Membrane Transport*. Springer-Verlag Berlin, Berlin; pp 33-68. doi: 10.1007/978-3-319-25304-6\_3
- Kane PM (1995). Disassembly and Reassembly of the Yeast Vacuolar H<sup>+</sup>-ATPase in Vivo. *J Biol Chem* 270(28): 17025-17032. PMID: 7622524
- Dechant R, Binda M, Lee SS, Pelet S, Winderickx J, Peter M (2010). Cytosolic pH is a second messenger for glucose and regulates the PKA pathway through V-ATPase. *EMBO J* 29(15): 2515-2526. doi: 10.1038/emboj.2010.138
- Diakov TT, Kane PM (2010). Regulation of vacuolar proton-translocating ATPase activity and assembly by extracellular pH. *J Biol Chem* 285(31): 23771-23778. doi: 10.1074/jbc.M110.110122
- Padilla-Lopez S, Pearce DA (2006). *Saccharomyces cerevisiae* lacking Btn1p modulate vacuolar ATPase activity to regulate pH imbalance in the vacuole. *J Biol Chem* 281(15): 10273-10280. doi: 10.1074/jbc.M510625200
- Dechant R, Peter M (2011). The N-terminal domain of the V-ATPase subunit 'a' is regulated by pH in vitro and in vivo. *Channels* 5(1): 4-8. doi: 10.4161/chan.5.1.13846
- Marshansky V (2007). The V-ATPase a2-subunit as a putative endosomal pH-sensor. *Biochem Soc Trans* 35(Pt 5): 1092-1099. doi: 10.1042/BST0351092
- Kawasaki-Nishi S, Bowers K, Nishi T, Forgac M, Stevens TH (2001). The amino-terminal domain of the vacuolar proton-translocating ATPase a subunit controls targeting and in vivo dissociation, and the carboxyl-terminal domain affects coupling of proton transport and ATP hydrolysis. *J Biol Chem* 276(50): 47411-47420. doi: 10.1074/jbc.M108310200
- Compton LM, Ikononov OC, Sbrissa D, Garg P, Shisheva A (2016). Active vacuolar H<sup>+</sup> ATPase and functional cycle of Rab5 are required for the vacuolation defect triggered by PtdIns(3,5)P<sub>2</sub> loss under PIKfyve or Vps34 deficiency. *Am J Physiol Cell Physiol* 311(3): C366-C377. doi: 10.1152/ajpcell.00104.2016
- Li SC, Diakov TT, Xu T, Tarsio M, Zhu W, Couoh-Cardel S, Weisman LS, Kane PM (2014). The signaling lipid PI(3,5)P<sub>2</sub> stabilizes V1–V0 sector interactions and activates the V-ATPase. *Mol Biol Cell* 25(8): 1251-1262. doi: 10.1091/mbc.E13-10-0563
- Banerjee S, Kane PM (2017). Direct interaction of the Golgi V-ATPase a-subunit isoform with PI(4)P drives localization of Golgi V-ATPases in yeast. *Mol Biol Cell* 28(19): 2518-2530. doi: 10.1091/mbc.E17-05-0316
- Davis-Kaplan SR, Compton MA, Flannery AR, Ward DM, Kaplan J, Stevens TH, Graham LA (2006). PKR1 encodes an assembly factor for the yeast V-type ATPase. *J Biol Chem* 281(42): 32025-32035. doi: 10.1074/jbc.M606451200
- Graham LA, Flannery AR, Stevens TH (2003). Structure and assembly of the yeast V-ATPase. *J Bioenerg Biomembr* 35(4): 301-312. PMID: 14635776
- Malkus P, Graham LA, Stevens TH, Schekman R (2004). Role of Vma21p in assembly and transport of the yeast vacuolar ATPase. *Mol Biol Cell* 15(11): 5075-5091. doi: 10.1091/mbc.E04-06-0514
- Ryan M, Graham LA, Stevens TH (2008). Voa1p functions in V-ATPase assembly in the yeast endoplasmic reticulum. *Mol Biol Cell* 19(12): 5131-5142. doi: 10.1091/mbc.E08-06-0629
- Brett CL, Kallay L, Hua Z, Green R, Chyou A, Zhang Y, Graham TR, Donowitz M, Rao R (2011). Genome-wide analysis reveals the vacuolar pH-stat of *Saccharomyces cerevisiae*. *PLoS One* 6(3): e17619. doi: 10.1371/journal.pone.0017619
- Orij R, Urbanus ML, Vizeacoumar FJ, Giaever G, Boone C, Nislow C, Brul S, Smits GJ (2012). Genome-wide analysis of intracellular pH reveals quantitative control of cell division rate by pH(c) in *Saccharomyces cerevisiae*. *Genome Biol* 13(9): 15. doi: 10.1186/gb-2012-13-9-r80
- Serrano R, Kielland-Brandt MC, Fink GR (1986). Yeast plasma membrane ATPase is essential for growth and has homology with (Na<sup>+</sup> + K<sup>+</sup>), K<sup>+</sup> and Ca<sup>2+</sup>-ATPases. *Nature* 319(6055): 689-693. doi: 10.1038/319689a0
- Portillo F (2000). Regulation of plasma membrane H<sup>+</sup>-ATPase in fungi and plants. *Biochim Biophys Acta* 1469(1): 31-42. PMID: 10692636
- Eraso P, Mazon MJ, Portillo F (2006). Yeast protein kinase Ptk2 localizes at the plasma membrane and phosphorylates in vitro the C-

- terminal peptide of the H<sup>+</sup>-ATPase. **Biochim Biophys Acta** 1758(2): 164-170. doi: 10.1016/j.bbame.2006.01.010
29. Estrada E, Agostinis P, Vandenheede JR, Goris J, Merlevede W, Francois J, Goffeau A, Ghislain M (1996). Phosphorylation of yeast plasma membrane H<sup>+</sup>-ATPase by casein kinase I. **J Biol Chem** 271(50): 32064-32072. PMID: 8943257
30. Mazon MJ, Eraso P, Portillo F (2015). Specific phosphoantibodies reveal two phosphorylation sites in yeast Pma1 in response to glucose. **FEMS Yeast Res** 15(5): fov030. doi: 10.1093/femsyr/fov030
31. Bouillet LE, Cardoso AS, Perovano E, Pereira RR, Ribeiro EM, Tropa MJ, Fietto LG, Tisi R, Martegani E, Castro IM, Brandao RL (2012). The involvement of calcium carriers and of the vacuole in the glucose-induced calcium signaling and activation of the plasma membrane H<sup>+</sup>-ATPase in *Saccharomyces cerevisiae* cells. **Cell Calcium** 51(1): 72-81. doi: 10.1016/j.ceca.2011.10.008
32. Yenush L, Merchan S, Holmes J, Serrano R (2005). pH-Responsive, posttranslational regulation of the Trk1 potassium transporter by the type 1-related Ppz1 phosphatase. **Mol Cell Biol** 25(19): 8683-8692. doi: 10.1128/MCB.25.19.8683-8692.2005
33. Huang CJ, Chang A (2011). pH-dependent Cargo Sorting from the Golgi. **J Biol Chem** 286(12): 10058-10065. doi: 10.1074/jbc.M110.197889
34. Fernandes AR, Sa-Correia I (2003). Transcription patterns of PMA1 and PMA2 genes and activity of plasma membrane H<sup>+</sup>-ATPase in *Saccharomyces cerevisiae* during diauxic growth and stationary phase. **Yeast** 20(3): 207-219. doi: 10.1002/yea.957
35. Schlessler A, Ulaszewski S, Ghislain M, Goffeau A (1988). A second transport ATPase gene in *Saccharomyces cerevisiae*. **J Biol Chem** 263(36): 19480-19487. PMID: 2904437
36. Eskes E, Deprez MA, Wilms T, Winderickx J (2017). pH homeostasis in yeast; the phosphate perspective. **Curr Genet**. Ahead of print doi: 10.1007/s00294-017-0743-2
37. Gournas C, Prevost M, Krammer EM, Andre B (2016). Function and Regulation of Fungal Amino Acid Transporters: Insights from Predicted Structure. **Adv Exp Med Biol** 892: 69-106. doi: 10.1007/978-3-319-25304-6\_4
38. Russnak R, Konczal D, McIntire SL (2001). A family of yeast proteins mediating bidirectional vacuolar amino acid transport. **J Biol Chem** 276(26): 23849-23857. doi: 10.1074/jbc.M008028200
39. Conrad M, Schothorst J, Kankipati HN, Van Zeebroeck G, Rubio-Teixeira M, Thevelein JM (2014). Nutrient sensing and signaling in the yeast *Saccharomyces cerevisiae*. **Fems Microbiol Rev** 38(2): 254-299. doi: 10.1111/1574-6976.12065
40. Sampaio-Marques B, Burhans WC, Ludovico P (2014). Longevity pathways and maintenance of the proteome: the role of autophagy and mitophagy during yeast ageing. **Microb Cell** 1(4): 118-127. doi: 10.15698/mic2014.04.136
41. Smets B, Ghillebert R, De Snijder P, Binda M, Swinnen E, De Virgilio C, Winderickx J (2010). Life in the midst of scarcity: adaptations to nutrient availability in *Saccharomyces cerevisiae*. **Curr Genet** 56(1): 1-32. doi: 10.1007/s00294-009-0287-1
42. Weidberg H, Moretto F, Spedale G, Amon A, van Werven FJ (2016). Nutrient Control of Yeast Gametogenesis Is Mediated by TORC1, PKA and Energy Availability. **PLoS Genet** 12(6): e1006075. doi: 10.1371/journal.pgen.1006075
43. Rolland F, De Winde JH, Lemaire K, Boles E, Thevelein JM, Winderickx J (2000). Glucose-induced cAMP signalling in yeast requires both a G-protein coupled receptor system for extracellular glucose detection and a separable hexose kinase-dependent sensing process. **Mol Microbiol** 38(2): 348-358. doi: 10.1046/j.1365-2958.2000.02125.x
44. Stracka D, Jozefczuk S, Rudroff F, Sauer U, Hall MN (2014). Nitrogen source activates TOR (target of rapamycin) complex 1 via glutamine and independently of Gtr/Rag proteins. **J Biol Chem** 289(36): 25010-25020. doi: 10.1074/jbc.M114.574335
45. Binda M, Peli-Gulli MP, Bonfils G, Panchaud N, Urban J, Sturgill TW, Loewith R, De Virgilio C (2009). The Vam6 GEF controls TORC1 by activating the EGO complex. **Mol Cell** 35(5): 563-573. doi: 10.1016/j.molcel.2009.06.033
46. Powis K, De Virgilio C (2016). Conserved regulators of Rag GTPases orchestrate amino acid-dependent TORC1 signaling. **Cell Discov** 2: 15049. doi: 10.1038/celldisc.2015.49
47. Zoncu R, Bar-Peled L, Efeyan A, Wang SY, Sancak Y, Sabatini DM (2011). mTORC1 Senses Lysosomal Amino Acids Through an Inside-Out Mechanism That Requires the Vacuolar H<sup>+</sup>-ATPase. **Science** 334(6056): 678-683. doi: 10.1126/science.1207056
48. Nicastro R, Sardu A, Panchaud N, De Virgilio C (2017). The Architecture of the Rag GTPase Signaling Network. **Biomolecules** 7(3): 48. doi: 10.3390/biom7030048
49. Thomas JD, Zhang YJ, Wei YH, Cho JH, Morris LE, Wang HY, Zheng XF (2014). Rab1A is an mTORC1 activator and a colorectal oncogene. **Cancer Cell** 26(5): 754-769. doi: 10.1016/j.ccell.2014.09.008
50. Kim A, Cunningham KW (2015). A LAPF/phafin1-like protein regulates TORC1 and lysosomal membrane permeabilization in response to endoplasmic reticulum membrane stress. **Mol Biol Cell** 26(25): 4631-4645. doi: 10.1091/mbc.E15-08-0581
51. Tanigawa M, Maeda T (2017). An In Vitro TORC1 Kinase Assay That Recapitulates the Gtr-Independent Glutamine-Responsive TORC1 Activation Mechanism on Yeast Vacuoles. **Mol Cell Biol** 37(14). doi: 10.1128/MCB.00075-17
52. Urban J, Soulard A, Huber A, Lippman S, Mukhopadhyay D, Deloche O, Wanke V, Anrather D, Ammerer G, Riezman H, Broach JR, De Virgilio C, Hall MN, Loewith R (2007). Sch9 is a major target of TORC1 in *Saccharomyces cerevisiae*. **Mol Cell** 26(5): 663-674. doi: 10.1016/j.molcel.2007.04.020
53. Swinnen E, Ghillebert R, Wilms T, Winderickx J (2014). Molecular mechanisms linking the evolutionary conserved TORC1-Sch9 nutrient signalling branch to lifespan regulation in *Saccharomyces cerevisiae*. **FEMS Yeast Res** 14(1): 17-32. doi: 10.1111/1567-1364.12097
54. Roelants FM, Torrance PD, Thorner J (2004). Differential roles of PDK1- and PDK2-phosphorylation sites in the yeast AGC kinases Ypk1, Pkc1 and Sch9. **Microbiology** 150(Pt 10): 3289-3304. doi: 10.1099/mic.0.27286-0
55. Voordeckers K, Kimpe M, Haesendonckx S, Louwet W, Versele M, Thevelein JM (2011). Yeast 3-Phosphoinositide-dependent Protein Kinase-1 (PDK1) Orthologs Pkh1-3 Differentially Regulate Phosphorylation of Protein Kinase A (PKA) and the Protein Kinase B (PKB)/S6K Ortholog Sch9. **J Biol Chem** 286(25): 22017-22027. doi: 10.1074/jbc.M110.200071
56. Crauwels M, Donaton MCV, Pernambuco MB, Winderickx J, deWinde JH, Thevelein JM (1997). The Sch9 protein kinase in the yeast *Saccharomyces cerevisiae* controls cAPK activity and is required for nitrogen activation of the fermentable-growth-medium-induced (FGM) pathway. **Microbiology** 143(Pt 8): 2627-2637. doi: 10.1099/00221287-143-8-2627
57. Lu JY, Lin YY, Sheu JC, Wu JT, Lee FJ, Chen Y, Lin MI, Chiang FT, Tai TY, Berger SL, Zhao Y, Tsai KS, Zhu H, Chuang LM, Boeke JD (2011). Acetylation of yeast AMPK controls intrinsic aging independently of caloric restriction. **Cell** 146(6): 969-979. doi: 10.1016/j.cell.2011.07.044

58. Roosen J, Engelen K, Marchal K, Mathys J, Griffioen G, Cameroni E, Thevelein JM, De Virgilio C, De Moor B, Winderickx J (2005). PKA and Sch9 control a molecular switch important for the proper adaptation to nutrient availability. *Mol Microbiol* 55(3): 862-880. doi: 10.1111/j.1365-2958.2004.04429.x
59. Smets B, De Snijder P, Engelen K, Joossens E, Ghillebert R, Thevissen K, Marchal K, Winderickx J (2008). Genome-wide expression analysis reveals TORC1-dependent and -independent functions of Sch9. *FEMS Yeast Res* 8(8): 1276-1288. doi: 10.1111/j.1567-1364.2008.00432.x
60. Yorimitsu T, Zaman S, Broach JR, Klionsky DJ (2007). Protein kinase A and Sch9 cooperatively regulate induction of autophagy in *Saccharomyces cerevisiae*. *Mol Biol Cell* 18(10): 4180-4189. doi: 10.1091/mbc.E07-05-0485
61. Pedruzzi I, Dubouloz F, Cameroni E, Wanke V, Roosen J, Winderickx J, De Virgilio C (2003). TOR and PKA signaling pathways converge on the protein kinase Rim15 to control entry into G(0). *Mol Cell* 12(6): 1607-1613. doi: 10.1016/s1097-2765(03)00485-4
62. Swinnen E, Wanke V, Roosen J, Smets B, Dubouloz F, Pedruzzi I, Cameroni E, De Virgilio C, Winderickx J (2006). Rim15 and the crossroads of nutrient signalling pathways in *Saccharomyces cerevisiae*. *Cell Div* 1:3. doi: 10.1186/1747-1028-1-3
63. Casal M, Cardoso H, Leao C (1996). Mechanisms regulating the transport of acetic acid in *Saccharomyces cerevisiae*. *Microbiology* 142 (Pt 6): 1385-1390. doi: 10.1099/13500872-142-6-1385
64. Bracey D, Holyoak CD, Coote PJ (1998). Comparison of the inhibitory effect of sorbic acid and amphotericin B on *Saccharomyces cerevisiae*: is growth inhibition dependent on reduced intracellular pH? *J Appl Microbiol* 85(6): 1056-1066. doi: 10.1111/j.1365-2672.1998.tb05271.x
65. Mollapour M (2008). Novel stress responses facilitate *Saccharomyces cerevisiae* growth in the presence of the monocarboxylate preservatives. *Yeast* 25(3): 169-177. doi: 10.1002/yea.1576
66. Orij R, Brul S, Smits GJ (2011). Intracellular pH is a tightly controlled signal in yeast. *Biochim Biophys Acta* 1810(10): 933-944. doi: 10.1016/j.bbagen.2011.03.011
67. Chan CY, Dominguez D, Parra KJ (2016). Regulation of Vacuolar H plus -ATPase (V-ATPase) Reassembly by Glycolysis Flow in 6-Phosphofructo-1-kinase (PFK-1)-deficient Yeast Cells. *J Biol Chem* 291(30): 15820-15829. doi: 10.1074/jbc.M116.717488
68. Kane PA (2006). The where, when, and how of organelle acidification by the yeast vacuolar H+-ATPase. *Microbiol Mol Biol Rev* 70(1): 177-191. doi: 10.1128/mmr.70.1.177-191.2006
69. Parra KJ, Kane PM (1998). Reversible association between the V-1 and V-0 domains of yeast vacuolar H+-ATPase is an unconventional glucose-induced effect. *Mol Cell Biol* 18(12): 7064-7074. PMID: 9819393
70. Wilms T, Swinnen E, Eskes E, Dolz-Edo L, Uwineza A, Van Essche R, Rosseels J, Zabrocki P, Cameroni E, Franssens V, De Virgilio C, Smits GJ, Winderickx J (2017). The yeast protein kinase Sch9 adjusts V-ATPase assembly/ disassembly to control pH homeostasis and longevity in response to glucose availability. *PLoS Genet* 13(6): e1006835. doi: 10.1371/journal.pgen.1006835
71. Orij R, Postmus J, Ter Beek A, Brul S, Smits GJ (2009). In vivo measurement of cytosolic and mitochondrial pH using a pH-sensitive GFP derivative in *Saccharomyces cerevisiae* reveals a relation between intracellular pH and growth. *Microbiology* 155(Pt 1): 268-278. doi: 10.1099/mic.0.022038-0
72. Ramos S, Balbin M, Raposo M, Valle E, Pardo LA (1989). The mechanism of intracellular acidification induced by glucose in *Saccharomyces cerevisiae*. *J Gen Microbiol* 135(9): 2413-2422. doi: 10.1099/00221287-135-9-2413
73. Dechant R, Saad S, Ibanez AJ, Peter M (2014). Cytosolic pH Regulates Cell Growth through Distinct GTPases, Arf1 and Gtr1, to Promote Ras/PKA and TORC1 Activity. *Mol Cell* 55(3): 409-421. doi: 10.1016/j.molcel.2014.06.002
74. Bond S, Forgac M (2008). The Ras/cAMP/Protein Kinase A Pathway Regulates Glucose-dependent Assembly of the Vacuolar (H+)-ATPase in Yeast. *J Biol Chem* 283(52): 36513-36521. doi: 10.1074/jbc.M805232200
75. Voss M, Vitavska O, Walz B, Wiczorek H, Baumann O (2007). Stimulus-induced phosphorylation of vacuolar H(+)-ATPase by protein kinase A. *J Biol Chem* 282(46): 33735-33742. doi: 10.1074/jbc.M703368200
76. Alzamora R, Thali RF, Gong F, Smolak C, Li H, Baty CJ, Bertrand CA, Auchli Y, Brunisholz RA, Neumann D, Hallows KR, Pastor-Soler NM (2010). PKA regulates vacuolar H+-ATPase localization and activity via direct phosphorylation of the a subunit in kidney cells. *J Biol Chem* 285(32): 24676-24685. doi: 10.1074/jbc.M110.106278
77. Williamson T, Adiamah D, Schwartz JM, Stateva L (2012). Exploring the genetic control of glycolytic oscillations in *Saccharomyces cerevisiae*. *BMC Syst Biol* 6(1): 108. doi: 10.1186/1752-0509-6-108
78. Chan CY, Parra KJ (2014). Yeast phosphofructokinase-1 subunit Pfk2p is necessary for pH homeostasis and glucose-dependent vacuolar ATPase reassembly. *J Biol Chem* 289(28): 19448-19457. doi: 10.1074/jbc.M114.569855
79. Lu M, Ammar D, Ives H, Albrecht F, Gluck SL (2007). Physical interaction between aldolase and vacuolar H+-ATPase is essential for the assembly and activity of the proton pump. *J Biol Chem* 282(34): 24495-24503. doi: 10.1074/jbc.M702598200
80. Lu M, Holliday LS, Zhang L, Dunn WA, Jr., Gluck SL (2001). Interaction between aldolase and vacuolar H+-ATPase: evidence for direct coupling of glycolysis to the ATP-hydrolyzing proton pump. *J Biol Chem* 276(32): 30407-30413. doi: 10.1074/jbc.M008768200
81. Dihazi H, Kessler R, Eschrich K (2003). Glucose-induced stimulation of the Ras-cAMP pathway in yeast leads to multiple phosphorylations and activation of 6-phosphofructo-2-kinase. *Biochemistry* 42(20): 6275-6282. doi: 10.1021/bi034167r
82. Seol JH, Shevchenko A, Shevchenko A, Deshaies RJ (2001). Skp1 forms multiple protein complexes, including RAVE, a regulator of V-ATPase assembly. *Nat Cell Biol* 3(4): 384-391. doi: 10.1038/35070067
83. Sardon AM, Tarsio M, Kane PM (2002). The RAVE complex is essential for stable assembly of the yeast V-ATPase. *J Biol Chem* 277(16): 13831-13839. doi: 10.1074/jbc.M200682200
84. Portillo F, Mazon MJ (1986). The *Saccharomyces cerevisiae* start mutant carrying the cdc25 mutation is defective in activation of plasma membrane ATPase by glucose. *J Bacteriol* 168(3): 1254-1257. doi: 10.1128/jb.168.3.1254-1257.1986
85. Ulaszewski S, Hilger F, Goffeau A (1989). Cyclic AMP controls the plasma membrane H+-ATPase activity from *Saccharomyces cerevisiae*. *FEBS Lett* 245(1-2): 131-136. doi: 10.1016/0014-5793(89)80206-6
86. dos Passos JB, Vanhalewyn M, Brandao RL, Castro IM, Nicoli JR, Thevelein JM (1992). Glucose-induced activation of plasma membrane H(+)-ATPase in mutants of the yeast *Saccharomyces cerevisiae* affected in cAMP metabolism, cAMP-dependent protein phosphorylation and the initiation of glycolysis. *Biochim Biophys Acta* 1136(1): 57-67. doi: 10.1016/0167-4889(92)90085-P
87. Mazon MJ, Behrens MM, Portillo F, Pinon R (1989). cAMP- and RAS-independent nutritional regulation of plasma-membrane H+-

- ATPase activity in *Saccharomyces cerevisiae*. **J Gen Microbiol** 135(6): 1453-1460. doi: 10.1099/00221287-135-6-1453
88. Mahmoud S, Planes MD, Cabedo M, Trujillo C, Rienzo A, Caballero-Molada M, Sharma SC, Montesinos C, Mulet JM, Serrano R (2017). TOR complex 1 regulates the yeast plasma membrane proton pump and pH and potassium homeostasis. **FEBS Lett** 591(13): 1993-2002. doi: 10.1002/1873-3468.12673
89. Bonangelino CJ, Catlett NL, Weisman LS (1997). Vac7p, a novel vacuolar protein, is required for normal vacuole inheritance and morphology. **Mol Cell Biol** 17(12): 6847. doi: 10.1128/MCB.17.12.6847
90. Bonangelino CJ, Nau JJ, Duex JE, Brinkman M, Wurmser AE, Gary JD, Emr SD, Weisman LS (2002). Osmotic stress– induced increase of phosphatidylinositol 3,5- bisphosphate requires Vac14p, an activator of the lipid kinase Fab1p. **J Cell Biol** 156(6): 1015-1028. doi: 10.1083/jcb.200201002
91. Dove SK, McEwen RK, Mayes A, Hughes DC, Beggs JD, Michell RH (2002). Vac14 controls PtdIns(3,5)P(2) synthesis and Fab1-dependent protein trafficking to the multivesicular body. **Curr Biol** 12(11): 885-893. doi: 10.1016/S0960-9822(02)00891-6
92. Bridges D, Fisher K, Zolov SN, Xiong T, Inoki K, Weisman LS, Saltiel AR (2012). Rab5 proteins regulate activation and localization of target of rapamycin complex 1. **J Biol Chem** 287(25): 20913-20921. doi: 10.1074/jbc.M111.334060
93. Jin N, Lang MJ, Weisman LS (2016). Phosphatidylinositol 3,5-bisphosphate: regulation of cellular events in space and time. **Biochem Soc Trans** 44(1): 177-184. doi: 10.1042/BST20150174
94. Jin Y, Weisman LS (2015). The vacuole/lysosome is required for cell-cycle progression. **eLife** 4: e08160. doi: 10.7554/eLife.08160
95. Rudge SA, Anderson DM, Emr SD (2004). Vacuole size control: regulation of PtdIns(3,5)P2 levels by the vacuole-associated Vac14-Fig4 complex, a PtdIns(3,5)P2-specific phosphatase. **Mol Biol Cell** 15(1): 24-36. doi: 10.1091/mbc.E03-05-0297
96. Stauffer B, Powers T (2015). Target of rapamycin signaling mediates vacuolar fission caused by endoplasmic reticulum stress in *Saccharomyces cerevisiae*. **Mol Biol Cell** 26(25): 4618-4630. doi: 10.1091/mbc.E15-06-0344
97. Young BP, Shin JJ, Oriji R, Chao JT, Li SC, Guan XL, Khong A, Jan E, Wenk MR, Prinz WA, Smits GJ, Loewen CJ (2010). Phosphatidic acid is a pH biosensor that links membrane biogenesis to metabolism. **Science** 329(5995): 1085-1088. doi: 10.1126/science.1191026
98. Purwin C, Nicolay K, Scheffers WA, Holzer H (1986). Mechanism of control of adenylate cyclase activity in yeast by fermentable sugars and carbonyl cyanide m-chlorophenylhydrazone. **J Biol Chem** 261(19): 8744-8749. PMID: 3522579
99. Thevelein JM (1991). Fermentable sugars and intracellular acidification as specific activators of the RAS-adenylate cyclase signalling pathway in yeast: the relationship to nutrient-induced cell cycle control. **Mol Microbiol** 5(6): 1301-1307. doi: 10.1111/j.1365-2958.1991.tb00776.x
100. Trevillyan JM, Pall ML (1979). Control of cyclic adenosine 3',5'-monophosphate levels by depolarizing agents in fungi. **J Bacteriol** 138(2): 397-403. PMID: 220213
101. Colombo S, Ma P, Cauwenberg L, Winderickx J, Crauwels M, Teunissen A, Nauwelaers D, de Winde JH, Gorwa MF, Colavizza D, Thevelein JM (1998). Involvement of distinct G-proteins, Gpa2 and Ras, in glucose- and intracellular acidification-induced cAMP signalling in the yeast *Saccharomyces cerevisiae*. **EMBO J** 17(12): 3326-3341. doi: 10.1093/emboj/17.12.3326
102. Rolland F, Wanke V, Cauwenberg L, Ma P, Boles E, Vanoni M, de Winde JH, Thevelein JM, Winderickx J (2001). The role of hexose transport and phosphorylation in cAMP signalling in the yeast *Saccharomyces cerevisiae*. **FEMS Yeast Res** 1(1): 33-45. doi: 10.1111/j.1567-1364.2001.tb00011.x
103. Longo VD, Shadel GS, Kaerberlein M, Kennedy B (2012). Replicative and chronological aging in *Saccharomyces cerevisiae*. **Cell Metab** 16(1): 18-31. doi: 10.1016/j.cmet.2012.06.002
104. Denoth Lippuner A, Julou T, Barral Y (2014). Budding yeast as a model organism to study the effects of age. **FEMS Microbiol Rev** 38(2): 300-325. doi: 10.1111/1574-6976.12060
105. Dakik P, Titorenko VI (2016). Communications between Mitochondria, the Nucleus, Vacuoles, Peroxisomes, the Endoplasmic Reticulum, the Plasma Membrane, Lipid Droplets, and the Cytosol during Yeast Chronological Aging. **Front Genet** 7(177). doi: 10.3389/fgene.2016.00177
106. Zhang N, Cao L (2017). Starvation signals in yeast are integrated to coordinate metabolic reprogramming and stress response to ensure longevity. **Curr Genet** 63(5): 839-843. doi: 10.1007/s00294-017-0697-4
107. Tsang F, Lin SJ (2015). Less is more: Nutrient limitation induces cross-talk of nutrient sensing pathways with NAD<sup>+</sup> homeostasis and contributes to longevity. **Front Biol** 10(4): 333-357. doi: 10.1007/s11515-015-1367-x
108. Arlia-Ciommo A, Piano A, Leonov A, Svistkova V, Titorenko VI (2014). Quasi-programmed aging of budding yeast: a trade-off between programmed processes of cell proliferation, differentiation, stress response, survival and death defines yeast lifespan. **Cell Cycle** 13(21): 3336-3349. doi: 10.4161/15384101.2014.965063
109. Madeo F, Zimmermann A, Maiuri MC, Kroemer G (2015). Essential role for autophagy in life span extension. **J Clin Invest** 125(1): 85-93. doi: 10.1172/JCI73946
110. Bitto A, Wang AM, Bennett CF, Kaerberlein M (2015). Biochemical Genetic Pathways that Modulate Aging in Multiple Species. **Cold Spring Harb Perspect Med** 5(11). doi: 10.1101/cshperspect.a025114
111. Ludovico P, Burhans WC (2014). Reactive oxygen species, ageing and the hormesis police. **FEMS yeast research** 14(1): 33-39. doi: 10.1111/1567-1364.12070
112. Wierman MB, Smith JS (2014). Yeast sirtuins and the regulation of aging. **FEMS yeast research** 14(1): 73-88. doi: 10.1111/1567-1364.12115
113. Delaney JR, Murakami C, Chou A, Carr D, Schleit J, Sutphin GL, An EH, Castanza AS, Fletcher M, Goswami S, Higgins S, Holmberg M, Hui J, Jelic M, Jeong KS, Kim JR, Klum S, Liao E, Lin MS, Lo W, Miller H, Moller R, Peng ZJ, Pollard T, Pradeep P, Pruett D, Rai D, Ros V, Schuster A, Singh M, et al. (2013). Dietary restriction and mitochondrial function link replicative and chronological aging in *Saccharomyces cerevisiae*. **Exp Gerontol** 48(10): 1006-1013. doi: 10.1016/j.exger.2012.12.001
114. Murakami C, Delaney JR, Chou A, Carr D, Schleit J, Sutphin GL, An EH, Castanza AS, Fletcher M, Goswami S, Higgins S, Holmberg M, Hui J, Jelic M, Jeong KS, Kim JR, Klum S, Liao E, Lin MS, Lo W, Miller H, Moller R, Peng ZJ, Pollard T, Pradeep P, Pruett D, Rai D, Ros V, Schuster A, Singh M, et al. (2012). pH neutralization protects against reduction in replicative lifespan following chronological aging in yeast. **Cell Cycle** 11(16): 3087-3096. doi: 10.4161/cc.21465
115. Mirisola MG, Longo VD (2012). Acetic acid and acidification accelerate chronological and replicative aging in yeast. **Cell Cycle** 11(19): 3532-3533. doi: 10.4161/cc.22042
116. Orlandi I, Ronzulli R, Casatta N, Vai M (2013). Ethanol and acetate acting as carbon/energy sources negatively affect yeast chronological aging. **Oxid Med Cell Longev** 2013: 802870. doi: 10.1155/2013/802870

117. Thomas KC, Hynes SH, Ingledew WM (2002). Influence of medium buffering capacity on inhibition of *Saccharomyces cerevisiae* growth by acetic and lactic acids. **Appl Environ Microbiol** 68(4): 1616-1623. doi: 10.1128/AEM.68.4.1616-1623.2002
118. Henderson KA, Hughes AL, Gottschling DE (2014). Mother-daughter asymmetry of pH underlies aging and rejuvenation in yeast. **eLife** 3:e03504. doi: 10.7554/eLife.03504
119. Smardon AM, Kane PM (2014). Loss of vacuolar H<sup>+</sup>-ATPase activity in organelles signals ubiquitination and endocytosis of the yeast plasma membrane proton pump Pma1p. **J Biol Chem** 289(46): 32316-32326. doi: 10.1074/jbc.M114.574442
120. Swiecilo A (2016). Cross-stress resistance in *Saccharomyces cerevisiae* yeast--new insight into an old phenomenon. **Cell Stress Chaperones** 21(2): 187-200. doi: 10.1007/s12192-016-0667-7
121. Calabrese EJ, Bachmann KA, Bailer AJ, Bolger PM, Borak J, Cai L, Cedergreen N, Cherian MG, Chiueh CC, Clarkson TW, Cook RR, Diamond DM, Doolittle DJ, Dorato MA, Duke SO, Feinendegen L, Gardner DE, Hart RW, Hastings KL, Hayes AW, Hoffmann GR, Ives JA, Jaworowski Z, Johnson TE, Jonas WB, Kaminski NE, Keller JG, Klaunig JE, Knudsen TB, Kozumbo WJ, et al. (2007). Biological stress response terminology: Integrating the concepts of adaptive response and pre-conditioning stress within a hormetic dose-response framework. **Toxicol Appl Pharmacol** 222(1): 122-128. doi: 10.1016/j.taap.2007.02.015
122. Giannattasio S, Guaragnella N, Corte-Real M, Passarella S, Marra E (2005). Acid stress adaptation protects *Saccharomyces cerevisiae* from acetic acid-induced programmed cell death. **Gene** 354:93-98. doi: 10.1016/j.gene.2005.03.030
123. Hoffmann GR, Moczula AV, Laterza AM, Macneil LK, Tartagliano JP (2013). Adaptive response to hydrogen peroxide in yeast: induction, time course, and relationship to dose-response models. **Environ Mol Mutagen** 54(6): 384-396. doi: 10.1002/em.21785
124. Mesquita A, Weinberger M, Silva A, Sampaio-Marques B, Almeida B, Leao C, Costa V, Rodrigues F, Burhans WC, Ludovico P (2010). Caloric restriction or catalase inactivation extends yeast chronological lifespan by inducing H<sub>2</sub>O<sub>2</sub> and superoxide dismutase activity. **Proc Natl Acad Sci USA** 107(34): 15123-15128. doi: 10.1073/pnas.1004432107
125. Casado C, Gonzalez A, Platara M, Ruiz A, Arino J (2011). The role of the protein kinase A pathway in the response to alkaline pH stress in yeast. **Biochem J** 438(3): 523-533. doi: 10.1042/BJ20110607
126. Mira NP, Becker JD, Sa-Correia I (2010). Genomic expression program involving the Haa1p-regulon in *Saccharomyces cerevisiae* response to acetic acid. **OMICS** 14(5): 587-601. doi: 10.1089/omi.2010.0048
127. Mira NP, Teixeira MC, Sa-Correia I (2010). Adaptive response and tolerance to weak acids in *Saccharomyces cerevisiae*: a genome-wide view. **OMICS** 14(5): 525-540. doi: 10.1089/omi.2010.0072
128. Ludovico P, Sousa MJ, Silva MT, Leao C, Corte-Real M (2001). *Saccharomyces cerevisiae* commits to a programmed cell death process in response to acetic acid. **Microbiology** 147(Pt 9): 2409-2415. doi: 10.1099/00221287-147-9-2409
129. Ludovico P, Rodrigues F, Almeida A, Silva MT, Barrientos A, Corte-Real M (2002). Cytochrome c release and mitochondria involvement in programmed cell death induced by acetic acid in *Saccharomyces cerevisiae*. **Mol Biol Cell** 13(8): 2598-2606. doi: 10.1091/mbc.E01-12-0161
130. Guaragnella N, Zdravlevic M, Antonacci L, Passarella S, Marra E, Giannattasio S (2012). The role of mitochondria in yeast programmed cell death. **Front Oncol** 2:70. doi: 10.3389/fonc.2012.00070
131. Pan Y (2011). Mitochondria, reactive oxygen species, and chronological aging: a message from yeast. **Exp Gerontol** 46(11): 847-852. doi: 10.1016/j.exger.2011.08.007
132. Aerts AM, Zabrocki P, Govaert G, Mathys J, Carmona-Gutierrez D, Madeo F, Winderickx J, Cammue BP, Thevissen K (2009). Mitochondrial dysfunction leads to reduced chronological lifespan and increased apoptosis in yeast. **FEBS Lett** 583(1): 113-117. doi: 10.1016/j.febslet.2008.11.028
133. Fabrizio P, Battistella L, Vardavas R, Gattazzo C, Liou LL, Diaspro A, Dossen JW, Gralla EB, Longo VD (2004). Superoxide is a mediator of an altruistic aging program in *Saccharomyces cerevisiae*. **J Cell Biol** 166(7): 1055-1067. doi: 10.1083/jcb.200404002
134. Hughes AL, Gottschling DE (2012). An early age increase in vacuolar pH limits mitochondrial function and lifespan in yeast. **Nature** 492(7428): 261-265. doi: 10.1038/nature11654
135. Laun P, Pichova A, Madeo F, Fuchs J, Ellinger A, Kohlwein S, Dawes I, Frohlich KU, Breitenbach M (2001). Aged mother cells of *Saccharomyces cerevisiae* show markers of oxidative stress and apoptosis. **Mol Microbiol** 39(5): 1166-1173. doi: 10.1111/j.1365-2958.2001.02317.x
136. Jazwinski SM (2015). Mitochondria to nucleus signaling and the role of ceramide in its integration into the suite of cell quality control processes during aging. **Ageing Res Rev** 23(Pt A): 67-74. doi: 10.1016/j.arr.2014.12.007
137. Traven A, Wong JM, Xu D, Sopta M, Ingles CJ (2001). Interorganellar communication. Altered nuclear gene expression profiles in a yeast mitochondrial dna mutant. **J Biol Chem** 276(6): 4020-4027. doi: 10.1074/jbc.M006807200
138. Hu J, Wei M, Mirzaei H, Madia F, Mirisola M, Amparo C, Chagoury S, Kennedy B, Longo VD (2014). Tor-Sch9 deficiency activates catabolism of the ketone body-like acetic acid to promote trehalose accumulation and longevity. **Ageing Cell** 13(3): 457-467. doi: 10.1111/ace1.12202
139. Petitjean M, Teste MA, Francois JM, Parrou JL (2015). Yeast Tolerance to Various Stresses Relies on the Trehalose-6P Synthase (Tps1) Protein, Not on Trehalose. **J Biol Chem** 290(26): 16177-16190. doi: 10.1074/jbc.M115.653899
140. Petitjean M, Teste MA, Leger-Silvestre I, Francois JM, Parrou JL (2017). A new function for the yeast trehalose-6P synthase (Tps1) protein, as key pro-survival factor during growth, chronological ageing, and apoptotic stress. **Mech Ageing Dev** 161(Pt B): 234-246. doi: 10.1016/j.mad.2016.07.011
141. Kyrjakov P, Beach A, Richard VR, Burstein MT, Leonov A, Levy S, Titorenko VI (2012). Caloric restriction extends yeast chronological lifespan by altering a pattern of age-related changes in trehalose concentration. **Front Physiol** 3:256. doi: 10.3389/fphys.2012.00256
142. Nakamura N, Matsuura A, Wada Y, Ohsumi Y (1997). Acidification of vacuoles is required for autophagic degradation in the yeast, *Saccharomyces cerevisiae*. **J Biochem** 121(2): 338-344. PMID: 9089409
143. Ruckenstuhl C, Netzberger C, Entfellner I, Carmona-Gutierrez D, Kickenweiz T, Stekovic S, Gleixner C, Schmid C, Klug L, Sorgo AG, Eisenberg T, Buttner S, Marino G, Koziel R, Jansen-Durr P, Frohlich KU, Kroemer G, Madeo F (2014). Lifespan Extension by Methionine Restriction Requires Autophagy-Dependent Vacuolar Acidification. **PLoS Genet** 10(5). doi: 10.1371/journal.pgen.1004347
144. Tang F, Watkins JW, Bermudez M, Gray R, Gaban A, Portie K, Grace S, Kleve M, Craciun G (2008). A life-span extending form of autophagy employs the vacuole-vacuole fusion machinery. **Autophagy** 4(7): 874-886. doi: 10.4161/auto.6556
145. Eisenberg T, Knauer H, Schauer A, Buttner S, Ruckenstuhl C, Carmona-Gutierrez D, Ring J, Schroeder S, Magnes C, Antonacci L, Fussi H, Deszcz L, Hartl R, Schraml E, Criollo A, Megalou E, Weiskopf D, Laun P, Heeren G, Breitenbach M, Grubeck-Loebenstien B, Herker E, Fahrenkrog B, Frohlich KU, Sinner F, Tavernarakis N, Minois N, Kroe-

- mer G, Madeo F (2009). Induction of autophagy by spermidine promotes longevity. **Nat Cell Biol** 11(11): 1305-1314. doi: 10.1038/ncb1975
146. Miller SB, Ho CT, Winkler J, Khokhrina M, Neuner A, Mohamed MY, Guilbride DL, Richter K, Lisby M, Schiebel E, Mogk A, Bukau B (2015). Compartment-specific aggregates direct distinct nuclear and cytoplasmic aggregate deposition. **EMBO J** 34(6): 778-797. doi: 10.15252/embj.201489524
147. Kaganovich D, Kopito R, Frydman J (2008). Misfolded proteins partition between two distinct quality control compartments. **Nature** 454(7208): 1088-1095. doi: 10.1038/nature07195
148. Song J, Yang Q, Yang J, Larsson L, Hao X, Zhu X, Malmgren-Hill S, Cvijovic M, Fernandez-Rodriguez J, Grantham J, Gustafsson CM, Liu B, Nystrom T (2014). Essential genetic interactors of SIR2 required for spatial sequestration and asymmetrical inheritance of protein aggregates. **PLoS Genet** 10(7): e1004539. doi: 10.1371/journal.pgen.1004539
149. Liu B, Larsson L, Caballero A, Hao X, Oling D, Grantham J, Nystrom T (2010). The polarisome is required for segregation and retrograde transport of protein aggregates. **Cell** 140(2): 257-267. doi: 10.1016/j.cell.2009.12.031
150. Liu B, Larsson L, Franssens V, Hao X, Hill SM, Andersson V, Hoglund D, Song J, Yang X, Oling D, Grantham J, Winderickx J, Nystrom T (2011). Segregation of protein aggregates involves actin and the polarity machinery. **Cell** 147(5): 959-961. doi: 10.1016/j.cell.2011.11.018
151. Liu Y, Liu N, Wu D, Bi Q, Meng S (2015). The longevity of *tor1Delta*, *sch9Delta*, and *ras2Delta* mutants depends on actin dynamics in *Saccharomyces cerevisiae*. **Cell Biosci** 5:18. doi: 10.1186/s13578-015-0008-z
152. Amberg D, Leadsham JE, Kotiadis V, Gourlay CW (2012). Cellular ageing and the actin cytoskeleton. **Subcell Biochem** 57:331-352. doi: 10.1007/978-94-007-2561-4\_15
153. Buttner S, Delay C, Franssens V, Bammens T, Ruli D, Zaunschirm S, de Oliveira RM, Outeiro TF, Madeo F, Buee L, Galas MC, Winderickx J (2010). Synphilin-1 enhances alpha-synuclein aggregation in yeast and contributes to cellular stress and cell death in a Sir2-dependent manner. **PLoS One** 5(10): e13700. doi: 10.1371/journal.pone.0013700
154. Hill SM, Hao X, Gronvall J, Spikings-Nordby S, Widlund PO, Amen T, Jorhov A, Josefson R, Kaganovich D, Liu B, Nystrom T (2016). Asymmetric Inheritance of Aggregated Proteins and Age Reset in Yeast Are Regulated by Vac17-Dependent Vacuolar Functions. **Cell Rep** 16(3): 826-838. doi: 10.1016/j.celrep.2016.06.016
155. Higuchi-Sanabria R, Pernice WM, Vevea JD, Alessi Wolken DM, Boldogh IR, Pon LA (2014). Role of asymmetric cell division in lifespan control in *Saccharomyces cerevisiae*. **FEMS Yeast Res** 14(8): 1133-1146. doi: 10.1111/1567-1364.12216
156. Peric M, Bou Dib P, Dennerlein S, Musa M, Rudan M, Lovric A, Nikolic A, Saric A, Sobocanec S, Macak Z, Raimundo N, Krisko A (2016). Crosstalk between cellular compartments protects against proteotoxicity and extends lifespan. **Sci Rep** 6:28751. doi: 10.1038/srep28751
157. Peric M, Lovric A, Saric A, Musa M, Bou Dib P, Rudan M, Nikolic A, Sobocanec S, Mikecin AM, Dennerlein S, Milosevic I, Vlahovick K, Raimundo N, Krisko A (2017). TORC1-mediated sensing of chaperone activity alters glucose metabolism and extends lifespan. **Aging Cell** 16(5): 994-1005. doi: 10.1111/acer.12623
158. Meusser B, Hirsch C, Jarosch E, Sommer T (2005). ERAD: the long road to destruction. **Nat Cell Biol** 7(8): 766-772. doi: 10.1038/ncb0805-766
159. Taylor EB, Rutter J (2011). Mitochondrial quality control by the ubiquitin-proteasome system. **Biochem Soc Trans** 39(5): 1509-1513. doi: 10.1042/BST0391509
160. Lang A, John Peter AT, Kornmann B (2015). ER-mitochondria contact sites in yeast: beyond the myths of ERMES. **Curr Opin Cell Biol** 35:7-12. doi: 10.1016/jceb.2015.03.002
161. Elbaz-Alon Y, Rosenfeld-Gur E, Shinder V, Futerman AH, Geiger T, Schuldiner M (2014). A dynamic interface between vacuoles and mitochondria in yeast. **Dev Cell** 30(1): 95-102. doi: 10.1016/j.devcel.2014.06.007
162. Honscher C, Mari M, Auffarth K, Bohnert M, Griffith J, Geerts W, van der Laan M, Cabrera M, Reggiori F, Ungermann C (2014). Cellular metabolism regulates contact sites between vacuoles and mitochondria. **Dev Cell** 30(1): 86-94. doi: 10.1016/j.devcel.2014.06.006
163. Belgareh-Touze N, Cavellini L, Cohen MM (2017). Ubiquitination of ERMES components by the E3 ligase Rsp5 is involved in mitophagy. **Autophagy** 13(1): 114-132. doi: 10.1080/15548627.2016.1252889
164. Bockler S, Westermann B (2014). Mitochondrial ER contacts are crucial for mitophagy in yeast. **Dev Cell** 28(4): 450-458. doi: 10.1016/j.devcel.2014.01.012
165. Bockler S, Westermann B (2014). ER-mitochondria contacts as sites of mitophagosome formation. **Autophagy** 10(7): 1346-1347. doi: 10.4161/auto.28981
166. Lakhani R, Vogel KR, Till A, Liu J, Burnett SF, Gibson KM, Subramani S (2014). Defects in GABA metabolism affect selective autophagy pathways and are alleviated by mTOR inhibition. **EMBO Mol Med** 6(4): 551-566. doi: 10.1002/emmm.201303356
167. Nazarko VY, Thevelein JM, Sibirny AA (2008). G-protein-coupled receptor Gpr1 and G-protein Gpa2 of cAMP-dependent signaling pathway are involved in glucose-induced pexophagy in the yeast *Saccharomyces cerevisiae*. **Cell Biol Int** 32(5): 502-504. doi: 10.1016/j.cellbi.2007.11.001
168. Natter K, Leitner P, Faschinger A, Wolinski H, McCraith S, Fields S, Kohlwein SD (2005). The spatial organization of lipid synthesis in the yeast *Saccharomyces cerevisiae* derived from large scale green fluorescent protein tagging and high resolution microscopy. **Mol Cell Proteomics** 4(5): 662-672. doi: 10.1074/mcp.M400123-MCP200
169. Bahmanyar S (2015). Spatial regulation of phospholipid synthesis within the nuclear envelope domain of the endoplasmic reticulum. **Nucleus** 6(2): 102-106. doi: 10.1080/19491034.2015.1010942
170. Fernandez-Murray JP, McMaster CR (2016). Lipid synthesis and membrane contact sites: a crossroads for cellular physiology. **J Lipid Res** 57(10): 1789-1805. doi: 10.1194/jlr.R070920
171. Malia PC, Ungermann C (2016). Vacuole membrane contact sites and domains: emerging hubs to coordinate organelle function with cellular metabolism. **Biochem Soc Trans** 44(2): 528-533. doi: 10.1042/BST20150277
172. Murley A, Sarsam RD, Toulmay A, Yamada J, Prinz WA, Nunnari J (2015). Ltc1 is an ER-localized sterol transporter and a component of ER-mitochondria and ER-vacuole contacts. **J Cell Biol** 209(4): 539-548. doi: 10.1083/jcb.201502033
173. Shai N, Schuldiner M, Zalckvar E (2016). No peroxisome is an island - Peroxisome contact sites. **Biochim Biophys Acta** 1863(5): 1061-1069. doi: 10.1016/j.bbamcr.2015.09.016
174. Tavassoli S, Chao JT, Young BP, Cox RC, Prinz WA, de Kroon AI, Loewen CJ (2013). Plasma membrane--endoplasmic reticulum contact sites regulate phosphatidylcholine synthesis. **EMBO Rep** 14(5): 434-440. doi: 10.1038/embor.2013.36



175. De Matteis MA, Rega LR (2015). Endoplasmic reticulum-Golgi complex membrane contact sites. *Curr Opin Cell Biol* 35:43-50. doi: 10.1016/j.ceb.2015.04.001
176. Henne WM, Liou J, Emr SD (2015). Molecular mechanisms of inter-organelle ER-PM contact sites. *Curr Opin Cell Biol* 35:123-130. doi: 10.1016/j.ceb.2015.05.001
177. Wolf W, Kilic A, Schrul B, Lorenz H, Schwappach B, Seedorf M (2012). Yeast Ist2 recruits the endoplasmic reticulum to the plasma membrane and creates a ribosome-free membrane microcompartment. *PLoS One* 7(7): e39703. doi: 10.1371/journal.pone.0039703
178. Yucel EB, Eraslan S, Ulgen KO (2014). The impact of medium acidity on the chronological life span of *Saccharomyces cerevisiae* - lipids, signaling cascades, mitochondrial and vacuolar functions. *FEBS J* 281(4): 1281-1303. doi: 10.1111/febs.12705
179. Mitrofanova D, Dakik P, McAuley M, Medkour Y, Mohammad K, Titorenko VI (2018). Lipid metabolism and transport define longevity of the yeast *Saccharomyces cerevisiae*. *Front Biosci (Landmark Ed)* 23:1166-1194. PMID: 28930594
180. Richard VR, Beach A, Piano A, Leonov A, Feldman R, Burstein MT, Kyryakov P, Gomez-Perez A, Arlia-Ciommo A, Baptista S, Campbell C, Goncharov D, Pannu S, Patrinos D, Sadri B, Svistkova V, Victor A, Titorenko VI (2014). Mechanism of liponecrosis, a distinct mode of programmed cell death. *Cell Cycle* 13(23): 3707-3726. doi: 10.4161/15384101.2014.965003
181. Eisenberg T, Buttner S (2014). Lipids and cell death in yeast. *FEMS Yeast Res* 14(1): 179-197. doi: 10.1111/1567-1364.12105
182. Halbleib K, Pesek K, Covino R, Hofbauer HF, Wunnicke D, Hanelt I, Hummer G, Ernst R (2017). Activation of the Unfolded Protein Response by Lipid Bilayer Stress. *Mol Cell* 67(4): 673-684 e678. doi: 10.1016/j.molcel.2017.06.012
183. Promlek T, Ishiwata-Kimata Y, Shido M, Sakuramoto M, Kohno K, Kimata Y (2011). Membrane aberrancy and unfolded proteins activate the endoplasmic reticulum stress sensor Ire1 in different ways. *Mol Biol Cell* 22(18): 3520-3532. doi: 10.1091/mbc.E11-04-0295
184. Obara K, Kihara A (2017). The Rim101 pathway contributes to ER stress adaptation through sensing the state of plasma membrane. *Biochem J* 474(1): 51-63. doi: 10.1042/BCJ20160580
185. Pineau L, Colas J, Dupont S, Beney L, Fleurat-Lessard P, Berjeaud JM, Berges T, Ferreira T (2009). Lipid-induced ER stress: synergistic effects of sterols and saturated fatty acids. *Traffic* 10(6): 673-690. doi: 10.1111/j.1600-0854.2009.00903.x
186. Pineau L, Ferreira T (2010). Lipid-induced ER stress in yeast and beta cells: parallel trails to a common fate. *FEMS Yeast Res* 10(8): 1035-1045. doi: 10.1111/j.1567-1364.2010.00674.x
187. Zhang M, Yu Q, Liang C, Zhang B, Li M (2016). Lipid homeostasis is involved in plasma membrane and endoplasmic reticulum stress in *Pichia pastoris*. *Biochem Biophys Res Commun* 478(2): 777-783. doi: 10.1016/j.bbrc.2016.08.024
188. Navarro-Tapia E, Nana RK, Querol A, Perez-Torrado R (2016). Ethanol Cellular Defense Induce Unfolded Protein Response in Yeast. *Front Microbiol* 7:189. doi: 10.3389/fmicb.2016.00189
189. Austriaco N (2012). Endoplasmic reticulum involvement in yeast cell death. *Front Oncol* 2:87. doi: 10.3389/fonc.2012.00087
190. Braun RJ, Zischka H (2008). Mechanisms of Cdc48/VCP-mediated cell death: from yeast apoptosis to human disease. *Biochim Biophys Acta* 1783(7): 1418-1435. doi: 10.1016/j.bbamcr.2008.01.015
191. Moir RD, Gross DA, Silver DL, Willis IM (2012). SCS3 and YFT2 link transcription of phospholipid biosynthetic genes to ER stress and the UPR. *PLoS Genet* 8(8): e1002890. doi: 10.1371/journal.pgen.1002890
192. Epstein S, Kirkpatrick CL, Castillon GA, Muniz M, Riezman I, David FP, Wollheim CB, Riezman H (2012). Activation of the unfolded protein response pathway causes ceramide accumulation in yeast and INS-1E insulinoma cells. *J Lipid Res* 53(3): 412-420. doi: 10.1194/jlr.M022186
193. Jadhav S, Russo S, Cottier S, Schneiter R, Cowart A, Greenberg ML (2016). Valproate Induces the Unfolded Protein Response by Increasing Ceramide Levels. *J Biol Chem* 291(42): 22253-22261. doi: 10.1074/jbc.M116.752634
194. Thibault G, Shui G, Kim W, McAlister GC, Ismail N, Gygi SP, Wenk MR, Ng DT (2012). The membrane stress response buffers lethal effects of lipid disequilibrium by reprogramming the protein homeostasis network. *Mol Cell* 48(1): 16-27. doi: 10.1016/j.molcel.2012.08.016
195. Liu M, Huang C, Polu SR, Schneiter R, Chang A (2012). Regulation of sphingolipid synthesis through Orm1 and Orm2 in yeast. *J Cell Sci* 125(Pt 10): 2428-2435. doi: 10.1242/jcs.100578
196. Fei W, Wang H, Fu X, Bielly C, Yang H (2009). Conditions of endoplasmic reticulum stress stimulate lipid droplet formation in *Saccharomyces cerevisiae*. *Biochem J* 424(1): 61-67. doi: 10.1042/BJ20090785
197. Bischof J, Salzmann M, Streubel MK, Hasek J, Geltinger F, Duschl J, Bresgen N, Briza P, Haskova D, Lejskova R, Sopjani M, Richter K, Rinnerthaler M (2017). Clearing the outer mitochondrial membrane from harmful proteins via lipid droplets. *Cell Death Discov* 3:17016. doi: 10.1038/cddiscovery.2017.16
198. Shpilka T, Elazar Z (2015). Lipid droplets regulate autophagosome biogenesis. *Autophagy* 11(11): 2130-2131. doi: 10.1080/15548627.2015.1093719
199. Shpilka T, Welter E, Borovsky N, Amar N, Mari M, Reggiori F, Elazar Z (2015). Lipid droplets and their component triglycerides and steryl esters regulate autophagosome biogenesis. *EMBO J* 34(16): 2117-2131. doi: 10.15252/embj.201490315
200. Velazquez AP, Tatsuta T, Ghillebert R, Drescher I, Graef M (2016). Lipid droplet-mediated ER homeostasis regulates autophagy and cell survival during starvation. *J Cell Biol* 212(6): 621-631. doi: 10.1083/jcb.201508102
201. Vevea JD, Garcia EJ, Chan RB, Zhou B, Schultz M, Di Paolo G, McCaffery JM, Pon LA (2015). Role for Lipid Droplet Biogenesis and Microlipophagy in Adaptation to Lipid Imbalance in Yeast. *Dev Cell* 35(5): 584-599. doi: 10.1016/j.devcel.2015.11.010
202. Deranieh RM, Shi Y, Tarsio M, Chen Y, McCaffery JM, Kane PM, Greenberg ML (2015). Perturbation of the Vacuolar ATPase: a novel consequence of inositol depletion. *J Biol Chem* 290(46): 27460-27472. doi: 10.1074/jbc.M115.683706
203. Hernandez A, Serrano-Bueno G, Perez-Castineira JR, Serrano A (2015). 8-Dehydrosterols induce membrane traffic and autophagy defects through V-ATPase dysfunction in *Saccharomyces cerevisiae*. *Biochim Biophys Acta* 1853(11 Pt A): 2945-2956. doi: 10.1016/j.bbamcr.2015.09.001
204. Teixeira V, Medeiros TC, Vilaca R, Ferreira J, Moradas-Ferreira P, Costa V (2016). Ceramide signaling targets the PP2A-like protein phosphatase Sit4p to impair vacuolar function, vesicular trafficking and autophagy in *Isc1p* deficient cells. *Biochim Biophys Acta* 1861(1): 21-33. doi: 10.1016/j.bbaliip.2015.10.004
205. Aerts AM, Zabrocki P, Francois IE, Carmona-Gutierrez D, Govaert G, Mao C, Smets B, Madeo F, Winderickx J, Cammue BP, Thevissen K (2008). Ydc1p ceramidase triggers organelle fragmentation, apoptosis and accelerated ageing in yeast. *Cell Mol Life Sci* 65(12): 1933-1942. doi: 10.1007/s00018-008-8129-8
206. Sherr GL, LaMassa N, Li E, Phillips G, Shen CH (2017). Pah1p negatively regulates the expression of V-ATPase genes as well as vacuolar

acidification. **Biochem Biophys Res Commun** 491(3): 693-700. doi: 10.1016/j.bbrc.2017.07.127

207. Fakas S, Qiu Y, Dixon JL, Han GS, Ruggles KV, Garbarino J, Sturley SL, Carman GM (2011). Phosphatidate phosphatase activity plays key role in protection against fatty acid-induced toxicity in yeast. **J Biol Chem** 286(33): 29074-29085. doi: 10.1074/jbc.M111.258798

208. Kliewe F, Kumme J, Grigat M, Hintze S, Schuller HJ (2017). Opi1 mediates repression of phospholipid biosynthesis by phosphate limitation in the yeast *Saccharomyces cerevisiae*. **Yeast** 34(2): 67-81. doi: 10.1002/yea.3215