

MEETING ABSTRACT

Open Access

Effectiveness of short-term heat acclimation on intermittent exercise in thermoneutral and hot environments

Fiona Nation, Matt Birkett, Damien Gleadall-Siddall, Rachel Burke, Christopher Towlson, James Bray, Grant Abt, Andrew Garrett*

From 15th International Conference on Environmental Ergonomics (ICEE XV)
Portsmouth, UK. 28 June - 3 July 2015

Introduction

It is well-established that repetition of heat stress exposure has been shown to facilitate adaptations to the heat but these protocols have tended to be of a fixed work intensity, continuous exercise, long-term in duration (>7 days) and use hydration. Secondly, there is limited information on the potential use of heat acclimation as a training method for human performance in thermoneutral conditions [1]. Therefore, the aims of this study were to investigate the effectiveness of short-term heat acclimation (STHA) for 5 days, using the controlled-hyperthermia technique with dehydration, on intermittent exercise in thermoneutral and hot environments.

Methods

Ten, healthy, active, moderately-trained males (Mean (SD); Age 25.6 (8.9) yrs; Height 180.7 (5.6) cm; Body Mass 83.2 (10.8) kg; $\dot{V}O_{2\max}$ 45.3 (6.5) mL.kg⁻¹.min⁻¹ and resting cardiac output 6.3 (1.8) L.min⁻¹), participated in a STHA programme. This consisted of 90 minutes dehydration heat acclimation (no fluid intake) for 5 consecutive days (39.5 °C; 60% rh), using the controlled-hyperthermia technique (~rectal temperature [T_{re}] 38.5 °C) [2]. The pre and post testing Exercise Stress Test (EST) consisted of 45 minutes of intermittent exercise (nine phases of 5 minutes). Including resting, walking, moderate and high intensity running) on a motorised, h/p Cosmos treadmill; and nine 6 second (s) maximal sprints on a Watt Bike, as a repeated, maximal sprint performance test. The EST was followed by a running, incremental test to exhaustion. This EST intermittent protocol was adapted from exercise

intensities of professional football players [3]. The EST was repeated in controlled conditions (C; 11 °C 45% rh); pre and post STHA in thermoneutral (TN; 11 °C 45% rh) and hot environments (H; 35 °C 45% rh). Data analysis was by paired t-test and reported as the mean differences with 95% confidence intervals (95%CI).

Results

Post STHA, in the H trial at 45-min there was a decrease in T_{re} by -0.2 °C (95%CI: -0.40 to 0.00 °C; $p = 0.03$), cardiac frequency (-3: -5 to -1 b.min⁻¹; $p = 0.01$) and RPE (-2: -3 to -1 units; $p = 0.01$). Mean average power (MavP) increased in sprints 7 (111: 25 to 197 W; $p = 0.01$) and 9 (240: 9 and 489 W; $p = 0.04$). The increase in Sprint 8 (87: -8 to 182 W; $p = 0.06$) and time to exhaustion (208: -24 to 439 s; $p = 0.06$) were close to significance. In the TN trial, increased MavP for sprint 9 (67: 2 to 131 W; $p = 0.04$) and time to exhaustion (43: 1 to 85 s; $p = 0.04$) were reported. There was limited change in the C conditions across all nine sprints ($p > 0.05$) and for time to exhaustion (-31: -72 to 11 s; $p = 0.12$).

Discussion

Short-term heat acclimation (5 days) with dehydration, using the controlled-hyperthermia technique, is effective for physiological adaptations during intermittent exercise in a hot environment. Furthermore, it has resulted in an increase in intermittent and endurance exercise performance in hot and thermoneutral conditions.

Conclusion

Short-term heat acclimation is effective for intermittent exercise in the heat. It may be a useful training method for human performance in thermoneutral conditions.

* Correspondence: a.garrett@hull.ac.uk

Department of Sport, Health and Exercise Science, Exercise Health and Human Performance Research Group, University of Hull, UK

Published: 14 September 2015

References

1. Corbett J, Neal RA, Lunt HC, Tipton MJ: **Heat acclimation and exercise performance under cooler conditions: a new hot topic.** *Sports Medicine* 2014, **44**:1323-1331.
2. Garrett AT, Goosens NG, Rehrer NJ, Patterson MJ, Harrison J, Sammut I, Cotter JD: **Short-term heat acclimation is effective and is not impaired by dehydration during acclimation.** *American Journal Human Biology* 2014, **26**:311-320.
3. Drust B, Atkinson G, Reilly T: **Future perspectives in the evaluation of the physiological demands of soccer.** *Sports Medicine* 2007, **37**(9):783-803.

doi:10.1186/2046-7648-4-S1-A120

Cite this article as: Nation *et al.*: Effectiveness of short-term heat acclimation on intermittent exercise in thermoneutral and hot environments. *Extreme Physiology & Medicine* 2015 **4**(Suppl 1):A120.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

