

The mean (SD) time taken in tip confirmation by using bedside ultrasound was 5.1(1.2) minutes, X-ray it was 28 (8.1) minutes ( $P<0.001$ ). The catheter tip was in an optimal position in the first attempt in 30 (91%) neonates after the ultrasound and confirmed by X-ray. In these three cases (9%) the tip of the PICC catheter was in the right atrium after first attempt confirmation. There was no inter-observer variation in the interpretation of the result.

Previous studies [4-8] have also shown that the mean time taken in confirmation of tip by using ultrasound is significantly less than standard care. The accuracy of ultrasound was also comparable with radiography. By using ultrasound, we can reduce radiation exposure, and ensure lesser handling of babies.

Bedside ultrasound is an accurate and time-efficient modality to guide the insertion and confirmation of the tip of the PICC line. However, training of neonatologists in ultrasound may be required before routine use of this modality.

*Contributors:* SS: collected data and wrote the article; HAV: planned study and supervised data collection; RS: statistical analysis and literature review; NKN: drafted and reviewed the final manuscript. All authors approved the final version of the manuscript.

*Funding:* None; *Competing Interest:* None stated

**SURJEET SINGH, HA VENKATESH,\* RAVI SWAMY AND N KARTHIK NAGESH**

*Department of Neonatology,  
Manipal Hospitals,  
Bengaluru, Karnataka, India.  
\*venkatveena46@gmail.com*

## REFERENCES

1. Ramasetu J. Complications of vascular catheters in the neonatal intensive care unit. *Clin Perinatol.* 2008;35: 199-222.
2. Diamond LK, Allen FH, Thomas WO. Erythroblastosis fetalis. VII. Treatment with exchange transfusion. *N Engl J Med.* 1951;244:39-49.
3. Sharma D, Farahbakhsh N, Tabatabaie SA. Role of ultrasound for central catheter tip localization in neonates: A review of the current evidence. *J Matern Fetal Neonatal Med.* 2019;32:2429-37.
4. Katheria AC, Fleming SE, Kim JH. A randomized controlled trial of ultrasound-guided peripherally inserted central catheters compared with standard radiograph in neonates. *J Perinatol.* 2013;33:791-4.
5. Tauzin L, Sigur N, Joubert C, Parra J, Hassid S, Moulies ME. Echocardiography allows more accurate placement of peripherally inserted central catheters in low birthweight infants. *Acta Paediatr.* 1992. 2013;102:703-6.
6. Saul D, Ajayi S, Schutzman DL, Horrow MM. Sonography for complete evaluation of neonatal intensive care unit central support devices: A pilot study. *J Ultrasound Med.* 2016;35:1465-73.
7. Telang N, Sharma D, Pratap OT, Kandraj H, Murki S. Use of real-time ultrasound for locating tip position in neonates undergoing peripherally inserted central catheter insertion: A pilot study. *Indian J Med Res.* 2017;145:373-6.
8. Jain A, McNamara PJ, Ng E, El-Khuffash A. The use of targeted neonatal echocardiography to confirm the placement of peripherally inserted central catheters in neonates. *Am J Perinatol.* 2012;29:101-6.

## Infantile Cardiac Beriberi in Rural North East India

Twenty eight exclusively breastfed infants presented between 1 July, 2017 and 30 June, 2018 with acute heart failure syndrome, with 23 (92%) showing dramatic clinical resolution of shock within 24 hours of receiving intravenous thiamine (100 mg) bolus. Our findings raise awareness for addressing this neglected nutritional disease in North East India.

**Keywords:** *Heart Failure, Infantile Beriberi, Thiamine.*

The cardiac form of infantile beriberi is a fulminant disease, affecting exclusively breastfed infants of mothers with thiamine deficiency. The classical description is a well thriving infant presenting in acute cardiac failure succumbing to the illness within four hours, if left untreated [1]. Laos has documented widespread thiamine deficiency in communities, causing a peak in infant mortality in the third month of life [2]. The overall infant mortality rates in the Karen refugee camp in Thailand

reduced from 183 to 78 per 1000 live births after early diagnosis and management of infantile beriberi [3]. We report on infantile beriberi as a preventable cause of death among infants from rural North East India.

The study was conducted in a charitable hospital in Karimganj district of Assam, which has an infant mortality rate of 69 per 1000 live births in 2012-13 (National average, 42/1000 live births). A retrospective review of medical records was conducted for all infants who were discharged between 1 July, 2017 and 30 June, 2018 with a diagnosis of infantile beriberi. Infantile beri beri was diagnosed when an otherwise well, exclusively breastfed infant presented with a thiamine responsive acute cardiac failure syndrome [1].

A total of 28 infants with a mean (SD) age 69 (29.1) days and weight 3.84 (1.26) kg from rural Assam and Tripura were diagnosed with infantile beriberi during the study duration. The commonest complaints were short history of vomiting, breathlessness and poor feeding. All infants presented in a critically ill state with prolonged capillary refill time (93%), tachycardia (93%), seizures (36%) and severe respiratory distress (92%). The capillary blood gas of all infants showed severe high anion-gap metabolic acidosis (**Table 1**).

**Table I Laboratory Abnormalities in Infants With Cardiac Beriberi (N=28)**

Variable	Value, median (IQR)
Hemoglobin (g/dL)	8.2 (7.7-9.9)
Leukocyte count (per mm <sup>3</sup> )	20510 (15410-27030)
Platelet count (per mm <sup>3</sup> )	297000 (15410-27030)
Blood sugar (mg/dL)	115 (70-206)
pH	6.9 (6.7-7.1)
HCO <sub>3</sub> (mmol/L)	5.2 (4.8-7.8)
Base excess	-26.6 (-23.05 to -29.0)
Anion gap	42.25 (30.9-43.0)

Of these, 25 (89%) received 100 mg of intravenous thiamine bolus within one hour of admission, followed by 100 mg intravenously for a minimum of seven days till discharge. Infants also received other treatment modalities as per the pediatric protocol for treatment of shock in the hospital. 20 infants (71%) required inotropes. Twenty three infants (92%) showed dramatic recovery, with features of shock resolving within 24 hours, and were initiated on breast feeds within two days. All 14 infants (50%) requiring invasive ventilation could be weaned within 60 hours, with 12 infants (86%) being extubated in the first 24 hours. There was a rapid improvement in the capillary blood gas measurements within 4-8 hours of bolus thiamine, with mean (SD) pH improving from 6.9 (0.22) to 7.35 (0.11) and mean (SD) base excess from -24.3 (6.69) to -3.8 (5.69).

The three infants who did not receive parenteral thiamine died within half an hour of admission while of those who received parenteral thiamine, two infants died. One of these due to a ventilator-associated adverse event. All infants who survived were discharged after a mean (SD) in-patient stay of 7 (2.81) days. They were prescribed 10 mg per day of thiamine supplements; 17 babies (74%) were subsequently reviewed and found to be well.

The presentation of infants in this case series corroborates with the classical description of infantile cardiac beriberi in literature, including a cohort from Kashmir [5]. Moreover, we have previously documented peripheral neuropathy with or without cardiomyopathy among peripartum women [6]. Since the clinical manifestation of beriberi in infants reflects poor maternal stores, communities in North East India are at risk populations, explaining the prevalence of this nutritional disease.

All infants presented in an acute critically ill state and none had documented fever or history of fluid loss to account for shock. The dramatic therapeutic response to parenteral thiamine administration in 92% of infants along with the rapid clinical deterioration of the three infants who did not receive it, favors the diagnosis of cardiac beriberi. X-ray done on five babies revealed cardiomegaly at admission, which disappeared after five days of thiamine. The diagnosis could have been strengthened using echocardiography, and determining RBC transketolase activity, which were not available at our institute. Subsequent to the study, echocardiography in other such infants has demonstrated features of pulmonary hypertension, which responded after thiamine administration.

Although infantile beriberi was believed to have been eliminated from India [7], there is emerging evidence to suggest that beriberi still continues to be a cause of preventable infant mortality among Indian children [4,8]. In places with high infant mortality and peak age of deaths at three months of age, beriberi needs to be considered in the differential diagnoses of infants presenting with unexplained shock.

Being a fatal but preventable and easily treatable disease, these observations on infantile beriberi require a strong public health response. Education campaigns and thiamine supplementation in pregnant and post partum women are possible strategies. Prospective studies using data from population surveys and nutritional assessments to identify the factors contributing to the epidemic in these high-risk populations are being planned with the National Institute of Nutrition, Government of India.

*Ethical Clearance:* Local research committee of MCL General Hospital; 27 June, 2017.

*Contributions:* ST: RMK conceptualized and designed the study, developed protocol; ST: collected and analyzed data, reviewed literature and prepared initial manuscript; RMK & VAI: Manuscript review; VAI: performed echocardiogram on some babies included in study. All the authors approved the final version of the manuscript.

*Funding:* None; *Competing interests:* None stated.

SHAJIN THANKARAJ,<sup>1\*</sup> ROSHINE M KOSHY<sup>2</sup> AND VIJAYANAND ISMAVEL<sup>3</sup>

Departments of <sup>1</sup>Pediatrics, <sup>2</sup>Medicine and <sup>3</sup>Pediatric Surgery, Makunda Christian Leprosy and General Hospital, Assam, India. \*mail4shajin@gmail.com

## REFERENCES

1. Department of Nutrition for Health and Development, World Health Organization. Thiamine deficiency and its prevention and control in major emergencies. WHO, 1999.
2. Barennes H, Sengkhamyong K, Rene JP, Phimmasane M. Beriberi (thiamine deficiency) and high infant mortality in northern Laos. PLoS Negl Trop Dis. 2015;9:3.
3. Luxemburger C, White NJ, ter Kuile F, Singh HM, Allier-Frachon I, Ohn M, *et al.* Beri-beri: the major cause of infant mortality in Karen refugees. Trans Royal Soc Trop Med Hyg. 2003;97:251-5.
4. Qureshi UA, Sami A, Atlatf U. Thiamine responsive acute life threatening metabolic acidosis in exclusively breast fed infants. Nutrition. 2016;32:213-6.
5. Koshy RM, Ismavel VA, Sharma H, Jacob PM. Suspected thiamine deficiency presenting as peripheral neuropathy among peripartum women in a hospital in rural Assam: A neglected public health problem. CHRISMED J Health and Research. 2018;5:178-81.
6. Nazir M, Lone R, Charoo BA. Infantile thiamine deficiency: New insights into an old disease. Indian Pediatr. 2019;56:673-81.
7. Bhat JI, Rather HA, Ahangar AA, Qureshi UA, Dar P, Ahmed QI, *et al.* Shoshin beriberi-thiamine responsive pulmonary hypertension in exclusively breastfed infants: A study from northern India. Indian Heart J. 2016;69:4.