

ORIGINAL ARTICLE

NR4A1 counteracts JNK activation incurred by ER stress or ROS in pancreatic β -cells for protection

Ze-qing Pu¹ | Dong Liu¹ | Hanse Pablick Patherny Lobo Mouguegue¹ | Cheng-wen Jin¹ | Esha Sadiq¹ | Dan-dan Qin¹  | Tian-fu Yu¹ | Chen Zong¹ | Ji-cui Chen² | Ru-xing Zhao³ | Jing-Yu Lin⁴ | Jie Cheng⁴ | Xiao Yu^{4,7}  | Xia Li¹ | Yu-chao Zhang⁵ | Yuan-tao Liu⁵ | Qing-bo Guan⁶ | Xiang-dong Wang^{1,7}

¹Department of Cell Biology, Shandong University School of Medicine, Jinan, China

²Blood Transfusion Department, Qilu Hospital of Shandong University, Jinan, China

³Department of Endocrinology, Qilu Hospital of Shandong University, Jinan, China

⁴Department of Physiology, Shandong University School of Medicine, Jinan, China

⁵Department of Endocrinology, Qingdao Municipal Hospital, Qingdao, China

⁶Department of Endocrinology, Shandong Provincial Hospital, Affiliated to Shandong University, Jinan, China

⁷Key Laboratory of Protein Sciences for Chronic Degenerative Diseases in Universities of Shandong (Shandong University), Jinan, China

Correspondence

Xiang-dong Wang, Shandong University School of Medicine, Jinan, China.
Email: xdongw@sdu.edu.cn

Funding information

This work was supported by the National Natural Science Foundation of China (NSFC funding numbers are 81670704, 81370881, 81500631, 81471078, and 81641030; Shandong Provincial Major Innovation Projects: 2017CXGC1214; Provincial Natural Science Foundation of Shandong: BS2015YY011).

Abstract

Sustained hyperglycaemia and hyperlipidaemia incur endoplasmic reticulum stress (ER stress) and reactive oxygen species (ROS) overproduction in pancreatic β -cells. ER stress or ROS causes c-Jun N-terminal kinase (JNK) activation, and the activated JNK triggers apoptosis in different cells. Nuclear receptor subfamily 4 group A member 1 (NR4A1) is an inducible multi-stress response factor. The aim of this study was to explore the role of NR4A1 in counteracting JNK activation induced by ER stress or ROS and the related mechanism. qPCR, Western blotting, dual-luciferase reporter and ChIP assays were applied to detect gene expression or regulation by NR4A1. Immunofluorescence was used to detect a specific protein expression in β -cells. Our data showed that NR4A1 reduced the phosphorylated JNK (p-JNK) in MIN6 cells encountering ER stress or ROS and reduced MKK4 protein in a proteasome-dependent manner. We found that NR4A1 increased the expression of cbl-b (an E3 ligase); knocking down cbl-b expression increased MKK4 and p-JNK levels under ER stress or ROS conditions. We elucidated that NR4A1 enhanced the transactivation of cbl-b promoter by physical association. We further confirmed that cbl-b expression in β -cells was reduced in NR4A1-knockout mice compared with WT mice. NR4A1 down-regulates JNK activation by ER stress or ROS in β -cells via enhancing cbl-b expression.

KEYWORDS

cbl-b, ER stress, NR4A1 (Nur77), pancreatic β -cells, p-JNK, ROS

1 | INTRODUCTION

Although the aetiology of type 2 diabetes is complex, it is well accepted that insulin resistance leads to sustained hyperglycaemia or

hyperlipidaemia, which consequently results in cellular ER stress and/or ROS overproduction in pancreatic β -cells, whereas ER stress or ROS causes cell damage or even apoptosis.¹⁻⁶ Pancreatic β -cell loss plays an important role in the pathogenesis of type 2 diabetes. To reveal the

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2020 The Authors. *Journal of Cellular and Molecular Medicine* published by Foundation for Cellular and Molecular Medicine and John Wiley & Sons Ltd

mechanisms by which pancreatic β -cells counteracts the apoptotic effect from ER stress or ROS may provide clues for β -cell protection.

NR4A1 is an inducible multi-stress response factor.⁷⁻⁹ It belongs to the NR4A family, and its molecular structure includes a transcriptional activation domain, a DNA-binding domain and a putative ligand-binding domain.¹⁰ So far, no nature ligand has been found in this family. Therefore, NR4A1 belongs to orphan nuclear receptors.¹¹ NR4A1 acts as a transcriptional factor to enhance the expression of some genes without ligand binding.^{12,13} NR4A1 participates in various physiological activities, including cell proliferation, differentiation and apoptosis.¹⁴⁻¹⁸ ER stress or ROS-inducing NR4A1 expression in pancreatic β -cells has been documented.^{4,5} Controversy still exists regarding the role of NR4A1 in apoptosis. Some studies showed that it had anti-apoptotic effects,^{15,19-24} whereas others showed it was pro-apoptotic.^{16,18,25-27} A possible explanation for the above controversial conclusions may be accounted by the cellular location of NR4A1. When NR4A1 remains in the nucleus as a transcriptional factor, it enhances the expression of some anti-apoptotic factors, such as WT1 and survivin^{4,5,24}; if NR4A1 is translocated into the cytoplasm, it associates with bcl2 and results in cell apoptosis.²⁷

JNKs, also known as stress-activated protein kinases, form an important subgroup of the mitogen-activated protein kinase (MAPK) superfamily and have three isoforms (JNK1, JNK2 and JNK3). Upon activation by upstream kinases, the activated JNK has diverse cellular functions, including cell proliferation and differentiation under physiological conditions. The role of activated JNK on pro-apoptosis has been described as in cell type-dependent manner.²⁸ Cellular stresses, such as ER stress and oxidative stress (ROS), incur high level or sustained JNK activation, which further triggers apoptotic pathways in different cells.²⁹⁻³⁹ Our preliminary data showed that either TG (thapsigargin, an ER stress inducer) or H₂O₂ (hydrogen peroxide, a kind of ROS) increased the accumulation of phosphorylated JNK in pancreatic β -cells, whereas overexpressing NR4A1 was able to reduce JNK phosphorylation. NR4A1 is a transcription factor. How can a transcript factor regulate JNK phosphorylation? To answer this question, we used pancreatic mouse insulinoma 6 cells (MIN6 cells) and NR4A1 KO mice to explore the underlying mechanism.

2 | MATERIALS AND METHODS

2.1 | Cell culture and reagents

MIN6 cells were obtained from ATCC and were cultured under the same conditions as described in the reference.⁴ NR4A1 overexpression cell lines (designated as OV cells) and control cell lines (designated as NC cells) were generated and maintained as described previously.⁴ cbl-b knockdown cells (designated as KD-cbl-b cells) or the control cell clone (designated as CON-cbl-b cells) were cultured in the same medium as NC/OV cells plus G418 at 1 mg/mL.

DMEM was purchased from M&C GENE technology. Foetal bovine serum (FBS) was purchased from Gibco. TG was purchased

from Sigma-Aldrich. H₂O₂ was purchased from Sinopharm Chemical Reagent. Puromycin and G418 (sulphate) were purchased from InvivoGen. MG132 was purchased from MedChemExpress.

2.2 | Mouse islet purification

NR4A1 knockout (KO) mice generated from C57BL/6J mice and wild-type (WT) mice were purchased from Cyagen Biosciences. The genotypes of the new-born mice were verified with PCR according to the manual from Cyagen Biosciences. Mouse islets were isolated with collagenase P (purchased from Roche Applied Science) as described previously.⁴⁰ After separation, the islets were deposited by Hank's Balanced Salt Solution. Finally, the islets were selected under stereoscopic microscope. Animal experiments were carried out according to the Principles of Laboratory Animal Care by NIH. All experiment procedures were approved by the Animal Care and Use Committee of Shandong University School of Basic Medical Sciences. The Animal Care and Use Committee of Shandong University School of Basic Medical Sciences approved that all the experiment procedures fully complied with the Helsinki Declaration.

2.3 | Lentiviral infection and stable cell line selection

Lentivirus encoding full-length NR4A1 and control lentivirus were constructed and generated by GenePharma. MIN6 cells were infected with recombinant lentivirus encoding NR4A1 or control lentiviral vector, and stable cells were selected under puromycin selection.

Lentivirus encoding shRNA targeting to cbl-b and control lentivirus were constructed in Obio Technology, and the targeting sequence was 5'-AACACAGACGCCATGATTTGC-3'.⁴¹ OV cells were infected with cbl-b shRNA lentivirus or control scramble shRNA lentivirus, and the stable cells were selected under G418 drug pressure.

2.4 | Plasmid construction

Mice genomic DNA was extracted as previously described.⁴² Mice cDNA library was obtained from WT C57BL/6J mouse liver with ReverTra Ace qPCR RT Kit (Toyobo). We designed a pair of primers with HindIII and XbaI restriction sites at 5' and 3' to amplify the cDNA of MKK4 and cloned the PCR product into pFLAG-CMV™-2 vector (Sigma). The sequence information of the primer is listed below:

Primer F: 5'-CCCAAGCTTATGGTCCACAAACCAAGTGGG-3'

Primer R: 5'-GGGTCTAGATCAGTCGACATACATGGGCGAGC-3'.

We designed four pairs of primers with KpnI and XhoI restriction sites at 5' and 3', respectively, to amplify four different lengths of the cbl-b promoter (1987, 1440, 981 and 485 bp) from mice genomic DNA. And then, the PCR products were cloned into pGL3-basic luciferase reporter vector (Promega).

The pairs of primers used for the four different lengths of cbl-b promoter are listed below:

Primer F (-2008 bp): 5'-CCC GGTACC GTCAGCCTACATAAAC ACTATC-3'

Primer R (-21 bp): 5'-CCC CTCGAG TCGAACCAATCCTGAGA-3'.

The primers used are 1440, 981 and 458 bp. They all shared the same reverse primer:

Primer F (-1454 bp): 5'-CCC GGTACC CCAGACTTGTACTTTAC TGC-3'

Primer F (-995 bp): 5'-CCCGGTACC TCACTGTAAGCGTTACCA-3'

Primer F (-499 bp): 5'-CCC GGTACC TATTTGGTGCTCTCCT-3'

Primer R (-14 bp): 5'-CCC CTCGAG TTGAGTCTTGAACCAA-3'.

All constructed plasmids were sequenced by Sangon Biotech Co., Ltd., to verify these sequences were completely aligned with the original DNA sequences from NCBI.

2.5 | Quantitative real-time PCR assay

Total RNA was extracted and purified from cultured cells by using RNAiso Plus (Takara) and from mouse islets using a RNeasy Mini Kit (Qiagen). qPCR was performed as previously described.⁴ According to the Cq value of the target gene obtained from the qPCR machine, the relative expression multiple of the target gene was calculated with a relative quantitative method for $2^{-\Delta\Delta Cq}$, and the numerical value was further corrected by using the housekeeping gene 18s rRNA as the internal control. All experiments were repeated three times, and each sample was assayed in triplicate. The primers used for PCR are listed in Table 1.

2.6 | Western blot analysis

Protein samples were obtained, and Western blot analysis was performed according to the methods as previously described.⁴³ The antibodies used in the experiment are as follows: rabbit antibodies, such as anti-p-JNK (1:1000), anti-JNK (1:1000), anti-MKK4 (1:1000) and anti-cbl-b (1:1000), were purchased from ProteinTech Group, Inc; anti-GAPDH antibody (1:30 000) was purchased from Zsfg Bio; anti-NR4A1 antibody (1:1000) were purchased from Affinity Biosciences and Abcam; and mouse anti-beta actin (1:5000) and rabbit anti- α -Tubulin (1:5000) were purchased from Bioworld Technology.

2.7 | Immunoprecipitation (IP)

Equal numbers of OV cells or NC cells were transfected with equal amount of pFLAG-CMV²-MKK4 and pUb-HA (a plasmid for over-expressing Ub with HA-tag from Dr Chengjiang Gao's Lab, Shandong University School of Basic Medical Sciences). After 48 hours

TABLE 1 Primers for real-time quantitative PCR

Genes	Sequence (5'→3')
18S rRNA F	CGCGGTTCTATTTTGTGGT
18S rRNA R	AGTCGGCATCGTTTATGGTC
NR4A1 F	ATGCCTCCCCTACCAATCTTC
NR4A1 R	CACCAGTTCCTGGAACCTTGGGA
MKK4 F	AGGACTTGAAAGACCTTGGAGA
MKK4 R	TATCTGCCCACTTGGTTTGTG
MKK7 F	ATGGAGAGCATCGAGATTGACC
MKK7 R	TTGATTCTGCCTGATAACGCT
Cbl-b F	GGTCGCATTTTGGGGATTATTGA
Cbl-b R	TTTGGCACAGTCTTACCCTTT
TRAF2 F	AGAGAGTAGTTCCGGCCTTTCC
TRAF2 R	GTGCATCCATCATTGGGACAG
Smurf1 F	AGCATCAAGATCCGTCTGACA
Smurf1 R	CCAGAGCCGTCCACAACAAT
JNK F	GTGGAATCAAGCACCTTCACT
JNK R	TCCTCGCCAGTCCAAAATCAA

post-transfection, the cells were harvested for immunoprecipitation as routinely used. The antibody used to pull down protein was anti-flag monoclonal antibody from ProteinTech Group. The beads with protein A and protein G were purchased from Solarbio Science & Technology. The pull-down samples were applied for Western blotting with 12CA5 (an anti-HA monoclonal antibody).

2.8 | Dual-luciferase reporter assays

Plasmids with different lengths of cbl-b promoter were co-transfected into cells with TK (thymidine kinase promoter-Renilla luciferase reporter) plasmid, or pGL3-basic vector was co-transfected into cells with TK plasmid as control. The luciferase activity was measured by a dual-luciferase reporter assay kit (Vazyme Biotech Co., Ltd) after 48 hours post-transfection.

2.9 | Chromatin immunoprecipitation (ChIP) analysis

MIN6 cells were infected with adenovirus encoding NR4A1-HA or control adenovirus. After 48 hours post-infection, the infected cells were applied for ChIP assay. ChIP analysis was accomplished with a ChIP Assay Kit (Beyotime) as previously described.⁴ The specific anti-HA monoclonal antibody (12CA5) was applied to pull down the DNA fragments associated with HA-tagged NR4A1.

A pair of primers, 5'-GCTGCTACTTTTTCAGTTCCCTTCTCG TTC-3'(F), and 5'-TCCACGGTACACAATGGCCC-3' (R), were designed to amplify the specific target sequence (-251 to -118 bp) of the cbl-b promoter (as indicated in Figure 6).




FIGURE 1 NR4A1 down-regulates JNK phosphorylation induced by TG or H_2O_2 in MIN6 cells. A-C, the relative NR4A1 mRNA and protein levels were determined by qPCR and Western blot in both OV (NR4A1 overexpression) and NC (control) cells. D-G, Western blotting exhibited the protein levels of p-JNK and JNK in response to 100 μ mol/L H_2O_2 at various time-points in both OV and NC cells. H-K, Western blotting exhibited the protein levels of p-JNK and JNK in response to 0.5 μ mol/L TG at various time-points in both OV and NC cells. Densitometric analyses of the blots were presented as histogram, and these data represented the means of three independent experiments; * $P < .05$, ** $P < .01$ and *** $P < .001$ vs ns; error bars indicate SD

2.10 | Immunofluorescence

The pancreatic tissues from WT mice or NR4A1 KO mice were fixed with paraformaldehyde (4%). The primary anti-insulin mice antibody (Santa Cruz Biotechnology, Inc) and the primary anti-cbl-b rabbit antibody (ProteinTech Group, Inc) were applied for immuno-staining. The overlapped colour (yellow) of insulin (red) with cbl-b (green) in islets of pancreatic tissue indicated cbl-b expression in pancreatic β -cells.

2.11 | Statistical analyses


Data were expressed as the mean \pm SD. Statistical analysis was done with t-test or analysis of variance (ANOVA) by using GraphPad Prism 5.0. A value of $P < .05$ (*) was considered a statistically significant difference.

3 | RESULTS

3.1 | NR4A1 down-regulates JNK phosphorylation induced by TG or H_2O_2 in MIN6 cells

Previously, we generated NR4A1 overexpression cells (OV cells) from MIN6 cells and the control cells (NC cells). The mRNA and protein expression levels of NR4A1 were confirmed in OV cells and NC cells (Figure 1A-C). Figure 1 showed that the phosphorylation level of JNK was increased in both OV cells and NC cells after they were treated with TG or H_2O_2 , but overall the JNK phosphorylation level was lower in OV cells compared with that in NC cells (Figure 1D,E,H,I), whereas the total JNK level was close between the two cells (Figure 1F,G,J,K). Overexpressing NR4A1 turned down the JNK phosphorylation in MIN6 cells rather than total JNK level.

FIGURE 2 NR4A1 down-regulates MKK4 protein level in MIN6 cells. A-C, the relative mRNA and protein levels of MKK4 and MKK7 were determined by qPCR and Western blot in both OV and NC cells. D and E, MKK4 and MKK7 were verified to be phosphorylated during the treatment with H₂O₂. F and G, the protein levels of MKK4 and MKK7 in response to 100 μmol/L H₂O₂ at various time-points in both OV and NC cells were determined by Western blot. H and I, the protein levels of MKK4 and MKK7 in response to 0.5 μmol/L TG at various time-points in both OV and NC cells were determined by Western blot. Densitometric analyses of the blots were presented as histogram, these data represented the means of three independent experiments; **P* < .05, ***P* < .01 and ****P* < .001 vs ns; error bars indicate SD



3.2 | NR4A1 down-regulates MKK4 protein level in MIN6 cells

To explore the mechanism by which NR4A1 attenuates JNK phosphorylation induced by ER stress or ROS, we searched online and found that there were two upstream kinase molecules specific for JNK, namely MKK4 and MKK7. The mRNA or protein level of MKK7 did not change between OV and NC cells (Figure 2A-C), but the protein level of MKK4 in OV cells was markedly reduced compared with NC cells (Figure 2A,B), whereas the mRNA level of MKK4 did not change (Figure 2C).

To verify MKK4 and MKK7 are two upstream kinase molecules for JNK activation, the MIN6 cells were treated with H₂O₂ for 30 minutes, the phosphorylated MKK4(p-MKK4) and phosphorylated MKK7(p-MMK7) were detected with the specific antibodies,

and the data showed that H₂O₂ increased the levels of both MKK4 and MKK7 phosphorylation levels. By comparison, the induced fold of p-MKK4 was higher than that of p-MMK7 (Figure 2D-E).

The OV cells and NC cells were treated with TG or H₂O₂ at different time-points, both MKK4 and MKK7 were examined in these samples, and we found that the MKK4 protein was maintained at a lower level in OV cells compared with that in NC cells during these series of treatments (Figure 2F-I).

3.3 | NR4A1 increases MKK4 ubiquitination and degradation in MIN6 cells

We tested whether overexpressing NR4A1 was able to enhance MKK4 ubiquitination level. In Figure 3, equal numbers of OV cells




FIGURE 3 NR4A1 increases MKK4 ubiquitination and degradation in MIN6 cells. **A**, both OV cells and NC cell were transfected with two plasmids to express Flag-MKK4 and HA-ub. The Western blotting results showed the different ubiquitination levels of Flag-MKK4 in OV and NC cells. **B** and **C**, OV cells or NC cells were treated with either 10 μ mol/L MG132 or DMSO for 6 h. The protein level of MKK4 was detected with Western blotting in four different conditions. Densitometric analyses of the blots were shown as histogram; the data represented the means of three independent experiments; * $P < .05$ vs ns; error bars indicate SD




FIGURE 4 NR4A1 enhances the expression of cbl-b in MIN-6 cells. **A**, the relative cbl-b mRNA levels were determined by real-time quantitative PCR in both OV and NC cells. **B** and **C**, the relative cbl-b protein levels were determined by Western blotting in both OV and NC cells. **D** and **E**, the protein levels of cbl-b in response to 100 μ mol/L H₂O₂ at various time-points in both OV and NC cells were assessed by Western blotting. **F** and **G**, the protein levels of cbl-b in response to 0.5 μ mol/L TG at various time-points in both OV and NC cells were assessed by Western blotting. Densitometric analyses of the blots are shown as histogram; the data represented the means of three independent experiments; * $P < .05$, ** $P < .01$ and *** $P < .001$ vs ns; error bars indicate SD

and NC cells were transfected with equal amount of Flag-MKK4 and HA-ubiquitin plasmids. The Flag-MKK4 protein was pulled down with anti-Flag monoclonal antibody. The results in Figure 3A showed that more ubiquitin modified exogenous Flag-MKK4 in OV cells compared with that in NC cells. From the data above, it was reasonable

to conclude that more overexpressed exogenous Flag-MKK4 in OV cells was ubiquitinated compared with that in NC cells.

To further confirm that MKK4 was reduced by ubiquitination-based proteasome degradation, MG132 was applied to figure out the dynamics of MKK4 in OV cells. As expected, MG132 was

able to reverse the effect of NR4A1 down-regulating MKK4 protein level in MIN6 cells (Figure 3B,C).

3.4 | NR4A1 enhances the expression of cbl-b in MIN6 cells

Proteasome-based protein degradation depends on the abundance of the specific E3 ligase. Each E3 ligase has a specific target or substrate to be modified with ubiquitin. It was reported that NR4A1 was able to enhance the expression of three possible E3 ligases, which were TRAF2,¹⁹ Smurf1⁴⁴ and cbl-b (casitas B-lineage lymphoma b).⁴¹ Our data showed that OV cells had higher mRNA and protein levels of cbl-b (Figure 4A-C), whereas the mRNA levels of TRAF2 or Smurf1 did not increase in OV cells compared with NC cells

(Figure 4A). During TG or H₂O₂ treatment for different time-points, OV cells always maintained a higher level of cbl-b protein expression compared with NC cells (Figure 4D-G).

3.5 | Cbl-b expression negatively correlates with MKK4 protein level and p-JNK level upon H₂O₂ or TG treatment in NR4A1-overexpression cells

To further confirm that cbl-b modulates MKK4 protein expression in MIN6 cells, we applied RNA interference scheme to knock down cbl-b expression in OV cells. Figure 5A-C showed that knocking down cbl-b expression resulted in increased MKK4 protein level. Knocking down cbl-b expression resulted in about 20% and 90% increase in mRNA and protein levels of MKK4, respectively; therefore, the increased MKK4




FIGURE 5 Cbl-b expression negatively correlates with MKK4 protein level and p-JNK level upon H₂O₂ or TG treatment in OV cells. A, OV cells were infected with Lentivirus encoding shRNA targeting cbl-b or encoding scramble shRNA. Real-time quantitative PCR was exploited to test the mRNA levels of cbl-b and MKK4 in KD-cbl-b and CON-cbl-b. B and C, Western blotting was applied to detect the protein levels of cbl-b and MKK4 in KD-cbl-b and CON-cbl-b. D and E, the protein levels of p-JNK were exhibited in CON-cbl-b and KD-cbl-b cells in response to 100 μ mol/L H₂O₂ at 0 or 30 min. F and G, the protein levels of p-JNK were exhibited in CON-cbl-b and KD-cbl-b cells in response to 0.5 μ mol/L TG at 0 or 3 h. The data represented the means of three independent experiments; ***P* < .01 and ****P* < .001 vs ns; error bars indicate SD




FIGURE 6 NR4A1 enhances the transactivation of *cbl-b* promoter via physical association. A, a sketch applied to show the putative binding sites of NR4A1 in *cbl-b* promoter. B, a diagram for *cbl-b* promoters with different lengths was designed for luciferase reporter construction. C, the relative luciferase activity of *cbl-b* promoters with different lengths was detected in both OV and NC cells. D, ChIP analysis. Chromatin DNA fragments combined with exogenous NR4A1-HA were pulled down by a monoclonal antibody of HA; thereafter, a pair of primers was applied to amplify the fragment of the *cbl-b* promoter containing a NR4A1 (or NBRE) binding site from the ChIP product. The data represented the means of three independent experiments; ** $P < .01$ and *** $P < .001$ vs ns; error bars indicate SD

protein level was not largely because of MKK4 transcription. Western blotting showed that TG or H_2O_2 treatment resulted in increased p-JNK level in *cbl-b* KD cells compared with that in control cells (Figure 5D-G).

3.6 | NR4A1 enhances the transactivation of *cbl-b* promoter via physical association

It was reported that NR4A1 enhanced the *cbl-b* promoter transactivation in human cells.⁴¹ We tested whether NR4A1 directly regulates *cbl-b* transactivation in mouse cells regarding that humans and mice have some differences in promoter sequences.

The promoter sequence of *cbl-b* between -14 and -2008 has four putative NR4A1 binding sites as shown in Figure 6A. We cloned four *cbl-b* luciferase reporters with different DNA lengths as shown in Figure 6B. We transfected the four *cbl-b* luciferase reporter plasmids into OV and NC cells, the dual-luciferase assay results showed that NR4A1 was able to enhance the luciferase activity of the four *cbl-b* reporters with different lengths, and the shortest promoter sequence of *cbl-b* (-14 to -499) was no less effective than the longer promoters (-14 to -995, -14 to -1454 and -22 to -2008) (Figure 6C). Namely, NR4A1 might modulate the promoter sequence at -14 to -499.

To verify whether NR4A1 associates with *cbl-b* promoter, we infected MIN6 cells with adenovirus encoding NR4A1-HA, and after the infection, the cells were applied for ChIP assay. The ChIP products obtained were applied for PCR examination. As shown in Figure 6D, the primers designed were covered the putative binding site at -118 to

-251 bp within -14 to -499. The results in Figure 6D exhibited that after ChIP pulled down by anti-HA antibodies, the putative binding site at -118 to -251 bp was amplified successfully. These results indicated that NR4A1 can physically associate with *cbl-b* promoter.

3.7 | Confirmation of the impact of NR4A1 on *cbl-b* expression in NR4A1-knockout mice

To confirm the conclusion that NR4A1 enhances the expression of *cbl-b* in vivo, we tested the expression of *cbl-b* in NR4A1-KO mice. We purified the islets from WT mice or NR4A1-KO mice, and further extracted the RNA and protein for real-time PCR and Western blotting. The mRNA expression of *cbl-b* in NR4A1-KO mice was much lower than that in WT mice (Figure 7A), whereas the *cbl-b* protein level in NR4A1-KO mice was also lower than that in WT mice (Figure 7B).

The immuno-staining data showed that the colocalization of *cbl-b* with insulin was dramatically decreased in the islets of pancreatic tissue from NR4A1-KO mice compared with that from WT mice (Figure 7C), which indicated that *cbl-b* protein expression in pancreatic β -cells was significantly reduced in NR4A1-KO mice.

3.8 | Summary of the results

Figure 7D is a graphic model for the mechanism that NR4A1 protects β -cells from ER stress or ROS-induced JNK phosphorylation. In

β -cells, ROS or ER stress results in JNK phosphorylation and consequently results in apoptosis. ROS or ER stress also induces NR4A1 expression, whereas NR4A1 enhances the expression of cbl-b and further increases the degradation of MKK4, which results in reduced JNK phosphorylation. The fate of the pancreatic β -cells is possibly determined by the balance between p-JNK and cbl-b.

In short, NR4A1 reduces p-JNK level by indirectly down-regulating the availability of its upstream MKK4, therefore, reduces the possibility of pancreatic β -cell apoptosis.

4 | DISCUSSION

Hyperglycaemia and hyperlipidaemia result in ER stress and/or increased ROS in pancreatic β -cells, and cause β cells damage and apoptosis.¹⁻⁶ Pancreatic β -cell loss plays an essential role in the pathogenesis of type 2 diabetes. Our pancreatic β -cells are always challenged with hyperglycaemia and/or hyperlipidaemia after meal, but they are always able to survive over the subsequent cellular stress by an unknown "protective network" with some protective molecules. If the "protective network" is weak, consequently, the pancreatic β -cells will be damaged or even undergo apoptosis. To search the molecules for pancreatic β -cell protection under cellular stress is important for type 2 diabetes prevention.

It was reported that ER stress or ROS resulted in increased JNK activation or phosphorylation in various cells^{31-33,35} and activated JNK further triggered the signalling pathways to apoptosis.³⁶⁻³⁹ We confirmed that MIN6 cells were treated with TG or H₂O₂ resulted in increased JNK phosphorylation. Some researcher applied JNK phosphorylation inhibitor which partially reversed the ER stress or ROS-induced apoptosis in Bcr/Abl-positive chronic myelogenous leukaemia cells,⁴⁵ which indicated that JNK phosphorylation was critical for ER stress or ROS-induced apoptosis.⁴⁶

As for the mechanisms of ROS or ER stress induced JNK activation, many research articles reported that ROS induced ER stress, which has been popularly accepted.⁴⁷⁻⁴⁹ Previously, we tested the survival of pancreatic beta cell (MIN6 cells) in response to TG (ER stress inducer) or H₂O₂ (a ROS molecule) and by comparison we found H₂O₂ resulted in apoptosis in a shorter time duration compared with TG.^{4,5} As the induction of JNK activation by H₂O₂ was much quicker than that by TG, it was reasonable that we speculated that ROS induced JNK activation via ER stress and another unknown parallel pathway.

NR4A1 (synonym as Nur77) was a multi-factor response molecule and an anti-stress factor.⁷⁻⁹ NR4A1 might have different roles in different tissues or different organs. People applied NR4A1 KO mice to test the pathophysiological changes in different research areas. For metabolism, Chao L C reported that NR4A1 KO mice fed with high-fat diet were much easier to become obese compared with WT mice⁵⁰; Pei L reported that NR4A1 had a role in gluconeogenesis⁵¹; Zhan Y Y reported that nur77 regulates LKB1 localization and thus modulates AMPK activation.⁵² For immunity system, some reporters showed that NR4A1 might have negative effect on T cell activation.^{53,54}

Previously, we found that the pancreatic cells from NR4A1 KO mice islets were more sensitive to TG or H₂O₂-induced apoptosis compared with that from WT mice.^{4,5} We observed that high-fat diet feeding resulted in increased 8-OHdG (a senescence marker) level and apoptosis rate in pancreatic cells from NR4A1 KO mice compared with that from WT mice.⁵ It is known that pancreatic cell senescence or/and pancreatic cell loss results in diabetes. We further figured out that NR4A1 directly or indirectly enhances the expression of some anti-apoptotic proteins, such as survivin, WT1, BCL2, SOD1 and GPX1.^{4,5,42} These anti-apoptotic proteins plus some other NR4A1-up-regulated molecules might form a protective network to work together to protect pancreatic β cells. Without the NR4A1-based protective network, the pancreatic β cells would be in great risk or danger. We are still exploring other possible members of NR4A1-up-regulated protective network in β cells by exploiting pancreatic β cell line and NR4A1 KO mice.

We previously found that NR4A1 was inducible upon treatment with ER stress inducer (TG) or ROS (H₂O₂) in pancreatic β -cells.^{4,5,42} Recently, we found that overexpression of NR4A1 in MIN6 cells resisted the JNK phosphorylation induced by TG or H₂O₂.

To study the mechanism of NR4A1 down-regulating JNK phosphorylation, we analysed the key upstream kinases for JNK and we focused on two potential JNK kinases, MKK4 and MKK7 as reported. We further confirmed that MKK4 and MKK7 were two upstream kinases for JNK activation in MIN6 cells induced by H₂O₂ or TG. We compared the protein expression levels of the two molecules vs NR4A1 expression level and found that NR4A1 overexpression in MIN6 cells resulted in reduced expression of MKK4 rather than MKK7. As a transcription factor, usually NR4A1 enhances the expression of a gene. How could NR4A1 reduce the protein expression of MKK4? There should be two possible ways for the reduction in a specific protein expression besides the transcription activity altering: One is to interfere the protein translation, and the other is to modulate the stability of the protein. We could not rule out that NR4A1 is able to reduce MKK4 protein translation via some special paths. But we did predict that NR4A1 might have the effect on the stability of MKK4 protein via an indirect way. Most proteins' stability is closely related to their ubiquitination level and proteasome degradation, which lies in the protein level of substrate-specific E3 ligase. We verified that the loss of MKK4 protein by NR4A1 overexpression was proteasome-dependent by using a specific drug MG132.

Thus, we searched the literatures and found that the NR4A1 was able to enhance the expression of three E3 ligases.^{19,40,41} However, we only found cbl-b mRNA level was increased in NR4A1 overexpression MIN6 cells. We further checked the protein expression of cbl-b in NR4A1 overexpression cells (OV cells), as expected that overexpressing NR4A1 resulted in enhanced protein expression of cbl-b. To confirm cbl-b was able to target MKK4, we knocked down the cbl-b expression in NR4A1 overexpression cells (OV cells) and found knocking-down expression of cbl-b in OV cells, resulted in increased MKK4 protein level and increased p-JNK level upon treatment with TG or H₂O₂. It was clear that cbl-b protein expression was




FIGURE 7 Confirmation of NR4A1 impacting *cbl-b* expression in NR4A1-KO mice. A, the relative mRNA levels of *cbl-b* were determined by real-time quantitative PCR in wild-type mice and NR4A1 KO mice. B, the relative protein levels of *cbl-b* and NR4A1 in pancreatic islets from WT or NR4A1 KO mice were determined by Western blotting. C, the colocalization of *cbl-b* (green) with insulin (red) was dramatically decreased in the islets of pancreatic tissue from NR4A1-KO mice compared with that from WT mice. The optical magnification of the image was 300 \times , and the larger square on the left bottom of each image was the enlarged image of the smaller section of the islet (the smaller square indicated within the islet). D, A graphic model for the mechanism that NR4A1 protects β -cells from JNK phosphorylation induced by ER stress or ROS. The data represented the means of three independent experiments; *** $P < .001$ vs ns; error bars indicate SD

reversely correlated with the MKK4 protein level or p-JNK level. Our data indicated that *cbl-b* was a down-stream molecule of NR4A1 to down-regulate MKK4 protein level in MIN6 cells.

As for *cbl-b*, Keane et al firstly cloned this gene. *Cbl-b*, with homology to the *c-cbl* proto-oncogene, has a proline-rich domain, a nuclear localization signal, a C3HC4 zinc finger and a putative leucine zipper. Their data suggested that *cbl-b* gene encodes a protein, which can interact with signal transduction proteins to regulate their function or to be regulated by them.⁵⁵ As *cbl-b* protein was reportedly found to contain a highly conserved RING finger domain with E3 ligase catalytic activity, it has been predicted that its biological functions are mediated via protein ubiquitination in diverse cellular signalling pathways. Such as, *cbl-b*, as E3 ubiquitin ligase, negatively regulates CLR-mediated antifungal innate immunity.⁵⁶ Upon nerve growth factor (NGF) stimulation, recruitment of *Cbl-b* promotes

tropomyosin-related kinase A (TrkA) ubiquitylation and degradation. *Cbl-b* limits NGF-TrkA signalling to control the length of neurites.⁵⁷ In macrophages, *Cbl-b* suppresses saturated FA-induced Toll-like receptor 4 (TLR4) signalling by ubiquitination and degradation of TLR4. *Cbl-b* deficiency could exaggerate HFD-induced insulin resistance through saturated FA-mediated macrophage activation.⁵⁸ We first discovered that *cbl-b* was able to degrade MKK4 through the ubiquitination-proteasome pathway, and thus to attenuate JNK phosphorylation in response to ER stress or ROS.

We further studied the possible mechanism(s) by which NR4A1 may up-regulate *cbl-b* expression. It was reported that NR4A1 enhanced *cbl-b* promoter transactivation in human cells. To verify whether NR4A1 directly enhances *cbl-b* transactivation in mice β -cells, we exploited luciferase assay and ChIP assay. We further found NR4A1 was able to enhance the transactivation activity of

cbl-b promoter and NR4A1 was able to physically associate with *cbl-b* at some putative targeting sequence. What we detected is the binding sites of NR4A1 in *cbl-b* promoter regulatory element in mouse cells were not the same as people found in human cells as reported.

In addition to our in vitro study, we further found the mRNA level or protein level of *cbl-b* was markedly reduced in the islets from NR4A1-KO mice compared with that from WT mice, which was further confirmed that NR4A1 was a positive factor for *cbl-b* expression in mice pancreatic islets. As shown in insulin immuno-staining data, the majority of cells (about 85%-90%) in pancreatic islets were β -cells. Therefore, the mRNA and protein expression levels of *cbl-b* detected from islets reflected the status of pancreatic β -cells to a large extent. To further clarify NR4A1 impacting *cbl-b* protein expression in β -cells, we applied insulin and *cbl-b* double immuno-staining scheme to show the colocalization of the two molecules in β -cells. Our immune-staining data showed that the protein level of *cbl-b* in pancreatic β -cells of NR4A1-KO mice was markedly reduced compared with that of WT mice. This evidence confirmed that NR4A1 impacted the expression of *cbl-b* in mice pancreatic β -cells, which was in concert with what we found in vitro.

In summary, our data demonstrated that NR4A1 could repress cellular ER stress or ROS-induced JNK activation via enhancing *cbl-b* expression to degrade MKK4 in pancreatic β -cells. Namely, NR4A1 down-regulates p-JNK level through NR4A1/*cbl-b*↑/MKK4↓/p-JNK↓ pathway. Our study indicates NR4A1 might be a major player in the protective network of pancreatic β -cells, and our finding might provide clues for β -cell protection and diabetes prevention.

ACKNOWLEDGEMENTS

Thankfulness. We are much thankful to Dr Guang-wei Wei for his suggestions and for his help with the high-quality animal facility.

CONFLICT OF INTEREST

No potential conflict of interest relevant to this article exists.

AUTHOR CONTRIBUTIONS

Xiangdong Wang: Conceptualization (lead); Funding acquisition (lead); Investigation (lead); Resources (lead); Supervision (lead); Validation (lead); Visualization (equal); Writing-original draft (lead); Writing-review & editing (equal). **Zeqing Pu:** Data curation (equal); Formal analysis (equal); Methodology (equal); Software (equal); Visualization (equal); Writing-original draft (equal). **Dong Liu:** Data curation (equal); Formal analysis (equal); Methodology (equal); Software (equal). **Hanse Pablick Patherny Lobo Mouguegue:** Data curation (equal); Investigation (equal); Methodology (equal). **Chengwen Jin:** Data curation (equal); Investigation (equal); Methodology (equal); Software (equal). **Esha Sadiq:** Data curation (equal); Methodology (equal). **Dandan Qin:** Data curation (equal); Formal analysis (equal); Methodology (equal); Software (equal). **Tianfu Yu:** Data curation (equal); Formal analysis (equal); Methodology (equal); Software (equal). **Chen Zong:** Data curation (equal); Methodology (equal); Software (equal). **Jicui Chen:** Methodology (equal); Software (equal);

Visualization (equal). **Ruxing Zhao:** Data curation (equal); Formal analysis (equal); Funding acquisition (supporting); Resources (equal). **Jingyu Lin:** Methodology (equal). **Jie Cheng:** Methodology (equal). **xiao yu:** Investigation (supporting); Methodology (equal). **Xia Li:** Conceptualization (equal); Funding acquisition (equal); Investigation (equal); Validation (equal). **Yuchao Zhang:** Methodology (equal); Software (equal). **Yuantao Liu:** Conceptualization (equal); Writing-original draft (equal). **Bo Qing Guan:** Conceptualization (supporting); Investigation (equal).

DATA AVAILABILITY STATEMENT

All data generated or analysed during this study are included in this published article, and the original data are always available to the editor and readers upon request.

ORCID

Dan-dan Qin  <https://orcid.org/0000-0001-8055-3681>
Xiao Yu  <https://orcid.org/0000-0003-1119-8334>

REFERENCES

- Butler AE, Janson J, Bonner-Weir S, Ritzel R, Rizza RA, Butler PC. β -cell deficit and increased β -cell apoptosis in humans with type 2 diabetes. *Diabetes*. 2003;52:102-110. <https://doi.org/10.2337/diabetes.52.1.102>
- Biarnés M, Montolio M, Nacher V, Raurell M, Soler J, Montanya E. β -cell death and mass in syngeneically transplanted islets exposed to short- and long-term hyperglycemia. *Diabetes*. 2002;51:66-72. <https://doi.org/10.2337/diabetes.51.1.66>
- Maris M, Robert S, Waelkens E, et al. Role of the saturated nonesterified fatty acid palmitate in beta cell dysfunction. *J Proteome Res*. 2013;12:347-362. <https://doi.org/10.1021/pr300596g>
- Yu C, Cui S, Zong C, et al. The orphan nuclear receptor NR4A1 protects pancreatic β -cells from endoplasmic reticulum (ER) stress-mediated apoptosis. *J Biol Chem*. 2015;290:20687-20699. <https://doi.org/10.1074/jbc.M115.654863>
- Zong C, Qin D, Yu C, et al. The stress-response molecule NR4A1 resists ROS-induced pancreatic β -cells apoptosis via WT1. *Cell Signal*. 2017;35:129-139. <https://doi.org/10.1016/j.cellsig.2017.03.012>
- Prause M, Christensen DP, Billestrup N, Mandrup-Poulsen T. JNK1 protects against glucolipotoxicity-mediated beta-cell apoptosis. *PLoS One*. 2014;9:e87067. <https://doi.org/10.1371/journal.pone.0087067>
- Berriel Diaz M, Lemke U, Herzig S. Discovering orphans' sweet secret: NR4A receptors and hepatic glucose production. *Cell Metab*. 2006;4:339-340. <https://doi.org/10.1016/j.cmet.2006.10.005>
- Li X, Wang Z, Zheng Y, et al. Nuclear receptor Nur77 facilitates melanoma cell survival under metabolic stress by protecting fatty acid oxidation. *Mol Cell*. 2018;69(3):480-492.e7. <https://doi.org/10.1016/j.molcel.2018.01.001>
- Hu M, Luo Q, Alitongbieke G, et al. Celastrol-induced Nur77 interaction with TRAF2 alleviates inflammation by promoting mitochondrial ubiquitination and autophagy. *Mol Cell*. 2017;66(1):141-153.e6. <https://doi.org/10.1016/j.molcel.2017.03.008>
- Walker NPC, Perlmann T, Wang Z, et al. Structure and function of Nur1 identifies a class of ligand-independent nuclear receptors. *Nature*. 2003;423:555-560. <https://doi.org/10.1038/nature01645>
- Mohan HM, Aherne CM, Rogers AC, Baird AW, Winter DC, Murphy EP. Molecular pathways: the role of NR4A orphan nuclear receptors in cancer. *Clin Cancer Res*. 2012;18:3223-3228. <https://doi.org/10.1158/1078-0432.CCR-11-2953>

12. Kurland IJ, Pei L, Waki H, Tontonoz P, Vaitheesvaran B, Wilpitz DC. NR4A orphan nuclear receptors are transcriptional regulators of hepatic glucose metabolism. *Nat Med*. 2006;12:1048-1055. <https://doi.org/10.1038/nm1471>
13. Seo H, Chen J, González-Avalos E, et al. TOX and TOX2 transcription factors cooperate with NR4A transcription factors to impose CD8+ T cell exhaustion. *Proc Natl Acad Sci USA*. 2019;116:12410-12415. [10.1073/pnas.1905675116](https://doi.org/10.1073/pnas.1905675116)
14. Boddupalli CS, Nair S, Gray SM, et al. ABC transporters and NR4A1 identify a quiescent subset of tissue-resident memory T cells. *J Clin Invest*. 2016;126:3905-3916. <https://doi.org/10.1172/JCI85329>
15. Hanna RN, Carlin LM, Hubbeling HG, et al. The transcription factor NR4A1 (Nur77) controls bone marrow differentiation and the survival of Ly6C- monocytes. *Nat Immunol*. 2011;12:778-785. <https://doi.org/10.1038/ni.2063>
16. Cheng Z, Völkens M, Din S, et al. Mitochondrial translocation of Nur77 mediates cardiomyocyte apoptosis. *Eur Heart J*. 2011;32:2179-2188. <https://doi.org/10.1093/eurheartj/ehq496>
17. Freire P, Conneely O. NR4A1 and NR4A3 restrict HSC proliferation via reciprocal regulation of C/EBP alpha and inflammatory signaling. *Blood*. 2018;131:1081-1093. <https://doi.org/10.1182/blood-2017-07-795757>
18. Deutsch AJA, Rinner B, Wenzl K, et al. NR4A1-mediated apoptosis suppresses lymphomagenesis and is associated with a favorable cancer-specific survival in patients with aggressive B-cell lymphomas. *Blood*. 2014;123:2367-2377. <https://doi.org/10.1182/blood-2013-08-518878>
19. Mullican S, Zhang S, Konopleva M, et al. Abrogation of nuclear receptors Nr4a3 and Nr4a1 leads to development of acute myeloid leukemia. *Nat Med*. 2007;13:730-735. <https://doi.org/10.1038/nm1579>
20. Ke N, Claassen G, Yu D, et al. Nuclear hormone receptor NR4A2 is involved in cell transformation and apoptosis. *Cancer Res*. 2004;64:8208-8212. <https://doi.org/10.1158/0008-5472.CAN-04-2134>
21. Suzuki S, Suzuki N, Mirtsos C, et al. Nur77 as a survival factor in tumor necrosis factor signaling. *Proc Natl Acad Sci USA*. 2003;100(14):8276-8280. <https://doi.org/10.1073/pnas.0932598100>
22. Lee S, Lee S, Jin U, et al. The orphan nuclear receptor NR4A1 (Nur77) regulates oxidative and endoplasmic reticulum stress in pancreatic cancer cells. *Mol Cancer Res*. 2014;12:527-538. <https://doi.org/10.1158/1541-7786.mcr-13-0567>
23. Lee S, Li X, Hedrick E, et al. Diindolylmethane analogs bind NR4A1 and Are NR4A1 antagonists in colon cancer cells. *Mol Endocrinol*. 2014;28:1729-1739. <https://doi.org/10.1210/me.2014-1102>
24. Lee S, Abdelrahim M, Yoon K, et al. Inactivation of the orphan nuclear receptor TR3/Nur77 inhibits pancreatic cancer cell and tumor growth. *Cancer Res*. 2010;70:6824-6836. <https://doi.org/10.1158/0008-5472.CAN-10-1992>
25. Oh S, Oh J, Lee C, et al. Expression of Twist2 is controlled by T-cell receptor signaling and determines the survival and death of thymocytes. *Cell Death Differ*. 2016;23:1804-1814. <https://doi.org/10.1038/cdd.2016.68>
26. Yang H, Zhan Q, Wan YY. Enrichment of Nur77 mediated by retinoic acid receptor β leads to apoptosis of human hepatocellular carcinoma cells induced by fenretinide and histone deacetylase inhibitors. *Hepatology*. 2011;53:865-874. <https://doi.org/10.1002/hep.24101>
27. Chen J, Fiskus W, Eaton K, et al. Cotreatment with BCL-2 antagonist sensitizes cutaneous T-cell lymphoma to lethal action of HDAC7-Nur77-based mechanism. *Blood*. 2009;113:4038-4048. <https://doi.org/10.1182/blood-2008-08-176024>
28. Shen H, Liu Z. JNK signaling pathway is a key modulator in cell death mediated by reactive oxygen and nitrogen species. *Free Rad Biol Med*. 2006;40:928-939. <https://doi.org/10.1016/j.freeradbiomed.2005.10.056>
29. Win S, Than TA, Min RWM, Aghajan M, Kaplowitz N. c-Jun N-terminal kinase mediates mouse liver injury through a novel Sab (SH3BP5)-dependent pathway leading to inactivation of intramitochondrial Src. *Hepatology*. 2016;63:1987-2003. <https://doi.org/10.1002/hep.28486>
30. Seki E, Brenner DA, Karin M. A Liver full of JNK: signaling in regulation of cell function and disease pathogenesis, and clinical approaches. *Gastroenterology*. 2012;143:307-320. <https://doi.org/10.1053/j.gastro.2012.06.004>
31. Ibrahim SH, Gores GJ. Who pulls the trigger: JNK activation in liver lipotoxicity? *J Hepatol*. 2011;2012(56):17-19. <https://doi.org/10.1016/j.jhep.2011.04.017>
32. Czaja MJ. JNK regulation of hepatic manifestations of the metabolic syndrome. *Trends Endocrinol Metab*. 2010;21:707-713. <https://doi.org/10.1016/j.tem.2010.08.010>
33. Holzer R, Park E, Li N, et al. Saturated fatty acids induce c-Src clustering within membrane subdomains, leading to JNK activation. *Cell*. 2011;147:173-184. <https://doi.org/10.1016/j.cell.2011.08.034>
34. Liu J, Lin A. Role of JNK activation in apoptosis: a double-edged sword. *Cell Res*. 2005;15(1):36-42. <https://doi.org/10.1038/sj.cr.7290262>
35. Win S, Than TA, Le BHA, García-Ruiz C, Fernández-Checa JC, Kaplowitz N. Sab (Sh3bp5) dependence of JNK mediated inhibition of mitochondrial respiration in palmitic acid induced hepatocyte lipotoxicity. *J Hepatol*. 2015;62:1367-1374. <https://doi.org/10.1016/j.jhep.2015.01.032>
36. Taylor CA, Zheng Q, Liu Z, Thompson JE. Role of p38 and JNK MAPK signaling pathways and tumor suppressor p53 on induction of apoptosis in response to Ad-elf5A1 in A549 lung cancer cells. *Mol Cancer*. 2013;12:35. <https://doi.org/10.1186/1476-4598-12-35>
37. Konishi H, Fujiya M, Tanaka H, et al. Probiotic-derived ferrichrome inhibits colon cancer progression via JNK-mediated apoptosis. *Nat Commun*. 2016;7:12365. <https://doi.org/10.1038/ncomms12365>
38. Lin S, Hoffmann K, Gao C, Petruionis M, Herr I, Schemmer P. Melatonin promotes sorafenib-induced apoptosis through synergistic activation of JNK/c-jun pathway in human hepatocellular carcinoma. *J Pineal Res*. 2017;62:e12398. <https://doi.org/10.1111/jpi.12398>
39. Song IS, Jun SY, Na H, et al. Inhibition of MKK7-JNK by the TOR signaling pathway regulator-like protein contributes to resistance of HCC cells to TRAIL-induced apoptosis. *Gastroenterology*. 2012;143:1341-1351. <https://doi.org/10.1053/j.gastro.2012.07.103>
40. Yang L, Zhao L, Gan Z, et al. Deficiency in RNA editing enzyme ADAR2 impairs regulated exocytosis. *FASEB J*. 2010;24:3720-3732. <https://doi.org/10.1096/fj.09-152363>
41. Li X, Wei W, Huynh H, Zuo H, Wang X, Wan Y. Nur77 prevents excessive osteoclastogenesis by inducing ubiquitin ligase Cbl-b to mediate NFATc1 self-limitation. *eLife*. 2015;4:e07217. <https://doi.org/10.7554/eLife.07217>
42. Yang Y, Xie F, Qin D, et al. The orphan nuclear receptor NR4A1 attenuates oxidative stress-induced β cells apoptosis via up-regulation of glutathione peroxidase 1. *Life Sci*. 2018;203:225-232. <https://doi.org/10.1016/j.lfs.2018.04.027>
43. Gao W, Fu Y, Yu C, et al. Elevation of NR4A3 expression and its possible role in modulating insulin expression in the pancreatic beta cell. *PLoS One*. 2014;9:e91462. <https://doi.org/10.1371/journal.pone.0091462>
44. Lin H, Lin Q, Liu M, et al. PKA/Smurf1 signaling-mediated stabilization of Nur77 is required for anticancer drug cisplatin-induced apoptosis. *Oncogene*. 2014;33:1629-1639. <https://doi.org/10.1038/onc.2013.116>
45. Mao X, Rong YUC, Hua Li W, Xin LW. Induction of apoptosis by shikonin through a ROS/JNK-mediated process in Bcr/Abl-positive chronic myelogenous leukemia (CML) cells. *Cell Res*. 2008;18:879-888. <https://doi.org/10.1038/cr.2008.86>

46. Timmins JM, Ozcan L, Seimon TA, et al. Calcium/calmodulin-dependent protein kinase II links ER stress with Fas and mitochondrial apoptosis pathways. *J Clin Invest*. 2009;119:2925-2941. <https://doi.org/10.1172/jci38857>
47. Park S, Lim W, Bazer FW, Song G. Apigenin induces ROS-dependent apoptosis and ER stress in human endometriosis cells. *J Cell Physiol*. 2018;233(4):3055-3065. <https://doi.org/10.1002/jcp.26054>
48. Wang Y, Lee K, Lim M, Choi J. TRPV1 antagonist DWP05195 induces ER stress-dependent apoptosis through the ROS-p38-CHOP pathway in human ovarian cancer cells. *Cancers*. 2020;12(6):1702. <https://doi.org/10.3390/cancers12061702>
49. Lin M, Li L, Zhang Y, et al. Baicalin ameliorates H₂O₂ induced cytotoxicity in HK-2 cells through the inhibition of ER stress and the activation of Nrf2 Signaling. *Int J Mol Sci*. 2014;15:12507-12522. <https://doi.org/10.3390/ijms150712507>
50. Chao LC, Wroblewski K, Zhang Z, et al. Insulin resistance and altered systemic glucose metabolism in mice lacking Nur77. *Diabetes*. 2009;58:2788-2796. <https://doi.org/10.2337/db09-0763>
51. Pei L, Waki H, Tontonoz P, Vaitheesvaran B, Wilpitz DC. NR4A orphan nuclear receptors are transcriptional regulators of hepatic glucose metabolism. *Nat Med*. 2006;12:1048-1055. <https://doi.org/10.1038/nm1471>
52. Zhan Y, Chen Y, Zhang Q, et al. The orphan nuclear receptor Nur77 regulates LKB1 localization and activates AMPK. *Nat Chem Biol*. 2012;8:897-904. <https://doi.org/10.1038/nchembio.1069>
53. Liu X, Wang Y, Lu H, et al. Genome-wide analysis identifies NR4A1 as a key mediator of T cell dysfunction. *Nature*. 2019;567:525-529. <https://doi.org/10.1038/s41586-019-0979-8>
54. Chen J, López-Moyado IF, Seo H, et al. NR4A transcription factors limit CAR T cell function in solid tumours. *Nature*. 2019;567:530-534. <https://doi.org/10.1038/s41586-019-0985-x>
55. Keane MM, Rivero-Lezcano OM, Mitchell JA, Robbins KC, Lipkowitz S. Cloning and characterization of cbl-b: A SH3 binding protein with homology to the c-cbl proto-oncogene. *Oncogene*. 1995;10:2367-2377. [https://doi.org/10.1016/0360-3016\(95\)93133-R](https://doi.org/10.1016/0360-3016(95)93133-R)
56. Zhu L, Luo T, Xu X, et al. E3 ubiquitin ligase Cbl-b negatively regulates C-type lectin receptor-mediated antifungal innate immunity. *J Exp Med*. 2016;213:1555-1570. <https://doi.org/10.1084/jem.20151932>
57. Emdal KB, Pedersen A, Bekker-Jensen DB, et al. Temporal proteomics of NGF-TrkA signaling identifies an inhibitory role for the E3 ligase Cbl-b in neuroblastoma cell differentiation. *Sci Signal*. 2015;8:ra40. <https://doi.org/10.1126/scisignal.2005769>
58. Abe T, Hirasaka K, Kagawa S, et al. Cbl-b is a critical regulator of macrophage activation associated with obesity-induced insulin resistance in mice. *Diabetes*. 2013;62:1957-1969. <https://doi.org/10.2337/db12-0677>

How to cite this article: Pu Z-Q, Liu D, Lobo Mouguegue HPP, et al. NR4A1 counteracts JNK activation incurred by ER stress or ROS in pancreatic β -cells for protection. *J Cell Mol Med*. 2020;24:14171-14183. <https://doi.org/10.1111/jcmm.16028>