

RESEARCH ARTICLE

Open Access

The impact of dialysis therapy on older patients with advanced chronic kidney disease: a nationwide population-based study

Chia-Jen Shih^{1,2†}, Yung-Tai Chen^{2,3†}, Shuo-Ming Ou^{2,4,5}, Wu-Chang Yang^{2,4}, Shu-Chen Kuo^{2,6,7} and Der-Cherng Tarn^{2,4,5*} for the Taiwan Geriatric Kidney Disease Research (TGKD) Group

Abstract

Background: Older patients with advanced chronic kidney disease (CKD) face the decision of whether to undergo dialysis. Currently available data on this issue are limited because they were generated by small, short-term studies with statistical drawbacks. Further research is urgently needed to provide objective information for dialysis decision making in older patients with advanced CKD.

Methods: This nationwide population-based cohort study was conducted using Taiwan's National Health Insurance Research Database. Data from 2000 to 2010 were extracted. A total of 8,341 patients ≥ 70 years old with advanced CKD and serum creatinine levels > 6 mg/dl, who had been treated with erythropoiesis-stimulating agents were included. Cox proportional hazard models in which initiation of chronic dialysis was defined as the time-dependent covariate were used to calculate adjusted hazard ratios for mortality. The endpoint was all-cause mortality.

Results: During a median follow-up period of 2.7 years, 6,292 (75.4%) older patients chose dialysis therapy and 2,049 (24.6%) received conservative care. Dialysis was initiated to treat kidney failure a median of 6.4 months after enrollment. Dialysis was associated with a 1.4-fold increased risk of mortality compared with conservative care (adjusted hazard ratio 1.39, 95% confidence interval 1.30 to 1.49). In subgroup analyses, the risk of mortality remained consistently increased, independent of age, sex and comorbidities.

Conclusions: In older patients, dialysis may be associated with increased mortality risk and healthcare cost compared with conservative care. For patients who are ≥ 70 years old with advanced CKD, decision making about whether to undergo dialysis should be weighted by consideration of risks and benefits.

Keywords: Advanced chronic kidney disease, Dialysis, Conservative care, Older people

* Correspondence: dctarn@vghtpe.gov.tw

†Equal contributors

²School of Medicine, National Yang-Ming University, Taipei, Taiwan

⁴Division of Nephrology, Department of Medicine, Taipei Veterans General

Hospital, No. 201, Section 2, Shih-Pai Road, Taipei 11217, Taiwan

Full list of author information is available at the end of the article

Background

Older patients with chronic kidney disease (CKD) stage 5 comprise a rapidly growing emerging population that may face the dilemma of whether to undergo dialysis or receive conservative care [1]. Although patients with end-stage renal disease (ESRD) tend to choose dialysis therapy over conservative care, the use of dialysis to treat kidney failure peaks in patients who are 75 years old and declines thereafter [2,3]. The perceived financial burden of dialysis, as well as a high comorbidity rate, uncertainty about the treatment's long-term benefit, and sense of life completion and acceptance of death, leads many older patients to forego dialysis [4]. The most critical concern of older patients with CKD stage 5 is whether dialysis provides benefits such as increased life expectancy and improved functional ability at the end of life [5,6]. The new guidelines of the Renal Physicians Association and the American Society of Nephrology on the initiation and withholding or withdrawal of dialysis emphasize shared decision making with patients, their family members and physicians in charge of their care [7]. Although comprehensive physician-patient communication may help to achieve treatment goals and avoid unnecessary medical expenditure [8,9], independent and objective evidence of the comparative effectiveness of treatment options is still needed to guide older patients' decision making about dialysis.

To date, very few studies have investigated the risks and benefits of dialysis therapy for older patients with advanced CKD. Previous studies examining this issue found that dialysis conferred a significant but small advantage over conservative care in older patients [10-12]. However, the statistical power of these studies was limited due to small samples, short follow-up periods and limitations of analytical methodologies. Because initiation of chronic dialysis commonly occurs during follow up in patients with advanced CKD, the use of appropriate statistical methods to calculate the fraction of mortality attributable to dialysis is essential. Accordingly, in this nationwide population-based cohort study based on Taiwan's National Health Insurance Research Database (NHIRD), we utilized the initiation of chronic dialysis as a time-dependent covariate in Cox regression models to assess the real effect of dialysis on older patients with advanced CKD.

Methods

Data source

The present study was performed using Taiwan's NHIRD. Taiwan launched the national health insurance (NHI) program in 1995. This single-payer system covers approximately 99% of residents. In 1999, as part of the NHIRD project, the Bureau of National Health Insurance began to release patient data in electronic form for research

purposes. These de-identified secondary data include all registry and claims data, ranging from demographic data to detailed orders from ambulatory and inpatient care. Diseases are coded according to the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM). The accuracy of diagnoses registered in the NHIRD has been validated for several diseases, including acute kidney injury [13], chronic kidney disease [14-16], acute coronary syndrome [17], ischemic stroke [18] and diabetes [19]. Because the dataset consisted of de-identified secondary data, this study was exempted from full review by the Institutional Review Board of Taipei City Hospital (TCHIRB-1030407-W).

Study design

This population-based, observational, retrospective cohort study was performed to determine the association between chronic dialysis and mortality in older patients with advanced CKD. We identified all subjects ≥ 70 years old in 2000 and extracted all relevant data for these subjects for the study period of January 2000 to December 2010. These data included demographic characteristics, diagnosis and procedure codes, drug prescriptions and information about outpatient visits and hospital admissions. We further extracted data from January 1995 to December 1999 to ensure the availability of information for all individuals for at least five years preceding enrollment; this information was used to identify comorbidities.

Among all individuals ≥ 70 years old in January 2000 in Taiwan, patients with ICD-9-CM codes for CKD (016.0, 042, 095.4, 189, 223, 236.9, 250.4, 271.4, 274.1, 403-404, 440.1, 442.1, 446.21, 447.3, 572.4, 580-589, 590-591, 593, 642.1, 646.2, 753 and 984) and receiving erythropoiesis-stimulating agents (ESAs) were identified. The first date of prescription of ESAs was defined as the index date. We excluded patients with histories of cancer, patients receiving chronic dialysis or kidney transplantation before or during the 30 days after the index date and patients with follow-up periods < 30 days. According to NHI reimbursement regulations, patients with CKD, serum creatinine levels > 6 mg/dl (approximately equivalent to glomerular filtration rate [GFR] < 15 ml/min/1.73 m²), and hematocrit $< 28\%$ should receive ESAs to maintain a target hematocrit level not to exceed 36%. In addition, unlike the relative low prevalence of ESA utilization (less than 20%) among CKD or ESRD patients in the US [20,21], a report from the Taiwan Department of Health indicated that 85% of patients with advanced CKD stage 5 not yet requiring dialysis received ESAs therapy in 2012 [22], possibly due to convenient medical access and minimal financial barrier of health insurance access in Taiwan. The median hematocrit value at the initiation of dialysis was 24.2% (interquartile range 20.6% to 27.5%) in Taiwan [23]. Thus, the selected cohort in our study is most representative of

older patients with advanced CKD not yet requiring dialysis in Taiwan.

During the study period, the records of patients receiving chronic dialysis were extracted from the Registry of Catastrophic Illness. For catastrophic illnesses such as ESRD on chronic dialysis, the government registered the confirmed subjects, after strict verification. After successful certification, ESRD patients on chronic dialysis can be exempted from related medical expenses. Therefore, the application of catastrophic illness certificate for ESRD on dialysis required specialist nephrological reviews in supported medical records, examination reports and imaging studies after careful exclusion of the causes of acute renal failure [24].

Outcome measures

The endpoint was all-cause mortality. All subjects were followed until death or 31 December 2011.

Potential confounders

Baseline demographic data, including age, sex, economic status and urbanization level of the patients' places of residence, were collected. Patients' systemic health status was evaluated using the Charlson Comorbidity Index (CCI). Each increase in the CCI represents a stepwise increase in cumulative mortality, that is, a score of 0 is associated with a 99% 10-year survival rate and a score of 5 is associated with a 34% 10-year survival rate [25]. The effect of primary renal disease on survival rate was also taken into consideration. Data on other comorbidities that affect survival and are not included in the CCI, such as diabetes mellitus, hypertension, dyslipidemia, atrial fibrillation, valvular heart disease, parkinsonism, autoimmune disease and drug abuse, were also extracted. Prescribed medications that could confound mortality, such as antiplatelet agents, warfarin, angiotensin-converting enzyme inhibitors (ACEIs), angiotensin receptor blockers (ARBs), beta blockers, calcium channel blockers, diuretics, nitrate, statins, dipyridamole, steroids, estrogen or progesterone, non-steroidal anti-inflammatory drugs, selective serotonin re-uptake inhibitors, proton pump inhibitors and oral hypoglycemic drugs, were identified.

Statistical analysis

The baseline characteristics were first analyzed using descriptive statistics and then compared using Pearson χ^2 tests for categorical variables, and the independent *t*-test and Mann-Whitney *U*-test for parametric and nonparametric continuous variables, respectively. Cox regression models were used to calculate hazard ratios (HRs) and 95% confidence intervals (CIs) characterizing the risk of mortality after chronic dialysis. Because no patient was undergoing chronic dialysis at the time of enrollment, this variable was calculated as a time-dependent

covariate to ensure that patients were considered at risk only when they were receiving chronic dialysis. The models allowed patients to switch from one exposure group to another and were adjusted for the variables listed in Table 1. Subgroup Cox regression analyses were performed to examine the effects of age, sex, CCI, underlying disease and time cohort on the risk of mortality after chronic dialysis. Interaction effects were examined using the likelihood ratio test.

Because the baseline characteristics of patients receiving chronic dialysis differed substantially from those receiving conservative care, we also performed a propensity score-matched analysis. Propensity scores representing the likelihood of receiving chronic dialysis were calculated using logistic regression analysis, conditional on the baseline covariates listed in Table 1. For each subject who received chronic dialysis, we randomly selected one subject who received conservative care based on nearest neighbor matching without replacement using calipers of width equal to 0.1 standard deviation of the logit of the propensity score. Comparisons of mortality rates and healthcare costs between groups were performed. Microsoft SQL Server 2012 (Microsoft Corporation, Redmond, WA, USA) was used for data linkage, processing and sampling. Propensity score was calculated using SAS software (version 12.0; SAS Institute, Inc., Cary, NC, USA). All other statistical analyses were conducted using STATA statistical software (version 12.0; StataCorp., College Station, Texas, USA). Statistical significance was defined as $P < 0.05$.

Results

Characteristics of the study population

We identified 8,341 older patients with advanced CKD (estimated GFR < 15 ml/min/1.73 m²) who met the inclusion criteria between January 2000 and December 2010. The mean age was 79.4 (standard deviation (SD) 7.0) years. Most (55.3%) patients were female, and the mean CCI was 5.6 (SD 2.1). A plurality of patients had the following comorbidities: hypertension, diabetes mellitus or dyslipidemia. Common concomitant medications were calcium channel blockers, diuretics, ACEIs or ARBs and dipyridamole, followed by antiplatelet agents (Table 1). During the follow-up period, 75.4% of the study subjects ($n = 6,292$) received chronic dialysis, and the rest ($n = 2,049$) received conservative care.

Incidence rate and risk of mortality among older patients with advanced CKD

A total of 5,807 deaths occurred in older patients with advanced CKD during the follow-up period of 28,397 person-years. The overall mortality rate was 204.5 per 1,000 person-years. Compared with conservative care, chronic dialysis was associated significantly with a higher

Table 1 Baseline characteristics of patients

Parameters	All patients	During follow-up period		P
		Not receiving chronic dialysis	Receiving chronic dialysis	
Number of patients	8,341	2,049	6,292	
Male, number (%)	3,726 (44.7)	955 (46.6)	2,771 (44.0)	0.042
Age, mean (SD), years	79.4 (7.0)	82.0 (6.4)	78.6 (7.1)	<0.001
Follow-up period, mean (SD), days	1,026 (880)	478 (546)	1,205 (894)	<0.001
Day of chronic dialysis initiation, median (IQR)	-	-	192 (90 to 404)	
Age at chronic dialysis initiation, mean (SD), years	-	-	79.4 (7.1)	
Monthly income (US\$), number (%)				0.158
Dependent	3,395 (40.7)	831 (40.6)	2,564 (40.8)	
0 to 637	1,933 (23.2)	505 (24.7)	1,428 (22.7)	
637 to 1,400	2,955 (35.4)	703 (34.3)	2,252 (35.8)	
>1,400	58 (0.7)	10 (0.5)	48 (0.8)	
Urbanization level ^a , number (%)				0.150
1	4,210 (50.5)	1,021 (49.8)	3,189 (50.7)	
2	3,236 (38.8)	800 (39.0)	2,436 (38.7)	
3	740 (8.9)	178 (8.7)	562 (8.9)	
4	155 (1.9)	50 (2.4)	105 (1.7)	
Charlson Comorbidity Index ^b , mean (SD)	5.6 (2.1)	5.7 (2.2)	5.5 (2.1)	0.003
Primary renal disease, number (%)				
Diabetes	3,070 (36.8)	721 (35.1)	2,349 (37.3)	0.080
Glomerulonephritis	4,008 (48.0)	984 (48.0)	3,024 (48.0)	0.979
Secondary glomerulonephritis/vasculitis	86 (1.0)	26 (1.2)	60 (0.9)	0.220
Hypertension/large vessel disease	986 (11.8)	273 (13.3)	713 (11.3)	0.015
Cystic/hereditary/congenital disease	615 (7.3)	142 (6.9)	473 (7.5)	0.377
Miscellaneous conditions	186 (2.2)	54 (2.6)	132 (2.0)	0.152
Comorbidities, number (%)				
Diabetes mellitus	4,807 (57.6)	1,152 (56.2)	3,655 (58.1)	0.137
Hypertension	7,807 (93.6)	1,904 (92.9)	5,903 (93.8)	0.151
Dyslipidemia	3,972 (47.6)	890 (43.4)	3,082 (49.0)	<0.001
Atrial fibrillation	579 (6.9)	170 (8.3)	409 (6.5)	0.005
Valvular heart disease	1,698 (20.4)	472 (23.0)	1,226 (19.5)	0.001
Parkinsonism	617 (7.4)	191 (9.3)	426 (6.8)	<0.001
Autoimmune disease	1,190 (14.3)	309 (15.1)	881 (14.0)	0.225
Drug abuse	42 (0.5)	13 (0.6)	29 (0.5)	0.335
Concomitant medications, number (%)				
Antiplatelet agents ^c	1,948 (23.4)	507 (24.7)	1,441 (22.9)	0.087
Warfarin	59 (0.7)	10 (0.5)	49 (0.8)	0.173
ACE inhibitors or ARB	2,716 (32.6)	678 (33.1)	2,038 (32.4)	0.558
Beta blockers	380 (4.6)	83 (4.1)	297 (4.7)	0.207
Calcium channel blockers	4,868 (58.4)	977 (47.7)	3,891 (61.8)	<0.001
Diuretics	4,707 (56.4)	1,098 (53.4)	3,609 (57.4)	0.003
Nitrate	1,626 (19.5)	376 (18.4)	1,250 (19.9)	0.132
Statins	710 (8.5)	146 (7.1)	564 (9.0)	0.010
Dipyridamole	2,107 (25.3)	463 (22.6)	1,644 (26.1)	0.001

Table 1 Baseline characteristics of patients (Continued)

Steroids	690 (8.3)	189 (9.2)	501 (8.0)	0.072
Estrogen or progesterone	38 (0.5)	11 (0.5)	27 (0.4)	0.529
Non-steroidal anti-inflammatory drugs	1,248 (15.0)	312 (15.3)	936 (14.9)	0.699
Selective serotonin re-uptake inhibitors	102 (1.2)	28 (1.4)	74 (1.2)	0.496
Proton pump inhibitors	473 (5.7)	157 (7.7)	316 (5.0)	<0.001
Oral hypoglycemic drugs	1,843 (22.1)	414 (20.2)	1,429 (22.7)	0.018

^aUrbanization levels in Taiwan are divided into four strata according to Taiwan National Health Research Institute publications. Level 1 designates the most urbanized areas and level 4 designates the least urbanized areas. ^bThe Charlson Comorbidity Index (CCI) is used to determine overall systemic health. Each increase in CCI represents a stepwise increase in cumulative mortality. ^cIncluding aspirin, clopidogrel, ticlopidine, and cilostazol. ACE, angiotensin-converting enzyme; ARB, angiotensin II receptor blocker; IQR interquartile range; SD, standard deviation.

mortality risk (adjusted HR (aHR) 1.39, 95% CI 1.30 to 1.49, $P < 0.001$; Table 2). The cumulative incidence of mortality is illustrated in Figure 1. Other significant risk factors for mortality among older patients with advanced CKD included older age (aHR per age 1.07, 95% CI 1.06 to 1.07), male (aHR 1.11, 95% CI 1.05 to 1.17), higher CCI (aHR per score 1.08, 95% CI 1.07 to 1.10), parkinsonism (aHR 1.24, 95% CI 1.13 to 1.37), use of diuretic (aHR 1.14, 95% CI 1.07 to 1.20) or steroid (aHR 1.11, 95% CI 1.01 to 1.21) (Table 3).

In subgroup analysis, age ($P < 0.001$), hypertension ($P = 0.010$) and year of index date ($P = 0.027$) interacted significantly with the receipt of chronic dialysis. Compared with conservative care, chronic dialysis increased mortality risk among patients <80 years old, those without hypertension and those diagnosed with advanced CKD between 2000 and 2002 (Figure 2).

Healthcare expense in older patients with advanced CKD undergoing chronic dialysis versus conservative care

During the 28,397-person-year follow-up period, chronic dialysis was initiated for 6,292 patients at a median of 192 days after enrollment and a mean age of 79.4 (SD 7.1) years. Total healthcare costs, including total hospital and ambulatory visit costs, were included in our analysis. Healthcare utilization costs were retrieved primarily from NHI records, and also from insurance deductible data. Total healthcare cost categories were dialysis costs, laboratory and examination fees, medication costs, fees for procedures or surgery and nurse and physician fees. The mean healthcare cost for patients receiving chronic dialysis was US\$23,994 (95% CI \$23,616 to \$24,371) per person-year; this cost was US\$8,738 (95% CI \$8,409 to \$9,068) per person-year for the pre-dialysis period and US\$42,980 (95% CI \$37,050 to \$48,909) per person-year

for the period after initiation of chronic dialysis. The mean healthcare cost among the 2,049 patients who never received chronic dialysis was US\$18,252 (95% CI \$16,990 to \$19,514) per person-year (Table 4).

Propensity score-matched analysis

To further validate the study results, a propensity score analysis including 1,984 study subjects who received chronic dialysis and 1,984 matched patients who did not receive chronic dialysis was conducted. Potentially confounding baseline characteristics did not differ significantly between these groups [see Additional file 1: Table S1]. Compared with matched subjects who received conservative care, patients who received chronic dialysis had a significantly increased risk of mortality (HR 1.16, 95% CI 1.07 to 1.25, $P < 0.001$; Additional file 1: Table S2) and higher total healthcare cost (US\$26,273 versus US\$17,973 per person-year; $P < 0.001$; Additional file 1: Table S3).


Discussion

This nationwide population-based study provides novel evidence that dialysis therapy does not always provide a substantial survival advantage among older patients with advanced CKD. Most of these patients face the prospect of dialysis therapy for the remainder of their lives. Thus, an understanding of the survival benefit of dialysis for this population is important. Using the initiation of chronic dialysis as a time-dependent variable in a Cox regression model, we found that dialysis therapy was associated with a nearly 40% increase in mortality risk, in patients ≥ 70 years old compared with those receiving conservative care. These increases remained significant in propensity score-matched analysis.

Table 2 Hazard ratios for mortality according to receipt of chronic dialysis during follow up

Variables	Number of deaths	Exposure time, patient-years	Crude hazard ratio (95% CI)	P value	Adjusted hazard ratio ^a (95% CI)	P value
Not receiving chronic dialysis	1,699	13,071	(Referent)		(Referent)	
Receiving chronic dialysis ^b	4,108	15,326	1.35 (1.26 to 1.44)	<0.001	1.39 (1.30 to 1.49)	<0.001

^aAdjusted for all covariates listed in Table 1. ^bChronic dialysis was calculated as a discrete time-dependent covariate. CI, confidence interval.



The risk of approaching dialysis was found to exceed that of mortality in most older patients with estimated GFR <15 ml/min/1.73 m² [26]. Some authors have argued that older patients may have no real choice in dialysis decision making due to the lack of comprehensive insurance coverage, which may prohibit access to dialysis through implicit or explicit dialysis rationing due to limited medical resources and financial barriers [27-29]. Our study results, however, suggest that NHI coverage of dialysis expenses eliminates financial barriers for patients who may benefit from the treatment. Furthermore, nephrologists in Taiwan cannot legally withdraw or withhold dialysis without patient agreement, even when they believe that the treatment will have no benefit or that any benefit is outweighed by the burdens of treatment. In other words, patients and their families take active roles in dialysis decision making. Thus, our findings

Table 3 Significant risk factors for mortality among older patients with advanced chronic kidney disease

Variables	Adjusted ^a	
	Hazard ratio (95% CI)	P value
Age, per year	1.07 (1.06 to 1.07)	<0.001
Male	1.11 (1.05 to 1.17)	<0.001
Charlson comorbidity index score, per 1 point	1.08 (1.07 to 1.10)	<0.001
Receiving chronic dialysis ^b	1.39 (1.30 to 1.49)	<0.001
Dyslipidemia	0.86 (0.81 to 0.91)	<0.001
Parkinsonism	1.24 (1.13 to 1.37)	<0.001
Calcium channel blocker use	0.88 (0.83 to 0.93)	<0.001
Diuretic use	1.14 (1.07 to 1.20)	<0.001
Dipyridamole use	0.93 (0.88 to 0.99)	0.019
Steroid use	1.11 (1.01 to 1.21)	0.034

^aAdjusted for all ovariates in Table 1. ^bChronic dialysis was calculated as a discrete time-dependent covariate. CI, confidence interval.

provide unrestricted objective evidence that can be used to optimize the risk-benefit analysis of dialysis therapy in older patients with advanced CKD.

The survival of patients undergoing dialysis in Taiwan improved rapidly after the initiation of the NHI program in 1995. However, outcomes of older patients have not improved substantially despite public insurance benefits enabling free healthcare access and total coverage of medical expenses [30]. Furthermore, Wu and colleagues [31] found that incident dialysis has been associated with a 6.27- to 10.4-fold greater risk of mortality in patients ≥70 years old compared with those <30 years old, even after adjusting for CCI. Among US nursing home residents, initiation of dialysis was related to substantial functional decline and up to 60% mortality within one year [32]. The cost of dialysis care also increases with age and the number of comorbidities [33]. In our study, the average annual per-patient cost of dialysis was much higher than that of conservative care among older patients; dialysis increased the total cost by US\$5,742 per patient year. The healthcare cost was also significantly higher after initiation of chronic dialysis than in the pre-dialysis period (US\$42,980 *versus* US\$8,738 per patient year). In 2011, Medicare expenditures for dialysis in the US were \$71,630 to \$87,945 per patient year [1], at least double the costs calculated in the present study. Thus, ample opportunity exists to improve the outcomes of older patients after the initiation of dialysis therapy from a global standpoint. Alternatively, our findings support that robust conservative care was encouraged when considering opportunity cost principles. However, risk-benefit analysis in the present study is insufficient for decision making about dialysis for older patients; examination of this issue requires further long-term assessment.

Most previous small-scale studies [10-12,34,35] found that dialysis therapy conferred a modest (2 to 45.9 months) absolute survival advantage over conservative care in older patients, although this advantage was largely offset by high comorbidity rates. By contrast, our study demonstrated that dialysis therapy in older patients increased mortality risk by almost 40% compared with conservative care. This discrepancy may be due to differences in enrollment criteria and drawbacks of the statistical methods used in previous studies, such as the consideration of initiation of chronic dialysis as a dichotomous (non-time-dependent) variable in Cox proportional hazards models. This approach involves the examination of the relationship between survival and patient characteristics at the time of study enrollment, which may result in a false increase in survival rate in the dialysis group because patients who survived longer had an increased likelihood of receiving dialysis therapy. Several prognostic models have been developed to resolve this issue [36-38]. The definition of initiation of chronic dialysis as a time-dependent covariate in




Figure 2 Subgroup analyses of associations between chronic dialysis and mortality risk among older patients with advanced chronic kidney disease.

the present study allowed for the divergence of survival curves after this event had occurred. This type of model ensures that the risk of mortality increases only after dialysis initiation, which is more accurate and clinically relevant.

Our subgroup analyses showed that the risk of mortality was consistently elevated in older patients receiving dialysis therapy, regardless of age, sex, comorbidities or time cohort. Mortality risk was lower in patients >80 years old than in those 70 to 80 years old, implying that the impact of dialysis on increased mortality risk became less dominant with advanced age in patients with advanced CKD. The influence of dialysis on mortality in very old patients may be offset by competition between the risks of dialysis and death. The mortality risk associated with dialysis increased consistently, irrespective of specific comorbidities; however, the adjusted HR of mortality was lower in patients with, than in those without, hypertension. This result may be due to the vulnerability of older patients receiving dialysis

to intradialytic hypotension, such that those with higher blood pressure targets may have a survival benefit [39]. Moreover, the effect of dialysis on mortality risk was smaller in the 2009 to 2010 cohort than in the 2000 to 2002 cohort. This finding may be attributed to advances in medical therapy and improvement in dialysis care over time.

Our study has several strengths. First, we present the largest currently available database for older patients with advanced CKD and their dialysis outcomes, with an extended follow-up period. Second, previous studies found that late or no referral to a nephrologist may be associated not only with an increased risk of short-term mortality, but also with incomplete understanding of dialysis in decision making [8,34]. Subjects in our study were referred early to nephrologists, with a median interval of 192 days between enrollment and dialysis. This characteristic eliminated selection bias because patients were placed under nephrologists' care with the intent of initiating dialysis.

Table 4 Per-person and total costs attributable to advanced chronic kidney disease

Parameters	All patients	Not receiving chronic dialysis	Receiving chronic dialysis during follow-up period
Number of patients	8,341	2,049	6,292
Age at time of enrollment, mean (SD), years	79.4 (7.0)	82.0 (6.4)	78.6 (7.1)
Total follow-up period, median (IQR), days	759 (302 to 1537)	279 (124 to 613)	985 (475 to 1758)
Day of chronic dialysis initiation, median (IQR)	-	-	192 (90 to 404)
Age at chronic dialysis initiation, mean (SD), years	-	-	79.4 (7.1)
Costs during total follow-up period, \$US/person-years (95% CI)			
Total healthcare cost	22,583 (22,159 to 23,007)	18,252 (16,990 to 19,514)	23,994 (23,616 to 24,371)
Total hospital cost	10,980 (10,553 to 11,407)	14,448 (13,230 to 15,665)	9,850 (9,450 to 10,251)
Total ambulatory visit cost	11,603 (11,444 to 11,763)	3,805 (3,646 to 3,963)	14,143 (13,982 to 14,305)
Costs before chronic dialysis, \$US/person-years (95% CI)			
Total healthcare cost	-	-	8,738 (8,409 to 9,068)
Total hospital cost	-	-	4,450 (4,158 to 4,743)
Total ambulatory visit cost	-	-	4,287 (4,199 to 4,377)
Costs after chronic dialysis initiation, \$US/person-years (95% CI)			
Total healthcare cost	-	-	42,980 (37,050 to 48,909)
Total hospital cost	-	-	24,333 (18,460 to 30,208)
Total ambulatory visit cost	-	-	18,646 (18,309 to 18,984)

CI, confidence interval; IQR, interquartile range; SD, standard deviation.

Our results provide objective information to facilitate dialysis decision making among older patients, their family members and physicians; however, some limitations of this study should be acknowledged. First, decisions about whether to receive dialysis may depend on patient preferences (family support, financial constraints and the will to live), but examination of this complex process was beyond the scope of the current study. Second, information on several potential confounding factors, including obesity, nutritional condition, psychosocial function, performance status and indication for dialysis initiation, were not available in the NHIRD database. Estimated GFRs at dialysis initiation were also not recorded, whereas the Initiating Dialysis Early and Late (IDEAL) Study showed no significant survival difference between early initiation (estimated GFRs: 10.0 to 14.0 ml per minute) and late initiation (estimated GFRs: 5.0 to 7.0 ml per minute) of dialysis [40]. Third, our study was subject to the inherent limitations of its retrospective and observational design. However, randomized control trials investigating this issue are not possible because of the potential of violating medical ethics. Finally, we based the diagnosis of advanced CKD on ESA prescriptions, resulting in the exclusion of patients with advanced CKD who never received such prescriptions (that is, those who had no obvious renal anemia). The results of the study cannot be generalized to all older patients with advanced CKD.

Conclusions

Our findings provide novel information useful for the re-consideration of dialysis decision making that accounts for risk-benefit relationships in older patients with advanced CKD. For frail older patients with late-stage CKD, conservative care may be a viable treatment option, as dialysis may not prolong life expectancy.

Additional file

Additional file 1: Table S1. Baseline characteristics of patients after propensity-score matching. **Table S2.** Crude and adjusted hazard ratios for mortality according to receiving chronic dialysis during the follow-up period. **Table S3.** Per person and total costs attributable to advanced chronic kidney disease.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

DCT had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis. Study concept and design: CJS, YTC, SMO, DCT. Acquisition of data: CJS, YTC, SCK, DCT. Analysis and interpretation of data: CJS, YTC, SMO, DCT. Drafting of the manuscript: CJS, YTC, SMO, DCT. Statistical analysis: YTC, SCK, WCY. Obtained funding: DCT. Administrative, technical, or material support: SMO, SCK, WCY, DCT. Study supervision: DCT. All authors read and approved the final manuscript.

Acknowledgements

This study was supported by grants to Dr. D.C. Targ from the National Science Council (NSC 99-2314-B-010-002-MY3 and NSC 102-2314-B-010-004-MY3), Taipei Veterans General Hospital (V102 C-129), and Ministry of Education's aim for the

Top University Plan. This study was based in part on data from the NHIRD provided by the Bureau of NHI, Department of Health and managed by National Health Research Institutes. The interpretation and conclusions contained herein do not represent those of Bureau of NHI, Department of Health, and National Health Research Institutes.

Author details

¹Department of Medicine, Taipei Veterans General Hospital, Yuanshan Branch, Yilan, Taiwan. ²School of Medicine, National Yang-Ming University, Taipei, Taiwan. ³Department of Medicine, Taipei City Hospital Heping Fuyou Branch, Taipei, Taiwan. ⁴Division of Nephrology, Department of Medicine, Taipei Veterans General Hospital, No. 201, Section 2, Shih-Pai Road, Taipei 11217, Taiwan. ⁵Institutes of Physiology and Clinical Medicine, National Yang-Ming University, Taipei, Taiwan. ⁶National Institute of Infectious Diseases and Vaccinology, National Health Research Institutes, Miaoli County, Taiwan. ⁷Division of Infectious Diseases, Taipei Veterans General Hospital, Taipei, Taiwan.

Received: 27 July 2014 Accepted: 1 September 2014

Published online: 06 October 2014

References

1. Chu VH, Sexton DJ, Cabell CH, Reller LB, Pappas PA, Singh RK, Fowler VG Jr, Corey GR, Aksoy O, Woods CW: **Repeat infective endocarditis: differentiating relapse from reinfection.** *Clin Infect Dis* 2005, **41**:406–409.
2. Kurella M, Covinsky KE, Collins AJ, Chertow GM: **Octogenarians and nonagenarians starting dialysis in the United States.** *Ann Intern Med* 2007, **146**:177–183.
3. Hemmelgarn BR, James MT, Manns BJ, O'Hare AM, Muntner P, Ravani P, Quinn RR, Turin TC, Tan Z, Tonelli M, Alberta Kidney Disease Network: **Rates of treated and untreated kidney failure in older vs younger adults.** *JAMA* 2012, **307**:2507–2515.
4. Morton RL, Turner RM, Howard K, Snelling P, Webster AC: **Patients who plan for conservative care rather than dialysis: a national observational study in Australia.** *Am J Kidney Dis* 2012, **59**:419–427.
5. Smith AK, Walter LC, Miao Y, Boscardin WJ, Covinsky KE: **Disability during the last two years of life.** *JAMA Intern Med* 2013, **173**:1506–1513.
6. Morton RL, Snelling P, Webster AC, Rose J, Masterson R, Johnson DW, Howard K: **Factors influencing patient choice of dialysis versus conservative care to treat end-stage kidney disease.** *CMAJ* 2012, **184**:E277–E283.
7. Moss AH: **Revised dialysis clinical practice guideline promotes more informed decision-making.** *Clin J Am Soc Nephrol* 2010, **5**:2380–2383.
8. Kawaguchi T, Karoboyas A, Robinson BM, Li Y, Fukuhara S, Bieber BA, Rayner HC, Andreucci VE, Pisoni RL, Port FK, Morgenstern H, Akizawa T, Saran R: **Associations of frequency and duration of patient-doctor contact in hemodialysis facilities with mortality.** *J Am Soc Nephrol* 2013, **24**:1493–1502.
9. O'Hare AM, Rodriguez RA, Hailpern SM, Larson EB, Kurella Tamura M: **Regional variation in health care intensity and treatment practices for end-stage renal disease in older adults.** *JAMA* 2010, **304**:180–186.
10. Chandna SM, Da Silva-Gane M, Marshall C, Warwicker P, Greenwood RN, Farrington K: **Survival of elderly patients with stage 5 CKD: comparison of conservative management and renal replacement therapy.** *Nephrol Dial Transplant* 2011, **26**:1608–1614.
11. Murtagh FE, Marsh JE, Donohoe P, Ekbal NJ, Sheerin NS, Harris FE: **Dialysis or not? A comparative survival study of patients over 75 years with chronic kidney disease stage 5.** *Nephrol Dial Transplant* 2007, **22**:1955–1962.
12. Smith C, Da Silva-Gane M, Chandna S, Warwicker P, Greenwood R, Farrington K: **Choosing not to dialyze: evaluation of planned non-dialytic management in a cohort of patients with end-stage renal failure.** *Nephron Clin Pract* 2003, **95**:c40–c46.
13. Wu VC, Wu CH, Huang TM, Wang CY, Lai CF, Shiao CC, Chang CH, Lin SL, Chen YY, Chen YM, Chu TS, Chiang WC, Wu KD, Tsai PR, Chen L, Ko WJ, NSARF Group: **Long-term risk of coronary events after AKI.** *J Am Soc Nephrol* 2014, **25**:595–605.
14. Wu IW, Wang SY, Hsu KH, Lee CC, Sun CY, Tsai CJ, Wu MS: **Multidisciplinary predialysis education decreases the incidence of dialysis and reduces mortality—a controlled cohort study based on the NKF/DOQI guidelines.** *Nephrol Dial Transplant* 2009, **24**:3426–3433.
15. Chen YR, Yang Y, Wang SC, Chiu PF, Chou WY, Lin CY, Chang JM, Chen TW, Ferng SH, Lin CL: **Effectiveness of multidisciplinary care for chronic kidney disease in Taiwan: a 3-year prospective cohort study.** *Nephrol Dial Transplant* 2013, **28**:671–682.
16. Hwang SJ, Tsai JC, Chen HC: **Epidemiology, impact and preventive care of chronic kidney disease in Taiwan.** *Nephrology (Carlton)* 2010, **15**:3–9.
17. Wu CY, Chan FK, Wu MS, Kuo KN, Wang CB, Tsao CR, Lin JT: **Histamine2-receptor antagonists are an alternative to proton pump inhibitor in patients receiving clopidogrel.** *Gastroenterology* 2010, **139**:1165–1171.
18. Cheng CL, Kao YH, Lin SJ, Lee CH, Lai ML: **Validation of the National Health Insurance Research Database with ischemic stroke cases in Taiwan.** *Pharmacoepidemiol Drug Saf* 2011, **20**:236–242.
19. Lin CC, Lai MS, Syu CY, Chang SC, Tseng FY: **Accuracy of diabetes diagnosis in health insurance claims data in Taiwan.** *J Formos Med Assoc* 2005, **104**:157–163.
20. Brophy DF, Daniel G, Gitlin M, Mayne TJ: **Characterizing hospitalizations of end-stage renal disease patients on dialysis and inpatient utilization of erythropoiesis-stimulating agent therapy.** *Ann Pharmacother* 2010, **44**:43–49.
21. Maddux FW, Shetty S, del Aguila MA, Nelson MA, Murray BM: **Effect of erythropoiesis-stimulating agents on healthcare utilization, costs, and outcomes in chronic kidney disease.** *Ann Pharmacother* 2007, **41**:1761–1769.
22. Hsu TW, Liu JS, Hung SC, Kuo KL, Chang YK, Chen YC, Hsu CC, Tarng DC: **Renoprotective effect of renin-angiotensin-aldosterone system blockade in patients with predialysis advanced chronic kidney disease, hypertension, and anemia.** *JAMA Intern Med* 2014, **174**:347–354.
23. Hwang SJ, Yang WC, Lin MY, Mau LW, Chen HC, Taiwan Society of Nephrology: **Impact of the clinical conditions at dialysis initiation on mortality in incident haemodialysis patients: a national cohort study in Taiwan.** *Nephrol Dial Transplant* 2010, **25**:2616–2624.
24. Chao PW, Ou SM, Chen YT, Lee YJ, Wang FM, Liu CJ, Yang WC, Chen TJ, Chen TW, Li SY: **Acute appendicitis in patients with end-stage renal disease.** *J Gastrointest Surg* 2012, **16**:1940–1946.
25. Charlson ME, Pompei P, Ales KL, MacKenzie CR: **A new method of classifying prognostic comorbidity in longitudinal studies: development and validation.** *J Chronic Dis* 1987, **40**:373–383.
26. O'Hare AM, Choi AI, Bertenthal D, Bacchetti P, Garg AX, Kaufman JS, Walter LC, Mehta KM, Steinman MA, Allon M, McClellan WM, Landefeld CS: **Age affects outcomes in chronic kidney disease.** *J Am Soc Nephrol* 2007, **18**:2758–2765.
27. Lamping DL, Constantinovici N, Roderick P, Normand C, Henderson L, Harris S, Brown E, Gruen R, Victor C: **Clinical outcomes, quality of life, and costs in the North Thames Dialysis Study of elderly people on dialysis: a prospective cohort study.** *Lancet* 2000, **356**:1543–1550.
28. Chandna SM, Schulz J, Lawrence C, Greenwood RN, Farrington K: **Is there a rationale for rationing chronic dialysis? A hospital based cohort study of factors affecting survival and morbidity.** *BMJ* 1999, **318**:217–223.
29. Holley JL, Hines SC, Glover JJ, Babrow AS, Badzek LA, Moss AH: **Failure of advance care planning to elicit patients' preferences for withdrawal from dialysis.** *Am J Kidney Dis* 1999, **33**:688–693.
30. Yang WC, Hwang SJ, Taiwan Society of Nephrology: **Incidence, prevalence and mortality trends of dialysis end-stage renal disease in Taiwan from 1990 to 2001: the impact of national health insurance.** *Nephrol Dial Transplant* 2008, **23**:3977–3982.
31. Wu PH, Lin YT, Lee TC, Lin MY, Kuo MC, Chiu YW, Hwang SJ, Chen HC: **Predicting mortality of incident dialysis patients in Taiwan - a longitudinal population-based study.** *PLoS One* 2013, **8**:e61930.
32. Kurella Tamura M, Covinsky KE, Chertow GM, Yaffe K, Landefeld CS, McCulloch CE: **Functional status of elderly adults before and after initiation of dialysis.** *N Engl J Med* 2009, **361**:1539–1547.
33. Lin YT, Wu PH, Kuo MC, Lin MY, Lee TC, Chiu YW, Hwang SJ, Chen HC: **High cost and low survival rate in high comorbidity incident elderly hemodialysis patients.** *PLoS One* 2013, **8**:e75318.
34. Joly D, Anglicheau D, Alberti C, Nguyen AT, Touam M, Grunfeld JP, Jungers P: **Octogenarians reaching end-stage renal disease: cohort study of decision-making and clinical outcomes.** *J Am Soc Nephrol* 2003, **14**:1012–1021.
35. Carson RC, Juszczak M, Davenport A, Burns A: **Is maximum conservative management an equivalent treatment option to dialysis for elderly**

- patients with significant comorbid disease? *Clin J Am Soc Nephrol* 2009, **4**:1611–1619.
36. Murtaugh PA, Dickson ER, Van Dam GM, Malinchoc M, Grambsch PM, Langworthy AL, Gips CH: **Primary biliary cirrhosis: prediction of short-term survival based on repeated patient visits.** *Hepatology* 1994, **20**:126–134.
 37. Christensen E, Schlichting P, Andersen PK, Fauerholdt L, Schou G, Pedersen BV, Juhl E, Poulsen H, Tygstrup N: **Updating prognosis and therapeutic effect evaluation in cirrhosis with Cox's multiple regression model for time-dependent variables.** *Scand J Gastroenterol* 1986, **21**:163–174.
 38. Altman DG, De Stavola BL: **Practical problems in fitting a proportional hazards model to data with updated measurements of the covariates.** *Stat Med* 1994, **13**:301–341.
 39. Peters R, Beckett N, McCormack T, Fagard R, Fletcher A, Bulpitt C: **Treating hypertension in the very elderly: benefits, risks, and future directions, a focus on the hypertension in the very elderly trial.** *Eur Heart J* 2014, **35**:1712–1718.
 40. Cooper BA, Branley P, Bulfone L, Collins JF, Craig JC, Fraenkel MB, Harris A, Johnson DW, Kesselhut J, Li JJ, Li JJ, Luxton G, Pilmore A, Tiller DJ, Harris DC, Pollock CA, IDEAL Study: **A randomized, controlled trial of early versus late initiation of dialysis.** *N Engl J Med* 2010, **363**:609–619.

doi:10.1186/s12916-014-0169-3

Cite this article as: Shih et al.: The impact of dialysis therapy on older patients with advanced chronic kidney disease: a nationwide population-based study. *BMC Medicine* 2014 **12**:169.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

