

Identification of antibiotic resistance genes in the multidrug-resistant *Acinetobacter baumannii* strain, MDR-SHH02, using whole-genome sequencing

HUALIANG WANG^{1*}, JINGHUA WANG^{2*}, PEIJUAN YU³, PING GE², YANQUN JIANG⁴,
RONG XU², RONG CHEN² and XUEJIE LIU²

Departments of ¹Molecular Biology Laboratory and ²Microbiology Laboratory, Shanghai Centre for Clinical Laboratory, Shanghai 200126; ³Department of Clinical Laboratory, The Second Affiliated Hospital of Soochow University, Suzhou, Jiangsu 215004; ⁴Department of Clinical Laboratory, The Sixth People's Hospital Affiliated to Shanghai Jiaotong University School of Medicine, Shanghai 200233, P.R. China

Received January 5, 2016; Accepted December 7, 2016

DOI: 10.3892/ijmm.2016.2844

Abstract. This study aimed to investigate antibiotic resistance genes in the multidrug-resistant (MDR) *Acinetobacter baumannii* (*A. baumannii*) strain, MDR-SHH02, using whole-genome sequencing (WGS). The antibiotic resistance of MDR-SHH02 isolated from a patient with breast cancer to 19 types of antibiotics was determined using the Kirby-Bauer method. WGS of MDR-SHH02 was then performed. Following quality control and transcriptome assembly, functional annotation of genes was conducted, and the phylogenetic tree of MDR-SHH02, along with another 5 *A. baumannii* species and 2 *Acinetobacter* species, was constructed using PHYLIP 3.695 and FigTree v1.4.2. Furthermore, pathogenicity islands (PAIs) were predicted by the pathogenicity island database. Potential antibiotic resistance genes in MDR-SHH02 were predicted based on the information in the Antibiotic Resistance Genes Database (ARDB). MDR-SHH02 was found to be resistant to all of the tested antibiotics. The total draft genome length of MDR-SHH02 was 4,003,808 bp. There were 74.25% of coding sequences to be annotated into 21 of the Clusters of Orthologous Groups (COGs) of protein terms, such as 'transcription' and 'amino acid transport and metabolism'. Furthermore, there were 45 PAIs homologous to the sequence MDRSHH02000806. Additionally, a total of 12 gene sequences in MDR-SHH02 were highly similar to the sequences of antibiotic resistance genes in ARDB, including genes encoding aminoglycoside-modifying enzymes [e.g., *aac(3)-Ia*, *ant(2'')-Ia*, *aph33ib* and *aph(3')-Ia*],

β -lactamase genes (*bl2b_tem* and *bl2b_tem1*), sulfonamide-resistant dihydropteroate synthase genes (*sul1* and *sul2*), *catb3* and *tetb*. These results suggest that numerous genes mediate resistance to various antibiotics in MDR-SHH02, and provide a clinical guidance for the personalized therapy of *A. baumannii*-infected patients.

Introduction

Acinetobacter baumannii (*A. baumannii*) is a notable pathogen that causes hospital-acquired infections among immune-compromised patients, accounting for 5% of Gram-negative infections (1). Due to a strong resistance to desiccation and multiple antibacterial agents, the widespread dissemination of multidrug-resistant (MDR) *A. baumannii* strains has been a threat to hospitalized patients in recent years (2).

Antibiotic resistance determinants play pivotal roles in whether or not *A. baumannii* will flourish in the host (3). For instance, the expression of β -lactamase genes [e.g., oxacillinase (OXA)-235 gene, *bla*_{OXA-51} and *bla*_{TEM-1}] has been shown to be involved in antibiotic resistance (4-6). An 86-kb region AbaR resistant to heavy metal and antibiotics has been found in a MDR isolate AYE (7), indicating the important role of AbaR in the spread of *A. baumannii* in hospitals (8,9). Furthermore, other resistance determinants, such as macrolide (*msrA/msrB*), aminoglycoside (e.g., *aacC1*, *armA*, and *aphA1*) and tetracycline [e.g., tet (39), tet (A), and tet (B)] have been identified in various *A. baumannii* isolates (10).

Bacterial whole-genome sequencing (WGS) has enhanced our ability to evaluate antibiotic resistance determinants. For example, a AbaR-type genomic resistance island, AbaR22, has been identified in the MDR *A. baumannii* strain, MDR-ZJ06, via WGS (11). A whole-genome comparison detected 18 putative single nucleotide polymorphisms (SNPs) between 2 pre- and post-therapy *A. baumannii* isolates (12). Furthermore, 10 types of AbaR resistance islands were identified in 2 *A. baumannii* isolates using WGS (13). Despite increased research on antibiotic resistance determinants in *A. baumannii*, the molecular mechanisms of antibiotic resistance in MDR *A. baumannii* are

Correspondence to: Dr Hualiang Wang, Department of Molecular Biology Laboratory, Shanghai Center for Clinical Laboratory, 528 Hongshan Road, Pudong New Area, Shanghai 200126, P.R. China
E-mail: wanghuanliang@hotmail.com

*Contributed equally

Key words: *Acinetobacter baumannii*, antibiotic resistance gene, whole-genome sequencing

not yet fully understood, and various antibiotic resistance genes have not been detected.

In the present study, we applied WGS to obtain the whole genomic sequence of the MDR *A. baumannii* strain, MDR-SHH02, isolated from a patient with breast cancer. Furthermore, the antibiotic resistance of MDR-SHH02 to multiple antibiotics was determined, and potential antibiotic resistance genes in MDR-SHH02 were predicted. The results of our study may enhance our understanding of the molecular mechanisms of antibiotic resistance in MDR *A. baumannii*, and provide a clinical guidance for the therapy of *A. baumannii*-infected patients.

Materials and methods

Isolation and identification of *A. baumannii* strain. The clinical MDR *A. baumannii* strain, named MDR-SHH02, was isolated from the blood obtained from a 65-year-old woman with terminal-stage breast cancer at Shanghai Sixth People's Hospital Affiliated to Shanghai Jiaotong University, Shanghai, China. This patient had received a double mastectomy and nearby lymph node excision. After being discharged from the hospital, this patient was hospitalized again due to symptoms of fever, cough (lasting for days) and shortness of breath. During her second hospital administration, she received several antimicrobial therapies, including maxipime, impenem, methylprednisolone, levofloxacin and cefoperazone-sulbactam sodium. *A. baumannii* was positive in the blood culture and sputum culture. The results of the antimicrobial susceptibility test revealed that the *A. baumannii* strain was resistant to multiple commonly used antibiotics. This study was approved by the Shanghai Health and Family Planning Commission Foundation (Shanghai, China), and informed consent was obtained from the patient.

The isolated strain was inoculated onto blood agar plates and then incubated in an atmosphere of 5% CO₂ at 35°C for 48 h. Afterwards, this strain was identified using morphological and biochemical tests according to standard methods (14). Colonies with typical morphological and biochemical characteristics of *Acinetobacter* were cultivated on 5% sheep blood agar and identified using an automated Microscan® system (Dade Behring, Inc., West Sacramento, CA, USA). The *A. baumannii* strain was stored at -70°C in skim milk for further analyses.

Antibiotic resistance test for *A. baumannii* MDR-SHH02. According to the Clinical and Laboratory Standards Institute (CLSI) guidelines (15), disc diffusion assay (DDA) with dry wafers saturated by 19 types of antibiotics, including gentamicin (10 µg/wafer), tobramycin (30 µg/wafer), amikacin (30 µg/wafer), ampicillin-sulbactam (10/10 µg/wafer), ceftazidime (30 µg/wafer), ciprofloxacin (5 µg/wafer), levofloxacin (5 µg/wafer), imipenem (10 µg/wafer), meropenem (10 µg/wafer), piperacillin/tazobactam (100/10 µg/wafer), ticarcillin/clavulanic acid (75/10 µg/wafer), cefepime (30 µg/wafer), cefotaxime (30 µg/wafer), ceftriaxone (30 µg/wafer), doxycycline (30 µg/wafer), minocycline (30 µg/wafer), tetracycline (30 µg/wafer), piperacillin (100 µg/wafer) and trimethoprim-sulfamethoxazole (1.25/23.75 µg/wafer) (all from Oxoid, Ltd., Basingstoke, UK), were carried out using the Kirby-Bauer (KB) method, as previously described (16). Briefly, Mueller-Hinton agar (Oxoid, Ltd.) plates were overlaid with the inocula of the

clinical *A. baumannii* strain, and the turbidity of the inocula was equivalent to the 0.5 McFarland standard. Subsequently, dry wafers saturated by antibiotics were placed on the surface of the agar, and plates were placed in an atmosphere of 5% CO₂ at 35°C. Following 24 h of culture, the diameter of the inhibition zone around each wafer was measured according to the CLSI criteria (15). In this test, *Escherichia coli* ATCC 25922, ATCC 35218 and *Pseudomonas aeruginosa* ATCC 27853 obtained from the Clinical Laboratory Center of the Ministry of Health were used as reference strains.

DNA preparation, library construction and sequencing. The genomic DNA of *A. baumannii* MDR-SHH02 was extracted using a bacterial genomic DNA purification kit (Tiangen Biotech Co., Ltd., Beijing, China) according to the manufacturer's instructions. The Illumina sequencing library was then prepared using the Nextera™ DNA Sample Preparation kit (Illumina®-Compatible). Paired-end dual index 2x90 bp sequencing was fulfilled following the Illumina HiSeq 2000. Sequencing was performed by Beijing Genomics Institute (BGI; Shenzhen, China). The sequencing data were uploaded to the public database the National Center for Biotechnology Information (NCBI; <http://www.ncbi.nlm.nih.gov/>) under the BioProject PRJNA256112 with BioSample accession no. SAMN02991371.

Quality control. For the raw sequencing data, the reads were cleaned by removing the empty reads, adapter sequences and reads with n≥10% using the SeqPrep program (<https://github.com/jstjohn/SeqPrep>) and Sickle (<https://github.com/najoshi/sickle>). In addition, the reads were trimmed by discarding the reads containing >30% bases with a Q-value ≤20 in the 3' terminal, as well as reads with adaptor sequences (the length of overlapping sequences between adaptor and read was at least >15 bp, and the number of mismatch bases was <3 bp).

Genome assembly. The clean reads were assembled using the short oligonucleotide analysis package SOAPdenovo (version 2.04; <http://soap.genomics.org.cn/>). To determine whether the GC content has a significant effect on sequencing randomness or not, the GC content and average depth of the genomic sequence were calculated without repetition as a unit of 500 bp.

Genome annotation. Genes in the assembled genomic sequence were predicted using Glimmer 3.0 (<http://www.cbc.edu/software/glimmer/>) (17), which is a system for identifying genes in DNA sequences of microorganisms, particularly bacteria, archaea and viruses.

Furthermore, tRNA and rRNA (5S, 16S, and 23S rRNA) in the genomic sequence were searched using tRNAscan-SE (<http://lowelab.ucsc.edu/tRNAscan-SE/>) (18) and RNAmmer 1.2 (<http://www.cbs.dtu.dk/services/RNAmmer/>) (19), respectively.

Additionally, tandem repeat sequences and clustered regularly interspaced short palindromic repeats (CRISPR) in the genomic sequence were predicted using the Tandem Repeat Finder (<http://tandem.bu.edu/trf/trf.html>) and CRISPR Finder (<http://crispr.u-psud.fr/Server/>) software, respectively.

Insertion sequences (ISs) were characterized using the IS Finder database (<https://www-is.biotoul.fr/>), and the parameter *-e* was set as $1e-5$, identity set as 35%. Besides, protein domains associated with the genomic sequence were predicted using the InterPro database (<https://www.ebi.ac.uk/interpro/>), and the parameter was set as *-appl PfamA*.

Functional annotation of genes. Sequence alignment of the amino acid sequences of genes to the Cluster of Orthologous Groups (COGs) of proteins database (<http://www.ncbi.nlm.nih.gov/COG/>) (20) was performed using the Basic Local Alignment Search Tool (BLASTP; version 2.0) program from NCBI (E-value $\leq 10^{-4}$) (21). We also performed sequence alignment of the amino acid sequences of genes to the NCBI non-redundant (NR) database (E-value $\leq 10^{-10}$, identity score $\geq 35\%$, and coverage length $\geq 80\%$). If the amino acid sequence of a gene was aligned to multiple sequences in the databases, the optimal result was retained.

Construction of phylogenetic tree. Based on the NCBI 16S rRNA gene database, 16S rRNA gene sequences of 5 *A. baumannii* strains, including ATCC 17978, ATCC 19606, CIP 70.34, DSM 30007 and *A. baumannii* JCM 68415, as well as 2 species belonging to *Acinetobacter* (*A. haemolyticus* ATCC 17906 and *A. bereziniae* ATCC 17924), were used to construct the phylogenetic tree, along with MDR-SHH02. Briefly, multiple sequence alignment was performed using ClustalW-2.1 (22). Subsequently, the software package PHYLIP 3.695 (<http://evolution.genetics.washington.edu/phylip.html>), along with the bootstrap algorithm, was used to construct the maximum likelihood phylogenetic tree, and the phylogenetic tree was visualized by FigTree v1.4.2 (<http://tree.bio.ed.ac.uk/software/figtree/>).

Prediction of pathogenicity islands (PAIs). The Pathogenicity Island database (PAIDB; http://www.paidb.re.kr/about_paidb.php), which is a web-based user-friendly resource and widely used for detecting PAIs in newly sequenced genomes (23), was utilized to predict PAIs in the genomic sequence of MDR-SHH02.

Identification of antibiotic resistance genes. To identify potential antibiotic resistance genes in the genomic sequence of MDR-SHH02, sequence alignment of the protein sequences of antibiotic resistance genes in the Antibiotic Resistance Genes database (ARDB; <http://ardb.cbcb.umd.edu/>) (24) and MDR-SHH02 genomic sequence was conducted using BLASTP (E-value $\leq 10^{-10}$, identity score $\geq 90\%$, and coverage length $\geq 80\%$).

Results

Antibiotic-resistance of *A. baumannii* MDR-SHH02. The antibiotic-resistance assay revealed that the diameter of the inhibition zone for 17 types of antibiotics on MDR-SHH02 plates was 6 mm, apart from levofloxacin and minocycline (diameter, 10 mm) (Table I). According to the CLSI criteria, MDR-SHH02 was resistant to all of the tested antibiotics.

Assembly and annotation of the genomic sequence of *A. baumannii* MDR-SHH02. During the genome assembly,

Table I. The results of antibiotic-resistance assay for *A. baumannii* MDR-SHH02.

Antibiotic name	Diameter of inhibition zone on MDR-SHH02 plate (mm)	Antibiotic-resistance of MDR-SHH02
Ampicillin-sulbactam	6	R
Ceftazidime	6	R
Ciprofloxacin	6	R
Levofloxacin	10	R
Imipenem	6	R
Meropenem	6	R
Gentamicin	6	R
Tobramycin	6	R
Amikacin	6	R
Piperacillin/tazobactam	6	R
Ticarcillin/clavulanic acid	6	R
Cefepime	6	R
Cefotaxime	6	R
Ceftriaxone	6	R
Doxycycline	6	R
Minocycline	10	R
Tetracycline	6	R
Piperacillin	6	R
Trimethoprim-sulfamethoxazole	6	R

Diameter of inhibition zone < 15 mm is determined as drug-resistant (R); diameter of inhibition zone between 15 and 17 mm is determined as intermediate (I); and diameter of inhibition zone > 17 mm is determined as drug-sensitive (S). *A. baumannii*, *Acinetobacter baumannii*.

a total of 85 scaffolds were generated, and the scaffold N50 length was 131,822 bp. The total draft genome length of MDR-SHH02 was 4,003,808 bp, with 38.99% of GC content. There were 3,787 coding sequences, 62 tRNA sequences and 3 rRNA sequences in the genomic sequence. Moreover, 2 CRISPR and 36 tandem repeat sequences, as well as 29 ISs were predicted in the genomic sequence (Table II).

Furthermore, numerous protein domains were predicted in the genomic sequence of MDR-SHH02, such as the AcrB/AcrD/AcrF family (Table III).

Functional annotation of the genomic sequence of *A. baumannii* MDR-SHH02. According to the COG annotation, 74.25% (2,812/3,787) of coding sequences were annotated into 21 COG terms, which were divided into 3 categories: information storage of processing, cellular processes and signaling, and metabolism. Apart from the category of poorly characterized, most of the coding sequences were annotated into 'transcription' (number of coding sequences, 261) and 'amino acid transport and metabolism' (number of coding




Figure 1. The functional annotation of the genomic sequence of *Acinetobacter baumannii* (*A. baumannii*) MDR-SHH02 using the Cluster of Orthologous Groups (COGs) of proteins database. (A) The bar diagram displaying the COG categories. The x-coordinate represents the COG categories, and y-coordinate represents the number of matched genes; (B) the annotation circle displaying the distribution of COG categories, open reading frames (ORFs) and GC content in the genomic sequence of *A. baumannii* MDR-SHH02. The outermost circle (circle 1) displays the distribution of COG categories on positive strand, and each kind of color represents one COG term, which is annotated as that in (A). Circles 2-4 and 5-7 display the distribution of ORFs on positive and negative strands. Circle 8 displays the distribution of COG categories on negative strand. Circle 9 displays the GC content. Circle 10 displays the GC skew (G-C)/(G+C) (green bars represent positive values, purple bars represent negative values). Circle 11 displays the scale in kbp.

sequences, 250) (Fig. 1A). The distribution of COG categories in the genomic sequence of *A. baumannii* MDR-SHH02 is shown in Fig. 1B.

Analysis of the phylogenetic tree. Based on the 16S rRNA gene sequences of *A. baumannii* in NCBI, the phylogenetic

tree revealed that the genomic sequence of MDR-SHH02 was most similar to the 16S rRNA gene sequence of *A. baumannii* ATCC 17978 (Fig. 2).

Analysis of PAIs. During the process of bacterial infection, PAIs play pivotal roles in the evolution of pathogens and




Figure 2. Whole-genome phylogeny of the genomes of *Acinetobacter baumannii* (*A. baumannii*) MDR-SHH02 and 5 sequenced *A. baumannii* genomes. The phylogenetic tree was rooted with 2 genomes of *Acinetobacter* species.

Table II. The results of genome assembly and annotation of *A. baumannii* MDR-SHH02.

Feature	Statistics
Length of total draft genome length (bp)	4,003,808
No. of scaffolds	85
Length of scaffold N50 (bp)	131,822
GC content (%)	38.99
No. of coding sequences	3,787
No. of tRNAs	62
No. of rRNAs	3
No. of CRISPR	2
No. of tandem repeat sequences	36
No. of insertion sequences	29

A. baumannii, *Acinetobacter baumannii*; CRISPR, clustered regularly interspaced short palindromic repeats.

the development of diseases. In the genomic sequence of MDR-SHH02, a total of 45 PAIs homologous to the sequence MDRSHH02000806 were detected (Table IV). Most of the PAIs were previously identified from *Escherichia coli* [e.g., locus of enterocyte effacement (LEE)] and *Pseudomonas* (e.g., PAPI-1 and T-PAI).

Screening of antibiotic resistance genes. To reveal the genes relevant to the antibiotic resistance of MDR-SHH02, sequence alignment of the protein sequences of antibiotic resistance genes in ARDB and MDR-SHH02 genomic sequence was performed. Based on the selection criteria, a total of 12 gene

Table III. The top 10 predicted protein domains in the genomic sequence of *A. baumannii* MDR-SHH02.

Gene ID	Protein domain	P-value
MDRSHH02002231	AcrB/AcrD/AcrF family	<1.0E-300
MDRSHH02000985	Monomeric isocitrate dehydrogenase	<1.0E-300
MDRSHH02003056	AcrB/AcrD/AcrF family	<1.0E-300
MDRSHH02000840	AcrB/AcrD/AcrF family	<1.0E-300
MDRSHH02003594	AcrB/AcrD/AcrF family	<1.0E-300
MDRSHH02000667	AcrB/AcrD/AcrF family	<1.0E-300
MDRSHH02000153	Phosphoenolpyruvate carboxylase	2.70E-292
MDRSHH02000199	Urocanase	5.20E-287
MDRSHH02000831	Phosphoenolpyruvate carboxykinase	2.70E-279
MDRSHH02003134	AcrB/AcrD/AcrF family	5.50E-266

A. baumannii, *Acinetobacter baumannii*.

sequences (e.g., MDRSHH02002408, MDRSHH02000600 and MDRSHH02000597) in MDR-SHH02 were highly similar

Table IV. Pathogenicity islands homologous to a region in the genomic sequence of *A. baumannii* MDR-SHH02.

Gene ID	Start	End	Size (bp)	no. of ORFs	PAIs homologous to this region
MDRSHH02000806	332	21071	20740	18	PAPI-1 (<i>Pseudomonas aeruginosa</i> PA14) PAPI-2(C) (<i>Pseudomonas aeruginosa</i> C) PAPI-3(SG) (<i>Pseudomonas aeruginosa</i> SG17M) PAPI-5 (<i>Pseudomonas aeruginosa</i> PSE9) PPHGI-1 (<i>Pseudomonas syringae</i> pv. phaseolicola 1302A) SPI-7 (<i>Salmonella enterica</i> subsp. <i>enterica</i> serovar Typhi str. CT18) SPI-7 (<i>Salmonella enterica</i> subsp. <i>enterica</i> serovar Typhi Ty2) AbaR25 (<i>Acinetobacter baumannii</i> BJAB07104) AbaR26 (<i>Acinetobacter baumannii</i> BJAB0868) Hrp PAI (<i>Erwinia amylovora</i> 321) S-PAI (<i>Pseudomonas cichorii</i> 83-1) Hrp PAI (<i>Pseudomonas syringae</i> pv. tomato DC3000) Hrp PAI (<i>Pseudomonas syringae</i> pv. tomato str. DC3000) T-PAI (<i>Pseudomonas viridiflava</i> LP23.1a) T-PAI (<i>Pseudomonas viridiflava</i> PNA3.3a) S-PAI (<i>Pseudomonas viridiflava</i> RMX23.1a) S-PAI (<i>Pseudomonas viridiflava</i> ME3.1b) S-PAI (<i>Pseudomonas viridiflava</i> RMX3.1b) Not named (<i>Corynebacterium urealyticum</i> DSM 7109) LEE (<i>Citrobacter rodentium</i> DBS100) PAI I 536 (<i>Escherichia coli</i> 536) LEE (<i>Escherichia coli</i> E2348/69) LEE (<i>Escherichia coli</i> O157:H7 str. EDL933 ATCC43895) LEE (<i>Escherichia coli</i> 71074) LEE (<i>Escherichia coli</i> 83/39) LEE (<i>Escherichia coli</i> REPEC 84/110-1) LEE (<i>Escherichia coli</i> RW1374) LEE (<i>Escherichia coli</i> RDEC-1) LEE (<i>Escherichia coli</i> O157:H7 EDL933) LEE (<i>Escherichia coli</i> O157:H7 str. Sakai) LEE (<i>Escherichia coli</i> O103:H2 str. 12009) LEE (<i>Escherichia coli</i> O26:H11 str. 11368) LEE (<i>Escherichia coli</i> O111:H- str. 11128) LEE II (<i>Escherichia coli</i> 413/89-1) AGI-3 (<i>Escherichia coli</i> BEN2908) TAI (<i>Escherichia coli</i> O157:H7 EDL933) OI-122 (<i>Escherichia coli</i> O157:H7 EDL933) PAI I CFT073 (<i>Escherichia coli</i> CFT073) Not named (<i>Escherichia coli</i> UMN026) SESS LEE (<i>Salmonella enterica</i> subsp. <i>salamae</i> serovar Sofia S1296) SESS LEE (<i>Salmonella enterica</i> subsp. <i>salamae</i> serovar Sofia S1635) SHI-1 (<i>Shigella flexneri</i> 2a str. 301) SHI-1 (<i>Shigella flexneri</i> 2a str. 2457T) SRL (<i>Shigella flexneri</i> 2a YSH6000) Not named (<i>Yersinia pestis</i> CO92)

A. baumannii, *Acinetobacter baumannii*; ORF, open reading frame; PAIs, pathogenicity islands.

to the sequences of antibiotic resistance genes in ARDB, such as *aac(3)-Ia*, *aac(6')-Ib*, *ant(2'')-Ia* and *aph(3')-Ia*. According

to the antibiotics that were resistant by the 12 gene sequences, MDR-SHH02 was resistant to multiple antibiotics, such as

Table V. The potential gene sequences in *A. baumannii* MDR-SHH02 relevant to antibiotic resistance.

Gene ID	Resistance gene type from ARDB	Resistance gene class	Antibiotics
MDRSHH02002408	<i>aac(3)-Ia</i>	Aminoglycoside N-acetyltransferase, which modifies aminoglycosides by acetylation	Astromicin, gentamicin , sisomicin
MDRSHH02000600	<i>aac(6')-Ib</i>	Aminoglycoside N-acetyltransferase, which modifies aminoglycosides by acetylation	Amikacin , dibekacin, isepamicin, netilmicin, sisomicin, tobramycin
MDRSHH02000597	<i>ant(2'')-Ia</i>	Aminoglycoside O-nucleotidyltransferase, which modifies aminoglycosides by adenylation	Dibekacin, gentamicin , kanamycin, sisomicin, tobramycin
MDRSHH02000611	<i>ant(3'')-Ia</i>	Aminoglycoside O-nucleotidyltransferase, which modifies aminoglycosides by adenylation	Spectinomycin, streptomycin
MDRSHH02001946	<i>aph33ib</i>	Aminoglycoside O-phosphotransferase, which modifies aminoglycosides by phosphorylation	Streptomycin
MDRSHH02002406	<i>aph(3')-Ia</i>	Aminoglycoside O-phosphotransferase, which modifies aminoglycosides by phosphorylation	Gentamicin b, kanamycin, neomycin, paromomycin, lividomycin, ribostamycin
MDRSHH02001945	<i>aph(6)-Id</i>	Aminoglycoside O-phosphotransferase, which modifies aminoglycosides by phosphorylation	streptomycin
MDRSHH02000608	<i>bl2b_tem, bl2b_tem1</i>	Class A β -lactamase. This enzyme breaks the β -lactam antibiotic ring open and deactivates the molecule's antibacterial properties	Cephalosporin, penicillin, cephalosporin i, cephalosporin ii
MDRSHH02000599	<i>catb3</i>	Group B chloramphenicol, acetyltransferase which can inactivate chloramphenicol. Also referred to as xenobiotic acetyltransferase	Chloramphenicol
MDRSHH02000610	<i>sul1</i>	Sulfonamide-resistant dihydropteroate synthase, which can not be inhibited by sulfonamide	Sulfonamide
MDRSHH02001738	<i>sul2</i>	Sulfonamide-resistant dihydropteroate synthase, which can not be inhibited by sulfonamide	Sulfonamide
MDRSHH02001941	<i>tetb</i>	Major facilitator superfamily transporter, tetracycline efflux pump	Tetracycline

A. baumannii, *Acinetobacter baumannii*; ARDB, Antibiotic Resistance Genes Database; DDA, disc diffusion assay. The antibiotics in bold are resisted by *A. baumannii* MDR-SHH02 in the DDA.

gentamicin, amikacin, tobramycin, spectinomycin, streptomycin and neomycin (Table V), which was partly consistent with the aforementioned results of antibiotic-resistance assay.

Discussion

In the present study, we reported the draft genomic sequence of the clinical MDR *A. baumannii* strain, MDR-SHH02, and predicted one gene sequence homologous to 45 PAIs and 12 potential gene sequences relevant to antibiotic resis-

tance. The antibiotic-resistance assay and the high similarity between the 12 gene sequences in MDR-SHH02 and the sequences of antibiotic resistance genes in ARDB, revealed that MDR-SHH02 was resistant to multiple antibiotics.

According to the prediction of PAIs, the gene sequence MDRSHH02000806 was homologous to 45 PAIs, such as LEE and PAPI-1. LEE PAIs were previously identified from multiple enteropathogenic *Escherichia coli* strains, and they are highly conserved in gene order and nucleotide sequence (25,26). PAPI-1 was previously identified from the *P. aeruginosa*

strain, PA14, and it contributes directly and synergistically along with PAPI-2 to the virulence of PA14 (27). Therefore, the virulence of MDR-SHH02 may be due to the presence of MDRSHH02000806 homologous to these PAIs.

In this study, we discovered a set of gene sequences that were highly similar to the sequences of multiple genes encoding aminoglycoside-modifying enzymes (AMEs), including 2 aminoglycoside N-acetyltransferase genes [*aac(3)-Ia* and *aac(6')-Ib*], 2 aminoglycoside *O*-nucleotidyltransferase genes [*ant(2'')-Ia* and *ant(3'')-Ia*], and 3 aminoglycoside *O*-phosphotransferase genes [*aph33ib*, *aph(3')-Ia* and *aph(6)-Id*]. The expression of AMEs enables bacteria to catalyze the modification of amino and hydroxyl groups on sugar moieties, such as aminoglycosides (28), which is a major cause of aminoglycoside resistance in many bacteria (29). The majority of aminoglycoside-resistant *Acinetobacter* isolates have the ability of enzymatic modification of aminoglycosides by acetyltransferases, nucleotidyltransferases and/or phosphotransferases (30). Previous studies have reported the prevalence of multiple AME genes [e.g., *aac(3)-Ia*, and *aac(6')-Ib*] in a set of *A. baumannii* isolates that are resistant to various aminoglycosides (e.g., amikacin, gentamicin and tobramycin) (31,32). Besides, *Aph(6)-Id* and *ant(3'')-Ia* have been detected in the *A. baumannii* strain, MRSN 12227, which is resistant to various antibiotics, such as amikacin, tobramycin and cefotaxime (33). Another study reported that *ant(2'')-Ia* present in a group of *A. baumannii* isolates (62.6%) was associated with resistance to the tested aminoglycosides (amikacin, tobramycin and gentamicin) (34). Furthermore, in this study, we found that MDR-SHH02 was resistant to 19 antibiotics, such as several types of aminoglycosides (amikacin, gentamicin and tobramycin), indicating that the resistance of MDR-SHH02 to aminoglycosides likely resulted from the coding sequences highly similar to AME genes. However, *aph33ib* has not been previously detected in *A. baumannii* isolates, and it is worthy of further study. For example, following knockout and complementation of the gene sequence that is highly similar to *aph33ib*, the resistance of MDR-SHH02 to aminoglycosides is determined.

In this study, several gene sequences in MDR-SHH02 had a high similarity to class A β -lactamase genes (*bl2b_tem* and *bl2b_tem1*), group B chloramphenicol acetyltransferase gene (*catb3*), sulfonamide-resistant dihydropteroate synthase genes (*sull* and *sul2*) and tetracycline efflux pump gene (*tetb*). The gene *bl2b_tem* has been detected in *Staphylococcus aureus* (35), and *bl2b_tem1* was detected in a series of marine bacteria, such as *Pelagibacter*, *Polaribacter* and *Roseobacter* (36). However, there is no evidence to support that *bl2b_tem* and *bl2b_tem1* are carried by *A. baumannii* isolates. All other genes (*catb3*, *sull*, *sul2* and *tetb*) have been found in *A. baumannii* isolates (37-40).

Despite the aforementioned results, there were still several limitations to this study. The association of MDRSHH02000806 with the virulence of MDR-SHH02 needs to be validated in further studies. Besides, the associations between the 12 gene sequences similar to AME genes and the antibiotic resistance of MDR-SHH02 are also needs to be confirmed in further studies. We aim to conduct such experiments in our future studies.

In conclusion, this study fulfilled the draft genomic sequence of the clinical MDR *A. baumannii* strain, MDR-SHH02, and 12 gene sequences in MDR-SHH02 had a highly similarity

to the sequences of genes encoding AMEs [e.g., *aac(3)-Ia*, *ant(2'')-Ia*, *aph33ib* and *aph(3'')-Ia*], β -lactamase genes (*bl2b_tem* and *bl2b_tem1*), sulfonamide-resistant dihydropteroate synthase genes (*sull* and *sul2*), *catb3* and *tetb*. Of these genes, *aph33ib*, *bl2b_tem* and *bl2b_tem1* were potential new antibiotic resistance genes. Furthermore, the antibiotic-resistance assay revealed that MDR-SHH02 was resistant to multiple antibiotics, such as amikacin, gentamicin and tobramycin. These findings were expected to enrich the data of antibiotic resistance genes in MDR *A. baumannii*, and provide a clinical guidance for the personalized therapy of *A. baumannii*-infected patients.

Acknowledgements

This study was supported by a project supported by Shanghai Health and Family Planning Commission Foundation, China (grant no. 20134010) and a project supported by the Natural Science Foundation of Shanghai, China (grant no. 15ZR1436100).

References

- Lockhart SR, Abramson MA, Beekmann SE, Gallagher G, Riedel S, Diekema DJ, Quinn JP and Doern GV: Antimicrobial resistance among Gram-negative bacilli causing infections in intensive care unit patients in the United States between 1993 and 2004. *J Clin Microbiol* 45: 3352-3359, 2007.
- Visca P, Seifert H and Towner KJ: *Acinetobacter* infection - an emerging threat to human health. *IUBMB Life* 63: 1048-1054, 2011.
- Mortensen BL and Skaar EP: Host-microbe interactions that shape the pathogenesis of *Acinetobacter baumannii* infection. *Cell Microbiol* 14: 1336-1344, 2012.
- Higgins PG, Pérez-Llarena FJ, Zander E, Fernández A, Bou G and Seifert H: OXA-235, a novel class D β -lactamase involved in resistance to carbapenems in *Acinetobacter baumannii*. *Antimicrob Agents Chemother* 57: 2121-2126, 2013.
- Zander E, Nemeč A, Seifert H and Higgins PG: Association between β -lactamase-encoding *bla*_(OXA-51) variants and DiversiLab rep-PCR-based typing of *Acinetobacter baumannii* isolates. *J Clin Microbiol* 50: 1900-1904, 2012.
- Krizova L, Poirel L, Nordmann P and Nemeč A: TEM-1 β -lactamase as a source of resistance to sulbactam in clinical strains of *Acinetobacter baumannii*. *J Antimicrob Chemother* 68: 2786-2791, 2013.
- Fournier PE, Vallenet D, Barbe V, Audic S, Ogata H, Poirel L, Riche H, Robert C, Mangenot S, Abergel C, *et al*: Comparative genomics of multidrug resistance in *Acinetobacter baumannii*. *PLoS Genet* 2: e7, 2006.
- Ramírez MS, Vilacoba E, Stietz MS, Merquier AK, JERIC P, Limansky AS, Márquez C, Bello H, Catalano M and Centron D: Spreading of AbaR-type genomic islands in multidrug resistance *Acinetobacter baumannii* strains belonging to different clonal complexes. *Curr Microbiol* 67: 9-14, 2013.
- Šepūtienė V, Povilonis J and Sužiedėlienė E: Novel variants of AbaR resistance islands with a common backbone in *Acinetobacter baumannii* isolates of European clone II. *Antimicrob Agents Chemother* 56: 1969-1973, 2012.
- Taitt CR, Leski T, Stockelman MG, Craft DW, Zurawski DV, Kirkup BC, and Vora GJ: Antimicrobial resistance determinants in *Acinetobacter baumannii* isolates taken from military treatment facilities. *Antimicrob Agents Chemother* 58: 767-781, 2014.
- Zhou H, Zhang T, Yu D, Pi B, Yang Q, Zhou J, Hu S and Yu Y: Genomic analysis of the multidrug-resistant *Acinetobacter baumannii* strain MDR-ZJ06 widely spread in China. *Antimicrob Agents Chemother* 55: 4506-4512, 2011.
- Hornsey M, Loman N, Wareham DW, Ellington MJ, Pallen MJ, Turton JF, Underwood A, Gaulton T, Thomas CP, Doumith M, *et al*: Whole-genome comparison of two *Acinetobacter baumannii* isolates from a single patient, where resistance developed during tigecycline therapy. *J Antimicrob Chemother* 66: 1499-1503, 2011.

13. Li H, Liu F, Zhang Y, Wang X, Zhao C, Chen H, Zhang F, Zhu B, Hu Y and Wang H: Evolution of carbapenem-resistant *Acinetobacter baumannii* revealed through whole-genome sequencing and comparative genomic analysis. *Antimicrob Agents Chemother* 59: 1168-1176, 2015.
14. Atlas RM, Brown AE and Parks LC: *Laboratory Manual of Experimental Microbiology*. Mosby, St. Louis, MO, pp119-127, 1995.
15. Watts JL: *Performance Standards For Antimicrobial Disk and Dilution Susceptibility Tests for Bacteria Isolated From Animals: Approved Standard*. Clinical and Laboratory Standards Institute, 2008.
16. Bauer AW, Kirby WM, Sherris JC and Turck M: Antibiotic susceptibility testing by a standardized single disk method. *Am J Clin Pathol* 45: 493-496, 1966.
17. Delcher AL, Bratke KA, Powers EC and Salzberg SL: Identifying bacterial genes and endosymbiont DNA with Glimmer. *Bioinformatics* 23: 673-679, 2007.
18. Schattner P, Brooks AN and Lowe TM: The tRNAscan-SE, snoscan and snoGPS web servers for the detection of tRNAs and snoRNAs. *Nucleic Acids Res* 33 (Web Server): W686-W689, 2005.
19. Lagesen K, Hallin P, Rødland EA, Staerfeldt H-H, Rognes T and Ussery DW: RNAmmer: consistent and rapid annotation of ribosomal RNA genes. *Nucleic Acids Res* 35: 3100-3108, 2007.
20. Tatusov RL, Fedorova ND, Jackson JD, Jacobs AR, Kiryutin B, Koonin EV, Krylov DM, Mazumder R, Mekhedov SL, Nikolskaya AN, *et al*: The COG database: an updated version includes eukaryotes. *BMC Bioinformatics* 4: 41, 2003.
21. Mount DW: Using the basic local alignment search tool (BLAST). *CSH Protoc* 2007: pdb. top17, 2007.
22. Chenna R, Sugawara H, Koike T, Lopez R, Gibson TJ, Higgins DG and Thompson JD: Multiple sequence alignment with the Clustal series of programs. *Nucleic Acids Res* 31: 3497-3500, 2003.
23. Yoon SH, Park Y-K, Lee S, Choi D, Oh TK, Hur C-G and Kim JF: Towards pathogenomics: a web-based resource for pathogenicity islands. *Nucleic Acids Res* 35 (Database): D395-D400, 2007.
24. Liu B and Pop M: ARDB - Antibiotic Resistance Genes Database. *Nucleic Acids Res* 37 (Database): D443-D447, 2009.
25. Pósfai G, Koob MD, Kirkpatrick HA and Blattner FR: Versatile insertion plasmids for targeted genome manipulations in bacteria: isolation, deletion, and rescue of the pathogenicity island LEE of the *Escherichia coli* O157:H7 genome. *J Bacteriol* 179: 4426-4428, 1997.
26. Tauschek M, Strugnell RA and Robins-Browne RM: Characterization and evidence of mobilization of the LEE pathogenicity island of rabbit-specific strains of enteropathogenic *Escherichia coli*. *Mol Microbiol* 44: 1533-1550, 2002.
27. Harrison EM, Carter ME, Luck S, Ou H-Y, He X, Deng Z, O'Callaghan C, Kadioglu A and Rajakumar K: Pathogenicity islands PAPI-1 and PAPI-2 contribute individually and synergistically to the virulence of *Pseudomonas aeruginosa* strain PA14. *Infect Immun* 78: 1437-1446, 2010.
28. Towner KJ: *Acinetobacter*: an old friend, but a new enemy. *J Hosp Infect* 73: 355-363, 2009.
29. Ramirez MS and Tolmasky ME: Aminoglycoside modifying enzymes. *Drug Resist Updat* 13: 151-171, 2010.
30. Nemeč A, Dolžani L, Brisse S, van den Broek P and Dijkshoorn L: Diversity of aminoglycoside-resistance genes and their association with class 1 integrons among strains of pan-European *Acinetobacter baumannii* clones. *J Med Microbiol* 53: 1233-1240, 2004.
31. Cho YJ, Moon DC, Jin JS, Choi CH, Lee YC and Lee JC: Genetic basis of resistance to aminoglycosides in *Acinetobacter* spp. and spread of *armA* in *Acinetobacter baumannii* sequence group 1 in Korean hospitals. *Diagn Microbiol Infect Dis* 64: 185-190, 2009.
32. Ploy MC, Denis F, Courvalin P and Lambert T: Molecular characterization of integrons in *Acinetobacter baumannii*: description of a hybrid class 2 integron. *Antimicrob Agents Chemother* 44: 2684-2688, 2000.
33. Waterman PE, McGann P, Snesrud E, Clifford RJ, Kwak YI, Munoz-Urbizo IP, Tabora-Castellanos J, Milillo M, Preston L, Aviles R, *et al*: Bacterial peritonitis due to *Acinetobacter baumannii* sequence type 25 with plasmid-borne new delhi metallo- β -lactamase in Honduras. *Antimicrob Agents Chemother* 57: 4584-4586, 2013.
34. Akers KS, Chaney C, Barsoumian A, Beckius M, Zera W, Yu X, Guymon C, Keen EF III, Robinson BJ, Mende K, *et al*: Aminoglycoside resistance and susceptibility testing errors in *Acinetobacter baumannii*-calcoaceticus complex. *J Clin Microbiol* 48: 1132-1138, 2010.
35. Kostić T, Ellis M, Williams MR, Stedtfeld TM, Kaneene JB, Stedtfeld RD and Hashsham SA: Thirty-minute screening of antibiotic resistance genes in bacterial isolates with minimal sample preparation in static self-dispensing 64 and 384 assay cards. *Appl Microbiol Biotechnol* 99: 7711-7722, 2015.
36. Hatosy SM and Martiny AC: The ocean as a global reservoir of antibiotic resistance genes. *Appl Environ Microbiol* 81: 7593-7599, 2015.
37. Chang-Tai Z, Yang L, Zhong-Yi H, Chang-Song Z, Yin-Ze K, Yong-Ping L and Chun-Lei D: High frequency of integrons related to drug-resistance in clinical isolates of *Acinetobacter baumannii*. *Indian J Med Microbiol* 29: 118-123 2011.
38. Martí S, Fernández-Cuenca F, Pascual A, Ribera A, Rodríguez-Baño J, Bou G, Miguel Cisneros J, Pachón J, Martínez-Martínez L and Vila J; Grupo de Estudio de Infección Hospitalaria (GEIH): Prevalence of the *tetA* and *tetB* genes as mechanisms of resistance to tetracycline and minocycline in *Acinetobacter baumannii* clinical isolates. *Enferm Infecc Microbiol Clin* 24: 77-80, 2006 (In Spanish).
39. Post V and Hall RM: AbaR5, a large multiple-antibiotic resistance region found in *Acinetobacter baumannii*. *Antimicrob Agents Chemother* 53: 2667-2671, 2009.
40. Ageršø Y and Petersen A: The tetracycline resistance determinant Tet 39 and the sulphonamide resistance gene *sulII* are common among resistant *Acinetobacter* spp. isolated from integrated fish farms in Thailand. *J Antimicrob Chemother* 59: 23-27, 2007.