

Exploring Myocardial Infarction Knowledge, Attitudes, and Practice Among Mogadishu's Somali Residents

Mohamed Omar Hassan ¹, Said Abdirahman Ahmed ¹, Mohamed Sheikh Hassan ²,
Mohamed Abdullahi Mohamud ¹, Ahmed Elmi Abdi¹

¹Department of Cardiology, Mogadishu Somali Turkish Training and Research Hospital, Mogadishu, Somalia; ²Department of Neurology, Mogadishu Somali Turkish Training and Research Hospital, Mogadishu, Somalia

Correspondence: Mohamed Omar Hassan, Email drmohamedyalax@gmail.com

Introduction: Myocardial infarction and cardiovascular disease are significant global health issues, particularly in Somalia. The Somali population faces challenges due to armed conflict, limited education, and underdeveloped healthcare infrastructure. Cardiovascular risk factors like diabetes, hypertension, and obesity are prevalent, and Somali people face increased susceptibility.

Methods: This study examined Somalia residents' perceptions, attitudes, and behaviors towards myocardial infarction using a cross-sectional paper-based questionnaire. The sample consisted of 313 urban residents in Mogadishu, aged 20 and above. The questionnaire was written in Somali and was designed for the demographic. Knowledge, attitude, and practice scores were categorized into two groups, with associations between knowledge, attitude, and practice with sociodemographic and health characteristics assessed using Logistic Regression.

Results: This study analyzed data from A random sample of participants who arrived at the hospital for various reasons regarding their perceptions, attitudes, and responses towards myocardial infarction. The mean age was 42 ± 19 years, with 54% being male and 46% female. Hypertension was the most common comorbidity, followed by dyslipidemia and diabetes. Over half of the respondents had no formal education, and only 3.8% had a smoking history. About 5.7% had a previous history of myocardial infarction. The mean knowledge score was 11.07 ± 0.697 , with 53.35% of respondents not knowing enough about myocardial infarction. The majority of the patients showed a favorable attitude, but only a slight majority could recognize symptoms of myocardial infarction. The majority of the patients had inadequate practice and behavior, with 53.6% not practicing regularly or engaging in sports.

Conclusion: The research highlights gaps in Mogadishu's Somali community's knowledge and practices regarding myocardial infarction. It emphasizes the need for health education, primary care, and community involvement to improve cardiovascular health awareness and reduce MI incidence.

Keywords: myocardial infarction, Somalia, cardiovascular disease, knowledge, practice

Introduction

Myocardial infarction (MI) and cardiovascular disease (CVD) are prominent global health issues, which also pose a major burden within the Somali population. The Somali population encounters distinct obstacles as a result of the repercussions of armed conflict, limited access to education, and an underdeveloped healthcare infrastructure. Myocardial Infarction (MI), commonly known as a heart attack, is a critical medical condition characterized by the interruption of blood supply to a part of the heart muscle, leading to the death of heart tissue. It is a global health concern with significant variations in knowledge, attitude, and practices in different regions, including Africa and specifically Somalia.

Numerous researches have shed light on the significant frequency of cardiovascular risk factors within Somali communities, encompassing conditions such as diabetes, hypertension, and obesity.¹ These identified risk variables are known to play a significant role in the pathogenesis of myocardial infarction (MI) and cardiovascular disease (CVD). Furthermore, it has been observed that Somali immigrants and refugees residing in the United States face an elevated

susceptibility to obesity and associated cardiovascular health risks.² This observation indicates the necessity of implementing specific interventions aimed at addressing these risk factors and enhancing cardiovascular well-being within the Somali population.

While there is limited research specifically focused on MI in Somalia, studies conducted in other African countries and among Somali diaspora populations provide valuable insights into the knowledge, attitudes, and practices regarding these conditions. Furthermore, the prevalence of MI is unknown in Somalia but is presumed to be high.

The influence of socioeconomic factors on the development and management of cardiovascular illnesses is of paramount importance. Research undertaken in the sub-Saharan African region, encompassing adjacent nations such as Ethiopia and Tanzania, has shed light on the influence of low socioeconomic level on cardiovascular risk factors and outcomes.^{3,4} Understanding and managing cardiovascular illnesses requires a high level of health literacy. For instance, low health literacy has been linked to an increased risk of death in people with end-stage renal disease.⁵

Attitudes toward healthcare and preventative medicine also have an impact on the treatment of MI. Somali patients are less familiar with preventive healthcare practices and tend to seek medical attention only when they are sick.⁶ Aside from individual knowledge, attitude, and practice, the healthcare system is also important in preventing myocardial infarction. The value of evidence-based preventative strategies is emphasized in guidelines for the management of acute myocardial infarction.⁷ This study aims to analyze knowledge, attitudes, and behaviors about myocardial infarction among Mogadishu citizens, focusing on socio-economic issues and health literacy, to address healthcare obstacles and improve cardiovascular health outcomes within the Somali community.

Methods and Materials

Design and Settings

A cross-sectional survey in the form of paper-based questionnaire was prepared for this study and used for data collection from 1 June to 31 August to examine Somalia inhabitants' Perceptions, Attitudes, and behaviors towards Myocardial Infarction. A random sample of participants who arrived at our hospital Mogadishu Somali Turkish Training and Research Hospital for various reasons were referred to our cardiology clinic for the study, they were greeted by a resident doctor, consent was obtained, and they were supported in understanding and answering the questions.

Participants

People above the age of 20 who came to our hospital and accepted to participate in the study. Somali urban residents of Mogadishu were the intended participants; Somalis living outside of Mogadishu or Somalia were not eligible for this study. A sample of 313 urban residents in Mogadishu were collected. With an approximately 95% confidence level, the margin of error is approximately $\pm 5\%$.

Questionnaire

The questionnaire utilized was similar to other myocardial infarction questionnaires. It was, however, specifically created for our demographic. The questionnaire was written in Somali, and the writing was clear and concise. The process of creating the questionnaire began with the establishment of precise research goals pertaining to cardiovascular health within the community. The selection of important variables was guided by a review of the literature. The questionnaire was created using language that was considerate of cultural differences and underwent validity testing. Any problems with the clarity and cultural appropriateness of the questions were found and fixed in a pilot study with a diverse sample. As a result, the questionnaire is revised. By giving the updated questionnaire to a subset of participants twice, test-retest reliability is guaranteed. The final version is then utilized to collect data for the primary sample. The data was collected and assisted by a resident doctor who was not involved with any patient care, and the sample was random and was collected from participants who came our hospital for various reasons, whether they had risk factors, previous history of myocardial infarction or simply as a relative of a patient or those who came for a simple check-up. Everyone had an equal chance of being requested to participate for the study.

It takes around 7 minutes to complete the questions, and any query or difficulty in comprehending was cleared by a resident doctor who assisted the patient at every step of the process. Questions included demographic data, such as age, weight, height, education level. Past medical data were also collected, for instance, diabetes or hypertension history, and if there was any previous myocardial infarction. Knowledge, attitude, and practice questions were answered similarly with a binary yes or no, and each response was scored 1 and 0, respectively. In order to assess the reliability of the questionnaire, it was subjected to a pretesting process including a randomly selected sample. No notable language-related challenges were found throughout this procedure.

Statistical Analysis

The study employed IBM SPSS 24 for Windows. For continuous variables, descriptive statistics were reported as mean \pm standard deviation (SD), while for other variables, frequencies (%) were employed. The scores of the participants pertaining to knowledge, attitudes, and beliefs were presented and analyzed. The highest attainable scores for knowledge, attitude, and practice were 22, 9, and 7, correspondingly. Moreover, the overall knowledge, attitude and practice scores was categorized into two distinct groups. For example, knowledge was considered to be sufficient for any result above 11, while 11 or less was considered insufficient knowledge. Furthermore, for attitude and practice we similarly scored them into two categories. 5 and less was considered poor attitude, while above was assigned as good attitude. For practice, scores 4 and less were considered inadequate practice, in contrast scores above were acknowledged as adequate practice.

Associations between knowledge, attitude and practice with variable sociodemographic and health characteristics were assessed using Logistic Regression. For every analysis, we employed $p < 0.05$ to establish statistical significance.

Ethical Consideration

Due to the fact that our hospital is a research hospital an informed consent is obtained from every patient before giving them the questionnaire and assisting them in answering the questions. This study did not disclose any personal information. The study was approved by the research ethics committee of Mogadishu Somali Turkey Training and Research Hospital. The study was performed in line with the principles of the Declaration of Helsinki.

Results

Socio-Demographic Features

In this study the data collected from 313 participants in the hospital for various reasons regarding their perceptions, attitudes, and responses towards myocardial infarction were analyzed. The mean age was 42 ± 19 years of age. One hundred and seventy patient (54%) of the subjects were male, and 143 (46%) were female. Hypertension was the most common comorbidity in these subjects 64 (20%), followed by dyslipidemia 41 (13%) and diabetes 36 (11.5%). More than half of the respondents 163 (51%) did not have formal education. Only 12 (3.8%) had smoking history, and 14 (4.4%) had regular practice of Khat chewing. The mean weight of patients was 68 ± 27 kg. Eighteen (5.7%) of the respondents had previous history of myocardial infarction.

Knowledge Results

The mean knowledge score of the participants was 11.07 ± 0.697 , with a maximum score of 22. It was discovered that $n=167$, 53.35% of respondents did not know enough about myocardial infarction using the cut-off criteria (Figure 1). Out of those surveyed, 172 (or 55%) said that a clot-induced arterial blockage was the reason behind the myocardial infarction. Similarly, 187 participants, or 59.7%, acknowledged that MI may be lethal. Additionally, over 50% of the participants were aware that MI typically manifests as chest pain; only 131 (41.9%) were aware that MI might occasionally manifest without chest pain. Only 153 (48.9), 141 (45%), and 141 (45%) of the respondents—or fewer than half—identified hypertension, diabetes, and dyslipidemia as risk factors for MI when questioned about their potential contributions. Sedentary lifestyle was identified as a risk factor by 129 (41.2%) and obesity by 137 (43.8%). Moreover, the majority of participants did not regard weight




Figure 1 Overall knowledge of the study respondents.

loss, diabetes management, or hypertension control as helpful in the fight against myocardial infarction and cardiovascular disease. Surprisingly, 205 (65.5%) of our respondents acknowledged that MI is a medical emergency requiring prompt medical intervention. (Table 1)

Table 1 Responses to the Knowledge Questions Provided by the Participants, Detailing Their Knowledge About Cardiovascular Disease and Myocardial Infarction

Questions	Yes	No
Myocardial infarction is mostly caused by a clot obstructing the heart arteries.	172(55%)	141(45%)
Myocardial infarction is a dangerous medical condition that could be deadly.	187(59.7%)	126(40.3%)
Myocardial infarction can present in a similar fashion to gastric reflux.	137(43.8%)	176(56.2%)
Hypertension is a cause of MI	153(48.9%)	160(51.1%)
Myocardial infarction presents with central chest pain and shortness of breath.	171(54.6%)	142(45.4%)
Myocardial infarction can occur sometimes without any noticeable symptoms.	131(41.9%)	182(58.1%)
Myocardial infarction can present with back pain.	117(37.4%)	196(62.6%)
Age is a risk factor for cardiovascular disease (such as MI).	144(46%)	169(54%)
Smoking is a risk factor for cardiovascular disease (such as MI).	160(51.1%)	153(48.9%)
Khat usage is a risk factor for cardiovascular disease (such as MI).	141(45%)	172(55%)
Diabetes is a risk factor for cardiovascular disease (such as MI).	141(45%)	172(55%)
Being obese is a risk factor for cardiovascular disease (such as MI).	137(43.8%)	176(56.2%)
High blood cholesterol is a risk factor for cardiovascular disease (such as MI).	141(45%)	172(55%)
Being inactive is a risk factor for cardiovascular disease (such as MI).	129(41.2%)	184(58.8%)
Too much salt consumption is a risk factor for cardiovascular disease (MI).	137(43.8%)	176(56.2%)
Hypertension control is beneficial for preventing cardiovascular disease (such as MI).	137(43.8%)	176(56.2%)
Diabetes control is beneficial for preventing cardiovascular disease (such as MI).	132(42.2%)	181(57.8%)
Weight loss if overweight is beneficial for preventing cardiovascular disease (MI).	143(45.7%)	170(54.3%)
Exercise is beneficial for preventing cardiovascular disease (such as MI).	228(72.8%)	85(27.2%)
Myocardial infarction treatment requires continues medication usage.	218(69.6%)	95(30.1%)
PCI must be performed within 12 hours of chest pain onset.	182(58.1%)	131(41.9%)
If Myocardial infarction is not treated within time it will lead to life-threatening complications.	205(65.5%)	108(34.5%)

Attitude Results

The mean attitude score of the respondents was 5.086 ± 0.3756 . Using cut-off criteria, a slight majority of the patient showed favorable attitude. Almost 168 (53.7%) said they could recognize the symptoms of myocardial infarction. In contrast only 144 (46%) said they can recognize it in others. Furthermore, 154 (49.2%) believed MI is unpreventable, while 201 (64.2%) said they will accept a MI diagnose from a doctor. Lastly, 202 (64.5%) said they are ready to do whatever it takes to lower their risk of myocardial infarction. (Table 2)

Practice Results

The practice score average was 3.463259 ± 0.352 . Majority of the patient had inadequate practice and behavior according to the cut-off score. 168 (53.6%) of the respondents did not practice regularly or engage in any kind of sport. Only 119 (38%) of those questioned did not regularly check their blood pressure, sugar level or their cholesterol level. Majority of the patient did not also control their salt, sugar and fatty food intake. Noticeably, out of those who had diabetes and hypertension or dyslipidemia more than 75% of them did not comply with their medications. (Table 3)

Factors Related to Myocardial Infarction Knowledge, Attitudes, and Practices

In a logistic regression, the following predictor variables such as age, gender, educational status, and a history of diabetes, hypertension, and dyslipidemia were input as variables against the knowledge, attitude and practice scores to be analyzed. With the exception of one group, however, we did not find any significant associations between the various variables and the level of knowledge. It was discovered that participants who were male and had not received any formal education had a higher probability of having a lower knowledge, unfavorable attitude and inadequate practice level ($P < 0.05$).

Table 2 Responses to the Attitude Questions Provided by the Participants, Detailing Their Attitude Towards Cardiovascular Disease and Myocardial Infarction

Questions	Yes	No
I know the symptoms of a Myocardial infarction (MI) if I am having one.	168(53.7%)	145(46.3%)
I know the symptoms of a Myocardial infarction (MI) if someone else is having one.	144(46%)	169(54%)
Myocardial infarction (MI) is a dangerous condition which requires immediate medical attention	211(67.4%)	102(32.6%)
Many people have Myocardial infarction (MI) in our country	184(58.8%)	129(41.2%)
I believe a doctor if he diagnosed me with Myocardial infarction (MI)	201(64.2%)	112(35.8%)
I only trust a cardiologist to make the diagnose of Myocardial infarction (MI)	142(45.3%)	171(54.7%)
Myocardial infarction (MI) is a preventable disease	154(49.2%)	159(50.8%)
Myocardial infarction (MI) is a treatable condition in our country	182(58.1%)	131(41.9%)
I am ready to do anything to lower my risk of developing cardiovascular disease (such as MI)	202(64.5%)	111(35.5%)

Table 3 Responses to the Practice Questions Provided by the Participants, Detailing Their Behavior Towards Cardiovascular Disease and Myocardial Infarction

Questions	Yes	No
If you have chronic diseases (HTN/DM/DL) do you take your medication regularly	145(46.7%)	168(53.7%)
Do you do routine body exercises	145(46.4%)	168(53.6%)
Do you regularly do blood pressure, sugar level, cholesterol level check-ups	119(38%)	194(62%)
Do you control the amount of sugar in your food	147(47%)	166(53%)
Do you control the amount of salt in your food	155(49.5%)	158(50.5%)
Do you control the amount of fat or oil in your food	154(49%)	159(50.7%)
Do you control your weight	165(52.7%)	148(47.3%)

Discussion

In this discussion, we will compare Mogadishu's citizens' understanding, attitudes, and practices around MI with those of other parts of Africa and the globe. We will also investigate the causes of the differences and provide potential solutions. Numerous researches have looked at various groups' knowledge, attitudes, and practices about MI. For instance, research carried out in Tanzania discovered that just a small percentage of people there believed they were at risk of MI, and that everyone knew very little about the disease's symptoms.⁸ This result was similar to our population findings. The prevalence of MI is rising throughout Africa.⁹ However, a study that was carried out in three nations—the United States, Australia, and New Zealand—reported similar findings, with 56% of participants knowing enough about ACS symptoms.¹⁰ Moreover, MI cases are rising in South Africa which many considers the most developed country in Africa.¹¹ These data show that improved MI prevention, education and management strategies are required in Africa. The relationship between health literacy and the management and prevention of cardiovascular disease is one crucial factor. The scientific statement released by the American Heart Association highlights the basic need of health literacy for both primary and secondary cardiovascular disease prevention.¹² Another study also demonstrated that health literacy is connected to healthy behaviors and self-reported health in people with cardiovascular disease, which further validates this.¹³ These studies emphasize how critical it is to raise health literacy in order to increase understanding and encourage healthy habits related to MI. Somalia is believed to have a low health literacy due to years of civil war and poverty. According to a comprehensive review and meta-analysis by, attitudes toward Myocardial infarction have also been demonstrated to predict attendance at cardiac rehabilitation after an acute myocardial infarction.¹⁴ A small majority of our patients showed a favorable attitude towards myocardial infarction. This may be due to a popular perception that heart attack is deadly acute condition in Somali culture. For instance, Patients' perceptions of their myocardial infarction can differ; according to a study by, some consider it to be an acute heart attack instead of a sign of a long-term illness.¹⁵ Effective management of myocardial infarction necessitates a thorough understanding of and implementation of preventative techniques.¹⁶ Diabetes mellitus, hypertension, dyslipidemia, and metabolic syndrome are risk factors for myocardial infarction.^{17,18} Based on the cut-off score, most of our patients exhibited inadequate conduct and practice. 53.6% of the respondents, or 168 people, said they did not play any sport or exercised on a regular basis. Improvements in lifestyle, such as frequent exercise and physical activity, are essential for preventing cardiovascular disease. Physical activity has demonstrated benefits for cardiovascular well-being and can lower the risk of atherosclerotic cardiovascular disease.¹⁹ Furthermore, most of the patients with chronic diseases did not comply with their medications in contrast to Seventy percent of hypertensive people in a Tshwane, South Africa research reported complying with treatment, while more than 80% of our hypertensive patients did not comply.²⁰

It is important to think about the effects of telemedicine, mobile technology-based lifestyle in order to improve knowledge and practice. People have seen telemedicine as a possible way to improve health care and practice, especially for people with heart diseases.²¹ The use of mobile technology and smart devices has shown promise in changing the way people with a high risk of heart disease live, providing a practical way to enhance patient outcomes. These mobile technologies offer a way in which many people with the disease, risk factors or relatives of these groups can be educated.²² Moreover, the internet provides a vast and extensive platform to access an audience of individuals who may have been previously deemed hard to reach. Sharing articles and videos about myocardial infarction can enhance understanding and influence behaviors related to this serious condition. Nevertheless, deceptive information can also be spread online.²³ To summarize, effectively addressing knowledge about myocardial infarction (MI) in Mogadishu necessitates implementing customized interventions that take into account cultural subtleties. It is crucial to utilize health literacy and technology to develop efficient strategies for prevention and management.

Limitations of Our Study

In our study we faced several limitations. First, there may be a limit due to the sample size and representativeness. Our study may not have included a representative and varied sample of the Somali population due to restricted access, which could have resulted in selection bias. Second, response bias is introduced by our data collection methods, such as surveys and interviews, as participants may give socially acceptable replies. This could result in an underestimating of bad

practices or an overestimation of knowledge and positive attitudes. Another issue with self-reported data is the potential for recollection or social desirability biases, which could skew the responses and effect data quality. Finally, the breadth of our study may be constrained by a lack of funding and time, which could have an impact on the number of participants or data collection locations.

Conclusion

In conclusion, our research highlights the serious gaps in the Somali community of Mogadishu's knowledge, attitudes, and practices surrounding myocardial infarction (MI). To improve cardiovascular health awareness and management, the results highlight the critical need for focused health education, enhanced primary care, and community involvement. In order to close these gaps and guarantee that prompt and efficient interventions result in a healthier and better-informed Somali population, policymakers and healthcare professionals must work together. This will eventually lessen the incidence of MI in the area.

Author Contributions

Each author contributed significantly to the work reported, whether it was through conception, study design, execution, data acquisition, analysis, and interpretation, or in all of these areas; they all helped draft, revise, or critically review the article; they approved the final version that was published; they all agreed on the journal to which the article was submitted; and they all agreed to take responsibility for the work in its entirety.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Mohamed A, Lantz K, Ahmed Y, et al. An assessment of health priorities among a community sample of Somali adults. *J Immigrant Minority Health*. 2021;24(2):455–460. doi:10.1007/s10903-021-01166-y
2. Njeru J, Wieland M, Okamoto J, et al. Social networks and obesity among Somali immigrants and refugees. *BMC Public Health*. 2020;20(1). doi:10.1186/s12889-020-8315-7
3. Yuyun M, Sliwa K, Kengne A, Mocumbi A, Bukhman G. Cardiovascular diseases in sub-saharan Africa compared to high-income countries: an epidemiological perspective. *Global Heart*. 2020;15(1):15. doi:10.5334/gh.403
4. Cappuccio F, Miller M. Cardiovascular disease and hypertension in sub-saharan Africa: burden, risk and interventions. *Int Emerg Med*. 2016;11(3):299–305. doi:10.1007/s11739-016-1423-9
5. Cavanaugh K, Wingard R, Hakim R, et al. Low health literacy associates with increased mortality in ESRD. *J Am Soc Nephrol*. 2010;21(11):1979–1985. doi:10.1681/asn.2009111163
6. Pavlish C, Noor S, Brandt J. Somali immigrant women and the American health care system: discordant beliefs, divergent expectations, and silent worries. *Soc sci med*. 2010;71(2):353–361. doi:10.1016/j.socscimed.2010.04.010
7. Steg P, James S, Atar D, et al. Esc guidelines for the management of acute myocardial infarction in patients presenting with ST-segment elevation. *Revista Española De Cardiología*. 2013;33:548.
8. Hertz J, Madut D, Tesha R, et al. Knowledge of myocardial infarction symptoms and perceptions of self-risk in Tanzania. *Am Heart J*. 2019;210:69–74. doi:10.1016/j.ahj.2019.01.003
9. Mensah G. Ischaemic heart disease in Africa. *Heart*. 2008;94(7):836–843. doi:10.1136/hrt.2007.136523
10. Dracup K, McKinley S, Doering LV, et al. Acute coronary syndrome: what do patients know? *Arch Internal Med*. 2008;168(10):1049–1054. doi:10.1001/archinte.168.10.1049
11. Stassen W, Olsson L, Kurland L. The application of optimisation modelling and geospatial analysis to propose a coronary care network model for patients with ST-elevation myocardial infarction. *Afr J Emergency Med*. 2020;10:S18–S22.
12. Magnani J, Mujahid M, Aronow H, et al. Health literacy and cardiovascular disease: fundamental relevance to primary and secondary prevention: a scientific statement from the American heart association. *Circulation*. 2018;138(2). doi:10.1161/CIR.0000000000000579
13. Aaby A, Friis K, Christensen B, Rowlands G, Maingal H. Health literacy is associated with health behaviour and self-reported health: a large population-based study in individuals with cardiovascular disease. *Eur J Preventive Cardiol*. 2017;24:1880.
14. French D, Cooper A, Weinman J. Illness perceptions predict attendance at cardiac rehabilitation following acute myocardial infarction: a systematic review with meta-analysis. *J Psychosomatic Res*. 2006;61(6):757–767. doi:10.1016/j.jpsychores.2006.07.029
15. Brink E, Karlson B, Hallberg L. Readjustment 5 months after a first-time myocardial infarction: reorienting the active self. *Journal of Advanced Nursing*. 2006;53(4):403–411. doi:10.1111/j.1365-2648.2006.03737.x
16. Adhikari A, Bhandari B. Awareness and practice of prevention of myocardial infarction among hypertensive patients: cross sectional study in a tertiary cardiac centre of Nepal. *Nepalese Heart J*. 2017;14(1):25–28. doi:10.3126/njh.v14i1
17. Uppin A, Badiger R, Sharma G, Soni R, Parme S, Maddiri P. Assessment of incidence rate and prognosis of metabolic syndrome among acute myocardial infarction: a longitudinal study. *Int J Adv Med*. 2020;7(2):267. doi:10.18203/2349-3933.ijam20200077

18. Shepherd J, Cobbe S, Ford I, et al. Prevention of coronary heart disease with pravastatin in men with hypercholesterolemia. *N Engl J Med.* 1995;333(20):1301–1308. doi:10.1056/nejm19951116
19. Thompson P, Buchner D, Piña I, et al. Exercise and physical activity in the prevention and treatment of atherosclerotic cardiovascular disease. *Circulation.* 2003;107(24):3109–3116. doi:10.1161/01.cir.00000075
20. Mafutha G, Wright S. Compliance or non-compliance of hypertensive adults to hypertension management at three primary healthcare day clinics in Tshwane. *Curationis.* 2013;36(1). doi:10.4102/curationis.v36i1.52
21. Hayiroğlu M. Telemedicine: current concepts and future perceptions. *Anatolian J Cardiol.* 2019. doi:10.14744/anatoljcardiol.2019.12525
22. Tekkeşin A, Hayiroğlu M, Çinier G, et al. Lifestyle intervention using mobile technology and smart devices in patients with high cardiovascular risk: a pragmatic randomised clinical trial. *Atherosclerosis.* 2021;319:21–27. doi:10.1016/j.atherosclerosis.2020.12.020
23. Hayiroğlu Mİ, Çinier G, Keser N, et al. Evaluation of websites reached using Google in the modern digital era related to approach to cholesterol. *Türk Kardiyol Dern Ars.* 2020;48(6):576–584. doi:10.5543/tkda.2020.40306

Vascular Health and Risk Management

Dovepress

Publish your work in this journal

Vascular Health and Risk Management is an international, peer-reviewed journal of therapeutics and risk management, focusing on concise rapid reporting of clinical studies on the processes involved in the maintenance of vascular health; the monitoring, prevention and treatment of vascular disease and its sequelae; and the involvement of metabolic disorders, particularly diabetes. This journal is indexed on PubMed Central and MedLine. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/vascular-health-and-risk-management-journal>