



Prognostic Value of Arterial Blood Gas Indices Regarding the Severity of Traumatic Injury and Fractures of the Femur and Pelvis

Leila Mohsenian¹, Mohamad Karim Khoramian^{1*}, Sara Sadat Mazloom¹

¹Department of Emergency Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

*Corresponding author: Mohamad Karim Khoramian

Address: Department of Emergency Medicine, Shiraz University of Medical Sciences, Shiraz, Iran
Cellphone: +98-917-721 4895
e-mail: mkkhoramian@yahoo.com

Received: July 7, 2018

Revised: August 18, 2018

Accepted: September 16, 2018

► ABSTRACT

Objective: To evaluate the prognostic value of arterial blood gas (ABG) indices, especially base excess (BE), regarding the severity of traumatic injury.

Methods: A total number of 96 trauma patients with fractures of femur and pelvis were included in this study. Demographic characteristics and clinical information of samples were recorded. The results of ABG test and hemoglobin (Hb) were evaluated at two time intervals (on admission and 6 hours later). The outcome of the patients was evaluated at the end of the study (dead or alive). The ABG indices included O₂ saturation (O₂Sat), PH, PO₂, PCO₂, bicarbonate (HCO₃) and base excess (BE). The mean of the hemodynamic and ABG indices were compared between those who were discharged and mortality cases. The confounders were compensated using a multivariate logistic regression model.

Results: Overall 94 trauma patients with long and pelvic fractures were included. There were 69 (73.4%) men and 25 (26.6%) women among the patients with mean age of 37.43±20.07 years. The mortality rate was 10 (10.6%). The most common mechanism of injury was motorcycle accident in 41 (43.6%) and car collision in 43 (45.7%) patients. We found that mortality was significantly associated with amount of FFP transfusion ($p=0.005$), but was not associated with amount of transfused packed cells ($p=0.113$). We also found that mortality was associated with lower BP after 6 hours of admission ($p=0.001$), higher HR on admission ($p=0.036$), lower HR after 6 hours ($p=0.017$), lower O₂Sat on admission ($p<0.001$), higher PCO₂ after 6 hours and lower BE on admission ($p=0.025$).

Conclusion: The ABG indices including O₂Sat, BE and PCO₂ are considered prognostic determinants of outcome in trauma patients with long and pelvic fractures. These findings can be considered as fundamental studies to achieve new diagnostic methods.

Keywords: Prognosis; Injuries; Femur; Pelvis; Mortality; Arterial blood gas (ABG); hemodynamic indices.

Please cite this paper as:

Mohsenian L, Khoramian MK, Mazloom SS. Prognostic Value of Arterial Blood Gas Indices Regarding the Severity of Traumatic Injury and Fractures of the Femur and Pelvis. *Bull Emerg Trauma*. 2018;6(4):318-324. doi: 10.29252/beat-060408.

Introduction

Trauma is among the most common causes of disability and mortality worldwide [1-3]. Meanwhile, fractures of pelvis and long bones are commonly observed in middle aged people due to osteoporosis and falls and in young people usually due to severe hits [4, 5]. Patients with severe pelvic fractures account for 3% of total fractures [6]. The complications of bone fracture include infection, avascular necrosis, large blood vessel damage, damage to the nerves, damage to the muscles, damage to the tendons, osteoarthritis and in certain cases, shock, fat embolism, compartment syndrome, lower extremity deep vein thrombosis (DVT) and lung embolism [7-9]. It is worth noting that hemodynamic compromise in patients with unstable pelvic fractures associated with arterial bleeding occurs in less than 20% of cases [10-12].

Severity of trauma and development of systemic inflammatory response syndrome have been studied as effective factors by causing inflammation [5, 9, 13]. Recent studies have shown that decreased tissue perfusion, without clear clinical symptoms, led to an increase in the levels of lactate and carbonic acid in trauma patients and was introduced as a predictor of mortality in these patients, and its criterion is measuring serum lactic acid [14, 15]. However, it is not possible to measure this parameter in all trauma centers. On the other hand, BE of the blood is an index of shock and physiological impairment in trauma patients and is associated with blood lactate concentration and the severity of the injury [16, 17]. Factors that affect the base deficit include the use of bicarbonate, hypothermia, hypocapnia, heparin, ethanol and ketoacidosis [18, 19]. Base excess (BE) is the levels of base that is required for titration of one liter of full blood to reach pH 7.4 in the case of full saturation of blood with oxygen, at 37°C and PaCO₂=40 mmHg [16, 17]. Studies have shown that in cases of severe base deficit, most of the trauma patients suffer from shock due to bleeding. However, in the absence of shock, it seems that this criterion can be a sign of lactic acidosis [16-19]. Some studies have suggested that the mortality of trauma patients can be predicted by the levels of base deficit within the first 24 hours after the trauma [20, 21].

Post-traumatic management with rapid diagnosis and timely action such as resuscitation is vital in trauma patients [8, 13, 22]. Trauma is one of the major causes of death and disability in Iran. Not only bone fractures cause permanent defects and disability in patients, but also cause death in many cases due to severe bleeding (the pelvic floor and long bone) [23]. Therefore, detection of dependent and prognostic variables can be very important and prevent the complications of these fractures. According to previous studies, the arterial blood gas (ABG) changes seem to be appropriate predictor of severity of traumatic injury [24, 25]. Nevertheless,

no comprehensive study has been conducted in this regards. Thus, the present study aims to examine the prognostic relationship between ABG indices, including BE, and bicarbonate, and the severity of traumatic injury.

Materials and Methods

Study Population

In this prospective cohort study, all trauma patients with major bone fracture, defined as long and pelvic fractures, referred to Shahid Rajaie hospital, a Level-I trauma center affiliated with Shiraz University of Medical Sciences located in southern Iran during a 6-month period from March to September 2017 were included. We included adult patients (18-65 years) with fractures of femur or pelvis. The exclusion criteria were patients with small bone fracture or patients whose fracture occurred more than 24 hours ago. Those also excluded those with severe multi-organ injuries, those who passed away less than 48 hours after admission, patients with comorbidities such as diabetes mellitus, ischemic heart disease, chronic obstructive pulmonary disease (COPD) and malignancies. The study protocol was approved by the institutional review board (IRB) and medical ethics committee of Shiraz University of Medical Sciences. All the patients or their legal representatives provided their informed written consents before the study.

Study Protocol

All the included patients were evaluated by an emergency medicine resident on admission. The demographic variables, including age and gender were recorded; clinical information of the samples such as fracture type, fracture site, the mechanism of injury and the level of consciousness were recorded into a data gathering form. The mechanism of injury was classified to motorcycle accident, car accident, exposed burn, fall and gunshot injuries. The vital signs including blood pressure (BP), heart rate (HR) and respiration rate (RR) were recorded. All the patients gave 2 samples for ABG and Hb on admission. The ANG and Hb measurement was repeated 6 hours after the admission.

The ABG was analyzed on arterial blood by using the Nova Biomedical (pHOx Plus, USA). We measured the PH, oxygen saturation (O₂ Sat), PCO₂, HCO₃ and BE. The normal range of these parameter was considered as follow; PH: 7.35-7.45, BE: -2 - +2 mmol/L, HCO₃: 22-26 mEq/L, PCO₂: 35-45mmHg and O₂Sat: 95-100%. In addition, PH higher than 7.45 was considered alkalemia and lower than 7.35 was recorded as acidemia, PCO₂ higher than 45mmHg was considered as respiratory acidosis and lower than 35mmHg as respiratory alkalosis. HCO₃ higher than 26mEq/L was metabolic alkalosis and lower than 22 mEq/L was recorded as metabolic acidosis [26, 27]. Furthermore, the Sysmex kx21 device was

used to run blood tests and determine the levels of Hb. The outcome of patients, including the length of hospitalization, the transfer of patients, discharge, or mortality was also recorded.

Statistical Analysis

All the data were analyzed using the statistical package for social sciences (SPSS Inc., Chicago, Illinois, USA) version. All the data are presented as mean±SD and proportions as appropriate. Descriptive data were composed from the frequency distribution table, central indexes, distribution and percentages. The changes in the parametric variables of ABG within groups were compared using paired t-test while normal distribution was encountered. In parametric variables without normal distribution, Kruskal-Wallis test was used. The parametric variables were compared between two groups (dead and alive) using independent t-test. The proportions were compared with chi-square test. We also run a multivariate logistic regression model to compensate for confounders such as age, gender and injury mechanism. A 2-sided p-value of less than 0.05 was considered statistically significant.

Results

Overall a total number of 94 patients were included in the current study. There were 69 (73.4%) men and 25 (26.6%) women among the patients with mean age of 37.43±20.07 years. The most common mechanism of injury was motorcycle accident in 41 (43.6%) and car collision in 43 (45.7%) patients. The mean GCS on admission was 13.38±3.27. The demographic and baseline characteristic of the patients is summarized in Table 1. We found that 66 (70.2%) of the patients required full blood, and 17 (18.1%) fresh frozen plasma (FFP) transfusions. The mean packed cells and FFP transfusion were 3.23±3.5 and 1.0±2.5 units per patient, respectively. Overall 48 (51.1%) patients were admitted to the Emergency Room (ER) while 13 (13.8%) were transferred to the orthopedic room immediately. Finally, 33 (35.1%) patients were admitted to the intensive care unit (ICU).

As demonstrated in Figure 1 the hemodynamic variables and the hemoglobin changes were observed between admission and a 6-hour period. The BP increased significantly ($p=0.033$) and HR levels decreased significantly ($p<0.001$). The Hb level also

Table 1. Demographic characteristics and mechanism of injury in the studied samples based on gender.

Variable	Men (n=67)	Women (n=25)	Total (n=94)	p-value
Age (years)	35.75±17.22	41.44±26.42	37.43±20.07	0.01
GCS ^a	13.03±3.59	14.36±1.93	13.38±3.27	<0.001
Mechanism of Injury				
Motorcycle (%)	37 (39.4%)	4 (4.3%)	41 (43.6%)	<0.001
Car (%)	26 (27.7%)	17 (18.1%)	43 (45.7%)	
Fall (%)	5 (5.3%)	3 (3.2%)	8 (8.5%)	
Gunshot (%)	1 (1.1%)	0 (0%)	1 (1.1%)	
Expose Burning (%)	0 (0%)	1 (1.1%)	1 (1.1%)	

^aGCS: Glasgow Coma Scale




Fig. 1. Changes in hemoglobin and hemodynamic indices between the two measurement intervals. The level of hemoglobin decreased significantly (A) and the respiratory rate also decreased after 6 hours (B). The heart rate decreased significantly after 6 hours of admission (C) while the blood pressure increased in this time period (D).

decreased significantly after 6 hours of admission in the study population ($p<0.001$) (Figure 1). As demonstrated in Table 2, the mean PH was below normal either on admission and 6 hours later. In the same way the patients had lower levels of HCO_3^- , BE and O_2Sat either on admission and 6 hours later (Table 2).

Overall 84 (89.4%) patients were discharged in good condition while 10 (10.6%) patients died during the hospital course. Among the mortality cases were 8 (8.5%) men and 2 (2.1%) women. The mean GCS was significantly lowered in deceased patients compared to those who were discharged from the hospital (9.4 ± 5.5 vs. 13.8 ± 2.5 ; $p<0.001$). We also found that mortality

was significantly associated with amount of FFP transfusion ($p=0.005$), but was not associated with amount of transfused packed cells ($p=0.113$). We found that mortality was associated with lower BP after 6 hours of admission ($p=0.001$), higher HR on admission ($p=0.036$), lower HR after 6 hours ($p=0.017$), lower O_2Sat on admission ($p<0.001$), higher PCO_2 after 6 hours and lower BE on admission ($p=0.025$). The results are summarized in Table 3. We have also run a multivariate logistic regression analysis to compensate for confounders including the age, gender and mechanism of injury. All the variables remained statistically significant predictors of mortality after compensating for confounders.

Table 2. The changes in parameters if of the arterial blood gas (ABG) on admission and 6 hours later in 94 trauma patients with long and pelvic fractures.

Variable		Mean	t	Threshold	p-value
PH	On admission	7.34±0.088	-5.777	7.4	<0.001
	6-hour	7.32±0.086	-8.244		
O_2Sat	On admission	92.1±6.68	-7.838	97.5	<0.001
	6-hour	95.98±2.47	-5.968		
PCO_2	On admission	40.69±11.02	0.596	40	0.122
	6-hour	37.99±10.44	-1.814		
HCO_3^-	On admission	22.12±4.18	-4.087	24	<0.001
	6-hour	20.40±4.05	-8.128		
BE	On admission	-3.251±4.117	-7.491	0	<0.001
	6-hour	-4.531±5.748	-7.436		

Table 3. The correlation of changes in hemodynamic and arterial blood gas indices with the outcome of 94 trauma patients with long and pelvic fractures.

	Dead (n=10)	Alive (n=84)	p-value
Blood pressure (mmHg)			
On admission	118.1±26.1	119.12±18.4	0.148
6-hour	102.6±29.9	126.3±15.4	0.001
Respiratory rate (per min)			
On admission	17.33±3.32	19.02±3.43	0.686
6-hour	17.36±3.84	17.36±3.84	0.799
Hear rate (per min)			
On admission	105.9±39.3	103.7±19.1	0.036
6-hour	92.1±31.8	95.9±15.5	0.017
Hemoglobin (g/dL)			
On admission	13.26±1.5	12.75±2.5	0.182
6-hour	11.67±2.1	11.26±2.7	0.224
PH			
On admission	7.25±0.07	7.35±0.08	0.670
6-hour	7.20±0.09	7.33±0.07	0.141
O_2Sat (%)			
On admission	83.9±12.6	93.1±4.8	<0.001
6-hour	94.4±2.45	96.17±2.4	0.801
PCO_2 (mmHg)			
On admission	50.8±12.7	39.5±10.3	0.551
6-hour	47.9±16.8	37.1±9.2	0.007
HCO_3^- (mEq/L)			
On admission	23.8±5.22	21.9±4.17	0.213
6-hour	18.9±5.12	20.5±3.93	0.244
BE (mmol/L)			
On admission	-4.68±5.8	-3.09±3.8	0.025
6-hour	-8.72±5.4	-7.11±5.6	0.557

Discussion

In recent years, trauma has been one of the most important causes of disability and mortality worldwide [1-3]. This problem can be caused by several factors, the most important of which are motorcycle accidents, car accidents and falling down. Trauma can cause various complications. Rupture of organs such as spleen, and lung, rupture of major vessels and internal bleeding are some of the most important complications of trauma. The most important complication that can be caused by trauma is bone fracture, which may lead to organ defects or disability [7-9].

Fractures of major bones such as femur and pelvic bone are among the most severe and complex injuries of the skeletal system [7, 13]. Due to the proximity of these bones to blood vessels and nerves, and organs such as the intestine and bladder, these bones increase the risks caused by this fracture. After bone fracture, due to the contraction of strong muscles around the bone, the fracture edges cross each other, the fractured organ becomes shorter and the injuries increase [7-9, 11, 13].

For these reasons, bleeding in the pelvic floor and long bones causes shock and even death in some cases [28-30]. Thus, identifying the prognostic criteria of the patients' conditions can be very important and help prevent complications. It seems that ABG changes can be considered as good predictor of the severity of traumatic injury.

The results show a significant relationship between BP and HR and mortality rate in the study samples; the lack of increase in BP to normal levels in the early hours can be considered a prognostic factor. In addition, the results showed that among the ABG indices, O_2 Sat, BE and PCO_2 could be prognostic indices of the status of patients with major bone fractures. In this study, it was found that the mean O_2 Sat, BE and PCO_2 levels in samples who were deceased due to the fracture of femur and pelvis showed significant difference compared with the recovered samples; the mean on admission O_2 Sat and BE levels in deceased samples was significantly lower than that of recovered samples. Moreover, mean PCO_2 after 6 hours in deceased samples was higher than that of recovered samples. The results showed that PH in the study samples was significantly lower than normal levels and the samples had acidemia. In addition, the mean O_2 Sat, HCO_3 and BE levels in the first and second times showed that these factors were significantly lower than the normal range. Furthermore, the results show a significant decrease in BE and HCO_3 levels in the second time compared to the first one. These results indicate the prognostic power of ABG indices regarding the complications of major bone fractures such as shock and death.

As stated earlier, decreased tissue perfusion without clear clinical symptoms is one of the complications of fracture, which leads to an increase in the levels

of lactate and carbonic acid in trauma patients and is introduced as a predictor of mortality in these patients. Some studies suggest that the lack of BE was the same as the clinical measurement of metabolic acidosis, which becomes normal with proper resuscitation and control of bleeding, and can be used for monitoring the primary care of a trauma patient [31]. In addition, blood pressure can be a major contributor to the complications of trauma and concealed hemorrhages. In particular, one of the best indicators of hypotension is BE, which estimates metabolic acidosis. Callaway *et al.* in their study stated that mortality due to hypoperfusion may increase in trauma patients with initial BE lower than -6 mmol/L [32]. Moreover, according to the study of Peñasco *et al.*, low levels of BE and BP could be an important risk factor for patients with trauma; $BE < -6$ mmol/L increases the risk of death in patients with abnormal BP [33]. In a study by Kincaid *et al.*, it was found that initial BE showed the severity of injury and could predict mortality and the possibility of increased mortality in children was associated with base deficit and was a warning sign for potentially lethal injuries or uncompensated shock [34]. In a study by Ibrahim *et al.*, it was concluded that as BE decreases, blood pressure, mortality, injuries and complications constantly increase [20]. Similar to the results of the present study, the study of Mofidi *et al.* concluded that in patients with blunt abdominal trauma, BE is an important indicator for identifying intra-abdominal injuries and the lack of attention to it can increase mortality in patients. In addition, BE can be a prognostic factor for determining the need for more blood transfusions [35].

In a study by Toth *et al.*, it was argued that hypotension and low BE levels could be decisive factors for performing angiography, identifying vascular problems and controlling pelvic bleeding in patients with pelvic fracture [36]. Similar results were reported in the study by Jeroukhimov *et al.*, which stated that the rates of blood transfusions and the reduction of BE levels can be indicators for early angiography and emergency follow-up in patients with severe pelvic pain [37].

Similar to our results, the study of Neville *et al.* indicated that BE lower than -4 mmol/L increases the mortality rate. They concluded that high BE and lactate could be very important predictors of early death (in the first 24 hours) in trauma patients [21]. These findings were in line with the study of Jung *et al.*, [38].

Our results suggest that the BE, BP, O_2 Sat and HCO_3 parameters are useful in the initial evaluation of these patients and are consistent with previous studies. These findings suggest that these parameters can be predictors of the patient's condition. It is worth noting that our study results, similar to Neville *et al.*, show that $BE < -4$ mmol/L is a risk factor and increases the chance of death. However, some studies have reported this value to be lower than

-6mmol/L [21]. However, it is crystal clear that this factor can be cited as a prognostic factor. Ultimately, the limitations of this study were: not evaluating the cause of death or joint effects or other injuries. Moreover, underlying diseases were not specifically evaluated. Nonetheless, other factors that can affect BE at the time of admission, such as hemoglobin levels, were evaluated.

In conclusion, the results of this study showed that ABG indices such as O₂Sat, BE and PCO₂ could be predictors of the status of patients with major bone fractures. Furthermore, BP acts as a determining factor. In this study, the mean O₂Sat, BE and PCO₂ levels in samples who were deceased due to fractures of femur and pelvis showed significant difference compared with the recovered samples. These

findings can be considered as fundamental studies to achieve new diagnostic methods. It is suggested that supplementary studies be carried out to achieve reliable results.

Acknowledgement

The present article was extracted from the thesis written by Leila Mohsenian (No. 10640010535) and was financially supported by Shiraz University of Medical Sciences. We would like to acknowledge the editorial assistance of Diba Negar Research Institute for improving the English and style of the manuscript.

Conflicts of Interest: None declared.

References

1. Abedzadeh-Kalahroudi M, Razi E, Sehat M, Asadi-Lari M. Psychometric properties of the world health organization disability assessment schedule II -12 Item (WHODAS II) in trauma patients. *Injury*. 2016;**47**(5):1104-8.
2. Rivera JC, Glebus GP, Cho MS. Disability following combat-sustained nerve injury of the upper limb. *Bone Joint J*. 2014;**96-b**(2):254-8.
3. Turk DC. A diathesis-stress model of chronic pain and disability following traumatic injury. *Pain Res Manag*. 2002; (1):9-19.
4. Byun CS, Park IH, Oh JH, Bae KS, Lee KH, Lee E. Epidemiology of trauma patients and analysis of 268 mortality cases: trends of a single center in Korea. *Yonsei Med J*. 2015;**56**(1):220-6.
5. Haagsma JA, Graetz N, Bolliger I, Naghavi M, Higashi H, Mullany EC, et al. The global burden of injury: incidence, mortality, disability-adjusted life years and time trends from the Global Burden of Disease study 2013. *Inj Prev*. 2016;**22**(1):3-18.
6. Pehle B, Nast-Kolb D, Oberbeck R, Waydhas C, Ruchholtz S. Significance of physical examination and radiography of the pelvis during treatment in the shock emergency room. *Unfallchirurg*. 2003;**106**(8):642-8.
7. Oryan A, Monazzah S, Bigham-Sadegh A. Bone injury and fracture healing biology. *Biomed Environ Sci*. 2015;**28**(1):57-71.
8. Maheshwari, J. and V.A. Mhaskar, Essential orthopaedics. 2015: JP Medical Ltd.
9. Mears DC, Velyvis JH. Primary total hip arthroplasty after acetabular fracture. *Instr Course Lect*. 2001;**50**:335-54.
10. Cherry RA, Goodspeed DC, Lynch FC, Delgado J, Reid SJ. Intraoperative angioembolization in the management of pelvic-fracture related hemodynamic instability. *J Trauma Manag Outcomes*. 2011;**5**:6.
11. Fernandez-Palomo LJ. Pelvic fractures: early management of hemodynamically unstable patients. *Acta Ortop Mex*. 2012;**26**(5):275-81.
12. Jeroukhimov I, Ashkenazi I, Kessel B, Gaziantz V, Peer A, Altshuler A, et al. Selection of patients with severe pelvic fracture for early angiography remains controversial. *Scand J Trauma Resusc Emerg Med*. 2009;**17**:62.
13. Ehara S. Complications of skeletal trauma. *Radiol Clin North Am*. 1997;**35**(3):767-81.
14. Davis JW, Kaups KL, Parks SN. Effect of alcohol on the utility of base deficit in trauma. *J Trauma*. 1997;**43**(3):507-10.
15. Paladino L, Sinert R, Wallace D, Anderson T, Yadav K, Zehtabchi S. The utility of base deficit and arterial lactate in differentiating major from minor injury in trauma patients with normal vital signs. *Resuscitation*. 2008;**77**(3):363-8.
16. Park M, Azevedo LC, Maciel AT, Pizzo VR, Noritomi DT, da Cruz Neto LM. Evolutionary standard base excess and serum lactate level in severe sepsis and septic shock patients resuscitated with early goal-directed therapy: still outcome markers? *Clinics (Sao Paulo)*. 2006;**61**(1):47-52.
17. Wijaya R, Ng JH, Ong L, Wong AS. Can venous base excess replace arterial base excess as a marker of early shock and a predictor of survival in trauma? *Singapore Med J*. 2016;**57**(2):73-6.
18. Park M, Taniguchi LU, Noritomi DT, Liborio AB, Maciel AT, Cruz-Neto LM. Clinical utility of standard base excess in the diagnosis and interpretation of metabolic acidosis in critically ill patients. *Braz J Med Biol Res*. 2008;**41**(3):241-9.
19. Rose R, Kheirabadi BS, Klemcke HG. Arterial blood gases, electrolytes, and metabolic indices associated with hemorrhagic shock: inter- and intrainbred rat strain variation. *J Appl Physiol (1985)*. 2013;**114**(9):1165-73.
20. Ibrahim I, Chor WP, Chue KM, Tan CS, Tan HL, Siddiqui FJ, et al. Is arterial base deficit still a useful prognostic marker in trauma? A systematic review. *Am J Emerg Med*. 2016;**34**(3):626-35.
21. Neville AL, Nemtsev D, Manasrah R, Bricker SD, Putnam BA. Mortality risk stratification in elderly trauma patients based on initial arterial lactate and base deficit levels. *Am Surg*. 2011;**77**(10):1337-41.
22. Feng ZC, Chang P, Tao SH, Chen H. Continuous blood purification in the treatment of pediatric septic shock. *Zhonghua Er Ke Za Zhi*. 2006;**44**(8):579-8.
23. Forouzanfar MH, Sepanlou SG, Shahrzad S, Dicker D, Naghavi P, Pourmalek F, et al. Evaluating causes of death and morbidity in Iran, global burden of diseases, injuries, and risk factors study 2010. *Arch Iran Med*. 2014;**17**(5):304-20.
24. Dumont TM, Visioni AJ, Rughani AI, Tranmer BI, Crookes B. Inappropriate prehospital ventilation in severe traumatic brain injury increases in-hospital mortality. *J Neurotrauma*. 2010;**27**(7):1233-41.
25. Glenn TC, Kelly DF, Boscardin WJ,

- McArthur DL, Vespa P, Oertel M, et al. Energy dysfunction as a predictor of outcome after moderate or severe head injury: indices of oxygen, glucose, and lactate metabolism. *J Cereb Blood Flow Metab.* 2003;**23**(10):1239-50.
26. Sola A, Chow L, Rogido M. Pulse oximetry in neonatal care in 2005. A comprehensive state of the art review. *An Pediatr (Bare).* 2005;**62**(3):266-81.
 27. Interview P, out Asthma R. of Activity: Chronic Obstructive Pulmonary Disease (1189A) Total Number of Contact Hours: 3 Intended Level of Learner: Intermediate Purpose/Goal: The student will recognize the stages and treatment options for COPD.
 28. Fitzgerald CA, Morse BC, Dente CJ. Pelvic ring fractures: has mortality improved following the implementation of damage control resuscitation? *Am J Surg.* 2014;**208**(6):1083-90; discussion 9-90.
 29. Vaidya R, Scott AN, Tonnos F, Hudson I, Martin AJ, Sethi A. Patients with pelvic fractures from blunt trauma. What is the cause of mortality and when? *Am J Surg.* 2016;**211**(3):495-500.
 30. Wang H, Phillips JL, Robinson RD, Duane TM, Buca S, Campbell-Furtick MB, et al. Predictors of mortality among initially stable adult pelvic trauma patients in the US: Data analysis from the National Trauma Data Bank. *Injury.* 2015;**46**(11):2113-7.
 31. Ziglar MK. Application of base deficit in resuscitation of trauma patients. *Int J Trauma Nurs.* 2000;**6**(3):81-4.
 32. Callaway DW, Shapiro NI, Donnino MW, Baker C, Rosen CL. Serum lactate and base deficit as predictors of mortality in normotensive elderly blunt trauma patients. *J Trauma.* 2009;**66**(4):1040-4.
 33. Penasco Y, Gonzalez-Castro A, Rodriguez-Borregan JC, Llorca J. Base excess, a useful marker in the prognosis of chest trauma in the geriatric population. *Rev Esp Anestesiol Reanim.* 2017;**64**(5):250-6.
 34. Kincaid EH, Miller PR, Meredith JW, Rahman N, Chang MC. Elevated arterial base deficit in trauma patients: a marker of impaired oxygen utilization. *J Am Coll Surg.* 1998;**187**(4):384-92.
 35. Mofidi M, Hasani A, Kianmehr N. Determining the accuracy of base deficit in diagnosis of intra-abdominal injury in patients with blunt abdominal trauma. *Am J Emerg Med.* 2010;**28**(8):933-6.
 36. Toth G, De Bruyne B, Casselman F, De Vroey F, Pyxaras S, Di Serafino L, et al. Fractional flow reserve-guided versus angiography-guided coronary artery bypass graft surgery. *Circulation.* 2013;**CIRCULATIONAHA.113.002740.**
 37. Jeroukhimov I, Ashkenazi I, Kessel B, Gaziants V, Peer A, Altshuler A, et al. Selection of patients with severe pelvic fracture for early angiography remains controversial. *Scand J Trauma Resusc Emerg Med.* 2009;**17**:62.
 38. Jung J, Eo E, Ahn K, Noh H, Cheon Y. Initial base deficit as predictors for mortality and transfusion requirement in the severe pediatric trauma except brain injury. *Pediatr Emerg Care.* 2009;**25**(9):579-81.

Open Access License

All articles published by Bulletin of Emergency And Trauma are fully open access: immediately freely available to read, download and share. Bulletin of Emergency And Trauma articles are published under a Creative Commons license. Mandated authors will be offered CC-BY; all other authors will choose between CC-BY, CC-BY-NC and CC-BY-NC-ND.