

# Isolation, identification and computational studies on *Pseudomonas aeruginosa* sp. strain MPC1 in tannery effluent

Renganathan Senthil\*, Kanagamani Jini Angel, Ravi Malathi, Dhanapal Venkatesan

Department of Bioinformatics, Marudupandiyar Institutions, Thanjavur-613403, TamilNadu, India; Renganathan Senthil – Email: renganathansenthil@gmail.com; Phone: 04362-266791, 265014, 650902; Fax: 265191; \*Corresponding author

Received April 20, 2011; Accepted May 07, 2011; Published May 26, 2011

## Abstract:

A study about isolation, identification and analysis of bacteria in waste water. Here the tannery effluent used as a sample for the entire analysis. A bacterial strain, designated MPC1 was isolated from a waste water sample collected from tannery effluent, Trichy, India and identified using a molecular approach. On the basis of the bacterial 16s rRNA gene sequence phylogeny and comparison of this gene sequence with sequence in RNA sequence database, it is considered that isolate is closely related to members of the *Pseudomonas aeruginosa* Sp. Phylogenetic and molecular evolutionary analyses were conducted using MEGA. Identification of regulatory elements and Transcription Factor with their binding sites in 16S rRNA gene of *Pseudomonas aeruginosa mpc1* was performed using BPROM tool. The sequence of 16s rRNA (*Pseudomonas aeruginosa* sp MPC 1) is submitted to Genbank in NCBI database (Ac.No-JF708077).

**Keywords:** *Pseudomonas aeruginosa* sp. Strain MPC1, isolation, identification, *in silico* analysis, Heavy metal resistant bacteria.

## Background:

The pollution of the environment with toxic heavy metals is spreading throughout the world along with industrial progress. Microorganisms and microbial products can be highly efficient bioaccumulators of soluble and particulate forms of metals especially dilute external solutions. Microbe related technologies may provide an alternative or addition to conventional method of metal removal or metal recovery. Waste water is a significant environmental problem. Wastewaters from the industries and sewage sludge applications have permanent toxic effects to human and the environment [5, 6]. Cadmium (Cd) is nonessential but poisonous for plants, animals, and humans. Cadmium is one of the most toxic pollutants of the surface soil layer, released into the environment by mining and smelting activities, atmospheric deposition from metallurgical industries, incineration of plastics and batteries, land application of sewage sludge, and burning of fossil fuels. Lead (Pb) a major pollutant that is found in soil, water and air is a hazardous waste and is highly toxic to human, animals, plants and microbes [3]. Tanneries, oil refineries and metal industries are causing depletion of surface and ground water quality. The discharge of various sub processes of tanneries like bathing, pickling, tanning, dyeing and fat liquoring may cause water pollution severely. The pollution of a particular water body can always be link to an industry or sewage or agricultural runoff. Likewise, contamination of soil by the release of heavy metals as a result of industrial and anthropogenic activities is a threat to human health and ecosystem. Presence of heavy metals in soil is known to have potential toxic impact on environmental quality and on human health via ground water and

surface water [12, 15]. The genus *Pseudomonas* includes many species of environmental, clinical, agricultural, and biotechnological interest [9]. *Pseudomonas aeruginosa* is a ubiquitous environmental bacterium that is one of the top three causes of opportunistic human infections. A major factor in its prominence as a pathogen is its intrinsic resistance to antibiotics and disinfectants [1, 7].

## Methodology:

### Isolation of bacteria:

The bacteria were identified based on colony characteristics, gram staining methods and by various biochemical tests as given by Bergey's (1984) Manual of Determinative Bacteriology.

### 16s rRNA SEQUENCING:

Isolates are sub cultured on agar plate. A single colony was obtained from a fresh subculture and resuspended in master mixture. A 380-bp fragment of the 16S rRNA (universal primer) used as the target. For 50µl reaction, 1µl of 200mM dNTPs, 5µl of Reaction buffer (Tris with MgCl2) "10pM forward primer RW01 (FP) 5'ACTGGAGGAAGGTGGGGAT-3'" "10pM reverse primer DG74 (RP) 5'AGGAGGTGATCCAACCGCA-3'" 1.2u/µl of Tag Polymerase were used, amplification was carried out in Thermal Cycler PTC 200(MJ Research), and a standard three-step protocol used for all reactions, including 35 cycles for each reaction with an annealing temperature of 55°C and 10-min extension time.



## Conclusion:

The nucleotide frequencies are 0.234 (A), 0.213 (T/U), 0.238 (C), and 0.316 (G). The transition/transversion rate ratios are  $k_1 = 14.587$  (purines) and  $k_2 = 16.07$  (pyrimidines). The statistic is calculated for numbers of synonymous and nonsynonymous substitutions per site, respectively. The variance of the difference was computed using the bootstrap method. There were a total of 75 codons and positions counted by Nei-Gojobori method. The Tajima test statistic was estimated for the 7 strains to identify the segregating sites. The relationship between the two estimates of genetic variation at the DNA level, namely the number of segregating sites: 326 and the average number of nucleotide differences estimated: 4.839163 from pairwise comparison and mutation are occurred in random, is investigated. Upstream region of 16S rRNA gene of *Pseudomonas aeruginosa strain MPC 1*, Promoter position are -10 box at pos. 60 GGTTAGACT; Score 63; -35 box at pos.41 TGGTAA; Score 10. As well as secondary structure and plot have predicted, which reveals the fold information of 16s rRNA gene sequence.

## Acknowledgments:

Authors thank to Marudupandiyar Institutions, Thanjavur and Aravind Medical Research foundation, Madurai and Tannery Industry in Trichy were helped to sequence the bacteria.

## References:

- [1] Bennisar A *et al. BMC Microbiol.* 2010 **10**: 118 [PMID: 20409328]
- [2] Singh BR *et al. Bioinformation* 2009 **3**(9): 375 [PMID: 19759811]

- [3] Edward raja C *et al. International joint symposium on Geodisaster prevention and Geoenvironment in Asia Js-Fukuoka*, 2009
- [4] Gojobori T *et al. J Mol Evol.* 1982 **18**: 360 [PMID: 7120431]
- [5] Hamid AAA *et al. Biotechnology* 2010 **9**(1): 33
- [6] Clarridge JE 3rd. *Clin Microbiol Rev.* 2004 **17**(4): 840 [PMID: 15489351]
- [7] Nath K *et al. J Environ Biol.* 2009 **30**(2): 227 [PMID: 20121023]
- [8] Tamura K *et al. Mol Biol Evol.* 2007 **24**(8): 1596 [PMID: 17488738]
- [9] Kumar P *et al. International Journal of Oceans and Oceanography* 2009 **3**(1): 29
- [10] Kumar S & Dudley J. *Bioinformatics* 2007 **23**(14): 1713 [PMID: 17485425]
- [11] Kumar S *et al. Brief Bioinform.* 2004 **5**: 150 [PMID: 15260895]
- [12] Sahu RK *et al. J Environ Biol.* 2007 **28**(3): 685 [PMID: 18380096]
- [13] Saitou N & Nei M. *Mol Biol Evol.* 1987 **4**: 406 [PMID: 3447015]
- [14] Srivastava S *et al. Bioinformation* 2008 **3**(4): 173 [PMID: 19238242]
- [15] Marcheggiani S *et al. BMC Microbiol.* 2008 **8**: 171 [PMID: 18842132]
- [16] Tamura K *et al. Proc Natl Acad Sci U S A.* 2004 **101**: 11030 [PMID: 15258291]
- [17] Tamura K & Nei M. *Mol Biol Evol.* 1993 **10**: 512 [PMID: 8336541]
- [18] Thompson JD *et al. Nucleic Acids Res.* 1994 **22**: 4673 [PMID: 7984417]
- [19] Yang Z & Kumar S. *Mol Biol Evol.* 1996 **13**: 650 [PMID: 8676739]
- [20] Sun Y *et al. Nucleic Acids Res.* 2010 **38**(22): e205 [PMID: 20929878]

Edited by P Kanguane

Citation: Senthil *et al.* Bioinformation 6(5): 187-190 (2011)  
provided the original author and source are credited.

License statement: This is an open-access article, which permits unrestricted use, distribution, and reproduction in any medium, for non-commercial purposes,

## Supplementary material:

**Table 1:** Results observed from the morphological and biochemical test

S.No	Bacterial isolates	Colony morphology	Appearance	Gram reaction	Motility	Indole Test	Methyl Red Test	VP Test	Citrate test	Starch hydrolysis Test
1	Isolate 1	Light green color	Watery appearance	Rod -	+	-	-	-	+	-
2	Isolate 2	Yellow color	Colloidal	Cocci +	+	+	+	-	+	+
3	Isolate 3	White color	Gummy appearance	Cocci +	+	+	+	-	+	+
4	Isolate 4	White color	Colloidal	Cocci +	+	+	+	-	+	-
5	Isolate 5	White color	Colloidal	Cocci +	+	+	+	-	+	+
6	Isolate 6	Red color	Colloidal	Cocci +	+	-	+	-	+	-

**Table 2:** Maximum composite likelihood estimate of the pattern of nucleotide substitution

	A	T	C	G
A	-	1.23	1.38	26.67
T	1.36	-	22.16	1.83
C	1.36	19.79	-	1.83
G	19.78	1.23	1.38	-