



HHS Public Access

Author manuscript

Nat Commun. Author manuscript; available in PMC 2014 July 22.

Published in final edited form as:

Nat Commun. 2014 ; 5: 3156. doi:10.1038/ncomms4156.

Integrated Analysis of Germline and Somatic Variants in Ovarian Cancer

Krishna L. Kanchi^{1,*}, Kimberly J. Johnson^{1,2,*}, Charles Lu^{1,*}, Michael D. McLellan¹, Mark D.M. Leiserson³, Michael C. Wendl^{1,4,5}, Qunyuan Zhang^{1,4}, Daniel C. Koboldt¹, Mingchao Xie¹, Cyriac Kandoth¹, Joshua F. McMichael¹, Matthew A. Wyczalkowski¹, David E. Larson^{1,4}, Heather K. Schmidt¹, Christopher A. Miller¹, Robert S. Fulton^{1,4}, Paul T. Spellman⁶, Elaine R. Mardis^{1,4,7}, Todd E. Druley^{4,8}, Timothy A. Graubert^{7,9}, Paul J. Goodfellow¹⁰, Benjamin J. Raphael³, Richard K. Wilson^{1,4,7}, and Li Ding^{1,4,7,9,#}

¹The Genome Institute, Washington University in St. Louis, MO 63108, USA

²Brown School Master of Public Health Program, Washington University in St. Louis, St. Louis, MO 63130, USA

³Department of Computer Science, Brown University, Providence, RI 02912, USA

⁴Department of Genetics, Washington University in St. Louis, MO 63108, USA

⁵Department of Mathematics, Washington University in St. Louis, MO 63108, USA

⁶Oregon Health & Science University, Portland, OR 97239, USA

⁷Siteman Cancer Center, Washington University in St. Louis, MO 63108, USA

⁸Department of Pediatrics, Washington University in St. Louis, MO 63108, USA

⁹Department of Medicine, Washington University in St. Louis, MO 63108, USA

¹⁰The Ohio State University Comprehensive Cancer Center, The Ohio State University, Columbus, OH 43210, USA

Abstract

We report the first large-scale exome-wide analysis of the combined germline-somatic landscape in ovarian cancer. Here we analyze germline and somatic alterations in 429 ovarian carcinoma cases and 557 controls. We identify 3,635 high confidence, rare truncation and 22,953 missense

Users may view, print, copy, download and text and data- mine the content in such documents, for the purposes of academic research, subject always to the full Conditions of use: http://www.nature.com/authors/editorial_policies/license.html#terms

#Corresponding Author: Li Ding, Ph.D, The Genome Institute, Division of Oncology, Department of Medicine, Washington University School of Medicine, St. Louis, MO 63108, ding@genome.wustl.edu.

*These authors contributed equally to this work.

COMPETING FINANCIAL INTERESTS

The authors declare no competing financial interests.

CONTRIBUTIONS

L.D. and R.K.W. jointly supervised research. L.D., K.L.K., K.J.J., C.L., M.D.M., M.D.M.L., C.K., M.A.W., J.F.M., D.C.K., C.A.M., P.T.S., and B.J.R. analyzed the data. M.C.W. and Q.Z. performed statistical analysis. K.L.K., C.L., J.F.M., M.D.M., M.A.W., and L.D. prepared figures and tables. R.S.F. performed experiments. E.R.M. and D.E.L. contributed analysis tools. L.D., K.J.J., T.A.G., P.J.G., T.E.D., and B.J.R. conceived and designed the experiments. L.D. and K.J.J. wrote the manuscript. KLK, KJJ, CL, MDM contributed equally but due to restrictions on the number of first authors only KLK, KJJ, and CL are denoted as such.

variants with predicted functional impact. We find germline truncation variants and large deletions across Fanconi pathway genes in 20% of cases. Enrichment of rare truncations is shown in *BRCA1*, *BRCA2*, and *PALB2*. Additionally, we observe germline truncation variants in genes not previously associated with ovarian cancer susceptibility (*NF1*, *MAP3K4*, *CDKN2B*, and *MLL3*). Evidence for loss of heterozygosity was found in 100% and 76% of cases with germline *BRCA1* and *BRCA2* truncations respectively. Germline-somatic interaction analysis combined with extensive bioinformatics annotation identifies 237 candidate functional germline truncation and missense variants, including 2 pathogenic *BRCA1* and 1 *TP53* deleterious variants. Finally, integrated analyses of germline and somatic variants identify significantly altered pathways, including the Fanconi, MAPK, and MLL pathways.

INTRODUCTION

Ovarian cancer is diagnosed in ~22,000 women annually in the United States. The average five year survival is relatively poor at ~43%¹, which is primarily due to late-stage diagnosis. It is currently estimated that 20–25% of women have an inherited germline mutation that predisposes them to ovarian cancer.^{2,3} New strategies for the prevention and control of ovarian cancer will rely on a thorough understanding of the contributing genetic factors both at the germline and somatic levels.

High throughput sequencing technologies are rapidly expanding our understanding of ovarian cancer biology by providing comprehensive descriptions of genetic aberrations in tumors.⁴ The ability to rapidly sequence individual tumor and normal genomes allows for efficient discovery of candidate cancer-causing events and such work is already transforming risk assessment, diagnosis, and treatment. For example, targeted sequencing of 21 tumor suppressor genes in 360 cases of ovarian, peritoneal, fallopian tube, and synchronous ovarian/endometrial carcinomas recently revealed that 24% of cases harbored germline loss of function mutations in 1 of 12 genes: *BRCA1*, *BRCA2*, *BARD1*, *BRIP1*, *CHEK2*, *MRE11A*, *MSH6*, *NBN*, *PALB2*, *RAD50*, *RAD51C*, and *TP53*.³ In a different study, The Cancer Genome Atlas (TCGA) consortium analyzed somatic alterations in 316 serous ovarian carcinomas, identifying recurrent somatic *TP53* mutations in nearly all cases (96%) and finding recurrent somatic mutations in *NF1*, *BRCA1*, *BRCA2*, *RBI*, and *CDK12* in a minority of cases.⁴ Such work is deepening our understanding of genes involved in ovarian cancer.

Cancer genomics studies have most often focused on independent analyses of either somatic or germline mutations. However, studies that perform sequencing of matched tumor and normal samples have the advantage that data from the somatic and germline genomes can be ascertained and integrated to build a fuller picture of each genome's contribution to disease. In addition, the rapidly growing number of publicly available exome datasets from non-cancer populations now facilitates rare germline susceptibility variant discovery.

Here we describe the somatic and germline mutation spectrum in the tumor and normal exome data from 429 TCGA serous ovarian cancer patients. To identify likely pathogenic variants, we compare the frequency of germline mutations to those from a large control dataset of sequences of post-menopausal women from the Women's Health Initiative Exome

Sequencing Project (WHISP). We identify several novel candidate germline predisposition variants in known ovarian genes (e.g., *BRCA1*, *BRCA2*, *ATM*, *MSH3*, and *PALB2*) as well as several genes not previously associated with ovarian cancer (e.g., *ASXL1*, *RBI*, *NF1*, *CDKN2A*, and *EXO1*). We also characterize patterns of loss of heterozygosity in tumor suppressor genes, including *BRCA1*, *BRCA2*, *BRIP1*, *ATM*, *CHEK2*, and *PALB2*, and identify significantly mutated pathways, including Fanconi anemia, MAPK, and MLL. These results provide a foundation for future functional and clinical assessment of susceptibility variants in ovarian cancer.

RESULTS

Clinical Characteristics of samples

Of the 429 TCGA cases in this analysis, 90.2% were Caucasian ($n=387$), 4.9% were African American ($n=21$), 3.5% were Asian ($n=15$), and 0.5% ($n=2$) were American Indian/Alaska Native. Patients were diagnosed between 26–89 years (mean 59.4 ± 11.8 years), frequently at late stage (93% at stages 3–4), and 50.8% were deceased at the time of TCGA sample procurement (Table 1). Nineteen of twenty-three cases with unknown ethnicity information were assigned Caucasian ($n=17$) and African ancestry ($n=2$) using principal components analysis (Supplementary Fig. 1). We performed systematic germline variant and somatic mutation analyses for the sample set, as illustrated in Fig. 1.

Data for 614 samples from the National Heart Lung and Blood Institute (NHBLI) Women's Health Initiative Exome Sequencing Project (WHISP) was used for comparison of genetic variants to TCGA ovarian cancer cases. After extensive quality checks (Methods), 557 Caucasians with an average age of $63.3 \text{ years} \pm 7.8 \text{ years}$ (range 50–79 years) were selected as controls for downstream ovarian susceptibility variant analysis (Supplementary Data 1).

Somatic mutations and significantly mutated genes

We analyzed somatic mutations in 429 ovarian cancer cases. Of these, 142 were new TCGA cases and 287 cases were previously reported⁴; the remaining twenty-nine cases reported in that study⁴ did not meet our coverage requirement ($\geq 20\times$ coverage for at least 50% of target exons) and were excluded from this analysis. The average exome-wide coverage for the entire sample set was 68.1X with 99.5X and 96.5X average coverages for *BRCA1* and *BRCA2*, respectively. We identified 11,479 somatic mutations in the 142 new TCGA cases. All of these mutations were manually reviewed, resulting in a total of 27,280 mutations in 429 cases (Fig. 1 and Supplementary Data 2 and 3). After removing genes with low or no RNA expression evidence from RNA-seq data, the significantly mutated genes (SMGs) identified by MuSiC⁵ include those previously reported: *TP53*, *NF1*, *RBI*, *CDK12(CRKRS)*, and *BRCA1*⁴, as well as the new SMG, *KRAS* (Supplementary Table 1). *BRCA2* and *RB1CC1* were near significance. We also identified 4 *NRAS* mutations, 3 *NF2* mutations, and 3, 8, and 10 mutations in the known tumor suppressor genes: *ATR*, *ATM*, and *APC*, respectively. Somatic truncation mutations were also observed in histone modifier genes including: *ARID1A*, *ARID1B*, *ARID2*, *SETD2*, *SETD4*, *SETD6*, *JARID1C*, *MLL*, *MLL2*, and *MLL3* as well as the DNA excision repair gene *ERCC6* (Supplementary Data 3).

Germline variant landscapes and significant germline events

We identified germline truncation variants (nonsense, nonstop, splice site, and frameshift indels) in these 429 matched tumor-normal cases using multiple algorithms.^{6–8} After removal of common variants, reference sequence errors, and recurrent artifacts, a total of 3,635 high confidence, rare (<1% population minor allele frequency) germline truncation variants were identified in 2,214 genes, 115 of which are in 40 known cancer genes (Fig. 1, Supplementary Fig. 2, Supplementary Data 4 and Methods).⁹ These 115 variants were validated using genomic DNA or a source of whole genome amplified DNA that differed from that used for discovery (Supplementary Data 5). We used several approaches to identify known and potentially pathogenic germline missense variants in the Caucasian subset (Table 1, $n = 387$). Specifically, a total of 22,953 missense variants in 3,637 genes were predicted to be functionally deleterious by Condel¹⁰ and also had population minor allele frequencies (MAFs) <1% in Caucasian data from the 1000 Genomes, and the current cohorts (TCGA ovarian cancer cases and WHISP exome controls) (Fig. 1, Supplementary Data 6, and Supplementary Fig. 3). After limiting our analyses to genes with an average expression RPKM >0.5 (Methods), we identified 17,348 missense variants in a total of 2,810 genes in this subset. We processed on 557 WHISP samples using the same software tools and filtering strategies and identified 7,889 rare (<1% minor allele frequency in the population and cohort) truncation variants and 30,335 rare missense variants defined as functional by Condel and in expressed genes (Supplementary Data 7 and 8).

Finally, although we performed a genome-wide germline copy number analysis using SNP array data, our manual review of the results indicated many false positives with very few passing our review criteria. Therefore, we focused our analysis of copy number alterations on *BRCA1*, *BRCA2*, and *TP53*, coupled with extensive manual review. Here, three high confidence germline deletion events in *BRCA1* were identified in three cases (TCGA-36-2539, TCGA-31-1959, and TCGA-23-1028) (Fig. 2). Two cases (TCGA-31-1959 and TCGA-23-1028) developed ovarian cancer at younger ages (50 and 43 years, respectively); information regarding age of diagnosis for TCGA-36-2539 was not available.

We used a right tailed CAST¹¹ burden test $CAST_{\text{greater}}$ (personal communication, Qunyuan Zhang) to evaluate expressed genes (Methods) having significant enrichment of rare, potentially pathogenic missense variants in the TCGA Caucasian exomes versus the WHISP control group and, the test identified 24 genes that had significant enrichment ($P < 0.0002$, $CAST_{\text{greater}}$). As expected, *BRCA1* is one of the most significant genes on the list ($P = 1.40 \times 10^{-6}$, $CAST_{\text{greater}}$). A total of 9 unique *BRCA1* rare missense variants were detected in this ovarian cancer cohort; this list included two known pathogenic missense variants (R1699W and G1788V) and three singletons (V772A, L668F, and P1637L). It also included one known ovarian susceptibility gene (*FANCM*; $P = 4.04 \times 10^{-6}$, $CAST_{\text{greater}}$) as well as three cancer genes (*ARID1A*, *EGFR*, and *DNMT1*), not previously implicated in ovarian cancer (Supplementary Data 6 and 9). *ARID1A*, frequently mutated in endometrial cancer¹² and *EGFR*, a prominent oncogene involved in lung cancer¹³ and glioblastoma¹⁴, harbored 10 and 5 rare (<1% MAF) unique missense variants in this ovarian sample set, respectively. Several other known cancer genes (e.g., *CREBBP*, *ASXL1*, *EZH2*, and *BRIP1*) were also

found to be in the top 100 and with $P_{\text{CAST,greater}} < 0.0015$. The significance of other top genes such as *EEF2K* requires additional investigation using larger sample sets.

We next focused on comparison of rare germline truncations in cancer genes between TCGA Ovarian cases and the WHISP control set. Three known ovarian cancer susceptibility genes were significant at the right tailed CAST test $p < 0.05$ as a threshold (*BRCA1* ($P = 2\text{E-}08$), *BRCA2* ($P = 8.89\text{E-}06$), *PALB2* ($P = 0.042$)) and two other known ovarian cancer susceptibility genes were among the highest ranked genes although they did not reach significance (*CHEK2* ($P = 0.11$), and *BRIP1* ($P = 0.11$)) (Supplementary Table 2). A total of 66 cases had truncations in one of these genes (Supplementary Data 4 and 5). It is worth noting that we have identified truncation mutations in *USP6*, *ROPNIL*, and *RYRI*, although their involvements in cancer are unclear. In addition, three truncation variants (T1222fs, Q645*, and L258fs) were detected in *BLM* that has recently been linked to familial breast cancer.¹⁵ Q645* and L258fs were previously reported in BLMbase (<http://bioinf.uta.fi/BLMbase/>). The distribution of germline and somatic mutations in these genes is shown in Fig. 3. It is interesting to note that 11 cases had germline truncation variants in multiple cancer genes, including two cases with *BRCA1* and *BRCA2* variants (diagnosis ages 49 and 55 years), one case with *BRCA2* and *ERCC3* variants, one with *PALB2* and *ATM* variants, and one with *BLM* and *FANCD2* truncation variants. Finally, five cases had germline truncation variants in other genes on the cancer gene list, including: *ERCC2* ($n=1$), *TET2* ($n=1$), *FANCD2* ($n=2$), and *NF1* ($n=1$) while one case had a germline mutation in *RAD51B* which has recently been linked to breast cancer susceptibility¹⁶ and whose family members (*RAD50*, *RAD51C*, *RAD51D*) have previously been implicated in ovarian cancer susceptibility.¹⁷

When we combined missense and truncation variants in cancer genes for burden testing, known cancer susceptibility genes were among the most significant genes on the list (*BRIP1*^{3,18} and *BRCA1*). In addition, other established/suspected ovarian/breast cancer susceptibility genes were significant, including *BRCA2*², and *NF1*¹⁹; novel genes such as *ASXL1*, frequently mutated in myelodysplastic syndromes²⁰, myeloproliferative neoplasms²¹, and AML²²; *SETD2*, involved in clear cell renal cell carcinoma²³; and *MAP3K1*, a newly discovered breast cancer gene^{24,25} (Supplementary Data 10).

Germline variants that have been detected as somatically mutated in cancer might signal functional relevance of these variants. We compared our identified germline truncation and missense variants to those present in the COSMIC and OMIM databases to determine whether any were reported in other studies. Of the 3,635 exome-wide truncation variants, 84 and 10 germline variants matched precisely or within ± 5 amino acids to reported variants in COSMIC and OMIM, respectively (Supplementary Data 11). Further analysis of 535 missense variants from cancer genes, using the same criteria applied for truncations, identified 35 and 14 missense events in COSMIC and OMIM, respectively (Supplementary Data 11). For example, the *ASXL1* germline variant G1397S that we identified in 6 of 387 ovarian cancer cases versus 2 of 557 WHISP non-cases and the *ASXL1* germline variant G643V identified in 1 of 387 cases vs. 0 of 557 WHISP non-cases have previously been found to be somatically mutated in hematologic malignancies.^{26,27} Although there was not an exact match of the germline variant P333L in *TET2* in COSMIC (observed in 1 of 387

cases vs. 0 of 557 WHISP non-cases), a somatic frameshift mutation, P333fs, was reported by Metzeler *et al.*²⁸ Another kinase domain germline variant, D837N, in *EGFR* was absent in WHISP controls but found in 5/387 ovarian cancer cases with a position matching a reported somatic mutation (D837G) in COSMIC.²⁹

Germline and somatic interactions in ovarian cancer

Since familial cancer predisposition genes are also often somatically mutated in non-familial cases³⁰, we examined previously characterized somatic SMGs (and *BRCA2*) that met our expression criteria for putative germline functional variants (truncation and predicted deleterious missense) in the germline data of ovarian cancer cases. As expected, a high frequency of germline truncation variants was observed in *BRCA1* ($n=32$) and *BRCA2* ($n=25$). We observed one germline truncation variant in *NF1* (D290fs) in one case (age of diagnosis: 39 years). We similarly investigated somatically mutated protein tyrosine phosphatases and identified 8 germline truncation events in 4 genes (*PTPN13*, *PTPRM*, *PTPRR*, and *PTPRH*). Notably, 4 truncation events (two H942fs, one R199fs, and one T79fs) were found in *PTPRH*, a gene not previously linked to ovarian cancer (Fig. 3). Analysis of germline truncations in somatically mutated chromatin modifier genes also identified truncations in *SETD4* (Y129fs), *SETD6* (M264fs), *MLL3* (e14-2), *SMC5* (Q810fs), and *SMC6* (Y954*). This suggests a potential role for histone modifiers in ovarian susceptibility and motivates further study. Predicted functional germline missense variants having low frequencies were detected in several somatic SMGs, including *BRCA1* (germline missense $n=27$), *BRCA2* ($n=13$), *NF1* ($n=8$), *RBI* ($n=3$), and *TP53* ($n=1$) (Supplementary Table 3). The two patients having a germline V2148D variant in *NF1*, developed ovarian cancer at age 36 and 45 years.

We further investigated the interplay between germline variants (truncation and missense) and somatic mutations in ovarian cancer, discovering 18 patients with germline truncation variants and somatic mutations in the same gene (Supplementary Table 4). For instance, a patient with a germline frameshift mutation (M723fs) in *PALB2* also harbored a somatic nonsense mutation (Q378*) and another patient with a germline nonsense variant (Q153*) in *CDK5RAP1* acquired a somatic splice site mutation in that gene (e9-2). We also detected 8 patients with both germline missense and somatic mutations from the same cancer gene. This list includes 2 patients with *BRCA1* (Germline: R1347G and S1512I; somatic: E111* and G813fs), 1 *NF1* (germline: A2644G; somatic: I85fs), and 1 *TP53* (germline: G334R; somatic: P177R).

We investigated LOH in tumor samples for 535 missense variants in cancer genes and 2,214 genes having germline truncation variants (3,635) and found a total of 732 truncation variants (63 in cancer genes) that displayed LOH in the tumor samples (>20% increase of VAF over normal was used for defining LOH, considering the average 77% purity of the ovarian tumor cohort, false discovery rate = 22%, Supplementary Fig. 5 and Methods), suggesting their potential roles in ovarian cancer susceptibility (Figure 4a and 4b and Supplementary Data 12). Most notably, we observed at least a 20% increased VAF for 30/32 truncation mutations in *BRCA1* (all 32 having increased VAFs) and 13/25 in *BRCA2* (19 having increased VAFs) in the tumor samples when compared to the paired germline

samples (Figure 4c, and 4d). In *BRCA1*, 13 LOH events were associated with a loss of one copy in tumor (copy number segmentation mean = 1.5), while 9 LOH events were associated with a single copy number loss for *BRCA2*. We also identified 14 *BRCA1* and 4 *BRCA2* copy number neutral LOH events in tumor samples ($1.5 < \text{copy number segmentation mean} \leq 2.5$). A small number of cases carried germline truncation variants with clear evidence of somatic LOH (loss of the wild-type allele) in the tumor samples occurring in genes involved in cell-cycle checkpoint, Fanconi/DNA-repair pathways (e.g., *ATM*, *BRIP1*, *CHEK2*, *FANCA*, and *MSH3*), phosphatases (*PTPRH* and *PTPRM*), and a putative prostate cancer susceptibility gene, *ELAC2* (Figure 4e and Supplementary Data 12). This evidence suggests several additional genes may be associated with ovarian cancer susceptibility.

We examined LOH patterns indicating retained germline missense variants in *BRCA1*. Here we identified two known pathogenic missense variants, G1788V and R1699W³¹ (Supplementary Figure 4); *R1699W* has VAFs of 42% and 79% and G1788V has VAFs of 57% and 98% in the germline and tumor samples, respectively. For one variant of unknown significance (VUS), S1521I, evidence indicating loss of the variant allele in the tumor was present in 3/3 cases, suggesting that S1521I is not pathogenic, in agreement with the BIC classification. Evidence of LOH was inconsistent for R1347G and R841W with 2/6 and 1/4 cases demonstrating LOH respectively. Three VUS (V772A, P1637L, L668F) identified in single cases showed LOH. The case with the V772A in *BRCA1* was diagnosed with ovarian cancer at age of 49 years, however this case also carried a *BRCA1* truncation variant. The case with the V1637L variant in *BRCA1* also had a truncation in *BRCA2* and V1637L has previously been predicted to be functionally neutral.³² For *L668F* that occurred in one ovarian cancer case and was not observed in the WHISP dataset no other truncation mutations were observed. None of the *BRCA2* missense variants were classified as clinically important in the BIC *BRCA2* database.^{31,33} Evidence of LOH for retaining some germline *BRCA2* missense variants (S1172L, T2088I, K2434T, and A2951T) was observed (Figure 4d, Supplementary Figure 4, and Supplementary Data 13). **The case** harboring K2434T in *BRCA2* was diagnosed at age of 37 years, however, further work is needed to confirm the functional relevance of such rare germline variants. We expanded our LOH analysis for all rare missense variants across cancer genes (Methods) and identified a total of 114 instances having a greater than 20% increase of VAF in the tumor compared to the germline (Figure 4d and Supplementary Data 13).

We further employed germline-somatic interaction analyses and extensive bioinformatics annotations to identify truncation and missense variants with high likelihood of having functional relevance. Specifically, we examined five aspects of each germline variant (3,635 truncations and 535 missense): pfam annotation, COSMIC/OMIM proximity match, LOH status, somatic SMG status, and somatic mutation in the same gene. When limiting our candidates to variants meeting at least two of the five criteria, the numbers of variants with putative function decreased to 302 truncation and 56 missense events, respectively. In addition, we limited our high confidence variants to genes expressed in ovarian cancer (RSEM (RNA-Seq by Expectation-Maximization) > 0.5) and those that had a lower frequency in cases than WHISP non-cases thereby obtaining 222 putative functional variants

(181 truncations and 41 missense) (Table 2 and Supplementary Data 14). After removing variants suspected to be non-pathogenic based on previous published findings (*ATM* F1463C³⁴, *BRCA1* L668F and P1637L³², *PALB2* H1170Y³⁵, *SMO*³⁶ and *TSC2*^{37,38}), the missense list includes variants from several genes including the two known pathogenic *BRCA1* variants (G1788V and R1699W), four *BRIP1* variants, three *ATM* variants, four NF1 variants, and one *TP53* variant previously identified in breast cancer³⁹ (Table 2). Notably some of the cases with variants identified through this analysis also had truncation variants in known ovarian cancer predisposition genes suggesting an alternative explanation or interacting risk alleles. Our integrated analysis of germline and somatic variants identifies a set of known ovarian cancer susceptibility variants and prioritizes a set of variants without previous association with ovarian cancer susceptibility.

Significant pathways in ovarian cancer

We performed pathway analysis using PathScan statistical test⁴⁰ including both germline truncation variants and somatic mutations and identified the KEGG Fanconi Anemia DNA repair pathway as significant ($P = 4.2E-08$) along with MAPK, Cell cycle, and TP53 signaling pathways (Fig. 5a and Supplementary Data 15). RB/RAS pathways were previously reported to be involved in ovarian cancer.⁴ Germline and somatic mutations in the Fanconi Anemia pathway affected a total of 40 genes in 37% (157/429) cases. Additional rare mutations detected but not shown occurred in *APITD1*, *EME1*, *ERCC1*, *HES1*, *MLH1*, *PMS2CL*, *POLK*, *POLI*, *RAD51*, *REV3L*, *RMI1*, *RPA1*, *RPA2*, *RPA4*, *TELO2*, *TOP3A*, *TOP3B*, *USP1*, and *WDR48*.

We used HotNet⁴¹ to identify subnetworks of a genome-scale protein-protein interaction network containing genes with significant numbers of somatic and germline variants. HotNet identified two such subnetworks ($P < 0.01$): one consisting of DNA repair and Fanconi Anemia genes (Fig. 5a and Supplementary Table 5) that is mutated in 33.1% (142/429) of samples. We combined Fanconi genes from PathScan and HotNet analyses and determined that 40.8% (175/429) of ovarian cancer patients in this study have germline/somatic defects in the Fanconi pathway. As expected, we found that germline alterations in 47 Fanconi genes are significantly enriched in younger patients by a Wilcoxon Rank-Sum test (427 tumors with data, P -value=1.1878E-05, Fig. 5b).

A second subnetwork containing somatic mutations and germline variants in *EGFR*, *ERBB2*, *ERBB3*, and other genes is shown in (Fig. 5c and Supplementary Data 16). The frequency of somatic mutations in each of these genes is low ($< 1.3\%$), as is the frequency of germline variants ($< 0.3\%$). The significance of this subnetwork is thus derived from the combined analyses of somatic mutations, germline variants, and biological interactions among these proteins. Using more permissive parameters, HotNet identifies two additional subnetworks (See Methods), including a subnetwork containing *MLL*, *MLL3*, and *SETD1A* (Fig. 5d and Supplementary Data 16). Mutations in these histone methyltransferases have been previously reported in leukemias⁴², breast cancer²⁴, and renal carcinomas⁴³, but have not been widely reported in ovarian carcinoma.

DISCUSSION

We report here the first large-scale exome-wide analysis of the combined germline-somatic landscape of ovarian cancer. We used several analytic approaches to sift through millions of germline variants to discover both known and candidate cancer susceptibility genes and loss-of-function truncation and missense variants. As expected, we found enrichment of germline presumed loss-of-function truncation variants in the known ovarian cancer susceptibility genes, *BRCA1*, *BRCA2*, *BRIP1*, *CHEK2*, and *PALB2*. The average diagnosis age for patients with germline *BRCA1/BRCA2* truncation variants was 53.4 years, significantly younger than either patients with somatic *BRCA1/BRCA2* mutations (61.8 years, $n=32$, $P = 0.0002$, t-test) or the entire cohort (59.4 years, $n=427$, $P = 5.73E-06$, t-test). Interestingly, patients harboring germline *BRCA1/BRCA2* alterations have an average of 1.87 somatic mutations ($n=60$) in 127 SMGs from MuSiC analysis of 12 TCGA cancer types⁴⁴ (curated from doi:10.1038/nature12634) which is markedly lower than patients with somatic *BRCA1/BRCA2* mutations (2.84 somatic mutations, $n=32$, $P = 2.1E-05$ t-test). Further, likely loss-of-function truncation variants were detected in several other genes/gene family members and syndromes (NF1) that have previously been associated with breast and/or ovarian cancer susceptibility including *BLM*¹⁵, *FANCD2*⁴⁵, *NFI*^{19,46}, *RAD51B*^{47,48}, *FANCA*⁴⁹, *FANCB*, *FANCL*, *FANCM*, *ATRIP*, and *ATR*⁵⁰. Notably, loss-of-function variants were dispersed across a set of genes, in particular, previously reported members of the Fanconi pathway⁵¹ and some novel members.

The identification of pathogenic missense variants in high-throughput sequencing data is challenging due to the large number of rare variants of unknown significance and inherent uncertainties associated with *in silico* based functional prediction. To identify a set of known and likely pathogenic missense variants, we used several complementary strategies including LOH, COSMIC/OMIM proximity match, PFAM domain, and case/control allele frequency analyses. We first applied the LOH analysis to germline truncation variants in *BRCA1* and *BRCA2* and a small set of other tumor suppressor genes, demonstrating a strong tendency to induce LOH of the wild-type allele in the tumor. For example, clear evidence for LOH of *BRCA1* wild-type alleles in the tumor was present in virtually all cases, similar to previous reports.^{3,52} Further, our analysis identified two pathogenic missense variants (G1788V, R1699W) as well as three with uncertain pathogenicity (L668F, V772A, P1637L) that demonstrated clear evidence of LOH. However, we note that the single cases with V772A and P1637L variants each had a *BRCA1* truncation variant suggesting an alternative explanation for these findings. LOH was also observed for several *BRCA2* missense variants.

Evidence for pathogenicity was also demonstrated for a number of variants in cancer genes including two pathogenic *BRCA1*, three *ATM*, and four *BRIP1* missense variants that met at least two of the five criteria for classifying candidate pathogenic missense variants. These results emphasize that integration of both somatic and protein domain information can facilitate identification of a set of known and potentially pathogenic missense variants among thousands of rare missense variants that informs functional assessment of variants of unknown significance.

Significance analysis of germline truncation and missense variants nominated a set of genes including *ASXL1*, *MAP3K1*, and *SETD2* as candidate novel ovarian susceptibility genes. COSMIC somatic mutation matches to *ASXL1* germline missense variant (G1397S) coupled with evidence for LOH support a potential role for this variant in ovarian cancer susceptibility. In addition, common variation in *MAP3K1*, another member of the MAP3K family, has been associated with breast cancer susceptibility⁵³, was recently identified as a target of frequent somatic breast cancer mutations^{24,25}, and was significant based on the burden test.

Pathway and network analyses of the integrated collection of germline and somatic variants revealed pathways with significant enrichment of variants including the Fanconi anemia/DNA repair pathway, MAPK pathway, and histone methyltransferases. In most cases, the individual genes in these pathways are altered rarely by either germline or somatic variants, and it is only through the combined analysis of both types of variants across many genes that the alteration of these pathways becomes apparent. This further emphasizes the extensive genetic heterogeneity in serous ovarian carcinoma, as suggested by the relatively small number of genes found to be recurrently mutated by somatic mutations in TCGA study.⁴

We are mindful of limitations of TCGA and WHISP data for germline analyses and the analysis of rare variants in general including lack of family history information in TCGA cases that would further inform these results, exclusion of women with a prior malignancy that required systemic therapy from the TCGA case set that might lead to an underestimation of the frequency of germline susceptibility alleles in the population, lack of personal cancer history information in WHISP controls, differences in sequencing platforms used to generate the TCGA and WHISP exome sequence data, and detection of rare germline variants that are extremely rare/private and have no pathogenic significance. With respect to differences in sequencing platforms between the case and control datasets, more variants were called in the WHISP data than the TCGA data, which would reduce our ability to detect significantly higher frequencies of rare deleterious germline variants in TCGA cases compared to WHISP controls. In addition, it is worth noting that the WHISP controls were older on average than TCGA cases and were assembled for the purpose of examining genetic susceptibility to non-cancer outcomes. Therefore pathogenic germline variants would most likely be under-represented in this cohort, which would increase our ability to identify pathogenic variants in TCGA OV cancer cases.

In conclusion, this is the first large scale and comprehensive analysis of both germline and somatic exome variants in ovarian cancer. Our exome-wide analysis strongly supports and extends results from previous studies employing candidate gene approaches for discovery of ovarian cancer genes, and is in line with previous reports by identifying Fanconi anemia pathway genes as the most frequent targets of germline and somatic mutations. Our integrated analyses of somatic and germline data indicate additional genes and variants of potential importance in ovarian cancer susceptibility for further investigation. In addition, we emphasize that candidate variants and genes nominated by our study will require extensive experimental functional validation as well as replication in additional ovarian

cancer datasets. Functionally validated variants will have important implications for the development of screening strategies to evaluate ovarian cancer predisposition.

METHODS

Study population

We obtained approval from the database of Genotypes and Phenotypes (dbGaP) to access the exome sequence and clinical data from TCGA ovarian cancer cases for this study (document #3281 Discover germline cancer predisposition variants). We selected a total of 460 ovarian cancer cases (316 cases previously reported⁴ and 144 new ovarian cases) with their germline and tumor DNA sequenced by exome capture followed by next generation sequencing on Illumina or SOLiD platforms. Of the 460 cases, 429 met our inclusion criteria of 50% coverage of targeted exome having at least 20X coverage in both germline and tumor samples. 74% of targets reached 20X coverage for 80% of breadth. Population estimates of allele frequencies were obtained from a control group of 3,505 European individuals from the National Heart, Lung, and Blood Institute (NHLBI) exome data set (<https://esp.gs.washington.edu/drupal/>), and from 379 European, 246 African, 286 ASN, and 181 AMR descent individuals from the 1000 genomes project.⁵⁶ The global minor allele frequencies were obtained from dbSNP release 137, based on the 1000 Genomes phase 1 genotypes for 1094 individuals, released in May 2011.

Ancestry classification using PLINK

TCGA ovarian cancer cases were classified with respect to ancestry using their SNP array data⁴ and the multi-dimensional scaling (MDS) analysis program in PLINK. (<http://pngu.mgh.harvard.edu/~purcell.plink/>, version 1.07). Five clusters were used for MDS analysis. Twenty-three TCGA cases had unknown ethnicity information; we were able to assign ethnicity for 19 of these as Caucasian (n=17) and African American (n=2), respectively, using principal components analysis (Supplementary Fig. 1).

Control cohort

WHISP (Women's Health Initiative Exome Sequencing Project, part of the Women's Health Initiative (WHI)) data for 614 samples were downloaded from dbGaP (dbGaP Study Accession: phs000281.v4.p2), verified for file integrity, and then imported as BAM files into our data warehouse. The WHISP data was collected as part of the NHLBI Exome Sequencing Project that has the objective of detecting genetic variants related to heart, lung, and blood diseases as described at http://www.ncbi.nlm.nih.gov/projects/gap/cgi-bin/study.cgi?study_id=phs000281.v4.p2. Women included in WHISP were a subset of women who were part of the WHI⁵⁷. To minimize batch differences between the ovarian dataset and these controls, we processed imported samples through the same pipeline, including alignment to the GRCh37-lite reference sequence with BWA⁵⁸ v0.5.9 with parameters `-t 4 -q 5` and marking of duplicates by Picard v1.46. SNVs and indels were called using VarScan v2.2.9 (with parameters `--min-coverage 3 --min-var-freq 0.20 --p-value 0.10 --strand-filter 1 --map-quality 10`) with the false-positive filter⁵⁹ and GATK⁶⁰ v5336 (with parameters `-T IndelGenotyperV2 --window_size 300`). Variant calls were restricted to the ~34 Mbp CDS target region.⁴

To remove outliers in data quality, we required that WHISP samples have read mapping rates <80%, duplication rates <40%, and at least 10,000 SNVs called in the target region. The 557 WHISP samples that met these criteria had, on average, mapping rates of ~95%, duplication rates of ~9%, and ~18,000 SNVs called in the target region. 81% of targets have reached 20X coverage for 80% of breadth. These were used as controls in the downstream analysis.

Germline variant calling and filtering

Sequence data from paired tumor and germline samples were aligned independently to NCBI Build 36 of the human reference using BWA 0.5.9 and de-duplicated using Picard 1.29. Germline SNPs and indels were identified in paired BAMs using VarScan2 with the following parameters: min-coverage = 30, min-var-freq = 0.08, normal-purity = 1, p-value = 0.10, somatic-p-value = 0.001, and validation = 1. Additional germline SNPs were identified using Samtools (Version 0.1.7a (revision #599)) and additional germline indels were identified using GATK (Version 1.0 (revision 5336)). All predicted variants were filtered to remove false positives related to potential homopolymer artifacts (variants found in homopolymers having sequence length ≤ 5 were removed), strand specific sequence artifacts, ambiguously mapped data (average mapping quality difference between the reference supporting reads and variant supporting reads ≤ 30), and low quality data at the beginning and end of reads (variants supported exclusively by bases observed in first or last 10% of the reads). Variants having an allele frequency <8% were removed. Initial variant transcript annotation was based on a combined database, including NCBI Refseq (May 2009) and Ensembl (version 54). All variants were additionally annotated using (Version 2.2) of Ensembl variant effect predictor (VEP).⁶¹ Variants that occurred outside of tier 1 (coding exons, canonical splice sites, and RNA genes) and variants that did not change the amino acid sequence were not included in the downstream analysis. Putative variants with translational effect were filtered in the multistep process shown in Supplementary Figure 2 and described below. Variants were filtered if they either could not be mapped uniquely from NCBI build 36 to GRCh37, were protein altering in a rare transcript that was exclusive to either the NCBI or Ensembl database, or if they were nonsynonymous only in transcripts that lacked a valid open reading frame due to internal frame shifts, missing start codons, and/or missing stop codons. In addition, all variants were discarded from genes suspected to have pseudogenes or other paralogs missing from the human reference sequence, such as *PDE4DIP*, *CDC27*, *MUC4*, *DUX4*, and *XPC*. We additionally filtered variants that occurred exclusively in non-coding RNA genes, those that affected only predicted, hypothetical, or olfactory genes, those that had a frequency >1% in the Caucasian population in the NHLBI GO exomes sequence data, those exclusively within a transcript annotated as a pseudogene or processed pseudogene based on Ensembl release (64) annotation downloaded via Biomart, and lastly those that were reported as a validated somatic mutation in the same sample. Sequence data supporting all remaining germline truncational variants were visually examined with the Integrative Genomics Viewer⁶² and any data that appeared to be supported by potential sequencing, amplification, or alignment artifacts were discarded. Additional validated germline variants reported in *BRCA1*, *BRCA2* were recovered, followed by removal (filtering) of any remaining nonsynonymous germline variants that were recurrent at the same position in more than 2% of the cohort (more than 8 samples at

the same position). Finally, for the analysis of significantly mutated genes, genes not typically expressed in ovarian adenocarcinoma tumor samples were filtered if they had an average RPKM ≤ 0.5 . For the RNA-seq based gene expression analysis, we used the Pancan12 per-sample log2-RSEM matrix from doi:10.7303/syn1734155.1. A gene qualified as expressed if it had at least 3 reads in at least 70% of samples. For every gene, the average per-sample RSEM value was calculated across samples from the same tumor-type. The genes that had an average RSEM $< 0.5\%$ were considered to be low expressed genes. Of the 20,239 genes that had an expression value in ovarian cancer, 4,957 were low expressed genes.

Cancer gene list

The cancer gene list (Supplementary Data 17) was comprised of a total of 672 unique genes of interest that included 436 genes from the Sanger Cancer Gene⁹ list (<http://www.sanger.ac.uk/genetics/CGP/Census/> as downloaded on December 1, 2010), 41 uterine and endometrial cancer genes that we previously identified as having recurrent somatic mutations¹², and 50 genes that have been identified in genome wide association studies as containing common cancer susceptibility variants to ovarian or breast cancer (HugeNet, <http://www.cdc.gov/genomics/about/index.htm>). Of note, the 436 genes on the Sanger cancer gene list contained gene clusters (IGH@, IGK@, IGL@). Individual genes from these clusters were extracted. Any genes on the list that represented common fusion products of translocation or any gene that could not be identified based on Ensembl release 58 and the corresponding release of NCBI Refseq from the same time point were excluded. This process resulted in a total of 616 putative cancer related genes.

Validation of truncation variants in cancer genes

We designed validation PCR primers pairs using Primer3 and tailed the sequences with universal forward and reverse primer sites. Primer pairs for PCR were selected to favor products with an optimal size of 200 to 300 bp. (Supplementary Data 19 and 20) Larger or smaller products were allowed to avoid problematic sequences. Alternate sources of WGA-amplified or original source genomic DNA samples from tumor and normal pairs were amplified with PCR using a single primer pair and each individual PCR product was sequenced with BigDye Terminators using universal primers. Products were purified and then loaded on an ABI 3730. Resulting reads were base called using Phred and aligned to genomic sequence representative of the PCR products using Crossmatch. PolyScan⁶³ and PolyPhred⁶⁴ were used to identify SNPs and Indels. Predicted putative rare germline variants were visually reviewed using Consed to determine the exact position and sequence of indel events and eliminate false positives due to data quality, LOH in the tumor sample, artifacts resulting from sequence context, paralog amplification, or WGA or Illumina library generation or sequencing artifacts.

Missense germline variant analysis

Missense germline variants were filtered using the same methods (Supplementary Figure 3) previously described for germline truncations. To minimize the number of variants tied with ancestral origins, only missense germline variants from individuals classified as Caucasian by Plink were used for downstream significance testing. Missense germline variants were

further filtered to retain only those identified as deleterious by the Ensembl implementation of Condel, a software program that employs a weighted approach to calculate the functional impact of missense variants from scores calculated by SIFT⁶⁵ and PolyPhen-2.⁶⁶ We then removed missense germline variants that occurred at >1% frequency in the ovarian cancer cases and followed that by removing germline predicted missense variants that were better classified as somatic variants. Variants with population minor allele frequencies > 1% in NHLBI ESP GO Exomes or 1000 Genomes were also filtered. Remaining sites were annotated using the Ensembl VEP instance of Condel and remaining predicted deleterious variants were retained for burden analysis. Sites were further filtered to only retain expressed variants in cancer genes (as described above). In addition, we have performed internal unbiased validation of all rare variants identified in 11 cases using available whole genome sequencing data that were independently generated. It is worth noting that whole genome sequencing data for 2 cases were generated using the SOLiD platform, furnishing orthogonal validation of the variants discovered using Illumina sequencing data. (Supplementary Data 18)

We applied a modified version of the cohort allelic sums test (CAST)¹¹ to the final list of germline missense variants in the ovarian cancer dataset to determine the statistical significance of deleterious variants in genes that were over-represented in ovarian cases vs. control exomes from the WHISP. A one-tailed CAST test was used to identify only the genes with higher burden frequency in cases than in controls.

Germline copy number alterations analysis

Segmented copy number deletion events were extracted from GISTIC (10.1073/pnas.0710052104) analysis of Affymetrix 6.0 SNP array data for a total of 426 exome sequenced tumor-normal sample pairs with available array genotype data. Matched tumor and normal samples were processed in parallel to identify putative germline copy number variations (CNV) with overlapping deletion segments defined by 8 consecutive probes in both tumor and normal. Potentially truncating CNV deletion events in the 672 cancer-related genes list (Details at <http://goo.gl/zLk8i>) were extracted from the total list. Graphical plots were visually examined to identify and filter suspected artifacts and somatic copy-number events. All CNV deletion events were annotated to identify those overlapping coding exons and those that were intronic, intergenic, or affected UTR exons were removed. Matched tumor-normal exome capture BAMs were examined to identify any heterozygous SNPs refuting germline copy number deletions or, alternatively to identify coverage anomalies supporting the presence of germline deletion events. Finally, individual probe intensities were plotted and reviewed to remove additional artifacts.

Loss of heterozygosity analysis

Loss of heterozygosity (LOH) Analysis was performed by calculating the variant allele frequency of both SNV and short indels using our internally developed tool bam-readcount (<https://github.com/genome/bam-readcount>) for SNVs and Samtools mpileup⁶/VarScan⁷ for indels. Significance testing was done on the basis of generating an approximate empirical distribution of the actual population null distribution using a resampling method (bootstrapping with replacement). We corrected each case for tumor purity using

$$\text{VAF}_{\text{tumor,C}} = \frac{\text{VAF}_{\text{tumor,U}} - (1 - P_{\text{tumor}}) \cdot \text{VAF}_{\text{normal}}}{P_{\text{tumor}}} \quad (1)$$

where $\text{VAF}_{\text{tumor,C}}$ and $\text{VAF}_{\text{tumor,U}}$ are the corrected and uncorrected tumor variant allele fractions, respectively, P_{tumor} is tumor purity, and $\text{VAF}_{\text{normal}}$ is variant allele fraction in the normal. This equation is an algebraic consequence of assuming that foreign variant and reference reads in the tumor are proportional to their corresponding numbers in the normal sample. The distribution converged within 10^8 trials (Supplementary Figure S4) and this, in turn, agreed well with another distribution model obtained by full enumeration of all possible VAF differences within the data set. A threshold of 20, i.e. $P_{\text{tumor}} \times (\text{VAF}_{\text{tumor}} - \text{VAF}_{\text{normal}}) \geq 20\%$, was taken as significant and this threshold incurs a false-positive error rate of roughly $\alpha = 22\%$. The actual error rate may be slightly less because VAF differences above 50 are, strictly speaking, spurious and likely due to contamination in the normal.

Pathway analysis using HotNet

We applied HotNet⁶⁷ to identify subnetworks in a genome-scale protein-protein interaction network, each containing genes with significant numbers of somatic and germline aberrations. HotNet identifies a list of subnetworks, each containing at least s genes, and employs a two-stage statistical test to assess the significance of the list of subnetworks. We used HotNet version 1.1 and an interaction network from iRefIndex⁹⁶⁸ containing 212,746 interactions among 14,384 proteins, using parameter $t = 0.05$ to derive the influence graph. With parameter $\delta = 0.02$, we find 2 subnetworks (Supplementary Table S5), each containing at least 6 genes ($P = 0.0005$). With parameter $\delta = 0.02$, we find 4 subnetworks (Supplementary Data 16), each containing at least 4 genes ($P = 0.1555$).

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by the National Cancer Institute grant R01CA180006 to L.D. and National Human Genome Research Institute grants R01HG005690 to B.J.R. and U54HG003079 to R.K.W.

References

1. Howlader, NNA.; Krapcho, M.; Neyman, N.; Aminou, R.; Altekruse, SF.; Kosary, CL.; Ruhl, J.; Tatalovich, Z.; Cho, H.; Mariotto, A.; Eisner, MP.; Lewis, DR.; Chen, HS.; Feuer, EJ.; Cronin, KA., editors. SEER Cancer Statistics Review, 1975–2009 (Vintage 2009 Populations). National Cancer Institute; Bethesda, MD: Apr. 2012 http://seer.cancer.gov/csr/1975_2009_pops09/, based on November 2011 SEER data submission, posted to the SEER web site
2. Weissman SM, Weiss SM, Newlin AC. Genetic testing by cancer site: ovary. *Cancer J*. 2012; 18:320–7. [PubMed: 22846732]
3. Walsh T, et al. Mutations in 12 genes for inherited ovarian, fallopian tube, and peritoneal carcinoma identified by massively parallel sequencing. *Proc Natl Acad Sci U S A*. 2011; 108:18032–7. [PubMed: 22006311]
4. Integrated genomic analyses of ovarian carcinoma. *Nature*. 2011; 474:609–15. [PubMed: 21720365]

5. Dees ND, et al. MuSiC: Identifying mutational significance in cancer genomes. *Genome research*. 2012; 22:1589–98. [PubMed: 22759861]
6. Li H, et al. The Sequence Alignment/Map format and SAMtools. *Bioinformatics*. 2009; 25:2078–9. [PubMed: 19505943]
7. Koboldt DC, et al. VarScan 2: somatic mutation and copy number alteration discovery in cancer by exome sequencing. *Genome research*. 2012; 22:568–76. [PubMed: 22300766]
8. McKenna A, et al. The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome research*. 2010; 20:1297–303. [PubMed: 20644199]
9. Futreal PA, et al. A census of human cancer genes. *Nat Rev Cancer*. 2004; 4:177–83. [PubMed: 14993899]
10. Gonzalez-Perez A, Lopez-Bigas N. Improving the assessment of the outcome of nonsynonymous SNVs with a consensus deleteriousness score, Condel. *American journal of human genetics*. 2011; 88:440–9. [PubMed: 21457909]
11. Morgenthaler S, Thilly WG. A strategy to discover genes that carry multi-allelic or mono-allelic risk for common diseases: a cohort allelic sums test (CAST). *Mutation research*. 2007; 615:28–56. [PubMed: 17101154]
12. Kandoth C, et al. Integrated genomic characterization of endometrial carcinoma. *Nature*. 2013; 497:67–73. [PubMed: 23636398]
13. Ding L, et al. Somatic mutations affect key pathways in lung adenocarcinoma. *Nature*. 2008; 455:1069–75. [PubMed: 18948947]
14. Comprehensive genomic characterization defines human glioblastoma genes and core pathways. *Nature*. 2008; 455:1061–8. [PubMed: 18772890]
15. Thompson ER, et al. Exome sequencing identifies rare deleterious mutations in DNA repair genes FANCC and BLM as potential breast cancer susceptibility alleles. *PLoS Genet*. 2012; 8:e1002894. [PubMed: 23028338]
16. Thomas G, et al. A multistage genome-wide association study in breast cancer identifies two new risk alleles at 1p11.2 and 14q24.1 (RAD51L1). *Nat Genet*. 2009; 41:579–84. [PubMed: 19330030]
17. Wickramanyake A, et al. Loss of function germline mutations in RAD51D in women with ovarian carcinoma. *Gynecol Oncol*. 2012; 127:552–5. [PubMed: 22986143]
18. Catucci I, et al. Germline mutations in BRIP1 and PALB2 in Jewish high cancer risk families. *Fam Cancer*. 2012; 11:483–91. [PubMed: 22692731]
19. Seminog OO, Goldacre MJ. Risk of benign tumours of nervous system, and of malignant neoplasms, in people with neurofibromatosis: population-based record-linkage study. *Br J Cancer*. 2013; 108:193–8. [PubMed: 23257896]
20. Thol F, et al. Prognostic significance of ASXL1 mutations in patients with myelodysplastic syndromes. *Journal of clinical oncology: official journal of the American Society of Clinical Oncology*. 2011; 29:2499–506. [PubMed: 21576631]
21. Carbuccia N, et al. Mutations of ASXL1 gene in myeloproliferative neoplasms. *Leukemia: official journal of the Leukemia Society of America, Leukemia Research Fund, U K*. 2009; 23:2183–6.
22. Schnittger S, et al. ASXL1 exon 12 mutations are frequent in AML with intermediate risk karyotype and are independently associated with an adverse outcome. *Leukemia: official journal of the Leukemia Society of America, Leukemia Research Fund, U K*. 2013; 27:82–91.
23. Comprehensive molecular characterization of clear cell renal cell carcinoma. *Nature*. 2013; 499:43–9. [PubMed: 23792563]
24. Comprehensive molecular portraits of human breast tumours. *Nature*. 2012; 490:61–70. [PubMed: 23000897]
25. Ellis MJ, et al. Whole-genome analysis informs breast cancer response to aromatase inhibition. *Nature*. 2012; 486:353–60. [PubMed: 22722193]
26. Patnaik MM, et al. Mayo prognostic model for WHO-defined chronic myelomonocytic leukemia: ASXL1 and spliceosome component mutations and outcomes. *Leukemia: official journal of the Leukemia Society of America, Leukemia Research Fund, UK*. 2013
27. Mian SA, et al. Spliceosome mutations exhibit specific associations with epigenetic modifiers and proto-oncogenes mutated in myelodysplastic syndrome. *Haematologica*. 2013

28. Metzeler KH, et al. TET2 mutations improve the new European LeukemiaNet risk classification of acute myeloid leukemia: a Cancer and Leukemia Group B study. *Journal of clinical oncology: official journal of the American Society of Clinical Oncology*. 2011; 29:1373–81. [PubMed: 21343549]
29. Penzel R, et al. EGFR mutation detection in NSCLC--assessment of diagnostic application and recommendations of the German Panel for Mutation Testing in NSCLC. *Virchows Arch*. 2011; 458:95–8. [PubMed: 21057810]
30. Fearnhead NS, Wilding JL, Bodmer WF. Genetics of colorectal cancer: hereditary aspects and overview of colorectal tumorigenesis. *Br Med Bull*. 2002; 64:27–43. [PubMed: 12421723]
31. Szabo C, Masiello A, Ryan JF, Brody LC. The breast cancer information core: database design, structure, and scope. *Hum Mutat*. 2000; 16:123–31. [PubMed: 10923033]
32. Easton DF, et al. A systematic genetic assessment of 1,433 sequence variants of unknown clinical significance in the BRCA1 and BRCA2 breast cancer-predisposition genes. *American journal of human genetics*. 2007; 81:873–83. [PubMed: 17924331]
33. National Human Genome Research Institute. Breast Cancer Information Core, An Open Access On-Line Breast Cancer Mutation Data Base. 2013
34. Offit K, et al. Rare variants of ATM and risk for Hodgkin's disease and radiation-associated breast cancers. *Clin Cancer Res*. 2002; 8:3813–9. [PubMed: 12473594]
35. Hellebrand H, et al. Germline mutations in the PALB2 gene are population specific and occur with low frequencies in familial breast cancer. *Hum Mutat*. 2011; 32:E2176–88. [PubMed: 21618343]
36. Wang XD, et al. Mutations in the hedgehog pathway genes SMO and PTCH1 in human gastric tumors. *PLoS One*. 2013; 8:e54415. [PubMed: 23349881]
37. Jozwiak J, Jozwiak S, Grzela T, Lazarczyk M. Positive and negative regulation of TSC2 activity and its effects on downstream effectors of the mTOR pathway. *Neuromolecular medicine*. 2005; 7:287–296. [PubMed: 16391386]
38. Nellist M, et al. Distinct effects of single amino-acid changes to tuberlin on the function of the tuberlin-hamartin complex. *European journal of human genetics*. 2004; 13:59–68. [PubMed: 15483652]
39. Rath MG, et al. Prevalence of germline TP53 mutations in HER2+ breast cancer patients. *Breast cancer research and treatment*. 2013
40. Wendl MC, et al. PathScan: a tool for discerning mutational significance in groups of putative cancer genes. *Bioinformatics*. 2011; 27:1595–602. [PubMed: 21498403]
41. Vandin F, Upfal E, Raphael BJ. De novo discovery of mutated driver pathways in cancer. *Genome research*. 2012; 22:375–85. [PubMed: 21653252]
42. Thirman MJ, et al. Rearrangement of the MLL gene in acute lymphoblastic and acute myeloid leukemias with 11q23 chromosomal translocations. *The New England journal of medicine*. 1993; 329:909–14. [PubMed: 8361504]
43. Duns G, et al. Histone methyltransferase gene SETD2 is a novel tumor suppressor gene in clear cell renal cell carcinoma. *Cancer research*. 2010; 70:4287–91. [PubMed: 20501857]
44. Kandoth C, et al. Mutational landscape and significance across 12 major cancer types. *Nature*. 2013; 502:333–9. [PubMed: 24132290]
45. Barroso E, et al. FANCD2 associated with sporadic breast cancer risk. *Carcinogenesis*. 2006; 27:1930–7. [PubMed: 16679306]
46. Seminog OO, Goldacre MJ. Risk of benign tumours of nervous system, and of malignant neoplasms, in people with neurofibromatosis: population-based record-linkage study. *British journal of cancer*. 2013; 108:193–8. [PubMed: 23257896]
47. Golmard L, et al. Germline mutation in the RAD51B gene confers predisposition to breast cancer. *BMC Cancer*. 2013; 13:484. [PubMed: 24139550]
48. Wickramanyake A, et al. Loss of function germline mutations in RAD51D in women with ovarian carcinoma. *Gynecologic oncology*. 2012; 127:552–5. [PubMed: 22986143]
49. Solyom S, et al. Screening for large genomic rearrangements in the FANCA gene reveals extensive deletion in a Finnish breast cancer family. *Cancer Lett*. 2011; 302:113–8. [PubMed: 21236561]

50. Durocher F, et al. Mutation analysis and characterization of ATR sequence variants in breast cancer cases from high-risk French Canadian breast/ovarian cancer families. *BMC Cancer*. 2006; 6:230. [PubMed: 17010193]
51. Pennington KP, Swisher EM. Hereditary ovarian cancer: beyond the usual suspects. *Gynecologic oncology*. 2012; 124:347–53. [PubMed: 22264603]
52. Rzepecka IK, et al. High frequency of allelic loss at the BRCA1 locus in ovarian cancers: clinicopathologic and molecular associations. *Cancer Genet*. 2012; 205:94–100. [PubMed: 22469508]
53. Easton DF, et al. Genome-wide association study identifies novel breast cancer susceptibility loci. *Nature*. 2007; 447:1087–93. [PubMed: 17529967]
54. Fokkema IF, et al. LOVD v.2.0: the next generation in gene variant databases. *Hum Mutat*. 2011; 32:557–63. [PubMed: 21520333]
55. Stenson PD, et al. The Human Gene Mutation Database: building a comprehensive mutation repository for clinical and molecular genetics, diagnostic testing and personalized genomic medicine. *Hum Genet*. 2013
56. Abecasis GR, et al. An integrated map of genetic variation from 1,092 human genomes. *Nature*. 2012; 491:56–65. [PubMed: 23128226]
57. Hays J, et al. The Women's Health Initiative recruitment methods and results. *Ann Epidemiol*. 2003; 13:S18–77. [PubMed: 14575939]
58. Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics*. 2009; 25:1754–60. [PubMed: 19451168]
59. Koboldt DC, et al. VarScan 2: Somatic mutation and copy number alteration discovery in cancer by exome sequencing. *Genome Res*. 2012; 22:568–76. [PubMed: 22300766]
60. McKenna A, et al. The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res*. 2010; 20:1297–303. [PubMed: 20644199]
61. McLaren W, et al. Deriving the consequences of genomic variants with the Ensembl API and SNP Effect Predictor. *Bioinformatics*. 2010; 26:2069–70. [PubMed: 20562413]
62. Thorvaldsdottir H, Robinson JT, Mesirov JP. Integrative Genomics Viewer (IGV): high-performance genomics data visualization and exploration. *Brief Bioinform*. 2012
63. Chen K, et al. PolyScan: an automatic indel and SNP detection approach to the analysis of human resequencing data. *Genome research*. 2007; 17:659–66. [PubMed: 17416743]
64. Nickerson DA, Tobe VO, Taylor SL. PolyPhred: automating the detection and genotyping of single nucleotide substitutions using fluorescence-based resequencing. *Nucleic acids research*. 1997; 25:2745–51. [PubMed: 9207020]
65. Ng PC, Henikoff S. SIFT: Predicting amino acid changes that affect protein function. *Nucleic acids research*. 2003; 31:3812–4. [PubMed: 12824425]
66. Nakken S, Alseth I, Rognes T. Computational prediction of the effects of non-synonymous single nucleotide polymorphisms in human DNA repair genes. *Neuroscience*. 2007; 145:1273–9. [PubMed: 17055652]
67. Vandin F, Upfal E, Raphael BJ. Algorithms for detecting significantly mutated pathways in cancer. *Journal of computational biology: a journal of computational molecular cell biology*. 2011; 18:507–22. [PubMed: 21385051]
68. Razić S, Magklaras G, Donaldson IM. iRefIndex: a consolidated protein interaction database with provenance. *BMC bioinformatics*. 2008; 9:405. [PubMed: 18823568]




Figure 1. Overview of the integrated analysis of germline and somatic variants in 429 TCGA serous ovarian cases

A total of 27,280 somatic mutations were identified, including 6 SMGs (blue shaded area). germline variants included a total of 3 BRCA1 large-scale deletions, Following filtering of variants with >1% MAF in the population, TCGA ovarian cancer cases, and WHISP controls, a total of 3,635 truncation variants, and 22,953 missense variants (17,348 in expressed genes) remained for TCGA cases. For WHISP controls a total of 10,443 truncation and 30,335 missense variants (in expressed genes) remained. After applying the burden test using WHISP exome sequence data, a total of 6 and 24 genes were significantly enriched for truncation events and missense variants, respectively (orange shaded area). The germline/somatic interaction analysis (purple shaded area) that retained variants in expressed genes in ovarian cancer that met 2 out of 5 criteria identified a total of 237 candidate germline susceptibility variants. The pathway analysis identified three significant pathways involved in ovarian cancer pathogenesis, Fanconi, MAPK, and MLL.




Figure 2. Germline copy number variants in *BRCA1*

Shown are three germline copy number deletion variants affecting *BRCA1* in three ovarian tumor pairs. Normal samples appear above the corresponding tumor samples. Red lines indicate normalized copy number segments based on a minimum of eight probes and blue dots indicate individual probe intensities from Affymetrix 6.0 SNP arrays within the region.




Figure 3. Lollipop plots showing the distribution of germline truncation variants and somatic mutations
 Somatic mutations in *BRCA1*, *BRCA2*, *PALB2*, *CHEK2*, *BRIP1*, *BLM*, *MAP3K15*, and *PTPRH* are shown in blue and germline truncation variants are in orange. Two known pathogenic *BRCA1* germline missense variants are also shown (G1788V and R1699W).




Figure 4. Loss of heterozygosity analysis in tumor samples

(a) Scatter plot displaying variant allele frequencies for all germline truncation variants in normal and tumor samples. Truncation variants in *BRCA1* and *BRCA2* are highlighted in red and blue, respectively. (b) Scatter plot displaying variant allele frequencies for germline missense variants from cancer genes in normal and tumor samples. Germline missense variants in *BRCA1* and *BRCA2* are highlighted in red and blue, respectively. (c) VAFs for the 32 samples showing LOH truncation in *BRCA1*, (d) VAFs for 25 samples showing LOH in *BRCA2*, (e) VAFs in *ATM*, *BLM*, *BRIP1*, *CHEK2*, *ERCC2*, *FANCA*, and *PALB2*. Overall, 100% (32/32) and 76% (19/25) of respective germline *BRCA1* and *BRCA2* truncation

variants showed increased VAFs in the tumor. All germline truncation variants in *BRIP1* and *CHEK2* also showed increased VAFs in corresponding tumors.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 5a




Figure 5b




Figure 5. Significant pathways and subnetworks in ovarian cancer

(a) Oncoprint of genes with germline truncation variants and somatic mutations found in the Fanconi subnetwork identified as significant by HotNet. Genes in the iRefIndex database 58 are underlined. (b) The age distribution for patients with or without germline alterations in Fanconi genes (genes include: Panel a). The Horizontal red line indicates the median age of the group and the blue whiskers represent the age of the individual sample. (c) Oncoprint of genes with germline truncation variants and somatic mutations found in the MAPK subnetwork identified as significant by HotNet. Additional genes in the MAPK pathway with somatic mutations and/or germline truncation variants are included. (d) Oncoprint of

genes with germline truncation variants and somatic mutations found in a subnetwork including *MLL*, *MLL3*, and *SETD1A* identified as significant by HotNet. Additional chromatin modifiers with somatic mutations and/or germline truncation variants are included.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 1

Clinical Characteristics of TCGA cases.

	Category	No. (%)
Ethnicity ^a	Caucasians	387 (90.2)
	African American	21 (4.9)
	Asian	15 (3.5)
	American Indian	2 (0.5)
	Unknown	4 (0.9)
Survival	Living	207 (48.3)
	Deceased	218 (50.8)
	Unknown	4 (0.9)
Age	45	57 (13.3)
	46–69	267 (62.2)
	70	103 (24.0)
	Unknown	2 (0.5)
Stage	IA–IC	5 (1.2)
	IIA–IIC	20 (4.7)
	IIIA–IIIC	338 (78.8)
	IV	62 (14.5)
	Unknown	4 (0.9)

^aNumber assigned to each category after PCA analysis (Supplementary Fig. S1)

Table 2

Thirty-five known and candidate functional missense variants

These variants were identified using a combination of integrated germline and somatic analysis and bioinformatics annotation.

Gene	Annotation	LOVD ^a	BIC ^b	HGMD ^c	HGMD pheno ^d	Exome VAF ^e	Exome Reads	RNA VAF ^e	RNA Reads	Case Freq ^f	Contol Freq ^f	LOF ^g Fanconi
<i>ATM</i>	p.R2459G	NR	NA	NR	NR	91.43	105	NA	NA	1/387 (0.003)	0	
<i>ATM</i>	p.L480F	NR	NA	NR	NR	75.44	57	100	1	1/387 (0.003)	0	<i>BRCA1</i>
<i>ATM</i>	p.P1112A	NR	NA	NR	NR	92.25	129	NA	NA	1/387 (0.003)	0	<i>BLM/FANCD2</i>
<i>BRCA1</i>	p.R1699W	1-/?; 10 ?/?; 8+/?	Clinically Important	DM	Breast and Colorectal Cancer susceptibility	79.01	81	70	10	1/387 (0.003)	0	
<i>BRCA1</i>	p.G1788V	1-/?; 10 ?/?; 8+/?	Clinically Important	DM	Ovarian Cancer	98.16	217	95.65	46	1/387 (0.003)	0	
<i>BRCA1</i>	p.V772A	4-/?; 2 ?/?; 1 +/?	Unknown	DM	Breast Cancer	91.44	292	NA	NA	1/387 (0.003)	0	<i>BRCA1</i>
<i>BRCA2</i>	p.A1996T	NR	Unknown	NR	-	7.14	14	NA	NA	1/387 (0.003)	0	
<i>BRCA2</i>	p.T2088I	NR	NR	NR	-	94.64	56	100	3	1/387 (0.003)	0	
<i>BRCA2</i>	p.K2434T	NR	Unknown	NR	-	82.4	125	NA	NA	1/387 (0.003)	0	
<i>BRCA2</i>	p.F1241L	NR	NR	NR	-	13.46	52	NA	NA	1/387 (0.003)	0	
<i>BRPI</i>	p.N370S	NR	NA	NR	-	76.66	377	0	1	1/387 (0.003)	0	
<i>BRPI</i>	p.P47A	NR	NA	DM	Breast Cancer	97.71	436	100	12	1/387 (0.003)	1/557 (0.002)	
<i>BRPI</i>	p.A349P	1 +/?	NA	DM	Fanconi anemia	13.87	411	20	5	1/387 (0.003)	0	
<i>BRPI</i>	p.K703I	NR	NA	NR	-	88.29	205	100	2	1/387 (0.003)	0	<i>BRPI</i>
<i>CLTC</i>	p.R1498H	NR	NA	NR	-	93.06	72	98.15	379	1/387 (0.003)	1/557 (0.001)	
<i>ERCC2</i>	p.R616P	NR	NA	DM	Trichothio-dystrophy	75.25	101	NA	NA	3/387 (0.008)	0	
<i>ERCC2</i>	p.R616P	NR	NA	DM	Trichothio-dystrophy	57.89	95	NA	NA	3/387 (0.008)	0	
<i>ERCC2</i>	p.R616P	NR	NA	DM	Trichothio-dystrophy	53.97	63	48.39	31	3/387 (0.008)	0	
<i>ERCC2</i>	p.A635V	NR	NA	NR	-	44.44	54	58.25	103	2/387 (0.005)	2/557 (0.003)	<i>BRCA2</i>
<i>ERCC2</i>	p.A635V	NR	NA	NR	-	68.18	22	97.26	73	2/387 (0.005)	2/557 (0.003)	
<i>FRG1</i>	p.G76V	NR	NA	NR	-	70.64	235	90.28	247	1/387 (0.003)	0	<i>BRPI</i>
<i>HIP1</i>	p.T62M		NA	NR	-	69.33	75	88.89	27	1/387 (0.003)	0	<i>BRCA1</i>
<i>ITK</i>	p.R448H	NR	NA	NR	-	41.49	94	0	1	1/387 (0.003)	0	
<i>ITK</i>	p.R581W	NR	NA	NR	-	43.16	95	NA	NA	1/387 (0.003)	1/557 (0.002)	
<i>MYH9</i>	p.R1400W	NR	NA	DM?	Epstein syndrome	93.59	78	89.68	599	1/387 (0.003)	1/557 (0.002)	

Gene	Annotation	LOVD ^a	BIC ^b	HGMD ^c	HGMD pheno ^d	Exome VAF ^e	Exome Reads	RNA VAF ^e	RNA Reads	Case Freq ^f	Contol Freq ^f	LOF ^g Fanconi
<i>MYH9</i>	p.D507N	NR	NA	NR	-	86.96	115	NA	NA	1/387 (0.003)	1/557 (0.002)	
<i>NCKIPSD</i>	p.R677H		NA	NR	-	85.71	14	92.73	55	1/387 (0.003)	0	
<i>NF1</i>	p.V2148D	NR	NA	NR	-	41.67	12	0	61	2/387 (0.005)	0	
<i>NF1</i>	p.V2148D	NR	NA	NR	-	35.71	14	0	76	2/387 (0.005)	0	
<i>NF1</i>	p.A2644G	NR	NA	NR	-	8.28	145	10.84	83	1/387 (0.003)	0	
<i>NF1</i>	p.P1421L	NR	NA	NR	-	89.04	146	81.82	11	1/387 (0.003)	0	
<i>NF1</i>	p.R765H	NR	NA	NR	-	95.2	542	100	17	1/387 (0.003)	0	
<i>NOTCH2</i>	p.H2032N	NR	NA	NR	-	36.78	87	49.28	414	2/387 (0.005)	1/557 (0.002)	<i>BLM</i>
<i>NOTCH2</i>	p.H2032N	NR	NA	NR	-	86.59	82	92.95	241	2/387 (0.005)	1/557 (0.002)	
<i>RB1</i>	p.I831V	NR	NA	NR	-	44.16	77	NA	NA	1/387 (0.003)	0	
<i>RB1</i>	p.R656W	2 ?/?	NA	NR	-	39.95	388	58.86	157	1/387 (0.003)	1/557 (0.002)	
<i>RNF213</i>	p.P978L	NR	NA	NR	-	82.76	29	100	2	1/387 (0.003)	0	
<i>SLC447</i>	p.V824L	NR	NA	NR	-	85.71	35	100	5	1/387 (0.003)	0	
<i>TP53</i>	p.G334R	NR (IARC)	NA	NR	-	83.95	81	NA	NA	1/387 (0.003)	0	
<i>WAS</i>	p.E285Q	NR	NA	NR	E285X DM for Wiskott- Aldrich	76.47	51	81.08	37	1/387 (0.003)	0	

NR=Not reported, NA=Not available,

^a Leiden Open Variation Database (LOVD)⁵⁴ key: Numbers indicate number of LOVD reports. Path: Variant pathogenicity, in the format Reported/Concluded; '+-' indicating the variant is pathogenic, '+?' probably pathogenic, '-?' probably no pathogenicity, '?' effect unknown,

^b Breast Cancer Information Core (BIC)³¹ report (*BRCA1* and *BRCA2* only),

^c Human Gene Mutation Base (HGMD)⁵⁵ status reported pathogenicity (DM=disease causing mutation),

^d Human Gene Mutation Base (HGMD)⁵⁵ phenotype

^e VAF=Variant Allele Frequency

^f Freq =Frequency

^g Loss of function truncation mutations in Fanconi Pathway