

ORIGINAL ARTICLE

Gene-by-gene interactions associated with the risk of conotruncal heart defects

Chen Lyu¹ | Daniel M. Webber² | Stewart L. MacLeod³ | Charlotte A. Hobbs⁴ | Ming Li¹  the National Birth Defects Prevention Study

¹Department of Epidemiology and Biostatistics, Indiana University, Bloomington, IN, USA

²Department of Pathology & Immunology, Washington University at St Louis, Saint Louis, MO, USA

³Arkansas Children's Research Institute, Little Rock, AR, USA

⁴Rady Children's Institute for Genomic Medicine, San Diego, CA, USA

Correspondence

Ming Li, Department of Epidemiology and Biostatistics, School of Public Health - Bloomington, Indiana University, 1025 E. Seventh Street, SPH C108, Bloomington, IN 47405, USA.

Email: li498@indiana.edu

Funding information

Centers for Disease Control and Prevention, Grant/Award Number: FOA #DD09-001, FOA #DD13-003, PA #02081 and PA #96043; Centers for Birth Defects Research and Prevention, Grant/Award Number: NOFO #DD18-001; National Heart, Lung, and Blood Institute, Grant/Award Number: K01HL140333; Eunice Kennedy Shriver National Institute of Child Health and Human Development, Grant/Award Number: R03HD092854

Abstract

Background: The development of conotruncal heart defects (CTDs) involves a complex relationship among genetic variants and maternal lifestyle factors. In this article, we focused on the interactions between 13 candidate genes within folate, homocysteine, and transsulfuration pathways for potential association with CTD risk.

Methods: Targeted sequencing was used for 328 case-parental triads enrolled in the National Birth Defects Prevention Study (NBDPS). To evaluate the interaction of two genes, we applied a conditional logistic regression model for all possible SNP pairs within two respective genes by contrasting the affected infants with their pseudo-controls. The findings were replicated in an independent sample of 86 NBDPS case-parental triads genotyped by DNA microarrays. The results of two studies were further integrated by a fixed-effect meta-analysis.

Results: One SNP pair (i.e., rs4764267 and rs6556883) located in gene *MGST1* and *GLRX*, respectively, was found to be associated with CTD risk after multiple testing adjustment using simpleM, a modified Bonferroni correction approach (nominal *p*-value of 4.62e-06; adjusted *p*-value of .04). Another SNP pair (i.e., rs11892646 and rs56219526) located in gene *DNMT3A* and *MTRR*, respectively, achieved marginal significance after multiple testing adjustment (adjusted *p*-value of .06).

Conclusion: Further studies with larger sample sizes are needed to confirm and elucidate these potential interactions.

KEYWORDS

case-parental triads, conditional logistic regression, conotruncal heart defects, gene-by-gene interaction, meta-analysis, single nucleotide polymorphism

1 | INTRODUCTION

Congenital heart defects (CHDs) are the most common type of birth defects, affecting 48.9 million people globally in 2015 (Vos et al., 2016). The prevalence of disease is approximately 1% (Vos et al., 2015), contributing to ~300,000 infant deaths per year (Naghavi et al., 2015; Wang et al., 2016). Children

with CHDs are usually at increased risks of other health conditions, such as developmental disabilities (Limperopoulos et al., 2000; Razzaghi, Oster, & Reefhuis, 2015) and cognitive disorders (Shillingford et al., 2008). CHDs comprise various cardiac anomalies that are etiologically heterogeneous. Conotruncal heart defects (CTDs), such as tetralogy of Fallot, truncus arteriosus, double outlet right ventricle, transposition

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2019 The Authors. Molecular Genetics & Genomic Medicine published by Wiley Periodicals, Inc.

of the great arteries, pulmonary atresia, malalignment ventricle septal defect, and interrupted aortic arch type B, are a major and severe subtype of CHDs, accounting for 20%–30% of all CHD cases (Botto, Lin, Riehle-Colarusso, Malik, & Correa, 2007; Kuehl & Loffredo, 2005). With the advance of surgical therapy and medical care, the survival rate of CTD-affected infants has improved substantially over the past 50 years (Oster et al., 2013). However, those infants who survived may require repeated surgeries and continuing medical care into their adulthood (Gilboa et al., 2016), imposing enormous emotional and financial burden to their families and the health care system (McClung, Glidewell, & Farr, 2018; Simeone et al., 2014). Understanding the pathogenesis of CTDs is of crucial importance to reduce the public health impact of the disease.

Previous studies have identified a number of risk factors, such as maternal obesity (Brite, Laughon, Troendle, & Mills, 2014; Gilboa et al., 2010), diabetes (Lisowski et al., 2010; Stavsky et al., 2017), tobacco use (Alverson, Strickland, Gilboa, & Correa, 2011; Malik et al., 2008), and medication use (Li et al., 2014a, 2014b) during pregnancy. Genetic variants have also been identified for association with disease susceptibility. For example, the microdeletion on the long arm of chromosome 22 (i.e., DiGeorge syndrome) substantially increases the disease risk, and may account for ~12% of conotruncal malformations (Hacihamdioglu, Hacihamdioglu, & Delil, 2015; Liu et al., 2014). Other chromosomal abnormalities and copy number variations, such as 1q21, 8p23, and 11q25 (Edwards & Gelb, 2016; Thienpont et al., 2007), may also increase the risk of CTDs. In addition, previous studies have shown that multiple genes, such as *NKX2.5*, *GATA4*, *GATA6*, *TBX1*, *TBX5*, *CITED2*, *HAND2*, *NOTCH1*, *JAG1*, *ACTC1*, and *THRAP2*, are associated with the transcription factors, ligands-receptors, contractile and miscellaneous proteins that are involved in essential cardiac developmental processes (Hu et al., 2013; Huang et al., 2010; Li, Pu, Liu, Xu, & Xu, 2017; Wessels & Willems, 2010; Zhang, Hong, et al., 2018a). In recent years, genetic association studies have identified additional genes from various biological pathways, such as folic acid, homocysteine, and transsulfuration, for potential association with CTD risk (Hobbs et al., 2014; Shaw et al., 2005, 2009). However, these genes generally have small to intermediate effect sizes, and collectively may only explain a small proportion of CTDs cases (Fahed, Gelb, Seidman, & Seidman, 2013). The etiology of CTDs is highly complex, involving multiple genetic variants, maternal environmental exposures and lifestyle factors (Gelb & Chung, 2014; Pierpont et al., 2007).

To date, most studies have adopted a single-locus analysis strategy, by testing each single genetic variant for potential association with the disease. However, accumulating evidence has demonstrated that gene-by-gene interaction exists pervasively in biological pathways and may be a major

source of the unexplained disease heritability (Eichler et al., 2010; Moore, 2003). For example, an animal study analyzed the transcriptome data among mouse embryos with various combinations of wild-type and mutant genotypes. The results suggested that the *Tbx5/Osr1/Pcsk6* gene interaction was associated with atrial septal defects through the signaling network (Zhang et al., 2016). Studies have also shown that gene-by-gene interactions can alter the risk of CTDs among human populations. For example, physical interactions were found between particular combinations of *GATA4* and *TBX5* molecules, with varying frequencies between families with and without cardiac septal defects (Garg et al., 2003). It was also suggested that additional gene-by-gene interactions might exist between *MYH6* and *TBX5* (Chen et al., 2017), *MYH6* and *NKX2-5* (Granados-Riveron et al., 2012), and *ZFPM2* and *GATA4* (Zhang et al., 2014). These studies might have provided potential explanations in terms of the complex pathophysiological and etiological processes that result in CTDs, and also support that gene-by-gene interactions are a common mechanism contributing to the disease development.

In this study, we evaluate the potential gene-by-gene interactions among 13 candidate genes selected from three biological pathways (i.e., folate, homocysteine, and transsulfuration) that have altered metabolites in CHD-affected pregnancies (Hobbs, Cleves, Zhao, Melnyk, & James, 2005; Hobbs et al., 2006; Kapusta et al., 1999; Obermann-Borst et al., 2011). A total of 515 common variants from 13 genes were analyzed in two independent samples, including a sequencing study with 328 case-parental triads and a microarray study with 86 case-parental triads. The analysis results from two independent samples were further integrated by a fixed-effect meta-analysis.

2 | MATERIALS AND METHODS

2.1 | Ethics statement

The study was approved by the Institutional Review Boards at University of Arkansas for Medical Sciences and Indiana University Bloomington, and the National Birth Defects Prevention Study (NBDPS) with protocol oversight by the Centers for Disease Control and Prevention (CDC) Center for Birth Defects and Developmental Disabilities. All study subjects gave informed consent. For minors, informed written consent was obtained from their legal guardian for DNA collection.

2.2 | Study population

The National Birth Defects Prevention Study (NBDPS) is a large population-based, case-control study of birth defects. The detailed study design and procedure has been described

elsewhere (Gallagher et al., 2011; Rasmussen et al., 2002; Yoon et al., 2001). Briefly, since 1997, subjects have been recruited from the population-based birth defects surveillance system in ten states (i.e., AR, CA, GA, IA, MA, NC, NJ, NY, TX, and UT). The study covers an annual birth population of 482,000 (i.e., 10% of U.S. births). Case infants were identified if presenting at least 1 major birth defect. Only non-syndromic cases were included, and cases with known single gene defects or chromosomal abnormalities were excluded. Case information and medical records were obtained by trained specialists. All diagnostic tests were reviewed by pediatric cardiologists to ensure uniform criteria for diagnoses. Study information was collected from recruited mothers via computer-assisted telephone-interviews with detailed questions about environmental exposures. After the interview, a buccal cell collection kit was sent to collect biological samples of both biological parents and the infant. The collected buccal cells were sent back to NBDPS laboratories for DNA extraction and genotyping.

In our current study, we adopt the case-parental triad design. A case-parental triad is defined as a singleton live-born infant with CTDs and his/her biological parents. Only complete triads with genetic data available for three family members were included in our study. Subjects in the sequencing study was genotyped using Illumina's targeted sequencing technology, while those in the microarray study were genotyped in one of our earlier studies using Illumina® GoldenGate™ platform (Hobbs et al., 2014). The human reference genome is version GRCh37.p13.

2.3 | Sequencing study

The sequencing study included 328 case-parental triads. Each subject was sequenced for 13 targeted regions, including

exons, introns, and 1kb upstream and downstream regions for the following folate-related genes: *GCLC* (OMIM: 606857), *SHMT1* (OMIM: 182144), *NOS2A* (OMIM: 163730), *MTRR* (OMIM: 602568), *MTHFS* (OMIM: 604197), *GLRX* (OMIM: 600443), *DNMT3A* (OMIM: 602769), *SOD2* (OMIM: 147460), *MGST1* (OMIM: 138330), *GPX4* (OMIM: 138322), *TCN2* (OMIM: 613441), *MTHFD2* (OMIM: 604887), and *TYMS* (OMIM: 188350). These genes were selected for sequencing based on a preliminary analysis of data from an earlier study (Hobbs et al., 2014). The 13 selected genomic regions are described in Table 1. DNA was extracted in accordance with well-established NBDPS protocols. Extracted DNA (50 ng per sample) was prepared for targeted sequencing using Illumina Nextera hybridization enrichment. Target libraries were pooled at 48x and sequenced on a HiScan-SQ or NextSeq500.

2.4 | Microarray study

The microarray study included 86 case-parental triads that are non-overlapping subjects with the sequencing study. These subjects are NBDPS participants in one of our previous studies (Hobbs et al., 2014). Each subject was initially genotyped for about 1,500 genetic variants in 62 candidate genes, including the 13 genes that were included in the sequencing study. Illumina's GoldenGate Custom DNA microarray was used for genotyping. The detailed process of SNP selection, DNA extraction and genotyping can be found elsewhere (Hobbs et al., 2014).

2.5 | Quality control

Thirty-one samples from the sequencing study were also genotyped by Illumina HumanOmni 5M BeadChip as part

TABLE 1 Candidate genes or regions sequenced in this study^a

Genes	Region of SNPs	Total (kb)	Number of SNPs	Pathways
<i>DNMT3A</i>	chr2: 25,451,420–25,568,539	117.1	45	Homocysteine
<i>MTHFD2</i>	chr2: 74,423,130–74,449,302	26.2	10	Folate
<i>MTRR</i>	chr5: 7,846,305–7,908,359	62.0	46	Homocysteine
<i>GLRX</i>	chr5: 95,148,650–95,164,023	15.4	9	Transsulfuration
<i>GCLC</i>	chr6: 53,357,360–53,414,706	57.3	37	Transsulfuration
<i>SOD2</i>	chr6: 160,091,691–160,119,427	27.7	29	Transsulfuration
<i>MGST1</i>	chr12: 16,492,829–16,533,128	40.3	41	Transsulfuration
<i>MTHFS</i>	chr15: 80,130,515–80,198,784	68.3	78	Folate
<i>SHMT1</i>	chr17: 18,222,647–18,274,323	51.7	91	Folate
<i>NOS2A</i>	chr17: 26,075,524–26,131,931	56.4	31	Transsulfuration
<i>TYMS</i>	chr18: 655,999–680,838	24.8	34	Folate
<i>GPX4</i>	chr19: 1,100,976–1,109,213	8.2	10	Transsulfuration
<i>TCN2</i>	chr22: 31,001,072–31,029,898	28.8	54	Folate

^aHuman reference genome version is GRCh37.p13.

of another project (unpublished data). Genotype calls were compared at 196 overlapping sites, and the concordance rate across platforms was 99.0%, indicating high accuracy of variant calls. All samples were screened for Mendelian inconsistencies using PLINK (Purcell et al., 2007). A cell-line from the 1,000 Genome project (Siva, 2008) was sequenced as a positive control and was used as an external quality assurance (EQA) measure for NBDPS. In the sequencing study, all SNPs had a call rate greater than 95%, and all families and SNPs showed Mendelian error rates less than 5%. All genotypes with Mendelian inconsistency were set to missing.

2.6 | Imputation and SNP selection

The targeted sequencing provided genotypes for 9,402 variants in the sequencing study. However, only 74 of those variants were readily available in the microarray study. In order to maximize the overlap of variants between two studies, we further conducted genotype imputation for each study by using 1,000 Genome Project Phase 3 Genome Build b37 as reference panels. Software IMPUTE version 2 was used for the imputation with a “multi-population reference panels” option (Howie, Donnelly, & Marchini, 2009), which chooses a “custom” reference panel for each individual based on all available reference haplotypes. The advantages for such a strategy has been discussed elsewhere (Howie, Marchini, & Stephens, 2011). After the imputation and quality control, the sequencing study and the microarray study included 9,588 and 3,505 variants, respectively. A total of 2,945 variants overlapped between two studies. In this article, we only considered 515 common variants with at least 5% minor allele frequencies in both studies for interaction analysis. We excluded variants with an allele frequency of less than 5% due to the limited statistical power for detecting potential interactions among rare variants.

2.7 | Statistical methods

2.7.1 | Single-locus association test

We first evaluated the association between each genetic variant and CTD risk with a genotypic transmission disequilibrium test (TDT) (Schaid, 1996, 1999). Conditional logistic regression model was fitted assuming additive inheritance. The conditional probability of disease risk was modeled by contrasting affected case infants with the corresponding pseudo-controls whose allele combinations are theoretically possible but not observed based on Mendelian transmission of given parents' genotypes (Falk & Rubinstein, 1987). A major advantage of using matched pseudo-controls is its robustness to confounding causes of association, such as population stratification (Cordell, 2002). Such a strategy has been commonly used for studies of other birth defects, such as cleft

palate (Liu et al., 2018, 2019; Xiao et al., 2017). We used genotypic TDT instead of conventional allelic TDT to have consistent modeling with the interaction analysis described below. The genotypic TDT analysis was performed in R version 3.43 using Bioconductor package “trio” (Schwender et al., 2014).

2.7.2 | Gene-by-gene interaction

Our study included 13 candidate genes (Table 1), leading to a total of 78 gene-by-gene combinations. For each given gene-by-gene combination, we estimated the interaction effect for all possible SNP pairs within two respective genes. We denote p_i as the conditional probability of being a case infant rather than the pseudo-controls for the i -th family; and X_{Ai} and X_{Bi} to be the genotype of two SNPs from two respective genes, coded as minor allele counts. A conditional logistic regression model was fitted as follows:

$$\text{logit}(p_i) = \beta_1 \times X_{Ai} + \beta_2 \times X_{Bi} + \gamma \times X_{Ai} X_{Bi}$$

The model conducted matched case-control analysis by contrasting the affected infants with all possible pseudo-controls assuming additive inheritance. For single bi-allelic variant, two heterozygous parents can produce four inheritance patterns of genotypes, suggesting 1 case versus 3 matched pseudo-controls. Similarly, two bi-allelic variants lead to 16 two-SNP inheritance patterns. The ratio between cases and their pseudo-controls is 1:15. The interaction effect between two variants can thus be tested against the null hypothesis of $\gamma = 0$. The same model was fitted for both the sequencing study and the microarray study. The analysis was conducted in R version 3.43 using Bioconductor package “trio” (Schwender et al., 2014).

2.7.3 | Meta-analysis

We further conducted a fixed-effect meta-analysis in order to integrate the results from the sequencing and the microarray studies. Top associations were selected based on the significance of meta-analysis (i.e., p -values) and the consistency of testing results across two studies, that is, achieving nominal significance level in both studies and having the same direction of effect. The analysis was conducted in R version 3.43 using package “rmeta.”

2.7.4 | Multiple testing adjustment

A total of 118,757 SNP pairs were tested for possible interaction effects. Because of the strong linkage-disequilibrium (LD) structure within the same candidate genes (Fig. S1), the conventional Bonferroni correction would be overly conservative. Alternatively, we adopted a SimpleM method for estimating the

TABLE 2 Characteristics of case-parental triads in sequencing and microarray study

	Sequencing study (<i>n</i> = 328)	Microarray study (<i>n</i> = 86)	<i>p</i> -value
Maternal Age			
Mean (<i>SD</i>)	28.31 (6.00)	30.03 (6.21)	.022
Maternal race, <i>n</i> (%)			.604
Caucasian	233 (71.04%)	68 (79.07%)	
African American	19 (5.80%)	3 (3.49%)	
Hispanic	52 (15.85%)	10 (11.63%)	
Others	24 (7.32%)	5 (5.81%)	
Maternal BMI, <i>n</i> (%)			.161
Underweight	14 (4.27%)	2 (2.32%)	
Normal	147 (44.82%)	53 (61.73%)	
Overweight	81 (24.70%)	16 (18.60%)	
Obese	76 (23.17%)	15 (17.44%)	
Missing	10 (3.05%)	0 (0.00%)	
Maternal Education, <i>n</i> (%)			.466
0–11 years	35 (10.67%)	7 (8.14%)	
12 years	72 (16.77%)	19 (22.09%)	
13–15 years	100 (30.49%)	21 (24.42%)	
16 or more years	121 (36.89%)	36 (41.86%)	
Missing	0 (0.00%)	3 (3.49%)	
Family Income, <i>n</i> (%)			.004
< 10,000	34 (10.36%)	9 (10.46%)	
10,000 to 30,000	55 (14.78%)	15 (17.44%)	
30,000 to 50,000	41 (12.50%)	13 (15.12%)	
> 50,000	62 (18.90%)	46 (53.49%)	
Missing	136 (41.46%)	3 (3.49%)	
Folic Acid Supplementation, <i>n</i> (%)			.755
Maternal Alcohol Use, <i>n</i> (%)			.063
Maternal Smoke, <i>n</i> (%)			.211
Missing	1 (0.30%)	1 (1.16%)	

effective number of independent tests among highly correlated data (Gao, Starmer, & Martin, 2008) and applied a Bonferroni correction based upon this estimate. The details and advantages of SimpleM has been discussed elsewhere (Gao, Becker, Becker, Starmer, & Province, 2010). In particular, SimpleM uses principal component analysis to estimate the effective number of independent tests while accounting for the majority of genetic variation (e.g., 95%). All analyses were performed using R version 3.4.3.

3 | RESULTS

The demographic information of our study populations is summarized in Table 2. A total of 328 and 86 case-parent triads were included in the sequencing and the microarray studies, respectively. Most of the maternal characteristics, including race, body mass index, education level, folic acid supplementation, alcohol use and smoking status, were similar between two studies (i.e., *p*-values > .05). The average maternal age at delivery was 28.31 and 30.03 years in the sequencing and the microarray studies, respectively, which were statistically different but not of clinical importance. Family income levels were also statistically different between two studies, which may largely due to the high level of missing values in the sequencing study. In both studies, the most frequent maternal characteristics were Caucasian race, normal body mass index (BMI), and over 16 years of education. The majority of the families had more than \$50,000 annual household income. In terms of maternal lifestyle factors, more women were taking folic acid supplementation regularly during the pregnancy, and less women smoking or drinking.

3.1 | Single-locus association test

Each of 515 SNPs was examined using a genotypic TDT as described above. The testing *p*-values are shown in Figure 1. The multiple testing significance thresholds were calculated as $0.05/144 = 3.5 \times 10^{-4}$, where the effective number of independent tests estimated by SimpleM was 144. In addition, results for each gene were summarized in Table 3. None of the associations remained significant after multiple testing adjustment.

3.2 | Gene-by-gene interaction

Each SNP pair was examined using a conditional logistic regression model described above. Under the null hypothesis of no association, the testing *p*-values were expected to follow a uniform distribution, and the distribution was evaluated by a quantile-quantile (QQ) plot (Fig. S2). No early departure from the uniform distribution was found, with genomic inflation factors (λ) close to 1, suggesting no inflated type I error rate. The top 5 SNP pairs are summarized in Table 4. In particular, the interaction effect between rs4764267 and rs6556883, located in genes *MGST1* and *GLRX*, respectively, had an adjusted *p*-value of .04, which remained significant after multiple testing adjustment. The interaction effect and corresponding genotypic relative risks varied largely between two studies, which we attributed mainly to the small sample size of our microarray study. These two SNPs may form 9 possible




FIGURE 1 Barplot of single-locus genotypic TDT test from meta-analysis. No significant findings after multiple testing adjustment. Each bar represents the negative 10-based logarithm of p-value for a SNP, ordered by its physical location in the genome. The dashed line represents the threshold of statistical significance after multiple testing adjustment

two-SNP genotype combinations. We estimated the relative risks among these genotype combinations formed by rs4764267 and rs6556883 (Table 5). To illustrate the interaction effect of these two SNPs, we further plotted these estimated relative risks in Figure 2. When genotype GG was observed at SNP rs6556883 (i.e. red line), allele T at SNP rs4764267 was estimated to increase the disease risk compared to allele G. However, when genotype AG or AA was observed at SNP rs6556883 (i.e. green and blue lines), allele T at SNP rs4764267 was expected to decrease the disease risk but with varying effect size. Such a pattern remained consistent across the two studies. SNPs rs4764267 and rs6556883 were located within genes *MGST1* and *GLRX*, respectively. The other four SNP pairs in Table 4 were from the same gene pairs of *DNMT3A* and *MTRR*, and were in strong LD. We believe these four SNP pairs were representing the same gene-by-gene interaction. In particular, the SNP pair of rs11892646 and rs56219526 was marginally significant after multiple testing adjustments (p -value = .06). We presented the analysis results in Table 6, and illustrated the relative risk among genotype combinations in Figure 3.

4 | DISCUSSION

In this study, we conducted a two-phase investigation to evaluate the possible interactions among 13 candidate genes for association with CTD risk. One SNP pair (i.e., rs4764267 and rs6556883) was identified with a significant p -value after multiple testing adjustment, and another SNP pair (i.e., rs11892646 and rs56219526) was

marginally significant. All identified genes are functionally involved in the transsulfuration pathway, supporting its possible contribution to the genetic susceptibility of CTDs.

Our results are consistent with previous work. Gene *MGST1* encodes microsomal glutathione S-transferase 1, which contributes to the antioxidant system by catalyzing glutathione binding to toxic electrophilic compounds. Decreased expression of *MGST1* has been linked structural cardiac abnormalities in *Nos3* deficient mice (Campbell, Li, Biendarra, Terzic, & Nelson, 2015). *GLRX* encodes glutaredoxin-1, which catalyzes the glutathione regeneration and provides reactant for *MGST1*, and is also involved in the response to oxidative stress. Overexpression of *GLRX2* in transgenic mice has been found to attenuate doxorubicin (DOX)-induced heart defects (Diotte, Xiong, Gao, Chua, & Ho, 2009). Others and we have previously reported the association of polymorphisms in *MGST1* and *GLRX* with CHDs in other NBDPS samples although the molecular mechanism remains unclear (Chowdhury et al., 2012; Nembhard et al., 2017, 2018). Meanwhile, accumulated evidence suggests that the oxidative stress/transsulfuration pathway plays an essential role in cardiac pathogenesis. For example, others and we have shown differences in markers of oxidative stress, including glutathione level, among women with infants affected by CHDs compared to those with healthy infants (Hobbs et al., 2005). In addition, decreased activity of antioxidant enzyme such as glutathione peroxidase and lower levels of molecular antioxidants were observed among children affected by CHDs compared to healthy controls (Mukhopadhyay, Gongopadhyay, Rani, Gavel, &

TABLE 3 Results of single-locus genotypic TDT test for each gene^a

Gene	SNP	Chro	Position	Study	Estimate (SE)	95% CI	p-value	Adjusted p-value
<i>MTHFD2</i>	rs1723285	2	74,426,038	Sequencing	1.42 (0.15)	(1.06, 1.92)	2.04*10 ⁻²	.21
				Microarray	1.66 (0.22)	(1.07, 2.57)	2.42*10 ⁻²	
				Meta-analysis	1.49 (0.13)	(1.16, 1.91)	1.45*10 ⁻³	
<i>NOS2A</i>	rs2072324	17	26,116,896	Sequencing	0.65 (0.49)	(0.49, 0.86)	2.47*10 ⁻⁴	.23
				Microarray	0.76 (0.26)	(0.46, 1.27)	0.30	
				Meta-analysis	0.68 (0.12)	(0.53, 0.86)	1.63*10 ⁻³	
<i>MGST1</i>	rs10744119	12	16,522,636	Sequencing	0.68 (0.14)	(0.52, 0.90)	6.08*10 ⁻³	.99
				Microarray	0.90 (0.26)	(0.54, 1.51)	0.69	
				Meta-analysis	0.72 (0.12)	(0.57, 0.92)	9.06*10 ⁻³	
<i>MTRR</i>	rs1801394	5	7,870,973	Sequencing	1.40 (0.12)	(1.11, 1.77)	4.38*10 ⁻³	1.00
				Microarray	0.98 (0.22)	(0.63, 1.50)	0.91	
				Meta-analysis	1.29 (0.10)	(1.05, 1.58)	1.41*10 ⁻²	
<i>SOD2</i>	rs5746105	6	160,112,638	Sequencing	0.66 (0.12)	(0.52, 0.85)	1.05*10 ⁻³	1.00
				Microarray	1.23 (0.23)	(0.79, 1.92)	0.36	
				Meta-analysis	0.76 (0.11)	(0.62, 0.95)	1.47*10 ⁻²	
<i>TCN2</i>	rs4820886	22	31,016,539	Sequencing	0.81 (0.18)	(0.57, 1.14)	0.22	1.00
				Microarray	0.41 (0.34)	(0.21, 0.81)	1.01*10 ⁻²	
				Meta-analysis	0.70 (0.16)	(0.52, 0.96)	2.43*10 ⁻²	
<i>SHMT1</i>	rs8073885	17	18,273,850	Sequencing	0.67 (0.22)	(0.43, 1.05)	7.92*10 ⁻²	1.00
				Microarray	0.69 (0.39)	(0.32, 1.48)	0.34	
				Meta-analysis	0.68 (0.20)	(0.46, 0.99)	4.56*10 ⁻²	
<i>MTHFS</i>	rs35919462	15	80,131,692	Sequencing	1.43 (0.16)	(1.05, 1.94)	2.47*10 ⁻²	1.00
				Microarray	0.90 (0.26)	(0.54, 1.50)	0.70	
				Meta-analysis	1.26 (0.14)	(0.97, 1.64)	8.60*10 ⁻²	
<i>GCLC</i>	rs2300420	6	53,376,551	Sequencing	0.66 (0.20)	(0.44, 0.98)	3.81*10 ⁻²	1.00
				Microarray	1.17 (0.39)	(0.54, 2.52)	0.70	
				Meta-analysis	0.74 (0.18)	(0.52, 1.06)	9.66*10 ⁻²	
<i>TYMS</i>	rs2741184	18	679,660	Sequencing	1.36 (0.18)	(0.96, 1.92)	8.29*10 ⁻²	1.00
				Microarray	1.08 (0.27)	(0.63, 1.84)	0.78	
				Meta-analysis	1.27 (0.15)	(0.95, 1.69)	0.11	
<i>DNMT3A</i>	rs62131064	2	25,568,528	Sequencing	0.60 (0.25)	(0.37, 0.98)	4.28*10 ⁻²	1.00
				Microarray	1.22 (0.45)	(0.51, 2.95)	0.66	
				Meta-analysis	0.71 (0.22)	(0.46, 1.09)	0.12	
<i>GLRX</i>	rs871775	5	95,158,768	Sequencing	1.02 (0.19)	(0.70, 1.49)	0.92	1.00
				Microarray	2.58 (0.34)	(1.33, 5.03)	5.25*10 ⁻³	
				Meta-analysis	1.28 (0.17)	(0.92, 1.78)	0.14	
<i>GPX4</i>	rs1808194	19	1,102,323	Sequencing	0.92 (0.12)	(0.73, 1.18)	0.54	1.00
				Microarray	1.00 (0.26)	(0.60, 1.67)	1.00	
				Meta-analysis	0.94 (0.11)	(0.75, 1.17)	0.58	

^aEstimates are exponential of β coefficients, representing relative risks. SNPs were ordered based on adjusted p values.

Mishra, 2015; Rokicki, Strzalkowski, Klapcinska, Danch, & Sobczak, 2003). Moreover, animal studies showed that oxidative stress might explain the increased risk of CHDs among female mice with diabetes (Wang, Reece, & Yang, 2015; Wu et al., 2016). In terms of the genes *DNMT3A* and *MTRR*, evidence suggest their association with CHD, respectively, in both animal models (Deng, Elmore, Lawrance, Matthews, & Rozen, 2008; Feng et al., 2013) and population-based case-control studies (van Beynum et al., 2006; Shaw et al., 2009; Sheng et al., 2013). It is

also biologically plausible that *MTRR* and *DNMT3A* may interact functionally since they work sequentially in homocysteine-methionine cycle (Ly, Hoyt, Crowell, & Kim, 2012).

In our study, a few assumptions were made while evaluating the possible interactions. First, by contrasting the affected infants with pseudo-controls, we assumed that these theoretically possible but unobserved siblings were healthy individuals. Such an assumption is in general reasonable since CTDs are a relative rare

TABLE 4 Top 5 gene-by-gene interaction by conditional logistic regression^a

Gene Pair	SNP pair	Chro	Position	Study	Estimate (SE)	95% CI	p-value	Adjusted p-value ^b
<i>MGST1</i>	rs4764267	12	16,523,580	Sequencing	0.49 (0.18)	(0.34, 0.70)	1.23*10 ⁻⁴	.04**
<i>GLRX</i>	rs6556883	5	95,152,085	Microarray	0.24 (0.50)	(0.09, 0.65)	4.78*10 ⁻³	
				Meta-analysis	0.45 (0.17)	(0.32, 0.63)	4.62*10 ⁻⁶	
<i>MTRR</i>	rs56219526	5	7,891,210	Sequencing	2.66 (0.27)	(1.56, 4.53)	3.34*10 ⁻⁴	.06*
<i>DNMT3A</i>	rs11892646	2	25,494,474	Microarray	4.53 (0.52)	(1.62, 12.69)	4.03*10 ⁻³	
				Meta-analysis	2.98 (0.24)	(1.85, 4.78)	6.55*10 ⁻⁶	
<i>MTRR</i>	rs3776454	5	7,896,604	Sequencing	2.53 (0.27)	(1.49, 4.29)	5.96*10 ⁻⁴	.12
<i>DNMT3A</i>	rs11892646	2	25,494,474	Microarray	4.53 (0.52)	(1.62, 12.69)	4.03*10 ⁻³	
				Meta-analysis	2.85 (0.24)	(1.78, 4.57)	1.25*10 ⁻⁵	
<i>MTRR</i>	rs326125	5	7,888,001	Sequencing	2.37 (0.27)	(1.40, 4.03)	1.39*10 ⁻³	.22
<i>DNMT3A</i>	rs11892646	2	25,494,474	Microarray	4.50 (0.50)	(1.68, 12.05)	2.76*10 ⁻³	
				Meta-analysis	2.74 (0.24)	(1.72, 4.37)	2.31*10 ⁻⁵	
<i>MTRR</i>	rs10380	5	7,897,191	Sequencing	2.26 (0.24)	(1.40, 3.66)	8.98*10 ⁻⁴	.24
<i>DNMT3A</i>	rs11892646	2	25,494,474	Microarray	4.34 (0.52)	(1.56, 12.06)	4.87*10 ⁻³	
				Meta-analysis	2.54 (0.22)	(1.65, 3.93)	2.62*10 ⁻⁵	

^aEstimates are exponential of β coefficients, representing relative risks.

^bAdjusted p-values were marked with **if less than .05 and marked with *if less than .1.

TABLE 5 Relative risk for all possible genotype combinations of rs4764267 and rs6556883^a

rs4764267	rs6556883	Estimate (Sequencing)	Estimate (Microarray)
GG	GG	1.00 (ref)	1.00 (ref)
TG	GG	1.70 (1.23, 2.36)	1.73 (0.77, 3.91)
TT	GG	2.89 (1.51, 5.56)	3.01 (0.59, 15.28)
GG	AG	1.55 (1.04, 2.31)	4.38 (1.59, 12.08)
TG	AG	1.29 (0.83, 2.02)	1.84 (0.54, 6.24)
TT	AG	1.08 (0.54, 2.18)	0.78 (0.11, 5.40)
GG	AA	2.41 (1.08, 5.35)	19.19 (2.53, 145.84)
TG	AA	0.98 (0.49, 2.00)	1.96 (0.27, 14.20)
TT	AA	0.40 (0.14, 1.19)	0.20 (0.01, 4.67)

^aEstimates are relative risks and 95% confident intervals.

disease with around 0.1% prevalence in the general population (Zhang, Li, et al., 2018b). Second, additive mode of inheritance was assumed when estimating the genetic effects, which is commonly used in genetic association studies.

The current analysis has several strengths. First, it is robust to population stratification and other environmental confounders as a result of case-parent design and use of conditional logistic regression model. Second, it integrates data from two studies to replicate results and achieve greater power. Third, we have optimized statistical power by including only common variants in the analysis. The study is limited by modest sample size, which reduces



**FIGURE 2** Estimated relative risk of CTD among all genotypic combinations of rs4764267 and rs6556883: (a) is for sequencing study and (b) is for microarray study

TABLE 6 Relative risk for all possible genotype combinations of rs11892646 and rs56219526^a

rs11892646	rs56219526	Estimate (Sequencing)	Estimate (Microarray)
TT	GG	1.00 (ref)	1.00 (ref)
CT	GG	0.78 (0.52, 1.16)	0.52 (0.21, 1.29)
CC	GG	0.61 (0.27, 1.35)	0.27 (0.04, 1.66)
TT	AG	0.48 (0.32, 0.73)	0.79 (0.38, 1.65)
CT	AG	1.00 (0.60, 1.64)	1.86 (0.71, 4.85)
CC	AG	2.07 (0.82, 5.20)	4.35 (0.76, 24.80)
TT	AA	0.23 (0.10, 0.54)	0.63 (0.14, 2.74)
CT	AA	1.28 (0.53, 3.09)	6.69 (1.20, 37.34)
CC	AA	7.02 (1.34, 36.71)	70.95 (3.11, 1619.84)

^aEstimates are relative risks and 95% confident intervals.**FIGURE 3** Estimated relative risk of CTD among all genotypic combinations of rs11892646 and rs56219526: (a) is for sequencing study and (b) is for microarray study

statistical power for epistasis detection. However, this is the one of the largest studies to investigate the gene-by-gene interaction and the risk of conotruncal heart defects.

5 | CONCLUSION

We identified two gene-by-gene interactions that are potentially associated with conotruncal heart defects. The genes are involved in transsulfuration pathways. Further studies with larger sample sizes are needed to confirm the associations. While we provide a possible explanation of their interactions, these explanations are our speculations based on existing knowledge. Additional molecular studies are needed to elucidate the molecular mechanisms of this hypothesis.

ACKNOWLEDGMENTS

This study is supported, in part, by the National Heart, Lung and Blood Institute under award number K01HL140333, the Eunice Kennedy Shriver National Institute of Child Health and Human Development under award number R03HD092854, the Centers for Disease Control and Prevention cooperative agreements under PA #96043, PA #02081, FOA #DD09-001, FOA #DD13-003, and NOFO #DD18-001 to the Centers for Birth Defects Research and Prevention participating in the National Birth Defects Prevention Study (NBDPS) and/or the Birth Defects Study To Evaluate Pregnancy exposureS (BD-STEPS), , and by a grant from the University of Arkansas for Medical Sciences, Children's University Medical Group Fund. The contents of this manuscript are solely the responsibility of the authors and do not necessarily represent the official views of the National Institute of Health and the Centers for Disease Control and Prevention.

CONFLICT OF INTEREST

The authors confirm that no conflicts of interest to disclose.

AUTHOR CONTRIBUTION

CL and ML conceptualized the study, conducted the data analysis, and wrote the manuscript. DW, CH, and SM designed the sequencing study. CH and SM designed the microarray study. CH, SM, and NBDPS collected the samples and provided study oversight. All authors read and revised the manuscript.

ORCID

Ming Li  <https://orcid.org/0000-0003-0273-6217>

REFERENCE

Alverson, C. J., Strickland, M. J., Gilboa, S. M., & Correa, A. (2011). Maternal smoking and congenital heart defects in the

- Baltimore-Washington Infant Study. *Pediatrics*, 127(3), e647–653. <https://doi.org/10.1542/peds.2010-1399>
- Botto, L. D., Lin, A. E., Riehle-Colarusso, T., Malik, S., & Correa, A.; National Birth Defects Prevention Study (2007). Seeking causes: Classifying and evaluating congenital heart defects in etiologic studies. *Birth Defects Research Part A: Clinical and Molecular Teratology*, 79(10), 714–727. <https://doi.org/10.1002/bdra.20403>
- Brite, J., Laughon, S. K., Troendle, J., & Mills, J. (2014). Maternal overweight and obesity and risk of congenital heart defects in offspring. *International Journal of Obesity(London)*, 38(6), 878–882. <https://doi.org/10.1038/ijo.2013.244>
- Campbell, K. A., Li, X., Biendarra, S. M., Terzic, A., & Nelson, T. J. (2015). Nos3^{-/-} iPSCs model concordant signatures of in utero cardiac pathogenesis. *Journal of Molecular and Cellular Cardiology*, 87, 228–236. <https://doi.org/10.1016/j.yjmcc.2015.08.021>
- Chen, H. X., Zhang, X., Hou, H. T., Wang, J., Yang, Q., Wang, X. L., & He, G. W. (2017). Identification of a novel and functional mutation in the TBX5 gene in a patient by screening from 354 patients with isolated ventricular septal defect. *European Journal of Medical Genetics*, 60(7), 385–390. <https://doi.org/10.1016/j.ejmg.2017.04.011>
- Chowdhury, S., Hobbs, C. A., MacLeod, S. L., Cleves, M. A., Melnyk, S., James, S. J., ... Erickson, S. W. (2012). Associations between maternal genotypes and metabolites implicated in congenital heart defects. *Molecular Genetics and Metabolism*, 107(3), 596–604. <https://doi.org/10.1016/j.ymgme.2012.09.022>
- Cordell, H. J. (2002). Epistasis: What it means, what it doesn't mean, and statistical methods to detect it in humans. *Human Molecular Genetics*, 11(20), 2463–2468. <https://doi.org/10.1093/hmg/11.20.2463>
- Deng, L., Elmore, C. L., Lawrance, A. K., Matthews, R. G., & Rozen, R. (2008). Methionine synthase reductase deficiency results in adverse reproductive outcomes and congenital heart defects in mice. *Molecular Genetics and Metabolism*, 94(3), 336–342. <https://doi.org/10.1016/j.ymgme.2008.03.004>
- Diotte, N. M., Xiong, Y., Gao, J., Chua, B. H., & Ho, Y. S. (2009). Attenuation of doxorubicin-induced cardiac injury by mitochondrial glutaredoxin 2. *Biochimica Et Biophysica Acta*, 1793(2), 427–438. <https://doi.org/10.1016/j.bbamer.2008.10.014>
- Edwards, J. J., & Gelb, B. D. (2016). Genetics of congenital heart disease. *Current Opinion in Cardiology*, 31(3), 235–241. <https://doi.org/10.1097/HCO.0000000000000274>
- Eichler, E. E., Flint, J., Gibson, G., Kong, A., Leal, S. M., Moore, J. H., & Nadeau, J. H. (2010). Missing heritability and strategies for finding the underlying causes of complex disease. *Nature Reviews Genetics*, 11(6), 446–450. <https://doi.org/10.1038/nrg2809>
- Fahed, A. C., Gelb, B. D., Seidman, J. G., & Seidman, C. E. (2013). Genetics of congenital heart disease: The glass half empty. *Circulation Research*, 112(4), 707–720. <https://doi.org/10.1161/CIRCRESAHA.112.300853>
- Falk, C. T., & Rubinstein, P. (1987). Haplotype relative risks: An easy reliable way to construct a proper control sample for risk calculations. *Annals of Human Genetics*, 51(3), 227–233. <https://doi.org/10.1111/j.1469-1809.1987.tb00875.x>
- Feng, Y., Zhao, L. Z., Hong, L., Shan, C., Shi, W., & Cai, W. (2013). Alteration in methylation pattern of GATA-4 promoter region in vitamin A-deficient offspring's heart. *Journal of Nutritional Biochemistry*, 24(7), 1373–1380. <https://doi.org/10.1016/j.jnutbio.2012.11.005>
- Gallagher, M. L., Sturchio, C., Smith, A., Koontz, D., Jenkins, M. M., Honein, M. A., & Rasmussen, S. A. (2011). Evaluation of mailed pediatric buccal cytobrushes for use in a case-control study of birth defects. *Birth Defects Research Part A: Clinical and Molecular Teratology*, 91(7), 642–648. <https://doi.org/10.1002/bdra.20829>
- Gao, X., Becker, L. C., Becker, D. M., Starmer, J. D., & Province, M. A. (2010). Avoiding the high Bonferroni penalty in genome-wide association studies. *Genetic Epidemiology*, 34(1), 100–105. <https://doi.org/10.1002/gepi.20430>
- Gao, X., Starmer, J., & Martin, E. R. (2008). A multiple testing correction method for genetic association studies using correlated single nucleotide polymorphisms. *Genetic Epidemiology*, 32(4), 361–369. <https://doi.org/10.1002/gepi.20310>
- Garg, V., Kathiriyi, I. S., Barnes, R., Schluterman, M. K., King, I. N., Butler, C. A., ... Srivastava, D. (2003). GATA4 mutations cause human congenital heart defects and reveal an interaction with TBX5. *Nature*, 424(6947), 443–447. <https://doi.org/10.1038/nature01827>
- Gelb, B. D., & Chung, W. K. (2014). Complex genetics and the etiology of human congenital heart disease. *Cold Spring Harbor Perspectives in Medicine*, 4(7), a013953. <https://doi.org/10.1101/cshperspect.a013953>
- Gilboa, S. M., Correa, A., Botto, L. D., Rasmussen, S. A., Waller, D. K., Hobbs, C. A., ... Riehle-Colarusso, T. J.; National Birth Defects Prevention Study (2010). Association between prepregnancy body mass index and congenital heart defects. *American Journal of Obstetrics and Gynecology*, 202(1): 51. e1–51. e10. <https://doi.org/10.1016/j.ajog.2009.08.005>
- Gilboa, S. M., Devine, O. J., Kucik, J. E., Oster, M. E., Riehle-Colarusso, T., Nembhard, W. N., ... Marelli, A. J. (2016). Congenital Heart Defects in the United States: Estimating the Magnitude of the Affected Population in 2010. *Circulation*, 134(2), 101–109. <https://doi.org/10.1161/CIRCULATIONAHA.115.019307>
- Granados-Riveron, J. T., Pope, M., Bu'lock, F. A., Thornborough, C., Eason, J., Setchfield, K., ... Brook, J. D. (2012). Combined mutation screening of NKX2-5, GATA4, and TBX5 in congenital heart disease: Multiple heterozygosity and novel mutations. *Congenital Heart Disease*, 7(2), 151–159. <https://doi.org/10.1111/j.1747-0803.2011.00573.x>
- Hacihamdioglu, B., Hacihamdioglu, D., & Delil, K. (2015). 22q11 deletion syndrome: Current perspective. *The Application of Clinical Genetics*, 8, 123–132. <https://doi.org/10.2147/TACG.S82105>
- Hobbs, C. A., Cleves, M. A., Macleod, S. L., Erickson, S. W., Tang, X., Li, J., ... Malik, S.; National Birth Defects Prevention (2014). Conotruncal heart defects and common variants in maternal and fetal genes in folate, homocysteine, and transsulfuration pathways. *Birth Defects Research Part A: Clinical and Molecular Teratology*, 100(2), 116–126. <https://doi.org/10.1002/bdra.23225>
- Hobbs, C. A., Cleves, M. A., Zhao, W., Melnyk, S., & James, S. J. (2005). Congenital heart defects and maternal biomarkers of oxidative stress. *American Journal of Clinical Nutrition*, 82(3), 598–604. <https://doi.org/10.1093/ajcn.82.3.598>
- Hobbs, C. A., Malik, S., Zhao, W., James, S. J., Melnyk, S., & Cleves, M. A. (2006). Maternal homocysteine and congenital heart defects. *Journal of the American College of Cardiology*, 47(3), 683–685. <https://doi.org/10.1016/j.jacc.2005.11.013>
- Howie, B. N., Donnelly, P., & Marchini, J. (2009). A flexible and accurate genotype imputation method for the next generation of genome-wide association studies. *PLoS Genetics*, 5(6), e1000529. <https://doi.org/10.1371/journal.pgen.1000529>
- Howie, B., Marchini, J., & Stephens, M. (2011). Genotype imputation with thousands of genomes. *G3 (Bethesda)*, 1(6), 457–470. <https://doi.org/10.1534/g3.111.001198>

- Hu, Z., Shi, Y., Mo, X., Xu, J., Zhao, B., Lin, Y., ... Shen, H. (2013). A genome-wide association study identifies two risk loci for congenital heart malformations in Han Chinese populations. *Nature Genetics*, *45*(7), 818–821. <https://doi.org/10.1038/ng.2636>
- Huang, J. B., Liu, Y. L., Sun, P. W., Lv, X. D., Du, M., & Fan, X. M. (2010). Molecular mechanisms of congenital heart disease. *Cardiovascular Pathology*, *19*(5), e183–193. <https://doi.org/10.1016/j.carpath.2009.06.008>
- Kapusta, L., Haagmans, M. L., Steegers, E. A., Cuyper, M. H., Blom, H. J., & Eskes, T. K. (1999). Congenital heart defects and maternal derangement of homocysteine metabolism. *Journal of Pediatrics*, *135*(6), 773–774. [https://doi.org/10.1016/S0022-3476\(99\)70102-2](https://doi.org/10.1016/S0022-3476(99)70102-2)
- Kuehl, K. S., & Loffredo, C. A. (2005). Genetic and environmental influences on malformations of the cardiac outflow tract. *Expert Review of Cardiovascular Therapy*, *3*(6), 1125–1130. <https://doi.org/10.1586/14779072.3.6.1125>
- Li, B., Pu, T., Liu, Y., Xu, Y., & Xu, R. (2017). CITED2 Mutations in Conserved Regions Contribute to Conotruncal Heart Defects in Chinese Children. *DNA and Cell Biology*, *36*(7), 589–595. <https://doi.org/10.1089/dna.2017.3701>
- Li, M., Liu, Z., Lin, Y., Chen, X., Li, S., You, F., ... Zhang, Y. (2014a). Maternal influenza-like illness, medication use during pregnancy and risk of congenital heart defects in offspring. *The Journal of Maternal-Fetal & Neonatal Medicine*, *27*(8), 807–811. <https://doi.org/10.3109/14767058.2013.838950>
- Li, M., Liu, Z., Lin, Y., Chen, X., Li, S., You, F., ... Zhu, J. (2014b). Maternal influenza-like illness, medication use during pregnancy and risk of congenital heart defects in offspring. *The Journal of Maternal-Fetal & Neonatal Medicine*, *27*(8), 807–811. <https://doi.org/10.3109/14767058.2013.838950>
- Limperopoulos, C., Majnemer, A., Shevell, M. I., Rosenblatt, B., Rohlicek, C., & Tchervenkov, C. (2000). Neurodevelopmental status of newborns and infants with congenital heart defects before and after open heart surgery. *Journal of Pediatrics*, *137*(5), 638–645. <https://doi.org/10.1067/mpd.2000.109152>
- Lisowski, L. A., Verheijen, P. M., Copel, J. A., Kleinman, C. S., Wassink, S., Visser, G. H., & Meijboom, E. J. (2010). Congenital heart disease in pregnancies complicated by maternal diabetes mellitus. An international clinical collaboration, literature review, and meta-analysis. *Herz*, *35*(1), 19–26. <https://doi.org/10.1007/s00059-010-3244-3>
- Liu, A. P., Chow, P. C., Lee, P. P., Mok, G. T., Tang, W. F., Lau, E. T., ... Chung, B. H. (2014). Under-recognition of 22q11.2 deletion in adult Chinese patients with conotruncal anomalies: Implications in transitional care. *European Journal of Medical Genetics*, *57*(6), 306–311. <https://doi.org/10.1016/j.ejmg.2014.03.014>
- Liu, D., Schwender, H., Wang, M., Wang, H., Wang, P., Zhu, H., ... Beaty, T. H. (2018). Gene–gene interaction between MSX1 and TP63 in Asian case-parent trios with nonsyndromic cleft lip with or without cleft palate. *Birth Defects Research*, *110*(4), 317–324.
- Liu, D., Wang, M., Yuan, Y., Schwender, H., Wang, H., Wang, P., ... Beaty, T. H. (2019). Gene–gene interaction among cell adhesion genes and risk of nonsyndromic cleft lip with or without cleft palate in Chinese case-parent trios. *Molecular Genetics & Genomic Medicine*, *7*(10), <https://doi.org/10.1002/mgg3.872>
- Ly, A., Hoyt, L., Crowell, J., & Kim, Y. I. (2012). Folate and DNA methylation. *Antioxidants & Redox Signaling*, *17*(2), 302–326. <https://doi.org/10.1089/ars.2012.4554>
- Malik, S., Cleves, M. A., Honein, M. A., Romitti, P. A., Botto, L. D., Yang, S., & ... National Birth Defects Prevention Study (2008). Maternal smoking and congenital heart defects. *Pediatrics*, *121*(4), e810–816. <https://doi.org/10.1542/peds.2007-1519>
- McClung, N., Glidewell, J., & Farr, S. L. (2018). Financial burdens and mental health needs in families of children with congenital heart disease. *Congenital Heart Disease*, *13*(4), 554–562. <https://doi.org/10.1111/chd.12605>
- Moore, J. H. (2003). The ubiquitous nature of epistasis in determining susceptibility to common human diseases. *Human Heredity*, *56*(1–3), 73–82. <https://doi.org/10.1159/000073735>
- Mukhopadhyay, B., Gongopadhyay, A. N., Rani, A., Gavel, R., & Mishra, S. P. (2015). Free radicals and antioxidants status in neonates with congenital malformation. *Journal of Indian Association of Pediatric Surgeons*, *20*(4), 179–183. <https://doi.org/10.4103/0971-9261.161037>
- Naghavi, M., Wang, H. D., Lozano, R., Davis, A., Liang, X. F., Zhou, M. G., ... Colla, G. M. C. D. (2015). Global, regional, and national age-sex specific all-cause and cause-specific mortality for 240 causes of death, 1990–2013: A systematic analysis for the Global Burden of Disease Study 2013. *Lancet*, *385*(9963), 117–171. [https://doi.org/10.1016/S0140-6736\(14\)61682-2](https://doi.org/10.1016/S0140-6736(14)61682-2)
- Nembhard, W. N., Tang, X., Hu, Z., MacLeod, S., Stowe, Z., Webber, D.; National Birth Defects Prevention (2017). Maternal and infant genetic variants, maternal preconceptional use of selective serotonin reuptake inhibitors, and risk of congenital heart defects in offspring: Population based study. *BMJ*, *356*, j832. <https://doi.org/10.1136/bmj.j832>
- Nembhard, W. N., Tang, X., Li, J., MacLeod, S. L., Levy, J., Schaefer, G. B., & ... National Birth Defects Prevention Study (2018). A parent-of-origin analysis of paternal genetic variants and increased risk of conotruncal heart defects. *American Journal of Medical Genetics. Part A*, *176*(3), 609–617. <https://doi.org/10.1002/ajmg.a.38611>
- Obermann-Borst, S. A., van Driel, L. M., Helbing, W. A., de Jonge, R., Wildhagen, M. F., Steegers, E. A., & Steegers-Theunissen, R. P. (2011). Congenital heart defects and biomarkers of methylation in children: A case-control study. *European Journal of Clinical Investigation*, *41*(2), 143–150. <https://doi.org/10.1111/j.1365-2362.2010.02388.x>
- Oster, M. E., Lee, K. A., Honein, M. A., Riehle-Colarusso, T., Shin, M., & Correa, A. (2013). Temporal trends in survival among infants with critical congenital heart defects. *Pediatrics*, *131*(5), e1502–1508. <https://doi.org/10.1542/peds.2012-3435>
- Pierpont, M. E., Basson, C. T., Benson, D. W. Jr, Gelb, B. D., Giglia, T. M., Goldmuntz, E., ... Webb, C. L.; American Heart Association Congenital Cardiac Defects Committee, Council on Cardiovascular Disease in the Young (2007). Genetic basis for congenital heart defects: Current knowledge: A scientific statement from the American Heart Association Congenital Cardiac Defects Committee, Council on Cardiovascular Disease in the Young: Endorsed by the American Academy of Pediatrics. *Circulation*, *115*(23), 3015–3038. <https://doi.org/10.1161/CIRCULATIONAHA.106.183056>
- Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M. A., Bender, D., ... Daly, M. J. (2007). PLINK: A tool set for whole-genome association and population-based linkage analyses. *The American Journal of Human Genetics*, *81*(3), 559–575. <https://doi.org/10.1086/519795>
- Rasmussen, S. A., Lammer, E. J., Shaw, G. M., Finnell, R. H., McGehee, R. E. Jr, Gallagher, M., ... Murray, J. C.; National Birth Defects Prevention Study (2002). Integration of DNA sample collection

- into a multi-site birth defects case-control study. *Teratology*, 66(4), 177–184. <https://doi.org/10.1002/tera.10086>
- Razzaghi, H., Oster, M., & Reefhuis, J. (2015). Long-term outcomes in children with congenital heart disease: National Health Interview Survey. *Journal of Pediatrics*, 166(1), 119–124. <https://doi.org/10.1016/j.jpeds.2014.09.006>
- Rokicki, W., Strzalkowski, A., Klapcinska, B., Danch, A., & Sobczak, A. (2003). Antioxidant status in newborns and infants suffering from congenital heart defects. *Wiadomosci Lekarskie*, 56(7–8), 337–340.
- Schaid, D. J. (1996). General score tests for associations of genetic markers with disease using cases and their parents. *Genetic Epidemiology*, 13(5), 423–449. [https://doi.org/10.1002/\(SICI\)1098-2272\(1996\)13:5<423:AID-GEPI1>3.0.CO;2-3](https://doi.org/10.1002/(SICI)1098-2272(1996)13:5<423:AID-GEPI1>3.0.CO;2-3)
- Schaid, D. J. (1999). Likelihoods and TDT for the case-parents design. *Genetic Epidemiology*, 16(3), 250–260. [https://doi.org/10.1002/\(SICI\)1098-2272\(1999\)16:3<250:AID-GEPI2>3.0.CO;2-T](https://doi.org/10.1002/(SICI)1098-2272(1999)16:3<250:AID-GEPI2>3.0.CO;2-T)
- Schwender, H., Li, Q., Neumann, C., Taub, M. A., Younkin, S. G., Berger, P., ... Ruczinski, I. (2014). Detecting disease variants in case-parent trio studies using the bioconductor software package trio. *Genetic Epidemiology*, 38(6), 516–522. <https://doi.org/10.1002/gepi.21836>
- Shaw, G. M., Iovannisci, D. M., Yang, W., Finnell, R. H., Carmichael, S. L., Cheng, S., & Lammer, E. J. (2005). Risks of human conotruncal heart defects associated with 32 single nucleotide polymorphisms of selected cardiovascular disease-related genes. *American Journal of Medical Genetics. Part A*, 138(1), 21–26. <https://doi.org/10.1002/ajmg.a.30924>
- Shaw, G. M., Lu, W., Zhu, H., Yang, W., Briggs, F. B., Carmichael, S. L., ... Finnell, R. H. (2009). 118 SNPs of folate-related genes and risks of spina bifida and conotruncal heart defects. *BMC Medical Genetics*, 10(1), 49. <https://doi.org/10.1186/1471-2350-10-49>
- Sheng, W., Qian, Y., Wang, H., Ma, X., Zhang, P., Chen, L., ... Huang, G. (2013). Association between mRNA levels of DNMT1, DNMT3A, DNMT3B, MBD2 and LINE-1 methylation status in infants with tetralogy of Fallot. *International Journal of Molecular Medicine*, 32(3), 694–702. <https://doi.org/10.3892/ijmm.2013.1427>
- Shillingford, A. J., Glanzman, M. M., Ittenbach, R. F., Clancy, R. R., Gaynor, J. W., & Wernovsky, G. (2008). Inattention, hyperactivity, and school performance in a population of school-age children with complex congenital heart disease. *Pediatrics*, 121(4), e759–767. <https://doi.org/10.1542/peds.2007-1066>
- Simeone, R. M., Oster, M. E., Cassell, C. H., Armour, B. S., Gray, D. T., & Honein, M. A. (2014). Pediatric inpatient hospital resource use for congenital heart defects. *Birth Defects Research Part A: Clinical and Molecular Teratology*, 100(12), 934–943. <https://doi.org/10.1002/bdra.23262>
- Siva, N. (2008). 1000 Genomes project. In: Nature Publishing Group.
- Stavsky, M., Robinson, R., Sade, M. Y., Krymko, H., Zalstein, E., Ioffe, V., ... Levitas, A. (2017). Elevated birth prevalence of conotruncal heart defects in a population with high consanguinity rate. *Cardiology in the Young*, 27(1), 109–116. <https://doi.org/10.1017/S1047951116000202>
- Thienpont, B., Mertens, L., de Ravel, T., Eyskens, B., Boshoff, D., Maas, N., ... Devriendt, K. (2007). Submicroscopic chromosomal imbalances detected by array-CGH are a frequent cause of congenital heart defects in selected patients. *European Heart Journal*, 28(22), 2778–2784. <https://doi.org/10.1093/eurheartj/ehl560>
- van Beynum, I. M., Kouwenberg, M., Kapusta, L., den Heijer, M., van der Linden, I. J., Daniels, O., & Blom, H. J. (2006). MTRR 66A>G polymorphism in relation to congenital heart defects. *Clinical Chemistry and Laboratory Medicine*, 44(11), 1317–1323. <https://doi.org/10.1515/CCLM.2006.254>
- Vos, T., Allen, C., Arora, M., Barber, R. M., Bhutta, Z. A., Brown, A., ... Chen, A. Z. (2016). Global, regional, and national incidence, prevalence, and years lived with disability for 310 diseases and injuries, 1990–2015: A systematic analysis for the Global Burden of Disease Study 2015. *The Lancet*, 388(10053), 1545–1602. [https://doi.org/10.1016/S0140-6736\(16\)31678-6](https://doi.org/10.1016/S0140-6736(16)31678-6)
- Vos, T., Barber, R. M., Bell, B., Bertozzi-Villa, A., Biryukov, S., Bolliger, I., ... Dicker, D. (2015). Global, regional, and national incidence, prevalence, and years lived with disability for 301 acute and chronic diseases and injuries in 188 countries, 1990–2013: A systematic analysis for the Global Burden of Disease Study 2013. *The Lancet*, 386(9995), 743–800. [https://doi.org/10.1016/S0140-6736\(15\)60692-4](https://doi.org/10.1016/S0140-6736(15)60692-4)
- Wang, F., Reece, E. A., & Yang, P. (2015). Oxidative stress is responsible for maternal diabetes-impaired transforming growth factor beta signaling in the developing mouse heart. *American Journal of Obstetrics and Gynecology*, 212(5), 650 e651–611. <https://doi.org/10.1016/j.ajog.2015.01.014>
- Wang, H., Naghavi, M., Allen, C., Barber, R. M., Bhutta, Z. A., Carter, A., ... Coates, M. M. (2016). Global, regional, and national life expectancy, all-cause mortality, and cause-specific mortality for 249 causes of death, 1980–2015: A systematic analysis for the Global Burden of Disease Study 2015. *The Lancet*, 388(10053), 1459–1544. [https://doi.org/10.1016/S0140-6736\(16\)31012-1](https://doi.org/10.1016/S0140-6736(16)31012-1)
- Wessels, M. W., & Willems, P. J. (2010). Genetic factors in non-syndromic congenital heart malformations. *Clinical Genetics*, 78(2), 103–123. <https://doi.org/10.1111/j.1399-0004.2010.01435.x>
- Wu, Y., Reece, E. A., Zhong, J., Dong, D., Shen, W. B., Harman, C. R., & Yang, P. (2016). Type 2 diabetes mellitus induces congenital heart defects in murine embryos by increasing oxidative stress, endoplasmic reticulum stress, and apoptosis. *American Journal of Obstetrics and Gynecology*, 215(3), 366. e1–366. e10. <https://doi.org/10.1016/j.ajog.2016.03.036>
- Xiao, Y., Taub, M. A., Ruczinski, I., Begum, F., Hetmanski, J. B., Schwender, H., ... Marazita, M. L. (2017). Evidence for SNP-SNP interaction identified through targeted sequencing of cleft case-parent trios. *Genetic Epidemiology*, 41(3), 244–250. <https://doi.org/10.1002/gepi.22023>
- Yoon, P. W., Rasmussen, S. A., Lynberg, M. C., Moore, C. A., Anderka, M., Carmichael, S. L., ... Edmonds, L. D. (2001). The National Birth Defects Prevention Study. *Public Health Reports*, 116(Suppl 1): 32–40. <https://doi.org/10.1093/phr/116.S1.32>
- Zhang, E., Hong, N., Chen, S., Fu, Q., Li, F., Yu, Y., & Sun, K. (2018a). Targeted sequencing identifies novel GATA6 variants in a large cohort of patients with conotruncal heart defects. *Gene*, 641, 341–348. <https://doi.org/10.1016/j.gene.2017.10.083>
- Zhang, K. K., Xiang, M., Zhou, L., Liu, J., Curry, N., Heine Suner, D., ... Xie, L. (2016). Gene network and familial analyses uncover a gene network involving Tbx5/Osr1/Pcsk6 interaction in the second heart field for atrial septation. *Human Molecular Genetics*, 25(6), 1140–1151. <https://doi.org/10.1093/hmg/ddv636>
- Zhang, M., Li, F. X., Liu, X. Y., Hou, J. Y., Ni, S. H., Wang, J., ... Yang, Y. Q. (2018b). TBX1 loss-of-function mutation contributes to congenital conotruncal defects. *Experimental and Therapeutic Medicine*, 15(1), 447–453. <https://doi.org/10.3892/etm.2017.5362>
- Zhang, W., Shen, L., Deng, Z., Ding, Y., Mo, X., Xu, Z., ... Yi, L. (2014). Novel missense variants of ZFPM2/FOG2 identified in conotruncal

heart defect patients do not impair interaction with GATA4. *PLoS ONE*, 9(7), e102379. <https://doi.org/10.1371/journal.pone.0102379>

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

How to cite this article: Lyu C, Webber DM, MacLeod SL, Hobbs CA, Li M. Gene-by-gene interactions associated with the risk of conotruncal heart defects. *Mol Genet Genomic Med.* 2020;8:e1010. <https://doi.org/10.1002/mgg3.1010>