


CORRECTION

Correction: Circulating Angiopoietin-2 and Its Soluble Receptor Tie-2 Concentrations Are Related to Renal Function in Two Population-Based Cohorts

The *PLOS ONE* Staff

The first row of Table 2 is missing. Please view the corrected table below. The publisher apologizes for the error.



OPEN ACCESS

Citation: The *PLOS ONE* Staff (2017) Correction: Circulating Angiopoietin-2 and Its Soluble Receptor Tie-2 Concentrations Are Related to Renal Function in Two Population-Based Cohorts. *PLoS ONE* 12(4): e0176477. <https://doi.org/10.1371/journal.pone.0176477>

Published: April 19, 2017

Copyright: © 2017 The PLOS ONE Staff. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Table 2. Associations between markers of renal function and serum angiotensin-2 concentration or serum Tie-2 concentration.

	All subjects						Subjects without hypertension or diabetes mellitus					
	Angiotensin-2			Tie-2			Angiotensin-2			Tie-2		
	Beta	Stderr	P	Beta	Stderr	P	Beta	Stderr	P	Beta	Stderr	P
SHIP-1												
Creatinine	-1.88E-03	1.16E-03	0.10	-9.80E-04	6.02E-03	0.87	1.09E-03	8.80E-04	0.22	-3.80E-03	1.03E-02	0.71
Creatinine'	1.67E-06	5.04E-07	<.01				-	-	-	-	-	-
eGFR(crea)	-3.53E-03	9.95E-04	<.01	-1.15E-03	5.21E-03	0.83	-7.05E-04	6.59E-04	0.29	3.05E-03	7.71E-03	0.69
eGFR(crea)'	9.76E-07	3.09E-07	<.01	-	-	-	-	-	-	-	-	-
Cystatin C	0.39	0.04	<.01	5.84	1.26	<.01	0.35	0.08	<.01	3.67	0.97	<.01
Cystatin C'	-	-	-	-15.03	4.87	<.01	-	-	-	-	-	-
eGFR(cys)	-5.53E-03	8.00E-04	<.01	-9.24E-03	9.54E-03	0.33	-3.73E-03	8.44E-04	<.01	-3.78E-02	9.88E-03	<.01
eGFR(cys)'	3.58E-07	1.74E-07	0.04	-5.08E-06	2.07E-06	0.01	-	-	-			
log(uACR)	3.67E-02	7.10E-03	<.01	0.20	0.08	0.02	3.51E-02	1.23E-02	<.01	0.28	0.15	0.05
SHIP-Trend												
Creatinine	-2.39E-03	9.76E-04	0.01	9.99E-03	4.80E-03	0.04	-2.97E-03	1.54E-03	0.05	1.31E-02	7.47E-03	0.08
Creatinine'	1.64E-06	4.41E-07	<.01	-	-	-	2.37E-06	9.18E-07	0.01	-	-	-
eGFR(crea)	-3.48E-03	8.03E-04	<.01	-7.69E-03	3.78E-03	0.04	-2.42E-03	1.22E-03	0.05	-7.02E-03	5.16E-03	0.17
eGFR(crea)'	1.12E-06	2.40E-07	<.01	-	-	-	9.51E-07	4.22E-07	0.02	-	-	-
Cystatin C	0.25	0.12	0.03	1.18	0.54	0.03	0.05	0.20	0.81	2.53	0.90	<.01
Cystatin C'	1.67	0.86	0.05	-	-	-	5.81	2.27	0.01	-	-	-
eGFR(cys)	-4.79E-03	4.99E-04	<.01	-1.20E-02	5.53E-03	0.03	-7.76E-03	1.29E-03	<.01	-2.50E-02	9.19E-03	0.01
eGFR(cys)'	-	-	-	-	-	-	1.45E-06	5.98E-07	0.02	-	-	-
log(uACR)	5.47E-02	7.10E-03	<.01	-0.60	0.22	<.01	6.46E-02	1.26E-02	<.01	-1.01	0.39	<.01
log(uACR)'	-	-	-	9.52E-02	3.62E-02	<.01	-	-	-	0.23	0.09	0.01

eGFR = estimated glomerular filtration rate based on either serum creatinine [eGFR(crea)] or serum cystatin C concentration [eGFR(cys)]. *Models were adjusted for age, sex, waist circumference, smoking, total cholesterol, systolic and diastolic blood pressure and additionally in the whole population for diabetes mellitus type 2 and use of antihypertensive or antidiabetic medication. Creatinine', eGFR' and cystatin C' represent spline components, for more detail see method section.

<https://doi.org/10.1371/journal.pone.0176477.t001>

Reference

1. Hennings A, Hannemann A, Rettig R, Dörr M, Nauck M, Völzke H, et al. (2016) Circulating Angiotensin-2 and Its Soluble Receptor Tie-2 Concentrations Are Related to Renal Function in Two Population-Based Cohorts. PLoS ONE 11(11): e0166492. <https://doi.org/10.1371/journal.pone.0166492> PMID: 27893762