

## RESEARCH ARTICLE

# Systematic review and standardised assessment of Chinese cross-cultural adapted hip Patient Reported Outcome Measures (PROMs)

James Reeves Mbori Ngwayi<sup>1</sup>✉\*, Jie Tan<sup>1</sup>, Ning Liang<sup>1</sup>, Emmanuel Gildas Eric Sita<sup>2</sup>, Kenedy Uzoma Obie<sup>3</sup>, Daniel Edward Porter<sup>4</sup>✉

**1** School of Clinical Medicine, Tsinghua University, Beijing, China, **2** School of Clinical Medicine, Fujian Medical University, Fujian, China, **3** School of Clinical Medicine, Central South University, Changsha, China, **4** Department of Orthopedics, Beijing Huaxin Hospital, Clinical Medicine School, Tsinghua University, Beijing, China

✉ These authors contributed equally to this work.

\* [zipromax@163.com](mailto:zipromax@163.com)



## OPEN ACCESS

**Citation:** Ngwayi JRM, Tan J, Liang N, Sita EGE, Obie KU, Porter DE (2021) Systematic review and standardised assessment of Chinese cross-cultural adapted hip Patient Reported Outcome Measures (PROMs). PLoS ONE 16(9): e0257081. <https://doi.org/10.1371/journal.pone.0257081>

**Editor:** Sinan Kardeş, Istanbul University Istanbul Faculty of Medicine: Istanbul Universitesi Istanbul Tip Fakultesi, TURKEY

**Received:** December 16, 2020

**Accepted:** August 23, 2021

**Published:** September 20, 2021

**Copyright:** © 2021 Ngwayi et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data are within the paper and its [Supporting information](#) files.

**Funding:** The author(s) received no specific funding for this work.

**Competing interests:** The authors have declared that no competing interests exist.

## Abstract

### Purpose

To perform a systemic literature search to identify Chinese cross culturally adapted and new designed Patient Reported Outcome Measures (PROMs) used for hip assessment, then a standardized evaluation of available instruments in order to provide evidence of high-quality PROMs for clinical use and adoption in future hip registries.

### Methods

A Systematic Review of the following databases: PUBMED, CINAHL, EMBASE, CNKI was performed to identify relevant PROMs. Instruments underwent standardized assessment and scoring using the EMPRO tool by two independent reviewers. Inter-rater reliability was assessed using intra-class correlation coefficients (ICC).

### Results

2188 articles were retrieved, with seven articles fitting the inclusion criteria consisting of six hip PROMs. Five PROMs were cross culturally adapted and one was originally designed in Mandarin Chinese. Total scores (/100) after EMPRO evaluation: Osteoarthritis of Knee and Hip Quality of Life (OAKHQOL): 55; Copenhagen Hip and Groin Outcome Score (HAGOS): 52; International Hip Outcome Tool (SC-iHOT-33): 45; Hip Disability and Osteoarthritis Outcome Score (HOOS): 37; Questionnaire on the Perceptions and Functions of Patients about Total Hip Arthroplasty (QPFPTHA): 36; Oxford Hip Score (OHS): 35. ICC values were 0.73 for the SC-iHOT-33 and ranged between 0.83–0.93 for the other PROMs indicating good to excellent inter-rater agreement.

## Conclusion

Among the commonly used hip-specific PROMs found in arthroplasty registries, none of the Chinese adapted versions evaluated by EMPRO is currently rated acceptable for clinical use. Only OAKHQOL and HAGOS reached acceptability threshold. Further research on the attributes of cross-cultural adaptation, interpretability and burden assessment would be helpful.

## Introduction

Hip arthroplasty constitutes one of the most cost-effective interventions in medicine [1]; reported cost-effectiveness ratios based on quality of life years (QALY) for common procedures include: £1372 for hip arthroplasty; £2101/QALY for knee arthroplasty; £3129–6904/QALY for tonsillectomy with adenoidectomy; £4928/QALY for inguinal hernia repair; £13205/QALY for laparoscopic cholecystectomy [2]. Patient Reported Outcome Measures (PROMs) assist in QALY calculation and are typically clinical outcome tools which give a voice to patients, documenting their own perspective without clinician interpretation [3–5]. PROMs increasingly dominate clinical outcome appraisal following advocacy of the central role of patients in their own health care affairs. A variety of hip related PROMs have been developed and are already in worldwide use [6, 7]. Most were originally designed in English. Before their deployment in another linguistic setting they must undergo rigorous transcultural adaptation and translation following guidelines such as International Society for Pharmacoeconomics and Outcomes Research (ISPOR) [8]. As with the English language original versions, implementation of culturally adapted PROMs should follow a quality control process for clinical use. Instruments to assess methodological quality of Health-related Patient Reported Outcomes (HR-PROs) include the Consensus Based Standards for the Selection of Health Measurement Instruments (COSMIN) checklist [9, 10]; and the Evaluating the Measurement of Patient-Reported Outcomes (EMPRO) tool, capable of both qualitative and quantitative assessment of PROs [11].

Arthroplasty registries around the world have expanded their endpoints from failure events such as revision surgery, infection and dislocation to include subjective endpoints such as pain relief, increased functionality and quality of life, involving a shift towards adopting PROMs within the protocol for clinical assessment [12]. Although there are no national or provincial arthroplasty registries in mainland China, it is important to evaluate and document the quality of PROMs presently in use to provide appropriate tools for future registry development. More immediately, surgeons should be aware of the best quality PROMs available for use in their patient population. The aim of the study was to carry out a systematic literature search for Chinese cross-culturally adapted PROMs and new Chinese-designed PROMs for use in the Chinese patients with hip disorders and evaluation using the EMPRO assessment instrument.

## Methods

### Systematic literature review and identification of hip studies

A search was performed of the earliest records up to 22/08/2020 according to guidelines of the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (S1 Checklist) [13]. The following databases were used: PubMed/MEDLINE, EMBASE(OVID), CINAHL(EBSCO) and CNKI (mainland Chinese database). Sensitive filters composed of MeSH terms and keywords based on previously documented PROM search strategies [14–16] (S1 File), were further

tailored to fit the target body region and population (Chinese) then applied to the databases. Publication languages for the articles were English and Chinese.

Based on Population, Intervention, Comparison, Outcome (PICO) criteria the following inclusion criteria were adapted: 1. Cross culturally adapted and translated hip PROMS tested in the Mainland Chinese population. 2. Hip specific PROMs evaluating hip procedures 3. PROMs restricted to the Chinese Mainland in Mandarin Chinese.

Exclusion Criteria were: 1. Non-Mainland Chinese PROMs including Mandarin language studies from geographical Taiwan and Hong Kong. 2. Instruments tested on populations out-with Mainland China. 3. Articles not meeting inclusion criteria.

Screening was carried out as a three-step process (Titles, Abstract and Full texts), performed independently by two reviewers (NJRM, DEP). Outputs were compared and consensus reached. After full text screen and identification of suitable articles we manually reviewed the in-article reference lists for potential relevant articles missed during the electronic search. The PRISMA chart of the review process is illustrated in [Fig 1](#).

### The EMPRO tool


The Evaluation Measurement of Patient Reported Outcomes (EMPRO) tool was designed in 2008 by Valderas JM et al, aimed at qualitative assessment of the methodology and development process of PROs. It comprises 8 attributes, namely: conceptual and measurement model (7 items), cross-cultural and linguistic adaptations (3 items), reliability (8 items), validity (6 items), responsiveness to change (3 items), interpretability (3 items), burden (7 items), alternative modes of administration (2 items); in total 39 items [11]. Quantitative assessment is achieved via an inherent scoring system where each item is graded on a 4-point Likert scale, from 4 (strongly agree) to 1 (strongly disagree) with additional option boxes for “No information” and “Not applicable” where necessary.

The questionnaire terminates with a rationale for recommendation by the reviewer according to the following response scale: “Strongly recommended”, “Recommended with provisos or alterations”, “Would not recommend” and “Unsure”.

The EMPRO tool requires a license application via the portal [www.bibliopro.org](http://www.bibliopro.org), which is free to-use, and presently available in two languages (English and Spanish). EMPRO has been used before in the systematic assessment of orthopedic-specific shoulder PROMs [17].

### Standardized and systematic evaluation; scoring and analysis

After the systematic review, data pertaining to the measurement properties of the selected PROMs studies were extracted from relevant articles. A standardized assessment of the adequacy of their measurement properties was undertaken using the EMPRO tool. To avoid possible bias due to evaluating cross cultural adaptation across different populations, assessment was restricted to studies on Chinese populations [14]. According to recommendations from the designers, two reviewers (both clinicians with a background in PROMs research) performed the assessment. Both had also completed the online EMPRO training webinar (<https://www.isoqol.org/category/webinar/page/3/>). The assessment was carried out in two phases. The first phase consisted of each reviewer independently scoring article(s) supporting each cross-culturally adapted hip PROM for methodological attributes, as well as the article describing the original design of the PROM for the conceptual and measurement model assessment. The second phase which followed a consensus method recommended by the EMPRO designers involved discussions between reviewers on discrepancies to obtain a common score for each item. Reviewers were based in two continents and did not converse on scoring until the discussion phase.



**Fig 1. PRISMA flow chart—Systematic literature search.**

<https://doi.org/10.1371/journal.pone.0257081.g001>

Scoring of the methodological attributes was calculated based on developers' instructions; <https://www.isoqol.org/?s=Empro>. Specifically, attribute-specific scores are obtained by calculating the response mean of the applicable items when at least 50% of them are rated; and items check marked with the option "no information" are assigned a score of 1 representing the lowest possible score. The response means for each attribute are then linearly transformed to a range of 0 to 100 (worst to best). Global scores (based on metric properties) are only calculated when at least 3 attributes can be scored and attributes without information are imputed zero. Panoramic assessment (which includes all culture/language versions of the instrument) involves conceptual model, reliability, validity, responsiveness and interpretability while culture/language specific evaluation involves conceptual model, reliability, validity,

responsiveness and interpretability and cross-cultural adaptation. Global scores for each domain of at least 50 (/100) are considered acceptable based on previous assessments by Garin O et al. [18].

Data synthesis involves combining results from different studies on measurement properties of the instrument under consideration. Results of measurement properties can differ depending on sample characteristics and setting; thus (dis)similarities have to be appropriately addressed before deciding whether articles contain measurement property data which can be pooled [19]. After the first phase scoring by reviewers, agreement between them was assessed by using two-way, random, single unit, absolute agreement intra-class correlation coefficients ICC [20]. The degree of reviewer agreement was categorized based on Cicchetti (1994) ICC < 0.40 poor; 0.40 to 0.59 moderate; 0.60–0.74 good; 0.75–1.00 excellent [21].

Statistical analysis, attribute scores calculation and graphics were performed with Microsoft Excel 2003<sup>®</sup> (Microsoft, Redmond, WA, USA), Intraclass correlation (ICC) was calculated using SPSS<sup>®</sup> Version 20.0 software (SPSS Inc, Chicago, IL, USA).

## Results

The literature search procedure resulted in a total of 2188 papers. Articles per database were as follows: PubMed/MEDLINE 1671 (76.4%), EMBASE 331 (15.1%), CINAHL 147 (6.7%) and CNKI 39 (1.8%). A total of seven papers were retained at the end of the process representing six hip-specific instruments. Five of the instruments were cross-cultural adaptations: Copenhagen Hip and Groin Outcome Score (HAGOS); International Hip Outcome Tool (SC-iHOT-33); Osteoarthritis of Knee and Hip Quality of Life (OAKHQOL); Oxford Hip Score (OHS); Hip Disability and Osteoarthritis Outcome Score (HOOS); and one instrument was Chinese-designed: Questionnaire on the Perceptions and Functions of Patients about Total Hip Arthroplasty (QFPPTHA). All papers were published between 2010 and 2019, two questionnaires were unidimensional while four had dimensions ranging from 2–6 dimensions. One PROM was validated for both knee and hip arthroplasty (OAKHQOL). [Table 1](#) describes general characteristics of each Chinese Cross-cultural adaptation and translation of hip PROM.

Qualities of measurement property domains/attributes for each hip PROM are illustrated in [Table 2](#). Following precedent in other studies [17] the global score for each domain was transformed into a five-point scale (denoted—/ + / ++ / +++ / ++++). +: EMPRO score < 25; ++: EMPRO score 25–49; +++: EMPRO score 50–74; ++++: EMPRO score 75–100; -: EMPRO score not applicable or not calculable according to designer instructions.

The properties of each PROM were assessed and scored with the EMPRO tool, for each of the attributes as well as a total score for the PROM. All questionnaires had acceptable scores for the concept and measurement model except OHS which scored ++. Questionnaires were all evaluated for cultural and language adaptations except the QFPPTHA originally designed in simplified Chinese. The HAGOS and the OAKHQOL had acceptable scores while the others could not be calculated ([Fig 2](#)). All instruments had acceptable scores on assessment of reliability ([Fig 3](#)) and responsiveness except the QFPPTHA which scored ++ in these attributes. On evaluation of validity all scores were above threshold for acceptability ([Fig 4](#)); however, data was insufficient for calculating interpretability. Burden scores could not be calculated for HAGOS, and for all remaining PROMs scored below acceptability threshold. Total scores were calculated for each PROM; OAKHQOL and HAGOS had scores above acceptability threshold (54 and 52 respectively) while SC-iHOT-33 scored 45, HOOS scored 36, QFPPTHA scored 36 and OHS scored 35 ([Fig 5](#)).

**Table 1. General characteristics of Chinese cross-cultural adaptation and translation of hip PROMs.**

Questionnaire	Author—Year	Population Characteristic	Dimensions and (items)	Scale design
Copenhagen Hip and Groin Outcome Score (HAGOS) [22]	Shi Qi Cao et al (2018)	Developmental dysplasia of hip joint, osteonecrosis of the femoral head, hip osteoarthritis for THA	6 Domains; symptoms (7 items), pain (10 items), function in ADL (5 items), function in Sport/Rec (8 items), participation in physical activities (2 items), hip- and/or groin-related QoL (5 items).	Standardized from 0 (worst health status) to 100 best health status
		Mean age 64 years		
International Hip Outcome Tool (SC-iHOT-33) [23]	D.H. Li et al (2016)	Avascular necrosis, Hip dysplasia, Femoro-acetabular impingement syndrome for THA	4 Domains; symptoms and functional limitations (16 items, Sport/Rec activities (6 items) job-related concerns (4 items) social, emotional, and lifestyle concerns (7 items).	Standardized from 0 (worst health status) to 100 best health status
		Mean age 43 years		
Osteoarthritis of Knee and Hip Quality of Life (OAKHQOL) [24]	W. Wang et al. (2016)	Osteoarthritis for both hip and knee Arthroplasty	5 Domains; physical activities (16 items), mental health (13 items), pain (4 items), social support (4 items) and social activities (3 items). The rest are three independent items related to relationships, sexual activity and professional life.	Standardized on a scale from 0 (worst quality of life) to 100 (best quality of life)
		Mean age 63 years		
Oxford Hip Score (OHS) - 2 articles [25, 26]	Wei Zheng et al (2013)	Hip OA for THA	12-items	0 (worst) to 48 (best).
		Mean age 66 years		
	Xia Zhen-lan et al (2012)	THA Pre-operation, 1-year post THA and 3-years post THA		
		Mean age 55 years		
Hip disability and osteoarthritis outcome score (HOOS) [27]	X. Wei et al (2012)	Primary hip OA for THA	5 Domains; pain (10 items), other symptoms (10 items), function in (ADL (17 items), function in Sport/Rec (4 items), hip-related QoL (4 items).	Normalized scores, from 0 (indicating extreme symptoms) to 100 (indicating no symptoms),
		Mean age 51 years		
Questionnaire on the perceptions and functions of patients about Total Hip Arthroplasty [28]	Tang Hong Yuan et al (2010)	Femoral head necrosis, emoral neck fracture, hip OA, hip joint dysplasia, AS, Bone tumours, hip joint tuberculosis for THA	10 items	10 (worst) to 50 (best).
		Mean age 57 years		

THA: Total Hip Arthroplasty; OA: Osteoarthritis; AS: Ankylosing spondylitis; QOL: quality of life; ADL: activities of daily living; Sport/Rec: sports and recreation.

<https://doi.org/10.1371/journal.pone.0257081.t001>


**Table 2. Standardized assessment of measurement attributes of Chinese cross-culturally adapted PROMs using EMPRO.**

PROM	Concept and measurement model	Cultural and Language Adaptations	Reliability	Validity	Responsiveness	Interpretability	Burden	Alternative forms of Administration	Total Score /100
OAKHQOL	++++	+++	+++	++++	+++	-	+	-	54.84
HAGOS	+++	+++	+++	++++	+++	-	-	-	52.19
SC-iHOT-33	+++	-	+++	+++	+++	-	++	-	45.05
HOOS	+++	-	+++	+++	+++	-	++	-	36.98
QFPPTH	+++	-	++	+++	++	-	+	-	36.02
OHS	++	-	+++	+++	+++	-	+	-	35.19

Copenhagen Hip and Groin Outcome Score (HAGOS); International Hip Outcome Tool (SC-iHOT-33); Osteoarthritis of Knee and Hip Quality of Life (OAKHQOL); Oxford Hip Score (OHS); Hip disability and Osteoarthritis Outcome Score (HOOS); Questionnaire on the Perceptions and Functions of Patients about Total Hip Arthroplasty (QFPPTH);

+: EMPRO score < 25; ++: EMPRO score 25–49; +++: EMPRO score 50–74; ++++: EMPRO score 75–100; -: EMPRO score not applicable or not calculable.

<https://doi.org/10.1371/journal.pone.0257081.t002>




**Fig 2. Global attribute score for each PROM as evaluated by EMPRO.**

<https://doi.org/10.1371/journal.pone.0257081.g002>

Intra-class correlation coefficient (ICC) values for the assessment of each PROM are depicted in Table 3. The agreement between the two reviewers ranged between ICC = 0.83–0.93 except for the SC-iHOT-33 with ICC of 0.73.


## Discussion

Global EMPRO scores from the assessment exercise indicated two PROMs had total scores above threshold, with the highest being OAKHQOL, followed by HAGOS. Four instruments



**Fig 3. Cultural and language adaptation attribute score for each PROM as evaluated by EMPRO.**

<https://doi.org/10.1371/journal.pone.0257081.g003>




**Fig 4. Reliability attribute score for each PROM as evaluated by EMPRO.**

<https://doi.org/10.1371/journal.pone.0257081.g004>

could not be scored for cultural and language adaptation and differential item functioning and harmonization was often under documented.

Reliability, validity and responsiveness properties all had scores greater than threshold except for QPFPTH. We found that these domains were more thoroughly evaluated than the others. Although interpretability is not a measurement property as it does not refer to the inherent quality of an instrument, it is a measure of practicality for clinical and research use.



**Fig 5. Validity attribute score for each PROM as evaluated by EMPRO.**

<https://doi.org/10.1371/journal.pone.0257081.g005>



**Table 3. Inter-rater reliability for 7 articles describing PROMs.**

PROM	Intra-class correlation coefficient (ICC)
OHS(zheng et al)	.934
OHS(Xia et al)	.841
OAKHQOL	.856
HAGOS	.916
HOOS	.881
SC-iHOT-33	.732
QFPFTHA	.832

Copenhagen Hip and Groin Outcome Score (HAGOS); International Hip Outcome Tool (SC-iHOT-33); Osteoarthritis of Knee and Hip Quality of Life (OAKHQOL); Oxford Hip Score (OHS); Hip disability and Osteoarthritis Outcome Score (HOOS); Questionnaire on the Perceptions and Functions of Patients about Total Hip Arthroplasty (QFPFTHA);

<https://doi.org/10.1371/journal.pone.0257081.t003>

This property often receives less attention in studies of instrument quality [19]. All hip PROMs had limited data on this attribute and scores could not be assigned.

Except for OHS which had two studies on measurement properties, all other assessed hip PROMs were supported by a single paper. Opportunities for gaining improved scores may have been restricted by the limited data set for analysis. More holistic assessments can be made where several articles describe attribute data which can be pooled [17].

Arthroplasty registries use PROMs because recording a high rate of technical success after arthroplasty does not capture the variability in pain and function experienced by patients [12]. In 2016, up to 18 national arthroplasty registries used PROMs which are published in their reports [29]. Rolfson et al [30], identified that hip-specific PROMs used in arthroplasty registries include OHS, HOOS, HOOS short form (joint replacement), Harris Hip Score (strictly not a PROM since it has a clinician-assessed component), Western Ontario and McMaster Universities Arthritis Index (WOMAC) and University of California at Los Angeles (UCLA) Activity Score. According to the International society of Arthroplasty Registries (ISAR) Working Group there is no single best generic or hip-specific PROM [12] however it is self-evident that PROMs have to be developed in appropriate patient population and have good measurement properties. Measurements of quality have varied between researchers, for example with the OHS [15, 31]. Some of these differences seem to be due to the availability of new evidence in later publications, reflecting a dynamic process of accumulating evidence of PROM utility.

In this study, reviewers were able to recommend all evaluated PROMs, but '*with provisos and alterations*'. Among these, only OHS and HOOS are commonly used hip-specific PROMs in registries worldwide. However, these failed to achieve threshold for acceptability using EMPRO scoring criteria. We identified HAGOS and OAKHQOL as having acceptable global scores. However, HAGOS is designed specifically for younger individuals with hip disease, and OAKHQOL is not commonly used in arthroplasty registries [30]. Chinese patient groups used to test the culturally-adapted PROMs mainly had osteoarthritis, although rheumatoid arthritis, avascular necrosis and fracture patients were included. All studies which tested responsiveness used arthroplasty as the intervention. All cohorts were younger than might be expected in arthroplasty patient groups elsewhere; mean age of each cohort in this series ranged from 43 to 66 years (Table 1).

Limitations of this study include that it relies on a single instrument (EMPRO) to assess measurement qualities semi-quantitatively, so the weighting of attribute and items within

attributes has certain subjective features; however, this study is the first assessment of cross culturally adapted Chinese-language hip PROMs within a dynamic evaluative process. We used two experienced reviewers; more would have added further reliability to the assessment, however they achieved good-excellent ICC scores for inter-rater agreement.

## Recommendations and conclusions

Among the commonly used hip-specific PROMs found in arthroplasty registries, none of the Chinese adapted versions evaluated by EMPRO is currently rated acceptable for clinical use. Compared to other non-English language environments which have developed PROM initiatives such as BiblioPro, a Spanish language-based PROM online library [32], there are none presently in China. We recommend that further studies on measurement qualities of these PROMs should remain a priority and include older patient groups. Deficient domains include cultural and language adaptation, interpretability and burden. WOMAC and UCLA Activity Score are commonly found in registries, and have been culturally adapted for Chinese patients with knee osteoarthritis but not for the hip [33, 34]. Since Chinese patients face barriers to follow-up after surgery due to large travel distances and lack of resources, the properties of electronic or telephonic methods of administration should also receive attention. Similar to the conclusion of researchers in other non-English language settings [14], we also cannot recommend any single hip-specific Chinese-mainland adapted PROM at present.

## Supporting information

**S1 Checklist. PRISMA 2009 checklist.**  
(DOC)

**S1 File. Pubmed/Medline filter: Psychometric properties of specific PROMs questionnaires in the Chinese population.**  
(PDF)

## Author Contributions

**Conceptualization:** James Reeves Mbori Ngwayi, Daniel Edward Porter.

**Data curation:** James Reeves Mbori Ngwayi, Jie Tan, Ning Liang.

**Formal analysis:** James Reeves Mbori Ngwayi, Daniel Edward Porter.

**Funding acquisition:** James Reeves Mbori Ngwayi, Daniel Edward Porter.

**Investigation:** James Reeves Mbori Ngwayi, Jie Tan, Daniel Edward Porter.

**Methodology:** James Reeves Mbori Ngwayi, Daniel Edward Porter.

**Project administration:** James Reeves Mbori Ngwayi, Jie Tan, Daniel Edward Porter.

**Resources:** James Reeves Mbori Ngwayi, Ning Liang, Emmanuel Gildas Eric Sita, Kenedy Uzoma Obie.

**Software:** James Reeves Mbori Ngwayi, Jie Tan, Kenedy Uzoma Obie.

**Supervision:** James Reeves Mbori Ngwayi, Daniel Edward Porter.

**Validation:** James Reeves Mbori Ngwayi, Daniel Edward Porter.

**Visualization:** James Reeves Mbori Ngwayi, Emmanuel Gildas Eric Sita, Kenedy Uzoma Obie, Daniel Edward Porter.

**Writing – original draft:** James Reeves Mbori Ngwayi.

**Writing – review & editing:** James Reeves Mbori Ngwayi, Daniel Edward Porter.

## References

1. Crawford RW and Murray DW. Total hip replacement: indications for surgery and risk-factors for failure. *Ann. Rheum. Dis.* 1997; 56:455–457. <https://doi.org/10.1136/ard.56.8.455> PMID: 9306866
2. Chen TC, Wanniarachige D, Murphy S, Lockhart K, O'Mahony J. Surveying the Cost-Effectiveness of the 20 Procedures with the Largest Public Health Services Waiting Lists in Ireland: Implications for Ireland's Cost-Effectiveness Threshold. *Value Health.* 2018 Aug; 21(8):897–904. <https://doi.org/10.1016/j.jval.2018.02.013> PMID: 30098666. Epub 2018 Jun 12.
3. Balck F, Kirschner S, Jeszenszky C, Lippmann M, Gunther KP. Validity and reliability of the German version of the HSS Expectation Questionnaire on Hip Joint Replacement. *Z Orthop Unfall.* 2016; 154:606. <https://doi.org/10.1055/s-0042-111329> PMID: 27612312
4. Rissanen P, Aro S, Slätis P, Sintonen H, Paavolainen P. Health and quality of life before and after hip or knee arthroplasty. *J Arthroplasty* 1995; 10: 169–75. [https://doi.org/10.1016/s0883-5403\(05\)80123-8](https://doi.org/10.1016/s0883-5403(05)80123-8) PMID: 7798097
5. Räsänen P, Paavolainen P., Sintonen H, Koivisto AM, Blom M, Ryyänen OP, et al. Effectiveness of hip or knee replacement surgery in terms of quality-adjusted life years and costs. *Acta Orthopaedica*, 2007. 78: p. 108–115. <https://doi.org/10.1080/17453670610013501> PMID: 17453401
6. Rolfson O, Eresian Chenok K, Bohm E, et al. Patient-Reported Outcome Measures Working Group of the International Society of Arthroplasty Registries. Patient-reported outcome measures in arthroplasty registries. *Acta Orthop* 2016; 87:3–8. <https://doi.org/10.1080/17453674.2016.1181815> PMID: 27168175
7. What is Patient-centered Health Care? A Review of Definitions and Principles. 2nd ed. London: IAPO; 2007. International Alliance of Patients' Organizations; pp. 1–34.
8. Wild D, Grove A, Martin M, Eremenco S, McElroy S, Verjee-Lorenz A, et al. Principles of Good Practice for the Translation and Cultural Adaptation Process for Patient-Reported Outcomes (PRO) Measures: Report of the ISPOR Task Force for Translation and Cultural Adaptation. *Value in Health*, Volume 8. Number 2. 2005. <https://doi.org/10.1111/j.1524-4733.2005.04054.x> PMID: 15804318
9. Mokkink LB, Terwee CB, Patrick DL, Alonso J, Stratford PW, Knol DL, et al. The COSMIN checklist for assessing the methodological quality of studies on measurement properties of health status measurement instruments: an international Delphi study. *Qual Life Res* 2010; 19:539–49. <http://dx.doi.org/10.1007/s11136-010-9606-8> PMID: 20169472
10. Terwee CB, Mokkink LB, Knol DL, Ostelo RW, Bouter LM, de Vet HC. Rating the methodological quality in systematic reviews of studies on measurement properties: a scoring system for the COSMIN checklist. *Qual Life Res* 2012; 21:651–7. <http://dx.doi.org/10.1007/s11136-011-9960-1> PMID: 21732199
11. Valderas JM, Ferrer M, Mendivil J, Garin O, Rajmil L, Herdman M, et al. Development of EMPRO: a tool for the standardized assessment of patient-reported outcome measures. *Value Health* 2008; 11:700–8. <http://dx.doi.org/10.1111/j.1524-4733.2007.00309.x> PMID: 18194398
12. Wilson I, Bohm E, Lübbecke A, Lyman S, Overgaard S, Rolfson O, et al. Orthopaedic registries with patient-reported outcome measures. *EFORT Open Rev.* 2019 Jun 3; 4(6):357–367. <https://doi.org/10.1302/2058-5241.4.180080> PMID: 31210973.
13. Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gøtzsche PC, Ioannidis JPA, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *Journal of clinical epidemiology.* 2009. pp. e1–34. <https://doi.org/10.1016/j.jclinepi.2009.06.006> PMID: 19631507
14. Arias-de la Torre J, Puigdomenech E, Valderas JM, Evans JP, Martín V, Molina AJ, et al. Availability of specific tools to assess patient reported outcomes in hip arthroplasty in Spain. Identifying the best candidates to incorporate in an arthroplasty register. A systematic review and standardized assessment. *PLoS One.* 2019 Apr 1; 14(4):e0214746. <https://doi.org/10.1371/journal.pone.0214746> PMID: 30934024.
15. Harris K, Dawson J, Gibbons E, Lim C, Beard D, Fitzpatrick R, et al. Systematic review of measurement properties of patient-reported outcome measures used in patients undergoing hip and knee arthroplasty. *Patient Relat Outcome Meas.* 2016; Volume 7: 101–108. <https://doi.org/10.2147/PROM.S97774> PMID: 27524925
16. Terwee CB, Jansma EP, Riphagen II, de Vet HCW. Development of a methodological PubMed search filter for finding studies on measurement properties of measurement instruments. *Qual Life Res.* 2009; 18: 1115–1123. <https://doi.org/10.1007/s11136-009-9528-5> PMID: 19711195

17. Schmidt S et al. Evaluation of shoulder-specific patient-reported outcome measures: a systematic and standardized comparison of available evidence. *J Shoulder Elbow Surg* (2014) 23, 434–444 <https://doi.org/10.1016/j.jse.2013.09.029> PMID: 24406123
18. Garin O, Herdman M, Vilagut G, Ferrer M, Ribera A, Rajmil L, et al. Assessing health-related quality of life in patients with heart failure: a systematic, standardized comparison of available measures. *Heart Fail Rev*. 2014 May; 19(3):359–67. <https://doi.org/10.1007/s10741-013-9394-7> PMID: 23681849.
19. de Vet Henrica C. W., Terwee Caroline B., Mokkink Lidwine B., Knol Dirk L.; *Measurement in Medicine: A Practical Guide*; Cambridge University Press, 11 August 2011.
20. Hallgren KA. Computing Inter-Rater Reliability for Observational Data: An Overview and Tutorial. *Tutor Quant Methods Psychol*. 2012; 8(1):23–34. <https://doi.org/10.20982/tqmp.08.1.p023> PMID: 22833776.
21. Cicchetti DV. Guidelines, criteria, and rules of thumb for evaluating normed and standardized assessment instruments in psychology. *Psychol Assess* 1994; 6 (4):284–90. <http://doi.apa.org/getdoi.cfm?doi=10.1037/1040-3590.6.4.284>
22. Cao S, Cao J, Li S, Wang W, Qian Q, Ding Y. Cross-cultural adaptation and validation of the Simplified Chinese version of Copenhagen Hip and Groin Outcome Score (HAGOS) for total hip arthroplasty. *J Orthop Surg Res*. 2018 Nov 6; 13(1):278. <https://doi.org/10.1186/s13018-018-0971-2> PMID: 30400966.
23. Li DH, Wang W, Li X, Gao YL, Liu DH, Liu DL, et al. Development of a valid Simplified Chinese version of the International Hip Outcome Tool (SC-iHOT-33) in young patients having total hip arthroplasty. *Osteoarthritis Cartilage*. 2017 Jan; 25(1):94–98. <https://doi.org/10.1016/j.joca.2016.08.013> PMID: 27621215. Epub 2016 Sep 9.
24. Wang W, He C- R, Zheng W, Li J, and Xu W- D. Development of a valid simplified Chinese version of the Osteoarthritis of Knee and Hip Quality of Life (OAKHQOL) in patients with knee or hip osteoarthritis. *Journal of Evaluation in Clinical Practice*, (2016) 22: 53–61. <https://doi.org/10.1111/jep.12431> PMID: 26239829
25. Zheng W, Li J, Zhao J, Liu D, Xu W. Development of a valid simplified Chinese version of the Oxford Hip Score in patients with hip osteoarthritis. *Clin Orthop Relat Res*. 2014 May; 472(5):1545–51. <https://doi.org/10.1007/s11999-013-3403-y> PMID: 24310602. Epub 2013 Dec 6.
26. Zhen-lan Xia, Bing He, Mei-Xia Fan, Xue-Qin Liu. Reliability and Validity of Chinese Version of Oxford Hip Score. *Journal of Nursing(China)*. March, 2012, Vol. 19 No. 3A.
27. Wei X, Wang Z, Yang C, Wu B, Liu X, Yi H, et al. Development of a simplified Chinese version of the Hip Disability and Osteoarthritis Outcome Score (HOOS): cross-cultural adaptation and psychometric evaluation. *Osteoarthritis Cartilage*. 2012 Dec; 20(12):1563–7. <https://doi.org/10.1016/j.joca.2012.08.018> PMID: 22944526. Epub 2012 Aug 31.
28. Tang HY, Yang MY, Zhang LM. Development and evaluation of questionnaire on the perceptions and function of patients about total hip arthroplasty. *Chin J Mod Nurs*. November 2010. Vol16. No.33
29. Rolfson O, Bohm E, Franklin P, et al; Patient-reported outcome measures working group of the international society of Arthroplasty registries. Patient-reported outcome measures in arthroplasty registries Report of the Patient-Reported Outcome Measures Working Group of the International Society of Arthroplasty Registries Part II. Recommendations for selection, administration, and analysis. *Acta Orthop* 2016; 87:9–23. <https://doi.org/10.1080/17453674.2016.1181816> PMID: 27228230
30. Rolfson O, Rothwell A, Sedrakyan A, et al. Use of patient-reported outcomes in the context of different levels of data. *J Bone Joint Surg [Am]* 2011; 93:66–71. <https://doi.org/10.2106/JBJS.K.01021> PMID: 22262427
31. Alviar MJ, Olver J, Brand C, Tropea J, Hale T, Pirpiris M, et al. Do patient-reported outcome measures in hip and knee arthroplasty rehabilitation have robust measurement attributes? A systematic review. *J Rehabil Med*. 2011 Jun; 43(7):572–83. <https://doi.org/10.2340/16501977-0828> PMID: 21607295.
32. <https://www.bibliopro.org>
33. Symonds T, Hughes B, Liao S, Ang Q, Bellamy N. Validation of the Chinese Western Ontario and McMaster Universities Osteoarthritis Index in Patients From Mainland China With Osteoarthritis of the Knee. *Arthritis Care Res (Hoboken)*. 2015 Nov; 67(11):1553–60. <https://doi.org/10.1002/acr.22631> PMID: 26018634.
34. Cao S, Liu N, Li L, Lv H, Chen Y, Qian Q. Simplified Chinese Version of University of California at Los Angeles Activity Score for Arthroplasty and Arthroscopy: Cross-Cultural Adaptation and Validation. *J Arthroplasty*. 2017 Sep; 32(9):2706–2711. <https://doi.org/10.1016/j.arth.2017.03.057> PMID: 28483213. Epub 2017 Apr 13.