

Corrigendum

Corrigendum to “Predictive Value of Plasma MicroRNA-216a/b in the Diagnosis of Esophageal Squamous Cell Carcinoma”

Shuling Dong,^{1,2} Huiqing Yin,^{1,2} Cuicui Dong,^{1,2} Kaiyan Sun,^{1,2} Pin Lv,^{1,2} Weiwei Meng,³ Liang Ming,^{1,2} and Fucheng He^{1,2}

¹Department of Medical Laboratory, The First Affiliated Hospital of Zhengzhou University, Zhengzhou, Henan 450052, China

²Key Medical Laboratory of Henan Province, Zhengzhou, Henan 450052, China

³Department of Medical Laboratory, The Second Affiliated Hospital of Zhengzhou University, Zhengzhou, Henan 450052, China

Correspondence should be addressed to Liang Ming; mingliang3072@163.com and Fucheng He; hefucheng@126.com

Received 29 June 2016; Accepted 12 October 2016; Published 12 March 2017

Copyright © 2017 Shuling Dong et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

In the article titled “Predictive Value of Plasma MicroRNA-216a/b in the Diagnosis of Esophageal Squamous Cell Carcinoma” [1], there was an error in Table 1. The heading incorrectly referred to miRNA-718 expression instead of miRNA-216a/b. miRNA-718 expression is the subject of

a related article, Sun L, Dong S, Dong C, Sun K, Meng W, Lv P, Yin H, Ming L, He F: “Predictive value of plasma miRNA-718 for esophageal squamous cell carcinoma”. *Cancer Biomark.* 2016;16(2):265-73. doi: 10.3233/CBM-150564. The table should be corrected as follows.

TABLE 1: Clinicopathological characteristics of patients with ESCC.

Characteristics	n (%)	Plasma miRNA relative expression level (mean ± SD)			
		miR-216a	P value	miR-216b	P value
Age (year)			0.1082		0.2125
≤63	46 (38.3)	0.058 ± 0.048		0.079 ± 0.086	
>63	74 (61.7)	0.074 ± 0.054		0.099 ± 0.087	
Gender			0.5388		0.5912
Male	79 (65.8)	0.070 ± 0.055		0.094 ± 0.087	
Female	41 (34.2)	0.064 ± 0.048		0.085 ± 0.089	
Smoking			0.1342		0.6168
Never	52 (43.3)	0.060 ± 0.047		0.087 ± 0.079	
Ever	68 (56.7)	0.074 ± 0.056		0.095 ± 0.093	
Alcohol use			0.2378		0.5386
Never	40 (33.3)	0.060 ± 0.054		0.098 ± 0.087	
Ever	80 (66.7)	0.072 ± 0.051		0.088 ± 0.087	

TABLE 1: Continued.

Characteristics	n (%)	Plasma miRNA relative expression level (mean $\pm$ SD)			
		miR-216a	P value	miR-216b	P value
Tumor location			0.3136		0.1810
Upper esophagus	11 (9.2)	0.065 $\pm$ 0.061		0.106 $\pm$ 0.092	
Middle esophagus	70 (58.3)	0.074 $\pm$ 0.056		0.100 $\pm$ 0.096	
Low esophagus	39 (32.5)	0.058 $\pm$ 0.041		0.072 $\pm$ 0.066	
Histologic grade			0.8959		0.8308
Well differentiated	33 (27.5)	0.070 $\pm$ 0.059		0.109 $\pm$ 0.112	
Moderate differentiated	70 (58.3)	0.068 $\pm$ 0.052		0.089 $\pm$ 0.082	
Poor differentiated	17 (14.2)	0.063 $\pm$ 0.040		0.068 $\pm$ 0.030	
T stage			0.3206		0.6600
Tis-T1	36 (30.0)	0.075 $\pm$ 0.061		0.115 $\pm$ 0.114	
T2	28 (23.3)	0.086 $\pm$ 0.065		0.100 $\pm$ 0.090	
T3	41 (34.2)	0.056 $\pm$ 0.037		0.074 $\pm$ 0.065	
T4	15 (12.5)	0.050 $\pm$ 0.019		0.067 $\pm$ 0.031	
Lymph node metastasis			0.0845		0.0168
Negative	42 (35.0)	0.088 $\pm$ 0.068		0.134 $\pm$ 0.117	
Positive	78 (65.0)	0.057 $\pm$ 0.038		0.068 $\pm$ 0.054	
TNM stage			0.0054		0.0359
0-I	29 (24.2)	0.088 $\pm$ 0.062		0.134 $\pm$ 0.119	
II	28 (23.3)	0.086 $\pm$ 0.065		0.101 $\pm$ 0.090	
III	40 (33.3)	0.058 $\pm$ 0.035		0.072 $\pm$ 0.067	
IV	23 (19.2)	0.039 $\pm$ 0.022		0.060 $\pm$ 0.029	

References

- [1] S. Dong, H. Yin, C. Dong et al., "Predictive value of plasma MicroRNA-216a/b in the diagnosis of esophageal squamous cell carcinoma," *Disease Markers*, vol. 2016, Article ID 1857067, 8 pages, 2016.