Clinical Rheumatology (2022) 41:1937–1938
https://doi.org/10.1007/s10067-022-06102-5

CORRECTION

Correction to: Coagulation in gout: is there a link with disease activity?

Daisy Vedder[1,2] · Martijn Gerritsen[1] · Joost C. M. Meijers[3,4] · Michael T. Nurmohamed[1,2,5]

Published online: 4 March 2022

Correction to: Clinical Rheumatology
https://doi.org/10.1007/s10067-022-06047-9

In the original version of the above article, The figures and legends used were incorrect. The correct figures are presented as follows (Figs.1, 2, 3, and 4):

The original article has been corrected.

The original article can be found online at https://doi.org/10.1007/s10067-022-06047-9.

✉ Daisy Vedder
d.vedder@reade.nl

[1] Amsterdam Rheumatology & Immunology Center, Reade, Amsterdam, Netherlands

[2] Amsterdam Cardiovascular Sciences, Vrije Universiteit, Amsterdam, Netherlands

[3] Department of Experimental Vascular Medicine, University of Amsterdam, Amsterdam UMC, Amsterdam, Netherlands

[4] Department of Molecular Hematology, Sanquin Research, Amsterdam, Netherlands

[5] Department of Rheumatology, Amsterdam UMC, Amsterdam, the Netherlands



Fig. 1 Disease activity according the Gout Activity Score is associated with Velocity index



Fig. 2 Disease activity according the Gout Activity Score is associated with Peak thrombin levels



Fig. 3 A decrease in CRP was associated with a decrease in Peak thrombin



Fig. 4 A decrease in ESR was associated with a decrease in D-dimer