



Published in final edited form as:

*Nat Commun.* ; 6: 6610. doi:10.1038/ncomms7610.

## Sumoylation controls the timing of Tup1-mediated transcriptional deactivation

Chong Han Ng<sup>1,2</sup>, Akhi Akhter<sup>3</sup>, Nathan Yurko<sup>1</sup>, Justin M. Burgener<sup>3</sup>, Emanuel Rosonina<sup>3,4</sup>, and James L. Manley<sup>1,4</sup>

<sup>1</sup>Department of Biological Sciences, Columbia University, New York, NY, 10027, USA

<sup>3</sup>Department of Biology, York University, Toronto, ON, M3J 1P3, Canada

### Abstract

The small ubiquitin-like modifier (SUMO) is implicated in various cellular activities, including transcriptional regulation. We previously showed that the yeast activator Gcn4 becomes sumoylated during activation, facilitating its eventual promoter eviction and transcriptional shut off. Here we show that the corepressor Tup1 is sumoylated, at two specific lysines, under various stress conditions. Mutation of these sites has no effect on Tup1 recruitment or RNAP II promoter occupancy immediately following induction. However, Tup1 levels subsequently decrease, while RNAP II and transcription increase in Tup1 mutant cells. Consistent with this, a Tup1 mutant displaying increased sumoylation led to reduced transcription. We also show that coordinated sumoylation of Gcn4 and Tup1 enhances Gcn4 promoter eviction, and that multiple Tup1-interacting proteins become sumoylated after stress. Together, our studies provide evidence that coordinated sumoylation of Gcn4, Tup1, and likely other factors, dampens activated transcription by stabilizing Tup1 binding and stimulating Gcn4 and RNAP II removal.

### INTRODUCTION

The SUMO polypeptide is present in all eukaryotes and is highly conserved from yeast to humans. SUMO covalently modifies many proteins that participate in diverse cellular processes, including transcriptional regulation, subcellular localization, DNA repair and signal transduction<sup>1-4</sup>. Many SUMO substrates are transcriptional activators, repressors, co-activators or co-repressors. Two lines of evidence have linked SUMO modifications with transcriptional repression<sup>1,5</sup>. First, in many cases interfering with sumoylation of transcriptional regulators at promoter regions leads to transcriptional activation. Second, sumoylated proteins can be recruited into repressive environments in higher eukaryotes,

Users may view, print, copy, and download text and data-mine the content in such documents, for the purposes of academic research, subject always to the full Conditions of use:[http://www.nature.com/authors/editorial\\_policies/license.html#terms](http://www.nature.com/authors/editorial_policies/license.html#terms)

<sup>4</sup>Co-corresponding authors. jlm2@columbia.edu OR rosonina@yorku.ca, PHONE: 212-854-4647, FAX: 212-865-8246.

<sup>2</sup>Current address: Faculty of Information Science & Technology, Multimedia University, Jalan Ayer Keroh Lama, 75450 Bukit Beruang, Melaka, Malaysia

### AUTHOR CONTRIBUTIONS

C.H.N., A.A., N.Y. and J.M.B. designed and carried out experiments. C.H.N., E.R., and J.L.M. analyzed data and wrote the paper.

### CONFLICT OF INTEREST

The authors declare no competing financial interests.

such as PML nuclear bodies. However, more recent studies have shown that SUMO modification of promoter-bound factors also occurs during the process of gene activation, suggesting a possible positive role in transcriptional control, in yeast <sup>6-8</sup> as well as mammalian <sup>9,10</sup> cells.

Modulation of SUMO levels at gene promoters is emerging as an important aspect of transcriptional activation. For example, activation of several inducible genes in yeast caused not only accumulation of SUMO at promoter regions, but also recruitment of Ubc9, the SUMO E2 conjugating enzyme, indicating that activation involves sumoylation of promoter-bound factors. However, Ubc9 inactivation, while reducing sumoylation at the induced promoters, paradoxically resulted in increased transcription <sup>6</sup> and the presence of Ulp1, a SUMO protease, is important for optimal gene activation <sup>8</sup>. Providing an explanation for these observations, the reduced sumoylation brought about by destabilizing Ubc9 impaired the cell's ability to shut off activated transcription appropriately, indicating that SUMO can facilitate transcriptional deactivation. Subsequently, the transcriptional activator Gcn4 was identified as one of the promoter-associated SUMO substrates, and Gcn4 sumoylation was shown to be required for its efficient removal from target promoters following RNA polymerase II (RNAP II) recruitment <sup>7</sup>. A similar result has also been reported for the mammalian activator AP-1 <sup>9</sup>.

Transcriptional activation is invariably dependent on multiple cofactors in addition to the activator. For example, for Gcn4 these include SAGA, SWI/SNF and RSC chromatin remodeling complexes, the SRB/MED complex, the transcriptional regulator CCR4-NOT, and the repressive Cyc8/Tup1 complex <sup>11-13</sup>. Some of the subunits of these complexes, including for example Gcn5, Snf2 and Tup1, have been identified as SUMO substrates in large-scale proteomics studies <sup>14-17</sup>. However, whether sumoylation of these proteins contributes to their function in gene control, and if so how, remains mostly elusive.

Tup1 has been suggested to function as both a corepressor and coactivator. The protein, which is conserved throughout eukaryotes <sup>18</sup>, was one of the first to be characterized as a transcriptional corepressor <sup>19,20</sup>, forming a complex with Cyc8 to mediate repression of diverse sets of genes under various stress conditions <sup>21</sup>. However, several studies have found that Tup1 remains associated with promoters of target genes after activation <sup>13,22,23</sup>. For example, Tup1 is bound to many glucose-repressed genes even after glucose repression is removed <sup>23</sup>. Tup1 has also been shown to play a role in recruitment of SAGA, SWI/SNF and Mediator to promoters <sup>13,24-26</sup>. Therefore, it has been proposed that Tup1 may switch from acting as a corepressor to a coactivator during transcriptional activation <sup>25,27,28</sup>. Consistent with this, Tup1 binds to the *ARG1* promoter after activation, and *ARG1* induction is reduced in a *tup1* null mutant strain <sup>13</sup>. However, there is still lack of direct evidence to support this notion, and several studies suggest that Tup1 may continue to act as a corepressor after gene activation. During galactose derepression, for example, the *GALI* gene is induced more quickly in a *tup1* null mutant <sup>29</sup>. Similarly, the *Drosophila* Tup1 ortholog Groucho was recently implicated in transcriptional attenuation of active genes <sup>30</sup>.

Here we show that Tup1 facilitates transcriptional deactivation in a manner enhanced by sumoylation. We first demonstrate that Tup1 is sumoylated at two specific lysine residues

under various stress conditions. By mutating these sites, we then show that Tup1 sumoylation does not affect its initial recruitment to promoters upon activation, but prolongs its association with the promoters, dampens transcription, and facilitates eventual removal of RNAP II and the Mediator component Gal11. Consistent with this, we identify a Tup1 mutant with enhanced sumoylation that results in reduced transcription. Although sumoylation of Tup1 and Gcn4 has opposite effects on their association with target gene promoters, modification of both proteins results in a net repressive effect on activated transcription. Interestingly, mutation of SUMO sites in both Tup1 and Gcn4 is found to prolong Gcn4 promoter occupancy, suggesting cooperation between sumoylated proteins. These results and others show that Tup1 sumoylation is part of a coordinated SUMO-mediated mechanism of controlling transcription levels and enabling gene deactivation.

## RESULTS

### Tup1 is sumoylated under various stress conditions

To extend our studies on the role of sumoylation during gene activation in yeast, we examined how SUMO affects the coactivator/corepressor Tup1. To show that Tup1 is a *bona fide* SUMO substrate, and to determine whether Tup1 sumoylation is affected by stress conditions, wild-type yeast expressing 6HA-tagged Tup1 (Tup1-HA) in its normal chromosomal location were treated with 100 J m<sup>-2</sup> UV, 1.0 mM H<sub>2</sub>O<sub>2</sub>, nitrogen starvation, carbon starvation, or amino acid starvation (induced by sulfometuron-methyl [SM], 0.5 μg ml<sup>-1</sup> <sup>31</sup>). Tup1-HA was immunoprecipitated (IPed) from cell extracts with an HA antibody and sumoylated forms of Tup1 were detected by western blot using an antibody that recognizes yeast SUMO (known as Smt3). We detected sumoylated Tup1-HA from yeast grown in normal, unstressed conditions, but higher levels were found in yeast grown in each of the tested stress conditions (Fig. 1a). These results confirm and extend previous global proteomic studies suggesting that Tup1 is a SUMO substrate <sup>14–17</sup>. *Drosophila* Groucho is also known to be sumoylated <sup>32,33</sup>, suggesting that Tup1 sumoylation is evolutionarily conserved.

To facilitate analysis of the function of Tup1 sumoylation, we next mapped its sumoylation site(s). Using the SUMOsp 2.0 sumoylation site prediction program <sup>34</sup>, three potential sumoylation sites, K229, K270 and K611, were predicted. SUMO western blot analysis of Tup1-HA IPs showed multiple bands, which may correspond to sumoylation at multiple sites (Fig. 1b, left panel). In agreement with this, we found that a single mutation, K270R (see also <sup>17</sup>), resulted in a major reduction, but not complete loss, of sumoylated Tup1, while another single mutation (K229R) caused a minor reduction after SM treatment. Combining both mutations (K229,270R) almost completely abolished Tup1 sumoylation, implying that K229 and K270 are the sumoylation sites in Tup1, in both untreated and SM treated cells (Fig. 1b, left panel). Consistent with this, a minor ~125 kDa form of Tup1-HA, detected when an HA western blot was overexposed, was not observed with the K229,270R mutant (Fig. 1b, right panel). Multiple higher molecular weight forms of Tup1, likely representing further modified isoforms, were also detected in the HA western blot (just below the 250 kDa marker; Fig. 1b, right panel), some of which also disappeared with the K229,270R mutations. This indicates that, in addition to sumoylation at K229 and K270, Tup1 is also

subject to additional modifications. Unexpectedly, mutation of the third predicted residue, K611, to Arg actually resulted in an increase of sumoylated Tup1 isoforms (Supplementary Fig. 1a). While the basis for this is unknown, one explanation is that K611 might normally be subject to other modification(s) that are inhibitory to sumoylation at other Lys residues. Alternatively, mutation of K611 may disrupt protein-protein interactions that naturally limit Tup1 sumoylation at K229 and K270. For example, K611 is located within a WD repeat domain that is capable of binding ubiquitin<sup>35</sup>.

The above data indicate that Tup1 is multi-sumoylated (modified by SUMO at multiple Lys residues). To determine whether some sumoylated isoforms were due to poly-sumoylation (SUMO chains forming at single modified Lys residues), we used a strain that expresses a form of Smt3 with Lys-to-Arg mutations at positions thought necessary for chain formation, *smt3-R11,15,19*, and another strain in which all Smt3 Lys were mutated to Arg, *smt3-allR*. These mutants are still capable of conjugating a single SUMO moiety on substrates, but the ability to form polySUMO chains is compromised (*smt3-R11,15,19*) or abolished (*smt3-allR*)<sup>36</sup>. In Western blot analysis of Tup1-HA IPs, the signal of sumoylated Tup1 was reduced slightly in the *smt3-R11,15,19* mutant, while only a single weak band was detected in the *smt3-allR* mutant (Supplementary Fig. 1b). This result suggests that a significant level of sumoylation associated with Tup1 derives from polySUMO chains.

We next set out to investigate phenotypes of mutant yeast strains deficient in Tup1 sumoylation. Since both K229 and K270 sites fall within the central domain which consists of the gene repression, H3/H4 binding and histone deacetylase Hda1 interaction domains<sup>37</sup>, we asked whether sumoylation of Tup1 affects its transcriptional repressive functions (Fig. 1c). Notably, Groucho sumoylation sites also fall within the central region required for HDAC interaction, and sumoylation enhances Groucho repressive activity likely through increased interaction with an HDAC<sup>33</sup>. While a *tup1* null mutant showed a mild growth defect in SM-containing media<sup>13</sup>, the Tup1-MT (K229,270R) mutant did not show any significant defect when grown on media containing SM (Fig. 1d). This result is consistent with the lack of a growth defect of a *gcn4* SUMO mutant grown in SM medium<sup>7</sup>. However, analysis of Tup1-WT and Tup1-MT (K229,270R) growth in liquid media revealed a growth delay for Tup1-MT (K229,270R) during exponential phase, although both strains reached the same density during stationary phase (Fig. 1e). Taken together, these results suggest that Tup1 sumoylation does not have a general, essential function in Tup1-mediated repression, or a direct role in activation of genes induced by amino acid starvation.

### Tup1 sumoylation enhances repression after gene activation

Our previous analyses demonstrated that Smt3 and Ubc9 proteins accumulate at multiple inducible gene promoters during gene activation<sup>6</sup> and that Gcn4 occupancy at target genes can be regulated by sumoylation<sup>7</sup>. To determine whether SUMO-deficient Tup1 also displays altered association with promoters of Gcn4 target genes, we conducted time course ChIP studies of Tup1-HA occupancy in Tup1-WT and Tup1-MT (K229,270R) strains during gene activation induced by SM. Figure 2a shows that both Tup1-WT and Tup1-MT (K229,270R) exhibited a similar initial increase in association with the *ARG1* and *CPA2* promoters after SM treatment. A similar Tup1-WT binding profile on the *ARG1* promoter

has been previously observed<sup>13</sup>. However, at 20 and 40 min after SM treatment, Tup1-MT (K229,270R) occupancy was significantly lower than Tup1-WT at the *ARG1* promoter, while a lower level of Tup1-MT (K229,270R) was detected at the *CPA2* promoter at 40 min. A time-course analysis of Tup1-WT sumoylation levels showed that Tup1 sumoylation peaked at 40 min after SM treatment, indicating that the greatest differential in Tup1-WT versus Tup1-MT (K229,270R) promoter occupancy was observed when Tup1 is normally most highly sumoylated (Supplementary Fig. 1c). To determine whether Tup1 associates specifically with promoter regions, we performed ChIP of Tup1-HA on the *ARG1* promoter, coding and downstream untranscribed regions (Supplementary Fig. 2a). We detected both Tup1-WT and Tup1-MT (K229,270R) mostly on the promoter region, and only after SM treatment, suggesting that Tup1 indeed associates with inducible gene promoter regions.

To extend these results, we examined Tup1-WT and Tup1-MT (K229,270R) promoter occupancy on two additional Tup1 target genes, *HIS4* and *SUC2*. The *HIS4* gene is also responsive to SM treatment<sup>31</sup>, and time-course ChIP (Supplementary Fig. 2b) revealed profiles for both Tup1 derivatives very similar to those observed with the *ARG1* and *CPA2* promoters, i.e., reduced Tup1-MT (K229,270R) occupancy at later times. In contrast, the *SUC2* promoter is responsive to low glucose starvation, but not to SM treatment, and no differences in Tup1-WT and Tup1-MT (K229,270R) occupancy were detected (Supplementary Fig. 2c). These results suggest that the effect of Tup1 sumoylation on Tup1 target genes is not universal and perhaps condition-dependent.

Reduced Tup1-MT (K229,270R) on SM-inducible target promoters was not due to lower expression levels of the mutant protein. On the contrary, in many cases, especially when strains were grown in nutrient-poor media, such as in the presence of proline as the main source of nitrogen, we detected significantly higher levels of Tup1-MT (K229,270R) than Tup1-WT (Supplementary Fig. 3a), suggesting that Tup1 sumoylation is involved in promoting Tup1 degradation. Similarly, sumoylation of *Drosophila* Groucho also promotes its degradation, via a ubiquitin-mediated pathway<sup>32</sup>. However, deletion of genes encoding SUMO-targeted ubiquitin ligases Slx5 or Rad18 did not result in increased Tup1-HA levels in cells grown under standard conditions, suggesting that sumoylation of Tup1 does not normally trigger its degradation through a ubiquitin pathway (Supplementary Fig 3b). Together, our results indicate that Tup1 is recruited to target gene promoters during activation independent of sumoylation, but preventing its sumoylation can, in a gene- and/or condition-dependent manner, result in lower occupancy at later time points.

In light of the above results, we next examined whether Tup1 sumoylation affects recruitment of RNAP II to the *ARG1* and *CPA2* promoters, again by time-course ChIP during SM induction. As with Tup1 itself, we did not observe any difference in RNAP II levels between Tup1-WT and Tup1-MT (K229,270R) strains at the beginning of the time course. However, increased levels of RNAP II were detected at both promoters in the Tup1-MT (K229,270R) strain at the 40-min time point (Fig. 2b). This is consistent with the reduced levels of promoter-associated Tup1-MT (K229,270R) allowing increased RNAP II recruitment. Conditional removal of Tup1 was previously shown to cause decreased histone H3 occupancy at several promoters<sup>26</sup>, and it was thus conceivable that defective Tup1 sumoylation might have a similar effect. We therefore compared histone H3 levels at the

above promoters in Tup1-WT and Tup1-MT (K229,270R) strains during induction. In both strains, H3 levels dropped during the time course, reflecting gene activation, but we detected no difference in H3 occupancy between the two strains (Fig. 2c).

Together, these results indicate that sumoylation regulates Tup1 occupancy on target gene promoters, as we observed previously with Gcn4<sup>7</sup>. However, whereas blocking Gcn4 sumoylation resulted in increased Gcn4 promoter occupancy, blocking Tup1 sumoylation had the opposite effect, causing reduced Tup1 persistence as well as increased RNAP II recruitment, which was not due to secondary effects on histone H3 levels.

To determine whether the reduced binding of SUMO-deficient Tup1 affects target gene expression, we performed RT-qPCR analysis of *ARG1* and *CPA2* transcripts over a time course after SM treatment with the Tup1-WT and Tup1-MT (K229,270R) strains (Fig. 2d). The mRNA levels of *ARG1* and *CPA2* peaked at the 40-min time point and decreased gradually at the 60 and 120 min time points. Similar to the ChIP analysis, we did not detect any significant initial difference in *ARG1* and *CPA2* mRNA levels between WT and MT (K229,270R) (first 10 min after SM treatment). However, both *ARG1* and *CPA2* mRNA levels were significantly higher in the Tup1-MT (K229,270R) strain at the 20, 40 and 60 min time points, correlating well with our observation that less Tup1-MT (K229,270R), and more RNAP II, was present at these promoters at later points during the time course. Strengthening these findings, we also found that a strain expressing the hyper-sumoylated Tup1 mutant (K611R) generated reduced *ARG1* mRNA levels compared to a WT strain (Supplementary Fig. 4). Taken together, our findings demonstrate that Tup1 sumoylation can affect target gene transcription in a negative manner, indicating a role in modulating, or dampening, activated transcription.

### Sumoylation of Tup1 and Gcn4 modulates Gcn4 promoter binding

Recent studies suggest that sumoylation can coordinately regulate multiple components of protein complexes, as opposed to regulating individual proteins independently, to achieve a biological effect. Consequently, stronger effects can be detected when sumoylation of several SUMO targets in the same complex is blocked<sup>38,39</sup>. In light of this, we examined the effect of combining SUMO site mutations in both Tup1 and Gcn4. Strains expressing either chromosomally tagged Tup1-WT or Tup1-K270R were generated and transformed with either Gcn4-WT or Gcn4 SUMO-deficient mutant (K50,58R) expression plasmids. Time-course ChIP analyses of Gcn4-Flag showed that the Tup1 SUMO mutation on its own slightly enhanced Gcn4-WT occupancy at the *ARG1* promoter following SM treatment (Fig. 3a). As we previously observed<sup>7</sup>, mutation of Gcn4 SUMO sites also resulted in increased Gcn4 occupancy on the *ARG1* promoter. Interestingly though, simultaneous mutation of SUMO sites on Tup1 and Gcn4 resulted in a considerably greater increase in Gcn4 promoter occupancy after SM treatment (Fig. 3a). Although individual SUMO site mutations on both Tup1 and Gcn4 resulted in increased *ARG1* transcript levels, combining the mutations did not result in a further increase (Fig. 3b), perhaps because maximal activation rates had already been achieved. These results nonetheless indicate that Tup1 sumoylation can affect promoter levels of Gcn4, and suggest that coordinated sumoylation of Tup1 and Gcn4, and

possibly other promoter-bound factors (see below), ensures efficient clearance of Gcn4 from target genes.

### Sumoylated Tup1 impairs recruitment of Gal11 but not Gcn5

We next wished to determine whether the repressive effect of Tup1 sumoylation involves additional proteins that function in *ARG1* activation. In addition to Tup1 and Gcn4, Gal11 and Gcn5 have been identified as SUMO targets<sup>14,16,40</sup>. Gcn4 interacts with Gal11 directly, and this interaction is important for efficient Gal11 recruitment to the *ARG1* promoter<sup>41</sup>. Moreover, depletion of Tup1 caused a significant increase in occupancy of both Gal11 and Gcn5 at several promoters<sup>26</sup>. We therefore investigated whether the repressive effect of sumoylated Tup1 reflects effects on binding of Gal11 and/or Gcn5. Strains expressing different combinations of chromosomally HA-tagged Tup1-WT or Tup1-K270R and Flag-tagged Gcn5-WT or Gal11-WT were generated. ChIP analysis confirmed that induction of *ARG1* by SM coincided with recruitment of both Gal11 and Gcn5 to the *ARG1* promoter (Fig. 3c). Whereas Gcn5-Flag occupancy was unaffected by the Tup1 SUMO site mutations, sharply elevated levels of Gal11-Flag were detected at the *ARG1* promoter in the Tup1-K270R strain. The magnitude of the effect (~3-fold) was significantly higher than the effect of Tup1 mutation on its own occupancy (~30% reduction in Tup1 occupancy; compare Fig. 3c with 2a), indicating that increased Gal11 levels were not simply a consequence of reduced Tup1 on the promoter. This result is consistent with our findings that impaired Tup1 sumoylation resulted in higher levels of RNAP II at the *ARG1* promoter. Taken together, our data point to a coordinated effect of Tup1 sumoylation on the *ARG1* promoter that enhances deactivation of activated transcription.

### Tup1-interacting factors are sumoylated after SM treatment

We next wished to investigate further the possibility that the effects of Tup1 sumoylation on promoter deactivation involves a coordinated network of sumoylated proteins, i.e, a consequence of “protein-group sumoylation”<sup>39</sup>. To this end, we examined whether other proteins known to interact physically or genetically with Tup1 are also SUMO targets, and whether their sumoylation is induced by SM treatment. These include histone H3<sup>42</sup>, Cti6, which relieves Tup1 transcriptional repression<sup>25</sup>, Gcn5<sup>43</sup>, Gal11<sup>24,26</sup> and Cyc8<sup>20,44,45</sup>. We generated strains expressing HA-tagged versions of each of these proteins from their chromosomal loci, and subjected them to SM treatment for 20 min. The HA-tagged proteins were IPed from whole-cell extracts and analyzed by SUMO western blotting. In every case, we detected sumoylation of the tested protein in untreated cells, but SM treatment resulted in significantly higher levels of sumoylation (Fig. 4a). This indicates that Tup1-associated proteins are coordinately sumoylated upon induction of target genes. This did not reflect a global increase in sumoylation induced by SM, as analysis of whole-cell lysates, prepared from three different yeast strains, by SUMO western blot revealed at most very slight increases in total sumoylation after SM treatment (Supplementary Fig. 5a), suggesting that the increase in sumoylation of Tup1-associated proteins was part of a specific response during gene activation by SM. Supporting this conclusion, we observed no increase in sumoylation of Tfg1, a subunit of transcription factor TFIIF, after SM treatment (Supplementary Fig. 5b), even though Tfg1 is a known SUMO target<sup>14–16</sup>

Gcn5 and histone H3 have previously been identified as SUMO substrates, and both sumoylated forms play negative roles in transcription regulation<sup>40,46</sup>. More recently, desumoylation of Cyc8 has been implicated in derepression of *GAL1* transcription<sup>8</sup>. In studies mentioned above, DNA damage was shown to induce sumoylation of multiple proteins involved in the DNA repair pathway, creating a synergistic response to DNA damage and promoting protein-protein interactions<sup>38,39</sup>. Likewise, we suggest that simultaneous sumoylation of Tup1, Gcn4 and associated proteins during gene activation functions to ensure subsequent transcriptional deactivation.

We next wished to investigate whether Tup1 sumoylation affects interactions with any of its network of associated factors. To this end, we examined whether sumoylation of Tup1 affects its association with histone H3. Reciprocal co-IP experiments with strains expressing Flag-tagged Hht1 (histone H3) and either HA-tagged Tup1-WT or Tup1-MT (K270R) were performed. Tup1-HA was IPed from whole-cell extracts with an HA antibody, and sumoylated forms of Tup1 and co-IPed Flag-Hht1 were detected by western blot using an Smt3 antibody and an Flag antibody, respectively (Fig. 4b). We detected increased association of Flag-Hht1 with Tup1-WT after SM treatment. However, association was strongly reduced in the Tup1-MT strain. Similarly, in the reciprocal Flag IP more Tup1-WT than Tup1-MT co-IPed with Flag-Hht1 after SM treatment (Fig. 4c). These results indicate that Tup1 sumoylation enhances its interaction with H3, likely contributing to Tup1-mediated gene deactivation. Interestingly, an Smt3 western blot of the Flag-Hht1 IP revealed fewer sumoylated species associated with Hht1 when Tup1 sumoylation was impaired (Fig. 4c). This supports the idea that Tup1 sumoylation is important for coordinating SUMO modifications of additional components of Tup1/Hht1-containing complexes.

## DISCUSSION

Transcription initiation by RNAP II is an immensely complex process involving numerous multi-subunit protein factors. While the molecular mechanisms of inducible gene activation following stimulation are well-characterized, it has been much less clear how transcription is returned to basal levels, or how genes are shut off, following activation. Our previous analyses demonstrated that sumoylation of Gcn4 is required for its efficient removal from promoters following transcription activation<sup>7</sup>, and similar findings with the mammalian activator AP-1 suggest this mechanism is evolutionarily conserved<sup>9</sup>. However, removal or inactivation of an activator can be insufficient to stop transcription entirely or to return a gene to an inactive state, and indeed Tup1 becomes ineffective under such conditions<sup>47</sup>. Thus, while our data has established important roles for SUMO-mediated clearance of Gcn4 and stabilization of Tup1 binding in promoter deactivation, additional mechanisms must be at play to insure efficient deactivation of induced genes when their products are no longer needed.

Our results showing that multiple Tup1-associated proteins display enhanced sumoylation upon induction point to a role for a “protein-group” sumoylation mechanism, as has been observed for several other nuclear processes involving complex protein assemblies<sup>39</sup>. As we have previously observed promoter-associated sumoylation during induction of other types of genes, such as *GAL1* and *STL1*<sup>6</sup>, we suggest that sumoylation of distinct protein

complexes found on different activated promoters is a general mechanism of regulating transcription. In any event, our study has demonstrated that sumoylation plays an important role in fine-tuning gene expression during activation and subsequent deactivation, and that SUMO-dependent modulation of Tup1 repression activity plays an important role in this process.

## MATERIALS AND METHODS

### Yeast strains and plasmids

All yeast strains used are listed in Supplementary Table 1. The chromosomally 6HA- and 3HA-tagged yeast strains (CHNM1 to 8 and YAA set) were generated by homologous recombination under the selection of minimal media without tryptophan<sup>48</sup>. The presence of the tagged alleles was verified by colony PCR and Western blot analysis using anti-HA antibodies. The pTUP1-6HA plasmid was generated by first amplifying *TUP1* genomic DNA including 1 kb upstream and downstream of the *TUP1* ORF and subsequently cloning the PCR product using the *URA3*-marked CEN vector pRS316. For generation of integrative mutagenic alleles, a site-directed mutagenesis PCR-based homologous recombination approach was employed<sup>49</sup>, while for generation of the mutagenic plasmid, a highly efficient site-directed, ligase-independent mutagenesis (SLIM) approach was used<sup>50</sup>. The details of the cloning procedure are available upon request. Primer sequences are listed in Supplementary Table 2.

### Western blot and immunoprecipitation

Yeast cultures were grown in minimal medium lacking isoleucine and valine to OD<sub>600</sub> of 0.6–0.7. Induction of Tup1 target genes, *ARG1* and *CPA2* by amino acid starvation was achieved by treatment with sulfometuron methyl (SM) from Sigma at a final concentration of 0.5  $\mu\text{g ml}^{-1}$  for 20 min (unless otherwise specified). Cells were lysed by agitation with glass beads in IP buffer (150 mM NaCl, 50 mM Tris pH 8.0, 0.1% NP-40, 10% glycerol) plus 2.5  $\text{mg ml}^{-1}$  N-ethylmaleimide. IPs and washes were performed in the same buffer, and 1  $\mu\text{g}$  of the appropriate antibody was added for overnight IP with Protein-G sepharose beads. Proteasomal inhibition<sup>51</sup> was performed by addition of 20  $\mu\text{M}$  MG132 or mock-treatment (DMSO) for 1 hr before harvesting for western blot analysis.

### Chromatin Immunoprecipitation

Exponentially growing yeast were treated (as indicated in Fig. 1a), or mock-treated, then cross-linked with 1% formaldehyde for 20 min, before quenching with 0.36 M glycine. Samples were pelleted by centrifugation and washed once with ice-cold TBS (20 mM Tris-HCl, pH 7.5, 150 mM NaCl), then in ChIP buffer (0.05 M HEPES-KOH, pH 7.5, 150 mM NaCl, 1 mM EDTA, 1% Triton X-100, 0.1% sodium deoxycholate, 0.1% SDS). Pelleted samples were re-suspended in 600  $\mu\text{L}$  of ChIP buffer and 400  $\mu\text{L}$  of glass beads were added before they were subjected to three rounds of 1-min bead-beating, with 1 min breaks in between. Lysates were recovered and sonicated to shear chromatin into fragments with an average length of approximately 500 bp. Samples were then micro-centrifuged, and NaCl was added to the supernatants to adjust the final concentration to 275 mM. Immunoprecipitations were then performed overnight, with 1  $\mu\text{g}$  of appropriate antibody and

Protein G-sepharose beads added to the samples. Beads were then washed first in ChIP buffer with 275 mM NaCl, then ChIP buffer with 500 mM NaCl, then in an additional wash buffer (10 mM Tris-HCl, pH 8, 0.25 M LiCl, 1 mM EDTA, 0.5% NP-40, 0.5% sodium deoxycholate), and finally in TE buffer (10 mM Tris-HCl, pH 8, 1 mM EDTA).

Immunoprecipitated chromatin was eluted by incubation of beads in ChIP Elution Buffer (50 mM Tris-HCl, pH 7.5, 10 mM EDTA, 1% SDS) at 65°C for ten minutes. Samples were micro-centrifuged and the supernatant was treated with Proteinase K for 1 hr at 42°C, before cross-links were reversed by heating to 65°C for at least 4 hr. DNA was recovered by phenol-chloroform extraction and ethanol precipitation, then analyzed by PCR. Primer sequences are listed in Supplementary Table 2. For quantification, a two-tailed Student's *t*-test was applied with *P*-values indicated in parentheses above paired bars within graphs for statistical data analysis.

### Antibodies

Antibodies used for western blot, immunoprecipitation and ChIP assay were anti-Flag M2 (Sigma, cat. F1804, WB – 1:1000), rabbit anti-HA (abm, cat. G166, WB – 1:1000), mouse anti-HA (Covance, cat. MMS101P, WB – 1:1000), rabbit anti-H3 (abm, cat. ab1791, WB – 1:2000), anti-GADPH (Sigma, cat. G9545, WB - 1:2000), rabbit anti-Smt3 (Santa Cruz, cat. sc-28649, WB – 1:1000) and rabbit anti-Rpb1 (Santa Cruz, cat. sc-25758). The uncropped scans of the Figure 1a, 1b, 4a, 4b and 4c are included in Supplementary Fig. 6. Quantitative real time PCR was performed using gene-specific primers and the primer sequences are listed in Supplementary Table 2.

### Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

### Acknowledgments

We thank Elizabeth Miller, Silvere Pagant and Kyle Stahmer for sharing yeast strains, and members of the labs of J.L.M. and Songtao Jia for technical help. This work was supported by Fulbright Malaysian Visiting Scholar Program 2011–2012 to C.H.N, an NSERC Discovery Grant to E.R., and grant GM097174 from the National Institutes of Health to J.L.M.


### References

1. Cubenas-Potts C, Matunis MJ. SUMO: a multifaceted modifier of chromatin structure and function. *Dev Cell*. 2013; 24:1–12. [PubMed: 23328396]
2. Flotho A, Melchior F. Sumoylation: a regulatory protein modification in health and disease. *Ann Rev Biochem*. 2013; 82:357–385. [PubMed: 23746258]
3. Gill G. Something about SUMO inhibits transcription. *Curr Opin Genet Dev*. 2005; 15:536–541. [PubMed: 16095902]
4. Johnson ES. Protein modification by SUMO. *Ann Rev Biochem*. 2004; 73:355–382. [PubMed: 15189146]
5. Garcia-Dominguez M, Reyes JC. SUMO association with repressor complexes, emerging routes for transcriptional control. *Biochim Biophys Acta*. 2009; 1789:451–459. [PubMed: 19616654]
6. Rosonina E, Duncan SM, Manley JL. SUMO functions in constitutive transcription and during activation of inducible genes in yeast. *Gene Dev*. 2010; 24:1242–1252. [PubMed: 20504900]

7. Rosonina E, Duncan SM, Manley JL. Sumoylation of transcription factor Gcn4 facilitates its Srb10-mediated clearance from promoters in yeast. *Gene Dev.* 2012; 26:350–355. [PubMed: 22345516]
8. Texari L, Dieppois G, Vinciguerra P, Contreras MP, Groner A, Letourneau A, Stutz F. The nuclear pore regulates *GALI* gene transcription by controlling the localization of the SUMO protease Ulp1. *Mol Cell.* 2013; 51:807–818. [PubMed: 24074957]
9. Tempe D, Vives E, Brockly F, Brooks H, De Rossi S, Piechaczyk M, Bossis G. SUMOylation of the inducible (c-Fos:c-Jun)/AP-1 transcription complex occurs on target promoters to limit transcriptional activation. *Oncogene.* 2013; 33:921–927. [PubMed: 23396363]
10. Liu HW, Zhang J, Heine GF, Arora M, Gulcin Ozer H, Onti-Srinivasan R, Huang K, Parvin JD. Chromatin modification by SUMO-1 stimulates the promoters of translation machinery genes. *Nucleic Acids Res.* 2012; 40:10172–10186. [PubMed: 22941651]
11. Swanson MJ, Qiu H, Sumibcay L, Krueger A, Kim SJ, Natarajan K, Yoon S, Hinnebusch AG. A multiplicity of coactivators is required by Gcn4p at individual promoters *in vivo*. *Mol Cell Biol.* 2003; 23:2800–2820. [PubMed: 12665580]
12. Govind CK, Yoon S, Qiu H, Govind S, Hinnebusch AG. Simultaneous recruitment of coactivators by Gcn4p stimulates multiple steps of transcription *in vivo*. *Mol Cell Biol.* 2005; 25:5626–5638. [PubMed: 15964818]
13. Kim SJ, Swanson MJ, Qiu H, Govind CK, Hinnebusch AG. Activator Gcn4p and Cyc8p/Tup1p are interdependent for promoter occupancy at *ARG1 in vivo*. *Mol Cell Biol.* 2005; 25:11171–11183. [PubMed: 16314536]
14. Wohlschlegel JA, Johnson ES, Reed SI, Yates JR 3rd. Global analysis of protein sumoylation in *Saccharomyces cerevisiae*. *J Biol Chem.* 2004; 279
15. Zhou W, Ryan JJ, Zhou H. Global analyses of sumoylated proteins in *Saccharomyces cerevisiae*. Induction of protein sumoylation by cellular stresses. *J Biol Chem.* 2004; 279:32262–32268. [PubMed: 15166219]
16. Denison C, Rudner AD, Gerber SA, Bakalarski CE, Moazed D, Gygi SP. A proteomic strategy for gaining insights into protein sumoylation in yeast. *Mol Cell Proteomics.* 2005; 4:246–254. [PubMed: 15542864]
17. Wykoff DD, O’Shea EK. Identification of sumoylated proteins by systematic immunoprecipitation of the budding yeast proteome. *Mol Cell Proteomics.* 2005; 4:73–83. [PubMed: 15596868]
18. Courey AJ, Jia S. Transcriptional repression: the long and the short of it. *Gene Dev.* 2001; 15:2786–2796. [PubMed: 11691830]
19. Keleher CA, Redd MJ, Schultz J, Carlson M, Johnson AD. Ssn6-Tup1 is a general repressor of transcription in yeast. *Cell.* 1992; 68:709–719. [PubMed: 1739976]
20. Tzamarias D, Struhl K. Functional dissection of the yeast Cyc8-Tup1 transcriptional co-repressor complex. *Nature.* 1994; 369:758–761. [PubMed: 8008070]
21. Smith RL, Johnson AD. Turning genes off by Ssn6-Tup1: a conserved system of transcriptional repression in eukaryotes. *Trends Biochem Sci.* 2000; 25:325–330. [PubMed: 10871883]
22. Profit M, Struhl K. Hog1 kinase converts the Sko1-Cyc8-Tup1 repressor complex into an activator that recruits SAGA and SWI/SNF in response to osmotic stress. *Mol Cell.* 2002; 9:1307–1317. [PubMed: 12086627]
23. Buck MJ, Lieb JD. A chromatin-mediated mechanism for specification of conditional transcription factor targets. *Nat Genet.* 2006; 38:1446–1451. [PubMed: 17099712]
24. Han SJ, Lee JS, Kang JS, Kim YJ. Med9/Cse2 and Gal11 modules are required for transcriptional repression of distinct group of genes. *J Biol Chem.* 2001; 276:37020–37026. [PubMed: 11470794]
25. Papamichos-Chronakis M, Petrakis T, Ktistaki E, Topalidou I, Tzamarias D. Cti6, a PHD domain protein, bridges the Cyc8-Tup1 corepressor and the SAGA coactivator to overcome repression at *GALI*. *Mol Cell.* 2002; 9:1297–1305. [PubMed: 12086626]
26. Wong KH, Struhl K. The Cyc8-Tup1 complex inhibits transcription primarily by masking the activation domain of the recruiting protein. *Gene Dev.* 2011; 25:2525–2539. [PubMed: 22156212]
27. Conlan RS, Gounalaki N, Hatzis P, Tzamarias D. The Tup1-Cyc8 protein complex can shift from a transcriptional co-repressor to a transcriptional co-activator. *J Biol Chem.* 1999; 274:205–210. [PubMed: 9867831]


28. Han BK, Emr SD. Phosphoinositide [PI(3,5)P<sub>2</sub>] lipid-dependent regulation of the general transcriptional regulator Tup1. *Gene Dev.* 2011; 25:984–995. [PubMed: 21536737]
29. Gligoris T, Thireos G, Tzamarias D. The Tup1 corepressor directs Htz1 deposition at a specific promoter nucleosome marking the *GAL1* gene for rapid activation. *Mol Cell Biol.* 2007; 27:4198–4205. [PubMed: 17387147]
30. Kaul A, Schuster E, Jennings BH. The Groucho co-repressor is primarily recruited to local target sites in active chromatin to attenuate transcription. *Plos Genet.* 2014; 10:e1004595. [PubMed: 25165826]
31. Jia MH, Larossa RA, Lee JM, Rafalski A, Derosé E, Gonye G, Xue Z. Global expression profiling of yeast treated with an inhibitor of amino acid biosynthesis sulfometuron methyl. *Physiol Genomics.* 2000; 3:83–92. [PubMed: 11015603]
32. Abed M, Barry KC, Kenyagin D, Koltun B, Phippen TM, Delrow JJ, Parkhurst SM, Orian A. Degringolade, a SUMO-targeted ubiquitin ligase, inhibits Hairy/Groucho-mediated repression. *EMBO J.* 2011; 30:1289–1301. [PubMed: 21343912]
33. Ahn JW, Lee YA, Ahn JH, Choi CY. Covalent conjugation of Groucho with SUMO-1 modulates its corepressor activity. *Biochem Biophys Res Commun.* 2009; 379:160–165. [PubMed: 19101520]
34. Ren J, Gao X, Jin C, Zhu M, Wang X, Shaw A, Wen L, Yao X, Xue Y. Systematic study of protein sumoylation: Development of a site-specific predictor of SUMOsp 2.0. *Proteomics.* 2009; 9:3409–3412. [PubMed: 19504496]
35. Pashkova N, Gakhar L, Winistorfer SC, Yu L, Ramaswamy S, Piper RC. WD40 repeat propellers define a ubiquitin-binding domain that regulates turnover of F box proteins. *Mol Cell.* 2010; 40:433–443. [PubMed: 21070969]
36. Bylebyl GR, Belichenko I, Johnson ES. The SUMO isopeptidase Ulp2 prevents accumulation of SUMO chains in yeast. *J Biol Chem.* 2003; 278:44113–44120. [PubMed: 12941945]
37. Malave TM, Dent SY. Transcriptional repression by Tup1-Ssn6. *Biochem Cell Biol.* 2006; 84:437–443. [PubMed: 16936817]
38. Cremona CA, Sarangi P, Yang Y, Hang LE, Rahman S, Zhao X. Extensive DNA damage-induced sumoylation contributes to replication and repair and acts in addition to the *mec1* checkpoint. *Mol Cell.* 2012; 45:422–432. [PubMed: 22285753]
39. Psakhye I, Jentsch S. Protein group modification and synergy in the SUMO pathway as exemplified in DNA repair. *Cell.* 2012; 151:807–820. [PubMed: 23122649]
40. Sterner DE, Nathan D, Reindle A, Johnson ES, Berger SL. Sumoylation of the yeast Gcn5 protein. *Biochemistry.* 2006; 45:1035–1042. [PubMed: 16411780]
41. Jedidi I, Zhang F, Qiu H, Stahl SJ, Palmer I, Kaufman JD, Nadaud PS, Mukherjee S, Wingfield PT, Jaroniec CP, Hinnebusch AG. Activator Gcn4 employs multiple segments of Med15/Gal11 including the KIX domain to recruit mediator to target genes *in vivo*. *J Biol Chem.* 2010; 285:2438–2455. [PubMed: 19940160]
42. Edmondson DG, Smith MM, Roth SY. Repression domain of the yeast global repressor Tup1 interacts directly with histones H3 and H4. *Gene Dev.* 1996; 10:1247–1259. [PubMed: 8675011]
43. Islam A, Turner EL, Menzel J, Malo ME, Harkness TA. Antagonistic Gcn5-Hda1 interactions revealed by mutations to the Anaphase Promoting Complex in yeast. *Cell Div.* 2011; 6:13. [PubMed: 21651791]
44. Varanasi US, Klis M, Mikesell PB, Trumbly RJ. The Cyc8 (Ssn6)-Tup1 corepressor complex is composed of one Cyc8 and four Tup1 subunits. *Mol Cell Biol.* 1996; 16:6707–6714. [PubMed: 8943325]
45. Redd MJ, Arnaud MB, Johnson AD. A complex composed of tup1 and ssn6 represses transcription *in vitro*. *J Biol Chem.* 1997; 272:11193–11197. [PubMed: 9111019]
46. Nathan D, Ingvarsdottir K, Sterner DE, Bylebyl GR, Dokmanovic M, Dorsey JA, Whelan KA, Krsmanovic M, Lane WS, Meluh PB, Johnson ES, Berger SL. Histone sumoylation is a negative regulator in *Saccharomyces cerevisiae* and shows dynamic interplay with positive-acting histone modifications. *Gene Dev.* 2006; 20:966–976. [PubMed: 16598039]

47. Biggar SR, Crabtree GR. Chemically regulated transcription factors reveal the persistence of repressor-resistant transcription after disrupting activator function. *J Biol Chem.* 2000; 275:25381–25390. [PubMed: 10801867]
48. Knop M, Siegers K, Pereira G, Zachariae W, Winsor B, Nasmyth K, Schiebel E. Epitope tagging of yeast genes using a PCR-based strategy: more tags and improved practical routines. *Yeast.* 1999; 15:963–972. [PubMed: 10407276]
49. Toulmay A, Schneiter R. A two-step method for the introduction of single or multiple defined point mutations into the genome of *Saccharomyces cerevisiae*. *Yeast.* 2006; 23:825–831. [PubMed: 16921548]
50. Chiu J, Tillett D, Dawes IW, March PE. Site-directed, Ligase-Independent Mutagenesis (SLIM) for highly efficient mutagenesis of plasmids greater than 8kb. *J Microbiol Meth.* 2008; 73:195–198.
51. Liu C, Apodaca J, Davis LE, Rao H. Proteasome inhibition in wild-type yeast *Saccharomyces cerevisiae* cells. *Bio Techniques.* 2007; 42:158, 160–162.




**Figure 1. Tup1 is sumoylated primarily at K270 with a minor site at K229**

(a) HA IPs of untreated cells (control, SC medium), or cells treated with 100 J m<sup>-2</sup> UV, H<sub>2</sub>O<sub>2</sub> (1.0 mM for 20 mins), nitrogen starvation (-N for 60 mins), sulfometuron methyl (SM) (0.5 µg ml<sup>-1</sup> for 20 mins), or carbon starvation (-C for 60 mins), were analyzed by HA and yeast SUMO (Smt3) immunoblots. (b) Left panel: Analysis of HA IPs obtained from untreated control and SM-induced cells expressing either Tup1-6HA WT or mutants plasmids. Right panel: Analysis of overexposed HA immunoblot and Smt3 immunoblot of induced cells expressing Tup1-HA WT and mutants plasmids after HA IP. (c) Diagram of Tup1 protein domains with probable sumoylation sites indicated. (d) A *tup1Δ* strain was transformed with plasmids expressing either vector, wt Tup1-HA or mutant Tup1-K229,270R. Transformed strains were serially diluted and spotted on medium lacking Val and Ile (left), or the same medium supplemented with 0.5 µg/ml SM. (e) Growth curve analysis of Tup1-WT and Tup1-MT (K229,270R). A *tup1Δ* strain was transformed with plasmids expressing either wt Tup1-6HA or mutant Tup1-K229,270R-6HA. Overnight culture were inoculated into fresh SC-Ura medium to OD<sub>600</sub> of 0.05 and incubated at 30°C with constant shaking. The absorbance OD<sub>600</sub> of yeast culture was measured at the indicated time points. Analyses were performed four times, with standard deviation (SD) indicated with error bars.




**Figure 2. Tup1 sumoylation enhances its association with promoters of *ARG1* and *CPA2* while reducing their transcription**

(a) Tup1-HA ChIP analysis of Tup1 wild-type (WT) or K229,270R mutant (MT) on *ARG1* and *CPA2* promoters at indicated times after SM induction. (b) RNAP II ChIP analysis of Tup1 wild-type (WT) or K229,270R mutant (MT) on *ARG1* and *CPA2* promoters at indicated times after SM induction. (c) H3 ChIP analysis of Tup1 wild-type (WT) or K229,270R mutant (MT) on *ARG1* and *CPA2* promoters at indicated times after SM induction. (d) mRNA levels were determined by RT-PCR for *ARG1* and a control housekeeping gene, *ACT1*, at the indicated times after exposure to SM. Analysis was performed with strains expressing Tup1 WT or MT. Values were normalized to *ARG1* RNA levels in Tup1-WT at 0 min. Data are represented as mean  $\pm$  SD of three independent experiments. *P*-values (refer to Materials and Methods for calculation) are indicated in parentheses above paired bars where relevant.



**Figure 3. Sumoylation of both Tup1 and Gcn4 enhances Gcn4 eviction from the induced *ARG1* promoter**

(a) Gcn4-Flag WT and K50,58R MT ChIP analysis of Tup1 wild-type (WT) or K270R mutant on *ARG1* promoter at indicated times after SM induction. (b) mRNA levels were determined by RT-PCR for *ARG1* and *ACT1* at the indicated times after exposure to SM. Analysis was performed in strains expressing Gcn4-WT, or K50,58R (MT) and Tup1 WT, or K270R, as indicated. Values were normalized to *ARG1* mRNA levels in strain expressing Gcn4-WT and Tup1-WT at 0 min. (c) Gal11-WT and Gcn5-WT Flag ChIP analysis of Tup1 wild-type (WT) or K270R mutant on *ARG1* promoters at indicated times after SM induction. Data are represented as mean  $\pm$  SD of three independent experiments.



**Figure 4. Multiple Tup1- and transcription-related proteins are sumoylated during amino acid starvation**

(a) Hht1-HA, Gcn5-HA, Cti6-HA, Gal11-HA and Cyc8-HA IPs of untreated (control), or SM-treated cells were analyzed by HA and yeast SUMO (Smt3) western blots. (b) Analysis of HA IPs obtained from untreated control and SM-induced cells expressing Hht1-WT Flag and either Tup1-WT HA or K270R HA. The asterisk indicates the HA antibody light chain. (c) Analysis of Flag IPs obtained from untreated control and SM-induced cells expressing Hht1- WT Flag and either Tup1 WT-HA or K270R-HA.